

Fluorescent leukotriene B₄: potential applications

Alan Sabirsh,^{1,*} Anders Wetterholm,* Jesper Bristulf,[†] Hakon Leffler,[§] Jesper Z. Haeggström,* and Christer Owman[†]

Department of Medical Biochemistry and Biophysics,* Division of Physiological Chemistry II, Karolinska Institute, Stockholm, Sweden; Department of Physiological Sciences,[†] Division of Molecular Neurobiology, Lund University, Lund, Sweden; and Department of Laboratory Medicine,[§] Division of Microbiology, Immunology, and Glycobiology, Lund University, Lund, Sweden

Abstract Leukotriene B₄ (LTB₄) is a potent lipid mediator of inflammation that acts primarily via a seven-transmembrane-spanning, G-protein-coupled receptor denoted BLT₁. Here, we describe the synthesis and characterization of fluorescent analogs of LTB₄ that are easy to produce, inexpensive, and without the disadvantages of a radioligand. Fluorescent LTB₄ is useful for labeling LTB₄ receptors for which no antibodies are available and for performing one-step fluorescence polarization assays conducive to high-throughput screening. We found that orange and green fluorescent LTB₄ were full agonists that activated the LTB₄ receptor BLT₁ with EC₅₀ values of 68 and 40 nM, respectively (4.5 nM for unmodified LTB₄). Flow cytometric measurements and confocal imaging showed that fluorescent LTB₄ colocalized with BLT₁. Fluorescence polarization measurements showed that orange fluorescent LTB₄ bound to BLT₁ with a K_d of 66 nM and that this binding could be displaced by unlabeled LTB₄ and other BLT₁-specific ligands. Fluorescent LTB₄ analogs were also able to displace tritiated LTB₄. Orange fluorescent LTB₄ binding to enhanced green fluorescent protein-tagged BLT₁ could be observed using fluorescence resonance energy transfer. **In addition to being a useful alternative to radiolabeled LTB₄, the unique properties of fluorescently labeled LTB₄ allow a variety of detection technologies to be used.**—Sabirsh, A., A. Wetterholm, J. Bristulf, H. Leffler, J. Z. Haeggström, and C. Owman. **Fluorescent leukotriene B₄: potential applications.** *J. Lipid Res.* 2005. 46: 1339–1346.

Supplementary key words G-protein coupled receptor • pharmacology • method

Leukotriene B₄ (LTB₄) is a proinflammatory lipid mediator generated from arachidonic acid, stored in phospholipids of the nuclear membrane, by the sequential action of phospholipase A₂, 5-lipoxygenase, and LTA₄ hydrolase (1). In humans, LTB₄ is produced primarily by leukocytes of the myeloid lineage, and as a potent chemoattractant, it is important for directing leukocytes toward infected or

inflamed tissues. Knockout experiments, deleting either LTA₄ hydrolase (2) or the high-affinity LTB₄ receptor BLT₁ (3, 4), have shown that in animals unable to produce or respond to LTB₄, death occurs after infections that would normally not be fatal and that this correlates with a lack of leukocyte infiltration into infected tissues. These data agree well with a recent report demonstrating that LTB₄, administered systemically to healthy volunteers promotes the production of antiviral cytokines (5). It has also been known for many years that LTB₄ plays a role in almost every pathophysiological condition involving inflammation, and this list has recently been expanded to include atherosclerosis as well (6–9). Recently, LTB₄ has also been found to modulate the behavior of lymphocytes, providing a link between the activation of the early innate immune system and the long-term adaptive immune responses (10).

The physiology of LTB₄ is complex, and this is partly because this mediator binds to several different receptor proteins in addition to BLT₁. Another G-protein-coupled receptor, BLT₂, also binds LTB₄, but with lower affinity (11–13). Unlike BLT₁, this receptor is expressed in many tissues, but its physiological role remains unclear. LTB₄ has also been shown to bind to and activate the vanilloid receptor VR1 (14) and the nuclear peroxisome proliferator-activating receptor α (15).

Radiolabeled LTB₄ is difficult to synthesize and is, at times, commercially unavailable, which is impeding molecular pharmacological studies of LTB₄ receptors. Furthermore, commercially available antibodies only recognize the human isoform of the LTB₄ receptor BLT₁ (16), making it impossible to assess LTB₄ receptor expression in other species. The current interest in LTB₄ as an immunomodulatory agent has also created a need for a method that would allow LTB₄ receptor expression to be easily evaluated in mouse tissues. A fluorescently labeled derivative of LTB₄, compatible with a range of fluorescence-based optical techniques, could potentially be useful for this pur-

Manuscript received 15 February 2005 and in revised form 17 March 2005.

Published, JLR Papers in Press, April 1, 2005.
DOI 10.1194/jlr.D500005-JLR200

¹ To whom correspondence should be addressed.
e-mail: alan.sabirsh@mbb.ki.se

Copyright © 2005 by the American Society for Biochemistry and Molecular Biology, Inc.

This article is available online at <http://www.jlr.org>

pose. In addition, a fluorescent ligand would permit real-time measurements to be made at true equilibrium using fluorescence polarization and fluorescence resonance energy transfer (FRET) techniques. These measurements are made in homogenous mixtures requiring no further handling, which is ideal for high-throughput screening. This also facilitates experiments involving soluble LTB₄ binding proteins that are not compatible with methodologies that use filtration-based techniques to measure binding. Finally, by carefully choosing the fluorophores used to label LTB₄, it is possible to use FRET to evaluate ligand binding to receptors.

Here, we present the synthesis of fluorescent LTB₄ molecules. We show that these ligands bind to and activate BLT₁ and that they can be used successfully to study BLT₁ receptors using a variety of techniques incompatible with radiolabeled LTB₄, including flow cytometry, confocal microscopy, fluorescence polarization, and FRET. This will allow LTB₄ physiology to be studied at equilibrium, over time, and in tissues or biochemical systems that have previously been inaccessible.

MATERIALS AND METHODS

Chemicals and reagents

All reagents were purchased from Sigma, unless stated otherwise. RP69698 was obtained from Rhône-Poulenc/Aventis (Vitry Sur Seine, Cedex, France). LTB₄ antiserum was obtained from Cayman Chemical (Ann Arbor, MI) and Assay Designs (Ann Arbor, MI). Hydroxamate, RB 3040, and Kelatorphan were generous gifts from Prof. B. P. Roques (UFR des Sciences Pharmaceutiques et Biologiques). All leukotrienes were from Biomol (Plymouth Meeting, PA).

Anti-BLT₁ receptor antibody 7B1 (16) was obtained from Serotec. The molar antibody concentration was calculated using the molecular weight of the appropriate (mouse) immunoglobulin isotype obtained from the crystal structure (17). The concentration of protein in the antibody solution was determined using a protein assay according to the manufacturer's instructions (BCA Protein AssayTM; Pierce, Rockford, IL). Antibody concentrations should be regarded as approximate because of uncertainties regarding their actual weight and the concentration of active antibody in antibody solutions.

Purified LTA₄ hydrolase was produced as described previously (18).

Labeling LTB₄ with Alexa FluorTM dyes

Fluorophore-labeled LTB₄ was produced using LTB₄-aminopropylamide (LTB₄-APA; Biomol), and amine reactive succinimidyl esters of Alexa FluorTM dyes (Molecular Probes, Eugene, OR) were dissolved in methanol to a concentration of 1 mM. LTB₄-APA was dried in a nitrogen atmosphere and redissolved in methanol to a concentration of 1.28 mM.

LTB₄-APA and Alexa FluorTM dye solutions were mixed at a ratio of 1:2 and incubated for 60 min at room temperature with continuous stirring. The LTB₄-APA-Alexa FluorTM product was isolated using reverse-phase HPLC with a Nova Pak C₁₈ column (Waters, Milford, MA) and a mobile phase consisting of methanol-water-acetate at a ratio of 60:40:0.08 (pH set to 6.8 using NaOH).

The fractions containing LTB₄-APA and fluorescent LTB₄ were collected, and the HPLC mobile phase was removed by diluting each sample 1:1 with water and then filtering through Supel

CleanTM LC-18 columns (Supelco/Sigma-Aldrich, Bellefonte, PA). Both LTB₄-APA and fluorescent LTB₄ were retained in the columns and were washed once with water before elution in methanol. The purified substances were dried in a nitrogen atmosphere and redissolved in methanol. Concentrations were determined using absorbance at 270 nm for LTB₄-APA (extinction coefficient = 50,000) and 365 nm (the Alexa FluorTM portion of the molecule absorbs light at 365 nm, whereas the unmodified LTB₄-APA does not). Using fluorescence emission, the concentration of fluorescent versions of LTB₄ could also be independently verified by comparison with a standard curve prepared using known quantities of each dye.

The various labeled versions of LTB₄ are referred to as "fluorescent LTB₄" or LTB₄-488 (for the green fluorescent LTB₄-APA-Alexa Fluor-488TM conjugate) and LTB₄-568 (for the orange fluorescent LTB₄-APA-Alexa Fluor-568TM conjugate) as necessary.

Cell culture and construction of HF1pBLT1 cell lines

Cultures of HeLa HF1pBLT1 luciferase reporter cells expressing BLT₁ or sham-transfected cells (HF1pSham) were established and maintained according to Kotarsky, Owman, and Olde (19). To create BLT₁ receptors that were C-terminally tagged with enhanced green fluorescent protein (EGFP), we used the mammalian expression vector pEAK12 (Edge Biosystems). This vector was used to construct the pEAK-HFTE vector containing a (His)₁₀ tag (H), a FLAG tag (F), and the recognition site for the tobacco etch virus (T) protease followed by EGFP (E). The HFTE cassette was assembled using PCR and cloned into pEAK12 between the *EcoRI* and *NotI* sites using standard methods. The open reading frame of the human BLT₁ receptor was generated using PCR with the following primers: 5 prime, ATATAAGCTTCCACCATGAACACTACATCTTCT (which includes a *HindIII* site and a Kozak consensus sequence); and 3 prime, CAGTGAATTCCCGTTCAGTTCGTTAACTTGAGAG (this primer adds an *EcoRI* site and removes the natural stop codon). BLT₁ was then cloned into pEAK-HFTE between *HindIII* and *EcoRI* to generate pEAK-BLTR-HFTE, in which the open reading frame of BLT₁ was in frame with, and upstream of, the HFTE cassette such that all three tags and the protease site were added to the C-terminal end of the expressed receptor protein. The integrity of the construct was confirmed by sequencing with BigDye (Applied Biosystems). HeLa HF1 cells were stably transfected with pEAK-BLT1-HFTE to produce HF1pBLT1-EGFP cells.

Neutrophils were purified from buffy coats using density gradient centrifugation (Lymphoprep; Axis-Shield, Oslo, Norway) after dextran sedimentation of erythrocytes. Purity and viability were determined using May-Grunwald-Giemsa staining and trypan blue exclusion, respectively. The resulting cell cultures contained at least 95% neutrophils, and the cells were at least 99% viable.

Membrane preparation

Cell cultures were grown on tissue culture plates until confluent and then chilled to 4°C. The cell growth medium was removed and the cells were rinsed once with ice-cold PBS, before ice-cold Tris-HCl buffer (50 mM Tris base, 5 mM MgCl₂, and 1 mM EGTA, pH 7.5) was added to each plate. The cells were then scraped off and homogenized using a motorized Teflon pestle.

The cell homogenates were centrifuged at 1,000 g for 10 min at 4°C. The pellet was discarded, additional buffer was added, and the supernatant was rehomogenized. The resulting homogenate was centrifuged at 100,000 g for 60 min at 4°C, and the supernatant was discarded, additional buffer was added, and the pellet was rehomogenized. The membrane protein concentration was determined (BCA protein assayTM), and the membrane isolates were portioned into aliquots before they were frozen at -80°C until use.

Confocal microscopy

For confocal microscopic studies, HF1pBLT1-EGFP and HF1pSham cells were seeded onto poly-D-lysine-coated cover slips at least 24 h before use. Just before staining, the cells were chilled to 4°C and rinsed with PBS. After addition of 100 nM LTB₄-568, the cells were incubated for 60 min at 4°C. The cells were then washed four times and fixed with 4% (w/v) paraformaldehyde for 5 min at room temperature. The expression of BLT₁-EGFP and LTB₄-568 binding were then examined using a TCS SP2 confocal laser-scanning microscope (Leica Microsystems, Wetzlar, Germany) equipped with an argon laser (488 nm emission line) and a green He/Ne laser (543 nm emission line), respectively.

Flow cytometry

Flow cytometric analysis of HF1pBLT1 and HF1pSham cells was performed using a FACSVantage™ (Becton Dickinson, Franklin Lakes, NJ). After staining with 100 nM LTB₄-568, fluorescence was analyzed using a FACSArray bioanalyzer (Becton Dickinson) equipped with a green diode laser emitting 532 nm light.

Radioligand binding assays

Radioligand binding assays were performed using opaque white 96-well filter plates with FC glass fiber filters (model MAFC-NOB, Multiscreen Assay System; Millipore, Bedford, MA). The plates were presoaked with binding buffer (0.02 M HEPES, 10 mM CaCl₂, and 10 mM MgCl₂·6H₂O, pH 7.5), which was then exchanged for 75 μl of binding buffer containing 1.0 nM [³H]LTB₄ and, if necessary, 2.0 μM unlabeled LTB₄ to determine nonspecific binding. The binding reaction was started by adding another 75 μl of binding buffer containing 1.0 μg of rehomogenized cell membrane and the appropriate concentration of any test substance. The membrane-ligand solution was incubated for 1 h at room temperature. The reaction was terminated by rapid filtration, and the filters were then washed three times with 200 μl volumes of ice-cold washing buffer (20 mM Tris-base and 0.5 g/l BSA). Excess washing buffer was removed by blotting, and the plates were dried at 40°C for 30 min. Twenty-five microliters of Microscint-O (Perkin-Elmer, Boston, MA) was then added to each well, and [³H]LTB₄ binding was evaluated using a MicroBeta scintillation counter (Perkin-Elmer).

FRET analysis of fluorescent LTB₄ binding to BLT₁

HF1pSham, HF1pBLT1, and/or HF1pBLT1-EGFP cells were seeded onto black, clear-bottomed 96-well plates at least 48 h before use. Just before use, the cells were chilled to 4°C and gently washed three times in an ice-cold buffer solution (135 mM NaCl, 4.6 mM KCl, 1.2 mM MgCl₂, 1.5 mM CaCl₂, 11 mM glucose, and 10 mM HEPES, pH 6.0, 7.5, or 9.0). Fifty microliters of cold buffer containing LTB₄-488 or LTB₄-568, or both, was added to appropriate wells, with or without excess unlabeled LTB₄. The cells were then incubated for 30 min at 4°C before they were washed three times with chilled buffer. Fluorescence and FRET were measured using a fluorometer (PolarStar™; BMG Labtech, Offenburg, Germany) chilled to <16°C. After fluorescence analysis, cell morphology was checked microscopically, the cells were counted, and the results were corrected for cell density. Cells on the same plate were also challenged with 100 nM LTB₄ to ensure that functional BLT₁ receptors were expressed. The resulting calcium release was visualized using FURA2-AM according to previously reported methods (20).

Fluorescence polarization assays of fluorescent LTB₄ binding

Fluorescence polarization assays of fluorescent LTB₄ binding to membrane preparations were performed in a total volume of 50 μl of binding buffer (0.02 M HEPES, 10 mM CaCl₂, and 10

mM MgCl₂·6H₂O, pH 7.5) on black, 96-well, low-volume (50 μl/well) plates from Molecular Devices. For competition experiments, membrane preparations were added to each well (at a concentration of 10 μg protein/well) together with the appropriate competitor or 1 μM LTB₄ and 10 nM fluorescent LTB₄. The membranes were incubated at room temperature for 1 h before fluorescent ligand binding was analyzed using a fluorometer (PolarStar™) equipped to measure fluorescence polarization. For each competitor or LTB₄, the concentration that inhibited half of the specific binding (IC₅₀) of fluorescent LTB₄ was determined when possible from competitive binding curves using nonlinear regression (Prism; GraphPad Software, San Diego, CA). For measurements of ligand binding to purified soluble proteins, the procedure was essentially the same as described above, and molar concentrations were calculated, after protein determination, using molar weights obtained from crystallographic structures.

For measurements of fluorescent LTB₄ association and dissociation, a final concentration of 10 nM fluorescent LTB₄ was added to membrane preparations as described above, and repeated polarization measurements were made over 30 min before adding a final concentration of 1 μM LTB₄ and measuring polarization for a further 30 min. Saturation experiments were performed using up to 100 nM fluorescent LTB₄ with or without 10 μM LTB₄. Results are presented in millipolars calculated as $(\parallel - \perp) / (\parallel + \perp) \times 1,000$, where \parallel is fluorescence emission polarized in parallel to the excitation source and \perp is perpendicularly polarized fluorescence emission.

Fluorescence assays of fluorescent LTB₄ binding

Fluorescent LTB₄ binding to whole cells was examined using cells growing on black, clear-bottomed, 96-well plates. The protocol was similar to that used for fluorescence polarization, except that the final incubation volume was 100 μl and the entire procedure was performed at 4°C with an incubation time of 30 min. After the incubation, the cells were washed four times with PBS Dulbecco's, and then all fluid was removed from the wells. Fluorescence was measured using a fluorometer (Fluostar™; BMG).

Luciferase assay of BLT₁ activity

The assay for agonist-induced luciferase production was performed according to Kotarsky, Owman, and Olde (19). This assay is used here to measure the relative potencies and efficacies of LTB₄ and its fluorescent counterparts. Briefly, HF1pBLTR1 cells were seeded as described above onto white, clear-bottomed, 96-well plates (Corning Costar, Corning, NY) and grown until they were 80–90% confluent. PBS solutions containing ligands were then added (in a volume not exceeding 10 μl) to the wells (results are reported as final ligand concentrations). After a further 16 h of incubation, the cells were washed once with PBS and lysed. Luciferase activity was measured using a 96-well luminometer (Lumistar™; BMG). The half-maximum effective ligand concentrations (EC₅₀) were determined from the luciferase assay concentration response curves using nonlinear regression (Prism), and results are presented as mean EC₅₀ values with 95% confidence intervals.

RESULTS

Synthesis of fluorescent LTB₄

The reaction between LTB₄-APA and the succinimidyl ester of the Alexa Fluor™ dye is simple (Fig. 1) and proceeds at room temperature, resulting in ~30% labeled LTB₄-APA after 1 h. The reaction mixture is slightly acidic (pH 6.0) and the yield was found to decrease significantly




Fig. 1. The synthesis of fluorescent leukotriene B₄ (LTB₄-FL) from LTB₄-aminopropylamide (LTB₄-APA) and Alexa Fluor™ fluorophores. The free amino group of LTB₄-APA hydrolyzes the succinimidyl ester to form mixed isomers of LTB₄ Alexa Fluor™ 568 or 488 (fluorescent LTB₄).

if the pH was increased using sodium hydroxide. The unreacted Alexa Fluor™ dye passed through the column immediately followed by fluorescent LTB₄ and unreacted LTB₄-APA. The two reaction components and the product could be detected using 270 nm absorption. Some additional reaction products were also detected, particularly after the Alexa Fluor™ dye was stored (after solvation) for more than 5 months at -80°C , but these contaminants were easily distinguished from fluorescent LTB₄.

Fluorescence polarization measurements

We used a PolarStar™ (BMG) to measure the fluorescent anisotropy (polarization) of labeled LTB₄, and this machine was found, under our experimental conditions, to be sensitive in a linear manner over concentrations ranging from >100 nM to 1 nM. Although the concentration of fluorescent LTB₄ did not affect the average fluorescence polarization over this range, concentrations less than 5 nM approached the noise floor of the assay and tended to produce erratic polarization values. Solutions with increasing concentrations of glycerol were also used to evaluate the dynamic range of fluorescence polarization values that could be obtained using fluorescent LTB₄. Fluorescent LTB₄ was found to be sensitive to increasing glycerol concentrations, although not as sensitive as the free Alexa Fluor™ fluorophores, which had a larger dynamic range. The polarity shift of fluorescent LTB₄ was not sensitive to the addition of unlabeled LTB₄.

Fluorescent LTB₄ as a BLT₁ agonist

The ability of fluorescent LTB₄ to activate HF1pBLT1 luciferase reporter cells was studied by comparing unmodified LTB₄ with labeled LTB₄. The EC₅₀ for LTB₄ was 4.5 nM (1.7–12) (means and 95% confidence intervals), whereas the EC₅₀ for LTB₄-568 was found to be ~ 15 times

higher at 68 nM (31–150), and the EC₅₀ for LTB₄-488 was found to be ~ 10 times higher at 40 nM (25–78). Both ligands were full agonists. The EC₅₀ for LTB₄ was not affected by the addition of free Alexa Fluor™ dye, even at concentrations as high as 10 μM .

Fluorescent LTB₄ and BLT₁ binding experiments

The binding of labeled LTB₄ to BLT₁ was examined using membranes prepared from HF1pBLT1 cells and measurements of fluorescence polarization. LTB₄-568 affinity for BLT₁ was calculated to be $K_d = 66$ nM (43–76) with a receptor concentration (Bmax) of 60 nM (31–101) according to the method of Prystay, Gosselin, and Banks (21). Accordingly, the association rate for LTB₄-568 was found to have an observed rate constant of 0.35 min^{-1} (0.33–0.37), so that specific binding of LTB₄-568 by BLT₁ was essentially complete after 20 min. LTB₄-568 is displaced quickly by unlabeled LTB₄, and it was impossible, for technical reasons, to make the first polarization measurement before a significant portion of the bound LTB₄-568 had been displaced. The specific LTB₄-568 binding could be displaced using unlabeled LTB₄, with an IC₅₀ of 110 nM (78–160), and also by the LTB₄ antagonist RP69698 [IC₅₀ = 1.60 μM (0.94–2.70)], the BLT₁-specific partial agonist U75302 [IC₅₀ = 2.10 μM (0.14–3.0)], and a BLT₁ antibody [IC₅₀ = 1.9 μM (0.1–3.6) approximate concentration; see Materials and Methods], but not by the chemically related LTC₄ or an isotype control antibody (Fig. 2). In competition experiments, there was no difference in the ability of unlabeled LTB₄ to displace labeled LTB₄, regardless of when the unlabeled ligand was added. No specific binding of fluorescent LTB₄ was observed when using sham-transfected HF1 cells or their membranes.

The interaction between fluorescent LTB₄ and BLT₁ was also examined using conventional nonequilibrium bind-




Fig. 2. Competition between LTB₄-568 and various ligands for BLT₁ binding. Displacement of LTB₄-568 (10 nM) binding to HF1pBLT1 cell membranes (10 μg/well) by unlabeled LTB₄, the LTB₄ antagonist RP69698, the BLT₁-specific partial agonist U75302, and the BLT₁ receptor antibody 7B1. Isotype control antibodies (Iso) and LTC₄ (which is not a ligand for BLT₁) did not displace LTB₄-568 binding. Data are presented as means ± SEM from triplicate wells in two separate experiments performed at 24°C.

ing experiments. Measuring fluorescence bound to whole cells expressing BLT₁, similar binding constants were obtained for LTB₄-568 affinity [$K_d = 64$ nM (27–102)] and LTB₄-568 displacement by LTB₄ [$IC_{50} = 75$ nM (36–156)]. The ability of fluorescently labeled LTB₄ to displace tritiated LTB₄ was also evaluated using membranes prepared from HF1pBLT₁ cells. Tritiated LTB₄ could be displaced using unlabeled LTB₄ with a calculated IC_{50} of 3 nM (2.1–4.6), by LTB₄-488 [$IC_{50} = 11$ nM (5–26)], LTB₄-568 [$IC_{50} = 22$ nM (12–38)], and LTB₄-APA [$IC_{50} = 180$ nM (91–360)].

Colocalization of fluorescent LTB₄ and EGFP-tagged BLT₁

Confocal microscopy of HF1pBLT1 cells revealed that the cell surface expression of EGFP-tagged receptors and LTB₄-568 fluorescence overlapped (Fig. 3C). Flow cytometric analysis of cells expressing a C-terminal EGFP-tagged BLT₁ receptor and surface labeled with BLT₁ antibodies indicated that more than 90% of the expressed receptors could be found on the cell surface (Fig. 3A). This same population of cells bound LTB₄-568 specifically, and this binding could be displaced using unlabeled LTB₄ (Fig. 3B). Sham-transfected cells did not specifically bind LTB₄-568, and cells exposed to only Alexa Fluor 568™ could not be differentiated from unlabeled cells (Fig. 3B). To demonstrate that fluorescent LTB₄ could also label immunologically relevant cells, we also labeled neutrophils with LTB₄-568 or with LTB₄-568 and excess unlabeled LTB₄ to measure nonspecific binding. Nonspecific binding represented only ~15% of the total binding under these conditions. In other words, LTB₄-568 labeling made the neutrophils approximately six times brighter than background, demonstrating that fluorescent LTB₄ can be used to detect LTB₄ receptors even in cell populations that are not overexpressing the receptors.

We also performed a FRET analysis of LTB₄-568 binding to EGFP-tagged BLT₁ receptors. In contrast to EGFP, LTB₄-568 is poorly excited by blue light (485 nm). When LTB₄-568 was added to membrane preparations from cells expressing EGFP-tagged BLT₁, the yellow-green EGFP fluorescence emission was able to stimulate LTB₄-568 molecules in close proximity, which reemitted this light as red-orange (610 nm) fluorescence (Fig. 4). LTB₄-568 does emit some fluorescence at 610 nm even when stimulated with 485 nm light, but the effect is much larger in the presence of BLT₁-EGFP, indicating that the ligand and the receptor are in close proximity. Membranes from HF1pSham cells did not exhibit this effect. The observed energy transfer could also be disrupted by adding excess unlabeled LTB₄.

Fluorescent LTB₄ binding to enzymes and carrier proteins

LTB₄-568 binding to putative carrier proteins was also examined, and it was found that LTB₄-568 could bind to albumin. This binding was unique to albumin, but it could not be reversed by adding excess quantities of unlabeled LTB₄ (Fig. 5).

Similarly, we found that both LTB₄-488 and LTB₄-568 could bind to LTA₄ hydrolase, the enzyme responsible for LTB₄ production in tissues (data not shown). Over a range of concentrations (1–100 nM), fluorescent LTB₄ bound to LTA₄ hydrolase but could not be displaced by excess unlabeled LTB₄ or various inhibitors (RB 3040, Kelatorphan, or the substrate mimic hydroxamate), indicating that the observed binding was probably nonspecific.

LTB₄ and a fluorescence polarization immunoassay

We attempted to measure the concentration of free unlabeled LTB₄ by allowing it to compete with labeled LTB₄ for binding to a polyclonal rabbit anti-LTB₄ peroxidase antibody (Cayman Chemical) and a polyclonal rabbit anti-LTB₄ antibody (Assay Designs). Fluorescent LTB₄ was not recognized by either antibody solution, even at antibody concentrations 100 times those recommended by the manufacturer. Therefore, it will be necessary to produce antibodies specific for fluorescent LTB₄ before a one-step fluorescence polarization immunoassay can be produced.

DISCUSSION

Synthesis and activity of fluorescent LTB₄

Our fluorescent LTB₄ analogs can reversibly bind to the LTB₄ receptor BLT₁ and function as full agonists. Fluorescent LTB₄ bound specifically to BLT₁ because this binding could be displaced, not only by unlabeled LTB₄ but also by the BLT₁-specific partial agonist U75302, the LTB₄ antagonist RP69698, and a BLT₁ antibody that has been shown to interfere with LTB₄ binding (22). LTB₄-568, however, does not bind very tightly to BLT₁, and there is a discrepancy between the values obtained using polarization measurements, made in homogenous solutions, and radioligand binding experiments. Polarization measurements, however, are made at true equilibrium and are dependent on a significant amount of ligand depletion, which makes




Fig. 3. Colocalization of LTB₄-568 and enhanced green fluorescent protein (EGFP)-tagged BLT₁. **A:** Flow cytometric analysis of HF1 cells expressing BLT₁ receptors that were C-terminally tagged with EGFP. The green fluorescence of the EGFP tag was used to measure the total number of receptors in each cell, and the red fluorescence of a BLT₁ antibody was used to measure surface expression. In combination, these measurements reveal that more than 90% of the receptor is on the cell surface (the percentage of double positive cells is given in the upper right quadrant of each scattergram). **B:** Representative curves showing that HF1pBLT1-EGFP cells bind 100 nM LTB₄-568 (grey shading) specifically. Also shown are curves depicting LTB₄-568 nonspecific binding (100 nM LTB₄-568 displaced using 10 μ M LTB₄) and HF1pSham cells, which exhibited the same level of nonspecific binding and a lack of specific binding. **C:** LTB₄-568 staining of HF1pBLT1-EGFP cells and HF1pSham cells observed using confocal microscopy. Green EGFP fluorescence from tagged receptors in the cell membranes colocalized with the orange fluorescence from LTB₄-568. **D:** Neutrophils isolated from buffy coats were stained with either 10 nM LTB₄-568 (to determine total binding) or LTB₄-568 together with 1 μ M LTB₄ (to determine nonspecific binding) before LTB₄-568 fluorescence was quantified using a plate fluorometer. The addition of unlabeled LTB₄ could displace almost 85% of the LTB₄-568 fluorescence. Data shown are means \pm SEM from quadruplicate wells and two separate donors.

the results difficult to compare directly with those of traditional nonequilibrium competition experiments. In spite of this apparent low affinity, LTB₄-568 was surprisingly difficult to displace. Ligands that typically displace tritiated LTB₄ at concentrations in the nanomolar range (such as LTB₄ itself) required higher concentrations to displace fluorescent LTB₄. This may be because the fluorophore portion of fluorescent LTB₄ participates in the ligand-receptor interaction. Excess free fluorophore, however, did not affect the binding or function of fluorescent LTB₄, so the mechanism appears to be more complex and may depend on interaction between epitopes from both the leukotriene and the fluorophore. This is supported by the observation that fluorescent LTB₄ binds more strongly to BLT₁ than LTB₄-APA. Baneres and Parello (23) have shown that LTB₄ binding induces the dimerization of purified BLT₁ receptors, which in turn increases BLT₁ affinity for LTB₄. However, preincubating the receptors with either fluores-

cent LTB₄ or LTB₄ before adding LTB₄ or fluorescent LTB₄, respectively, did not affect the results. Nor are the apparently low ligand affinities dependent on active receptor conformations (and presumably G-protein coupling), as even antagonists (both small molecules and antibody peptides) bound more weakly than expected and were insensitive to the order in which they were added.

Fluorescent versions of LTB₄ bind to and activate BLT₁ specifically, but they are somewhat less potent agonists compared with unmodified LTB₄. They are not partial agonists, however, because the addition of labeled LTB₄ to LTB₄ solutions did not affect receptor activation by LTB₄. Exposure to labeled LTB₄ could also provoke equally large responses, provided the concentrations used were high enough.

LTB₄, modified such that the hydroxyl portion of the carboxyl group is exchanged for aminopropylamide, has previously been used to chemically label receptors for LTB₄




Fig. 4. Fluorescence resonance energy transfer (FRET) from EGFP-tagged BLT₁ to LTB₄-568. Membranes prepared from cells expressing EGFP-tagged BLT₁ were illuminated with 485 nm light, and fluorescence emission at 520 or 610 nm was recorded. FRET between BLT₁-EGFP and LTB₄-568 is revealed as a shift from 520 nm fluorescence emission to 610 nm emission. LTB₄-568 was added to wells containing membranes (white bars), membranes plus an excess of unlabeled LTB₄ (light gray bars), no membranes (dark gray bars), or a buffer control (no fluorophores; black bars). After the addition of LTB₄-568, much more orange-red (610 nm) fluorescence was produced by membranes containing EGFP that were stimulated with 485 nm (blue) light. Data have been normalized to the green fluorescence of EGFP and are shown as means \pm SEM from quadruplicate wells in two separate experiments. The x axis shows excitation/emission wavelengths in nanometers.

(24), and this ligand retains agonistic activity. It also reacts readily with a variety of fluorophores that have been designed to include succinimidyl esters to facilitate labeling of the free amino groups found in proteins. Although we have focused on Alexa Fluor™ 488 and 568, any fluorophore coupled to a succinimidyl ester will react with similar chemistry, allowing the production of labeled LTB₄ molecules with emission maxima across the visible spectrum and into infrared.

The location of the fluorophore (Fig. 1) is fortuitous because in this position it allows the LTB₄ moiety to bind

to and interact with BLT₁. Receptor modeling, ligand docking experiments, and point mutation of the receptor protein (our unpublished data) also support this idea.

Using the unique optical properties of fluorescent LTB₄

Using the optical properties of fluorescent LTB₄, we could locate LTB₄ molecules and LTB₄ receptors in living cells using a variety of techniques incompatible with radiolabeled LTB₄, including flow cytometry, fluorescence histochemistry, and the aforementioned fluorescence polarization.

We also used fluorescently labeled LTB₄ to label membranes from cells expressing BLT₁ receptors that were tagged at the C-terminal end with EGFP and observed both physical overlap between the two fluorophores and resonance energy transfer between the fluorescent ligand and the tagged receptor. The red shift in EGFP fluorescence after LTB₄-568 binding could also be reversed by the addition of unlabeled LTB₄, implying that this is not attributable to random interactions between EGFP and LTB₄-568 molecules in close proximity. Although we did observe binding to purified samples of the cytosolic LTA₄ hydrolase (see below), no cytosolic staining was observed in preparations of intact cells. This may be because, unlike leukocytes, parenchymal cells do not express LTA₄ hydrolase protein, or because fluorescent LTB₄ does not cross intact cell membranes.

Fluorescent LTB₄ binding to water-soluble proteins

One advantage of using fluorescent polarization is that there is no need to separate unbound ligand from bound ligand before measuring ligand binding. This in turn means that measuring fluorescent LTB₄ binding to soluble proteins is, at least theoretically, relatively straightforward. Molecules that bind fluorescent LTB₄ are interesting because many represent possible drug targets and could be used as platforms for high-throughput screening. If the displacement of fluorescent LTB₄ from LTA₄ hydrolase by specific enzymatic inhibitors could be mea-





Fig. 5. LTB₄-568 binding to serum albumin. A: LTB₄-568 was added to solutions each containing one of three different polypeptides or the free amino acid glycine. Over a wide range of concentrations, only human serum albumin (HSA) bound LTB₄-568. B: LTB₄-568 (closed diamonds) was added to albumin solution with concentrations ranging from 1 μg/ml to 1 mg/ml, and binding was observed to increase in proportion to the albumin concentration. This binding, however, was not restricted to a site specific for the leukotriene moiety of LTB₄-568, because the addition of excess unlabeled LTB₄ (1 μM) displaced LTB₄-568 only slightly (closed squares). Data are presented as means \pm SEM from quadruplicate wells in two separate experiments performed at 24°C. mP, millipolars.

sured using fluorescence polarization, this would create a one-step homogenous assay for screening enzymatic inhibitors. The binding of labeled LTB₄ to serum albumin showed that fluorescent LTB₄ can bind to this important serum protein. Unlike fluorescent LTB₄ binding to BLT₁, however, binding to serum albumin could not be displaced by unlabeled LTB₄, indicating that there are not specific binding epitopes for the LTB₄ moiety. Free Alexa Fluor™ 568 dye was also unable to displace serum albumin-bound LTB₄-568, suggesting that it was not binding of the fluorophore alone that was responsible for LTB₄-568 binding. Similar results were obtained using LTA₄ hydrolase, the enzyme responsible for the enzymatic production of LTB₄ and polyclonal antibodies specific for LTB₄. Although we did observe binding to LTA₄ hydrolase, this binding was not specific and could not be displaced by unlabeled LTB₄ or various LTA₄ hydrolase inhibitors. The two different antibodies we examined, which were designed for use in immunoassays, did not recognize (bind to) our fluorescent LTB₄ analogs.

Conclusions

Fluorescent LTB₄ has several advantages over native LTB₄ and tritiated LTB₄. Fluorescent LTB₄ is not radioactive and thus is safer to handle. Fluorescently labeled LTB₄ is also significantly cheaper than tritiated LTB₄, assuming that tritiated LTB₄ can be purchased at all. Fluorescent LTB₄ can also be used to label receptors for LTB₄ in tissues that are incompatible with currently available antibodies (at present, only antibodies against the human BLT₁ receptor are commercially available). The fact that fluorescent labels can be easily detected optically makes these LTB₄ analogs compatible with a range of techniques, including flow cytometry, microscopy, polarization, and resonance energy transfer. This in turn makes it possible to study LTB₄ binding proteins in new ways, in new tissues with fewer steps, at higher throughput, even in homogeneous solutions. 

This study was financially supported by the AFA Health Foundation, the Swedish Research Council (03X-10350), Konung Gustav V:s 80 årsfond, and European Commission FP6 funding (LSHM-CT-2004-005033). This publication reflects only the authors' views. The European Commission is not liable for any use that may be made of information herein.

REFERENCES

1. Funk, C. D. 2001. Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science*. **294**: 1871–1875.
2. Byrum, R. S., J. L. Goulet, J. N. Snouwaert, R. J. Griffiths, and B. H. Koller. 1999. Determination of the contribution of cysteinyl leukotrienes and leukotriene B₄ in acute inflammatory responses using 5-lipoxygenase- and leukotriene A₄ hydrolase-deficient mice. *J. Immunol.* **163**: 6810–6819.
3. Haribabu, B., M. W. Verghese, D. A. Steeber, D. D. Sellars, C. B. Bock, and R. Snyderman. 2000. Targeted disruption of the leukotriene B₄ receptor in mice reveals its role in inflammation and platelet-activating factor-induced anaphylaxis. *J. Exp. Med.* **192**: 433–438.
4. Tager, A. M., J. H. Dufour, K. Goodarzi, S. D. Bercery, U. H. von Andrian, and A. D. Luster. 2000. BLTR mediates leukotriene B₄(4)-

- induced chemotaxis and adhesion and plays a dominant role in eosinophil accumulation in a murine model of peritonitis. *J. Exp. Med.* **192**: 439–446.
5. Flamand, L., P. Borgeat, R. Lalonde, and J. Gosselin. 2004. Release of anti-HIV mediators after administration of leukotriene B₄ to humans. *J. Infect. Dis.* **189**: 2001–2009.
6. Aiello, R. J., P. A. Bourassa, S. Lindsey, W. Weng, A. Freeman, and H. J. Showell. 2002. Leukotriene B₄ receptor antagonism reduces monocytic foam cells in mice. *Arterioscler. Thromb. Vasc. Biol.* **22**: 443–449.
7. Subbarao, K., V. R. Jala, S. Mathis, J. Suttles, W. Zacharias, J. Ahamed, H. Ali, M. T. Tseng, and B. Haribabu. 2003. Role of leukotriene B₄ receptors in the development of atherosclerosis: potential mechanisms. *Arterioscler. Thromb. Vasc. Biol.* **24**: 369–375.
8. Dwyer, J. H., H. Allayee, K. M. Dwyer, J. Fan, H. Wu, R. Mar, A. J. Lusis, and M. Mehrabian. 2004. Arachidonate 5-lipoxygenase promoter genotype, dietary arachidonic acid, and atherosclerosis. *N. Engl. J. Med.* **350**: 29–37.
9. Helgadóttir, A., A. Manolescu, G. Thorleifsson, S. Gretarsdóttir, H. Jónsdóttir, U. Thorsteinsdóttir, N. J. Samani, G. Gudmundsson, S. F. Grant, G. Thorgeirsson, et al. 2004. The gene encoding 5-lipoxygenase activating protein confers risk of myocardial infarction and stroke. *Nat. Genet.* **36**: 233–239.
10. Tager, A. M., S. K. Bromley, B. D. Medoff, S. A. Islam, S. D. Bercery, E. B. Friedrich, A. D. Carafone, R. E. Gerszten, and A. D. Luster. 2003. Leukotriene B₄ receptor BLT1 mediates early effector T cell recruitment. *Nat. Immunol.* **4**: 982–990.
11. Kamohara, M., J. Takasaki, M. Matsumoto, T. Saito, T. Ohishi, H. Ishii, and K. Furuichi. 2000. Molecular cloning and characterization of another leukotriene B₄ receptor. *J. Biol. Chem.* **275**: 27000–27004.
12. Tryselius, Y., N. E. Nilsson, K. Kotarsky, B. Olde, and C. Owman. 2000. Cloning and characterization of cDNA encoding a novel human leukotriene B₄ receptor. *Biochem. Biophys. Res. Commun.* **274**: 377–382.
13. Wang, S., E. Gustafson, L. Pang, X. Qiao, J. Behan, M. Maguire, M. Bayne, and T. Laz. 2000. A novel hepatointestinal leukotriene B₄ receptor. Cloning and functional characterization. *J. Biol. Chem.* **275**: 40686–40694.
14. Hwang, S. W., H. Cho, J. Kwak, S. Y. Lee, C. J. Kang, J. Jung, S. Cho, K. H. Min, Y. G. Suh, D. Kim, and U. Oh. 2000. Direct activation of capsaicin receptors by products of lipoxygenases: endogenous capsaicin-like substances. *Proc. Natl. Acad. Sci. USA.* **97**: 6155–6160.
15. Devchand, P. R., H. Keller, J. M. Peters, M. Vazquez, F. J. Gonzalez, and W. Wahli. 1996. The PPARalpha-leukotriene B₄ pathway to inflammation control. *Nature.* **384**: 39–43.
16. Pettersson, A., A. Boketoft, A. Sabirsh, N. E. Nilsson, K. Kotarsky, B. Olde, and C. Owman. 2000. First-generation monoclonal antibodies identifying the human leukotriene B₄ receptor-1. *Biochem. Biophys. Res. Commun.* **279**: 520–525.
17. Harris, L. J., E. Skaletsky, and A. McPherson. 1998. Crystallographic structure of an intact IgG1 monoclonal antibody. *J. Mol. Biol.* **275**: 861–872.
18. Rudberg, P. C., F. Tholander, M. Andberg, M. M. Thunnissen, and J. Z. Haeggstrom. 2004. Leukotriene A₄ hydrolase: identification of a common carboxylate recognition site for the epoxide hydrolase and aminopeptidase substrates. *J. Biol. Chem.* **279**: 27376–27382.
19. Kotarsky, K., C. Owman, and B. Olde. 2001. A chimeric reporter gene allowing for clone selection and high-throughput screening of reporter cell lines expressing G-protein-coupled receptors. *Anal. Biochem.* **288**: 209–215.
20. Sabirsh, A., J. Bristulf, and C. Owman. 2004. Exploring the pharmacology of the leukotriene B₄ receptor BLT1, without the confounding effects of BLT2. *Eur. J. Pharmacol.* **499**: 53–65.
21. Prystay, L., M. Gosselin, and P. Banks. 2001. Determination of equilibrium dissociation constants in fluorescence polarization. *J. Biomol. Screen.* **6**: 141–150.
22. Sabirsh, A., A. Pettersson, Å. Boketoft, K. Kotarsky, and C. Owman. 2003. Differential inhibition of receptor activation by two mouse monoclonal antibodies specific for the human leukotriene B₄ receptor, BLT1. *Int. Immunopharmacol.* **3**: 1829–1839.
23. Baneres, J. L., and J. Parello. 2003. Structure-based analysis of GPCR function: evidence for a novel pentameric assembly between the dimeric leukotriene B₄ receptor BLT1 and the G-protein. *J. Mol. Biol.* **329**: 815–829.
24. Goldman, D. W., L. A. Gifford, R. N. Young, T. Marotti, M. K. Cheung, and E. J. Goetzl. 1991. Affinity labeling of the membrane protein-binding component of human polymorphonuclear leukocyte receptors for leukotriene B₄. *J. Immunol.* **146**: 2671–2677.