

Substrate specificity of lysophosphatidic acid acyltransferase β —evidence from membrane and whole cell assays

David Hollenback,¹ Lynn Bonham, Laura Law, Eddie Rossnagle, Lisa Romero, Heather Carew, Christopher K. Tompkins, David W. Leung, Jack W. Singer, and Thayer White

Department of Biochemistry, Cell Therapeutics, Inc., Seattle, WA 98119

Abstract Membranes of mammalian cells contain lysophosphatidic acid acyltransferase (LPAAT) activities that catalyze the acylation of *sn*-1-acyl lysophosphatidic acid (lysoPA) to form phosphatidic acid. As the biological roles and biochemical properties of the six known LPAAT isoforms have yet to be fully elucidated, we have characterized human LPAAT- β activity using two different assays. In a membrane-based assay, LPAAT- β used lysoPA and lysophosphatidylmethanol (lysoPM) but not other lysophosphoglycerides as an acyl acceptor, and it preferentially transferred 18:1, 18:0, and 16:0 acyl groups over 12:0, 14:0, 20:0, and 20:4 acyl groups. The fact that lysoPM could traverse cell membranes permitted additional characterization of LPAAT- β activity in cells: PC-3 and DU145 cells converted exogenously added lysoPM and ¹⁴C-labeled 18:1 into ¹⁴C-labeled phosphatidylmethanol (PM). The rate of PM formation was higher in cells that overexpressed LPAAT- β and was inhibited by the LPAAT- β inhibitor CT-32501. In contrast, if lysoPM and ¹⁴C-labeled 20:4 were added to PC-3 or DU145 cells, ¹⁴C-labeled PM was also formed, but the rate was neither higher in cells that overexpressed LPAAT- β nor inhibited by CT-32501. We propose that LPAAT- β catalyzes the intracellular transfer of 18:1, 18:0, and 16:0 acyl groups but not 20:4 groups to lysoPA.—Hollenback, D., L. Bonham, L. Law, E. Rossnagle, L. Romero, H. Carew, C. K. Tompkins, D. W. Leung, J. W. Singer, and T. White. **Substrate specificity of lysophosphatidic acid acyltransferase β —evidence from membrane and whole cell assays.** *J. Lipid Res.* 2006. 47: 593–604.

Supplementary key words phosphatidic acid • lysophosphatidylmethanol • phosphatidylmethanol • acyl-CoA

Membrane-associated lysophosphatidic acid acyltransferase (LPAAT) activities, identified in bacteria, yeast, plant, and animal cells, catalyze the transfer of acyl groups from acyl-CoA to lysophosphatidic acid (lysoPA) to form phosphatidic acid (PA) (1–3). PA is a component of cell membranes and a key intermediate in the de novo synthesis of phosphoglycerides, the major components of

cell membranes, and triacylglycerol, the major form of energy storage in plants and animals. Models that suggest that PA can activate numerous cell functions, including proliferation, cytoskeletal organization, vesicle trafficking and the oxidative burst in neutrophils, have also gained wide acceptance.

Six human LPAAT isoforms (α – ζ), also known as 1-acyl-*sn*-glycerol-3-phosphate acyltransferase 1–6, have been cloned. Only the α and β isoforms, which share 34% sequence identity, have been shown to have significant LPAAT activity and to complement the growth defect in *plsC*, an *Escherichia coli* strain mutated in its sole LPAAT gene (4). These isoforms also contain two to four predicted transmembrane domains and two highly conserved motifs, H(X)₄D and EGTR, which are essential for the catalytic activity of a family of acyltransferases (5–7). Northern blot analysis of human LPAAT- α suggests that it has broad tissue distribution, whereas that of human LPAAT- β is more restricted, primarily to heart, liver, adipose, and pancreas (8–11). The inherited loss of function of LPAAT- β has also been linked to a rare form of congenital, generalized lipodystrophy, characterized by a nearly complete absence of adipose tissue from birth (12, 13).

Although membrane-associated LPAAT activities were first described by Kornberg and Pricer (14) in 1953, progress has been slow to understand their cellular function(s) and to separately define each isozyme's enzymatic properties. Significant steps could be made if these enzymes were solubilized from the membranes and purified to homogeneity. Unfortunately, attempts to solubilize LPAATs into detergent micelles have resulted in the immediate and complete loss of enzymatic activity. Thus, most information on LPAAT enzymatic properties comes from experiments using cell lysates or membrane preparations containing a recombinant LPAAT isozyme. When the

Abbreviations: GFP, green fluorescent protein; LPAAT, lysophosphatidic acid acyltransferase; lysoPA, lysophosphatidic acid; lysoPC, lysophosphatidylcholine; lysoPM, lysophosphatidylmethanol; PA, phosphatidic acid; PLD, phospholipase D; PM, phosphatidylmethanol.

¹To whom correspondence should be addressed.

e-mail: dhollenback@earthlink.net

Manuscript received 3 October 2005 and in revised form 16 December 2005.

Published, JLR Papers in Press, December 20, 2005.

DOI 10.1194/jlr.M500435-JLR200

Copyright © 2006 by the American Society for Biochemistry and Molecular Biology, Inc.

This article is available online at <http://www.jlr.org>

potential acyl donors 14:0-CoA, 16:0-CoA, 18:0-CoA, and 20:4-CoA were tested separately against lysates from COS 7 cells containing human recombinant LPAAT- β , all acyl donors were active. When the potential acyl acceptors *sn*-1-18:1 lysoPA, *sn*-1-acyl lysophosphatidylcholine (lysoPC), and lysoplatelet-activating factor (*sn*-1-alkyl lysoPC) were tested separately, only lysoPA was active (10). A separate study showed that 18:1-CoA was also an acceptable acyl donor (8). Studies of human or murine LPAAT- α expressed in JC201, COS 7, or Sf21 cells have shown that it has broad preference for acyl-CoAs and can use lysoPA but not lysoPC, lysophosphatidylethanolamine, lysophosphatidylinositol, or lysoplatelet-activating factor as an acyl acceptor (9, 15, 16). In spite of these efforts, the LPAATs' enzymatic properties need a more detailed characterization to fully understand the effect that LPAAT catalysis may have on biological processes requiring PA.

To this end, we studied the substrate specificity of LPAAT- β in more detail using Sf9 membrane preparations containing recombinant LPAAT- β and compared it with that of native Sf9 membranes and Sf9 membrane preparations containing recombinant LPAAT- α . We also developed a novel assay to estimate LPAAT- β activity in whole cells. This assay measured the intracellular formation of phosphatidylmethanol (PM) and took advantage of two reagents, lysophosphatidylmethanol (lysoPM) and LPAAT- β chemical inhibitors. LysoPM was shown to be an acyl acceptor for LPAAT- β but not LPAAT- α , and unlike lysoPA, it readily traversed cell membranes. The LPAAT- β chemical inhibitors show submicromolar potency against LPAAT- β but do not inhibit LPAAT- α even at 40 μ M and were used to help differentiate the activity of the two isozymes. We used this second assay to test whether the LPAAT- β acyl chain transfer specificity observed in the membrane assay was reciprocated in cells.

EXPERIMENTAL PROCEDURES

Materials

14 C-labeled 18:1, 14 C-labeled 20:4, 14 C-labeled 18:1-CoA, 3 H-labeled *sn*-1-18:1 lysoPA, and γ - 33 P-labeled ATP were from Perkin-Elmer; 33 P-labeled *sn*-1-18:1 lysoPA was prepared using *E. coli* diacylglycerol kinase (Calbiochem) to phosphorylate 1-monoolein (Doosan Serdery Research Laboratories) with γ - 33 P-labeled ATP as described (17); all other lipids were from Avanti Polar Lipids. Silica Gel 60 HP-TLC plates were from Analtech. Organic solvents were American Chemical Society grade or better from J. T. Baker or EM Science. The synthesis of compounds CT-32228 and CT-32212 (18) and of compound CT-32501 (19) were as described. Anti-LPAAT- α and anti-LPAAT- β antibodies have been described (20). All other chemicals were of reagent grade from Sigma.

Preparation of lysoPM

sn-1-18:1 lysoPM was synthesized from *sn*-1-18:1 lysoPA essentially as described (21). One gram of the protonated form of *sn*-1-18:1 lysoPA was dissolved in 50 ml of chloroform and mixed with an ethereal solution of diazomethane until a yellow color persisted. The solvent was evaporated, and the dimethylated lysoPA was mixed with 2.0 g of NaI in anhydrous 2-butanone and

then refluxed for 90 min. The solvent was removed, and the lysoPM was purified by flash chromatography as the sodium salt by eluting with chloroform-methanol-acetic acid-water (50:30:15:5). The solvent was evaporated, and the lysoPM was reconstituted in chloroform. This yielded 800 mg of lysoPM, whose structure and purity (>95%) were confirmed by TLC, LC-MS, and NMR.

Ectopic expression of LPAAT- β and LPAAT- α in PC-3 and DU145 cells

Cells were maintained at 37°C in a humidified atmosphere of 95% air and 5% CO₂. PC-3, a human prostate tumor cell line (American Type Culture Collection; CRL-1435), was cultured in Ham's F12K medium supplemented with 2 mM L-glutamine, 1.5 g/l sodium bicarbonate, and 10% fetal bovine serum. DU145, a human prostate tumor cell line (American Type Culture Collection; HTB-81), was cultured in Eagle's minimum essential medium supplemented with 2 mM L-glutamine, 1.5 g/l sodium bicarbonate, 0.1 mM nonessential amino acids, 1.0 mM sodium pyruvate, and 10% fetal bovine serum. For the overexpression of LPAATs in these cells, human LPAAT- β cDNA and human LPAAT- α cDNA were cloned separately into a modified version of the LXSN retroviral vector (Clontech) (22). The Moloney murine leukemia virus long terminal repeat drives LPAAT expression, and the SV40 promoter drives the selectable marker (*neo*) expression. As a further modification of the vector, a recognition site for cre recombinase (23) was introduced into the long terminal repeat regions. Control vectors containing alkaline phosphatase cDNA, green fluorescent protein (GFP) cDNA, or vector alone were also produced using the same method. The retroviral packaging cell lines for LPAAT- β , LPAAT- α , and control vectors were produced and characterized as described (24). Clones producing high-titer virus (0.5 – 1×10^6 G418-resistant colony-forming units/ml) were used for the transduction of PC-3 and DU145 cells in their respective media; clonal cell lines were subsequently generated. None of the transduced cells exhibited any gross morphological or proliferative alterations.

LPAAT activity assay in cell lysates

The magnitude of overexpression of each recombinant LPAAT was determined by assaying total endogenous LPAAT activity in cell lysates essentially as described (25). Briefly, transduced PC-3 and DU145 cells were harvested by trypsinization, suspended in 400 μ l of ice-cold PBS, and probe-sonicated with a Branson Sonifier at output = 2, duty cycle = 2 for 10 pulses. The samples were centrifuged at 1,500 *g* for 2 min at 4°C to obtain the supernatants, which were quantified for protein. The LPAAT activity in these supernatants was estimated during a 7 min incubation at 37°C in 12 \times 75 mm silanized, borosilicate glass test tubes containing 50 μ l of 50 mM HEPES (pH 7.5), 100 mM NaCl, 1 mM EDTA, 1 mg/ml fatty acid-free BSA, 0.2 mM 14 C-labeled 18:1-CoA (\sim 40,000 cpm/assay), 0.2 mM *sn*-1-18:1 lysoPA, and 0.005–0.03 mg/ml protein; the reaction was quenched with organic solvent. The 14 C-labeled PA was resolved by TLC on Silica Gel 60 plates, and the density of radioactive bands was quantitated by exposing the plate to a phosphor screen overnight and scanning the phosphor screen with a Storm 840 PhosphorImager (GE Healthcare Life Sciences) as described (26). The increase in total LPAAT activity achieved in PC-3 cells was typically 5- to 20-fold for LPAAT- α and 20- to 45-fold for LPAAT- β . One clone from PC-3 cells containing recombinant LPAAT- β was used frequently in this study and is referred to as PC-3/ β 1 (or β 1 in Fig. 3D). The increase in total LPAAT activity achieved in DU145 cells typically was 2- to 5-fold for LPAAT- α and 4- to 10-fold for LPAAT- β .

Whole cell PM formation assay

PM synthesis was estimated in cultured cells by quantitating the conversion of exogenously added lysoPM and ^{14}C -labeled fatty acid to ^{14}C -labeled PM during a 35 min incubation at 37°C . PC-3 and DU145 cells, containing a recombinant protein as indicated, were separately plated onto 96-well plates in 500 μl of the appropriate medium (see above) and grown to 70–80% confluence. On the day of the assay, the medium volume was adjusted to 96 μl . The lysoPM and ^{14}C -labeled fatty acid stocks were prepared separately by drying each solvent under a stream of nitrogen and resuspending the lysoPM in PBS and the ^{14}C -labeled fatty acid in PBS + 20 mg/ml fatty acid-free BSA. The assay was performed by incubating 2 μl of the lysoPM stock with the cells for 5 min, adding 2 μl of the ^{14}C -fatty acid stock, and incubating for an additional 30 min. The final concentrations were 50 μM *sn*-1-18:1 lysoPM and 0.2 μCi of ^{14}C -labeled 18:1 in a total volume of 100 μl , unless stated otherwise. Background PM levels were determined by adding PBS without lysoPM. The assay was terminated by removing the incubation mixture, washing the cells three times with ice-cold PBS, discarding the final wash, and then adding 0.25 ml of methanol. The plates were gently swirled, and the methanol was transferred to glass test tubes. This method extracted >98% of the PM from the wells (data not shown). Carrier 18:1-18:1 PM (4 μg), 1.0 ml of chloroform, 0.25 ml of methanol, and 0.25 ml of 0.2 M H_3PO_4 were added to each tube. The tubes were vortexed for 10 s and centrifuged for 2 min at 500 g to clarify the phases; the upper phase was discarded. The lower phase was dried under a stream of nitrogen, and the sample was resuspended in 100 μl of chloroform-methanol (2:1). ^{14}C -labeled PM was typically resolved by TLC in chloroform/methanol/ NH_4OH (65:30:4). To help confirm the identity of the PM, in some instances a two-solvent system was also used in which the TLC plates were first developed in chloroform/methanol/ NH_4OH (70:25:3), dried thoroughly, and then developed in chloroform-methanol-acetic acid-water (85:10:8:1). The density of radioactive bands was scanned and quantified with a Storm PhosphorImager as described above. Unless stated otherwise, ^{14}C -labeled PM formation is expressed as a percentage of the lane: the pixel volume of the PM region \div the pixel volume of the entire lane $\times 100$, where pixel volume is the signal intensity calculated with the Storm PhosphorImager and is directly proportional to the amount of radioactivity. This calculation provided a relative rate of esterification of radioactive fatty acid into lysoPM versus the entire lipid fraction and served as a loading control. For experiments using LPAAT- β chemical inhibitors, the inhibitors were diluted from a DMSO stock into PBS + 4 mg/ml BSA. The volume of medium was adjusted to 94 μl , and 2 μl of inhibitor stock or excipient alone was added to the cells 5 min before the addition of the lysoPM stock. The final concentration of DMSO was 0.01% (v/v) and did not affect the outcome of the assay (data not shown). IC_{50} values were determined using GraphPad Prism version 4.00 for Windows (GraphPad Software, San Diego CA; www.graphpad.com). Other changes in the assay are noted in the text and figure legends.

Expression of LPAAT- β and LPAAT- α in Sf9 cells

For the construction of Baculovirus expression vectors, the full-length human LPAAT- α and LPAAT- β cDNAs were amplified by PCR from the DNA templates pCE9.LPAAT- α and pCE9.LPAAT- β (8) using the primers 5'-TGATATCCGAAGAAGATCT-TATGGATTTGTGGCCAGGGGC-3' (olpa1F) and 5'-CAGGCTC-TAGATCACCCACGCCCCAGGCTTC-3' (olpa1R) for LPAAT- α and the primers 5'-TGATATCCGAAGAAGATCTTATGGAGC-TGTGGCCGTGTC-3' (olpb1F) and 5'-CAGGCTCTAGACTACT-GGGCCGGCTGCAC-3' (olpb1R) for LPAAT- β . The ~ 870 bp

fragments generated were reamplified by PCR using the primers 5'-CCTACGTCGACATGGAACAAAAATTGATATCCGAAGAA-GATC-3' (olp2F) and olpa1R for LPAAT- α and the primers olp2F and olpb1R for LPAAT- β . The ~ 890 bp fragments generated were then cleaved with *Sal*I and *Xba*I for insertion into pFastBacTM HTc vector (Invitrogen) between the *Sal*I and *Xba*I sites. The plasmids pFB.LPAAT- α and pFB.LPAAT- β were further modified by cleaving them with *Rsa*I and *Sal*I, converting the cohesive ends to blunt ends with Klenow DNA polymerase and deoxy-nucleoside triphosphate and recircularizing by ligase to remove the polyhistidine tag coding region. These plasmids were then transformed into *E. coli* DH10BacTM (Invitrogen) for the generation of recombinant Bacmid DNAs for transfection into Sf9 insect cells for the production of recombinant Baculovirus stocks using the protocol described in the Bac-to-Bac[®] Baculovirus Expression System (Invitrogen), a eukaryotic expression system for generating recombinant Baculovirus through site-specific transposition in *E. coli*. Viral stocks harvested from the transfected cells were then used to infect fresh insect cells for the subsequent expression of LPAAT- α and LPAAT- β proteins.

Preparation of membranes from Sf9 cells

All steps were performed at 0 – 4°C . Sf9 cell pellets ($\sim 10^8$ cells) containing recombinant LPAAT were thawed and resuspended in 5 ml of buffer A [20 mM HEPES, pH 7.5, 1 mM DTT, 1 mM EDTA, 20% (w/v) glycerol, 1 mM benzamide, 1 $\mu\text{g}/\text{ml}$ soybean trypsin inhibitor, and 1 $\mu\text{g}/\text{ml}$ pepstatin A] with 1 mM Pefabloc. The cells were lysed by sonication using a Branson Sonifier at output = 2, duty cycle = 2, 10 pulses each for 10 s. DTT was then added to 1 mM from a 1 M stock. The samples were centrifuged at 1,500 rpm for 5 min. The resulting supernatant was centrifuged at 100,000 g for 1 h. The resulting pellet, which contained the Sf9 membranes, was resuspended in buffer A + 1 mM DTT and stored at -70°C . LPAAT- β and LPAAT- α activities were stable in this form for >2 years.

LPAAT activity assay in Sf9 membranes

Enzyme assays were performed in 12×75 mm silanized, borosilicate glass test tubes at 37°C for 3–10 min. For the estimation of LPAAT activity in Sf9 membrane preparations, typical assay mixtures (300 μl) consisted of 251 μl of buffer B (50 mM HEPES, pH 7.5, 1 mM EDTA, and 100 mM NaCl), 25 μl of enzyme in buffer B + 10% glycerol, 12 μl of ^{14}C -labeled 18:1-CoA (40,000–100,000 cpm), and 12 μl of *sn*-1-18:1 lysoPA. To prevent the nonspecific binding of substrates to the tubes, stock solutions of each substrate were prepared separately in DMSO. The final concentration of DMSO in all assays was 8% (v/v) and did not affect LPAAT- β or LPAAT- α activity as determined in separate experiments (data not shown). At the end of the assay, 1 ml of chloroform/methanol/HCl (1:2:0.027), carrier PA and lysoPA (0.3 μg), 0.3 ml of chloroform, and 0.1 ml of water were added. The tubes were vortexed for 10 s and centrifuged for 2 min at 500 g to clarify the phases; the upper phase was discarded. The lower phase was washed with 1 ml of chloroform-methanol-water (10:150:100) and then dried under a stream of nitrogen. ^{14}C -labeled PA was resolved by TLC in chloroform-methanol-acetic acid-water (85:12.5:12.5:3) and quantitated using the Storm PhosphorImager as described above. Product formation was linear with time and protein concentration. V_{max} and K_m values were determined using GraphPad Prism software. If other potential acyl acceptors were used in the assay, the following TLC solvent systems were used: for lysophosphatidylinositol, lysophosphatidylserine, or lysophosphatidylglycerol, chloroform-methanol-acetic acid-water (85:12.5:12.5:3); for lysoPC and

lysophosphatidylethanolamine, chloroform-methanol-ammonia (65:40:5); for monoolein and diolein, hexane-diethylether-acetone (60:40:5); and for glycerol-3-phosphate, chloroform-methanol-acetic acid-water (70:30:4:6).

For competition experiments between two different acyl-CoA species, [^3H] *sn*-1-18:1 lysoPA, ^{14}C -labeled 18:1-CoA, and a second, unlabeled acyl-CoA were included in the reaction. After extraction and TLC, the PA band was scraped from the plate and extracted from the silica with 1.3 ml of chloroform-methanol-water (1:1:0.1). The extract was transferred to a glass scintillation vial and dried. Samples were counted in Ultima Gold scintillant (Perkin-Elmer) on a Packard Tri-Carb 1600 Scintillation Counter using the preset $^3\text{H}/^{14}\text{C}$ channels (pulse height discriminator settings of 0–12 for ^3H and 12–156 for ^{14}C) and the tSIE/AEC quench indicating parameter. The amount of 18:1 groups transferred equaled the moles of ^{14}C -labeled PA formed, and the amount of the second acyl group transferred equaled the

moles of ^3H -labeled PA formed minus the moles of ^{14}C -labeled PA formed. Relative, normalized velocities were calculated from the equation $v_1/v_2 = [S_1]/[S_2] \times [(k_{\text{cat}1}/K_{\text{m}1})/(k_{\text{cat}2}/K_{\text{m}2})]$, where v = initial velocity, $[S]$ = substrate concentration, and k_{cat} and K_{m} are catalytic constants for substrates 1 and 2 as indicated (27).

Other methods

Protein concentrations were assayed using the bicinchoninic acid method (Pierce) with BSA as the standard. NuPAGE (Invitrogen) was performed according to the manufacturer's instructions, and Western blotting with anti-LPAAT- β and anti-LPAAT- α monoclonal antibodies was performed as described (28). Error bars in the figures indicate standard deviations unless only two values were obtained, in which cases the standard errors of means are shown. Error bars that are not visible are smaller than the data point symbols.




Fig. 1. Acyl-CoA molecular species preferences of lysophosphatidic acid acyltransferase- β (LPAAT- β) and LPAAT- α . A: Preparations of Sf9 wild-type membranes or those containing recombinant LPAAT- β or LPAAT- α were analyzed by NuPAGE and Western blotting, as described in Experimental Procedures. Separate blots were incubated with either the anti-LPAAT- α antibody or the anti-LPAAT- β antibody as indicated. Positions of the molecular mass markers are indicated on the right side of the blot. B: The preferences of the LPAAT enzymes for different molecular species of acyl-CoA were determined by measuring the amount of phosphatidic acid (PA) formed in separate incubation assays (8 min, 37°C) containing buffer B, 8 μM ^{33}P -labeled *sn*-1-18:1 lysophosphatidic acid (lysoPA), 8 μM of the indicated molecular species of acyl-CoA, and 0.1 $\mu\text{g}/\text{ml}$ Sf9 membrane preparations; wild-type membranes or those containing recombinant LPAAT- β or LPAAT- α were used as indicated on the abscissa. Two experiments were performed that used all three membrane preparations, and a third experiment was performed that used only the Sf9 membrane preparations containing recombinant LPAAT- β . Results from one representative experiment are shown, and all three experiments gave similar results. Error bars represent standard errors of the means from duplicate samples. C: Sf9 membrane preparations containing recombinant LPAAT- β were incubated as described for B except with the indicated concentrations of 18:1-CoA. The experiment was performed in duplicate, and the results shown are representative of three experiments.

RESULTS

Expression of LPAAT- β and LPAAT- α in Sf9 cells

We expressed the LPAAT- β and LPAAT- α transgenes in Sf9 cells under the control of the SV40 promoter; membrane preparations were generated as described in Experimental Procedures. Membrane preparations from cells infected with LPAAT- β or LPAAT- α cDNA could catalyze the transfer of oleoyl groups from 18:1-CoA to *sn*-1-18:1 lysoPA at rates of 205 ± 36 nmol/min/mg protein ($n = 5$) and 115 ± 48 nmol/min/mg protein ($n = 4$), respectively, which are 110- and 60-fold higher than the rate observed using wild-type Sf9 membranes. Furthermore, when these membrane preparations were analyzed by Western blotting using anti-LPAAT- β and anti-LPAAT- α monoclonal antibodies, the anti-LPAAT- β antibody recognized a band at the expected molecular mass of 31 kDa only in membranes containing recombinant LPAAT- β , whereas the anti-LPAAT- α antibody recognized a band of similar molecular mass only in membranes containing recombinant LPAAT- α (Fig. 1A). We originally planned to solubilize and purify the LPAAT isozymes before studying their enzymatic properties. Unfortunately, this proved inefficient, as both activities were immediately and completely inactivated in the presence of detergents. On the other hand, it appeared that both isozymes were over-expressed in the membrane fractions and accounted for >98% of the total LPAAT enzymatic activity. Thus, it was possible to use the membrane preparations to separately study each isoform, without the concern for other interfering LPAAT activities in the assay.

Acyl donor and acceptor preferences of LPAAT- β

Several sets of experiments were performed to examine the acyl donor preference of LPAAT- β . One set examined the ability of LPAAT- β to use different molecular species of acyl-CoA. Sf9 membrane preparations containing recombinant LPAAT- β (see Experimental Procedures) were separately incubated with *sn*-1-18:1 lysoPA and different molecular species of acyl-CoA. The results revealed that LPAAT- β could use all acyl-CoAs tested (Fig. 1B). When the concentration of 18:1-CoA was varied in the assay, LPAAT activity was saturable, with an apparent V_{max} of 210 nmol/min/mg protein and an apparent K_m of 0.4 μ M (Fig. 1C). The catalytic site specificity for an enzyme is defined as the relative values of the specificity constant (V_{max}/K_m) for different substrates (27). As these values cannot be derived from assays using one concentration of one acyl donor, as shown in Fig. 1B, competition experiments were performed in which Sf9 membrane preparations containing recombinant LPAAT- β were incubated with 3 H-labeled *sn*-1-18:1 lysoPA, 14 C-labeled 18:1-CoA, and a second, unlabeled acyl-CoA in the same tube. The amount of each acyl group transferred to lysoPA was determined as described in Experimental Procedures, and the ratio of these two values represents the specificity of 18:1-CoA to the second acyl-CoA. The results shown in Table 1 suggest that, compared with 18:1-CoA, LPAAT- β displays approximately equal preference toward 16:0-CoA and 18:0-CoA, a moderate

TABLE 1. Competition between 18:1-CoA and selected acyl-CoAs in the LPAAT- β -catalyzed reaction

14 C-Labeled Acyl-CoA	12 C-Labeled Acyl-CoA	Relative Normalized Velocities (18:1-CoA \div acyl-CoA)
18:1	12:0	>18
18:1	14:0	>18
18:1	16:0	1.8 ± 0.9
18:1	18:0	1.0 ± 0.3
18:1	18:1	1.2 ± 0.4
18:1	18:2	8.5 ± 5.1
18:1	20:0	>18
18:1	20:4	>18

LPAAT- β , lysophosphatidic acid acyltransferase- β . The ability of LPAAT- β to catalyze the transfer of 18:1 acyl groups over other acyl groups was measured in a single reaction consisting of Sf9 membranes containing recombinant LPAAT- β , 6 μ M 3 H-labeled lysophosphatidic acid, 0.6 μ M 14 C-labeled 18:1-CoA, and either 0.6 μ M 20:0-CoA or 1.2 μ M of the indicated acyl-CoA for 7 min at 37°C; relative normalized velocities were calculated as described in Experimental Procedures. Results are from two experiments for 14:0-CoA and 20:0-CoA and from three experiments for 20:4-CoA (the same result was observed each time); results are averages \pm SD from four separate experiments for all others. The limit of detection in the assay was a ratio of 18.

preference (8.5-fold) compared with 18:2-CoA, and an at least 18-fold greater preference compared with 12:0-CoA, 14:0-CoA, 20:0-CoA, and 20:4-CoA.

Mouse LPAAT- α also catalyzes a CoA-dependent transacylase reaction, in which acyl groups are transferred from donor phospholipids to acceptor lysophospholipids in the presence of unesterified CoA, although at a much slower rate compared with its acyltransferase activity (15). When Sf9 membrane preparations containing recombinant human LPAAT- β were incubated with 3 H-labeled *sn*-1-18:1 lysoPA and unesterified CoA, no PA formation was detected. Furthermore, unesterified CoA did not affect the rate of LPAAT- β -catalyzed acylation of *sn*-1-18:1 lysoPA with 18:1-CoA (data not shown). Thus, LPAAT- β does not appear to display a CoA-dependent transacylase activity in this assay.

Experiments were also performed to elucidate the acyl acceptor preference of LPAAT- β . Sf9 membrane preparations containing recombinant LPAAT- β were incubated separately with 14 C-labeled 18:1-CoA and various *sn*-1-16:0, *sn*-1-18:0, or *sn*-1-18:1 lysophospholipids. The results showed that LPAAT- β transferred oleoyl groups to lysoPA; transfer to the other lysophospholipids was never consistently above background levels (Fig. 2A). LPAAT- β was also unable to use glycerol-3-phosphate, 1-monoolein, or dioleoylglycerol as an acyl acceptor (data not shown). When the concentration of *sn*-1-18:1 lysoPA was varied in the assay, LPAAT activity was saturable, with a representative V_{max} of 200 nmol/min/mg protein and an apparent K_m of 2.0 μ M (Fig. 2B). We then examined the nature of the head group specificity of LPAAT- β more closely using *sn*-1-18:1 lysoPM as a substrate, which represents a very small modification to the phosphate moiety of lysoPA but significantly affects the charge and hydrophobicity of the molecule. When *sn*-1-18:1 lysoPM was substituted for lysoPA in the LPAAT membrane assay, PM was formed at lysoPM concentrations similar to that of lysoPA. At 3 μ M




Fig. 2. Acyl acceptor preferences of LPAAT- β and LPAAT- α . **A:** The preferences of the LPAAT enzymes for different lysophospholipids were determined by measuring the amount of phospholipid formed in separate incubation assays (10 min, 37°C) containing buffer B, 5 μ M 14 C-labeled 18:1-CoA, 5 μ M of the indicated lysophospholipid, and 0.1 μ g/ml Sf9 membrane preparations: wild-type membranes (black bars) or those containing recombinant LPAAT- β (white bars) or LPAAT- α (striped bars). Two experiments were performed that used all three membrane preparations, and a third experiment was performed that used only the Sf9 membrane preparations containing recombinant LPAAT- β . Results from one representative experiment are shown, and all three experiments gave similar results. Error bars represent standard errors of the means from duplicate samples. Abscissa labels are as follows: oLPA, *sn*-1-18:1 lysoPA; sLPA, *sn*-1-18:0 lysoPA; pLPA, *sn*-1-16:0 lysoPA; oLPC, *sn*-1-18:1 lysophosphatidylcholine (lysoPC); sLPC, *sn*-1-18:0 lysoPC; oLPE, *sn*-1-18:1 lysophosphatidylethanolamine; oLPG, *sn*-1-18:1 lysophosphatidylglycerol; LPI, liver lysophosphatidylinositol (primarily the *sn*-1-18:0 species); oLPS, *sn*-1-18:1 lysophosphatidylserine. **B:** Sf9 membrane preparations containing recombinant LPAAT- β were incubated as described for A except with the indicated concentrations of *sn*-1-18:1 lysoPA. **C:** Membrane preparations from Sf9 wild-type cells (triangles) or cells that overexpressed LPAAT- β (squares) or LPAAT- α (circles) were incubated as described for A except with the indicated concentrations of *sn*-1-18:1 lysophosphatidylmethanol (lysoPM). **D:** Sf9 membrane preparations containing recombinant LPAAT- β were incubated as described for A except with 5 μ M *sn*-1-18:1 lysoPA (circles) or 5 μ M *sn*-1-18:1 lysoPM (squares) and with the indicated concentrations of CT-32228. Activities without inhibitor were 223 nmol PA formed/min/mg protein and 23 nmol phosphatidylmethanol (PM) formed/min/mg protein. The results shown in B–D are representative of at least two separate experiments performed in duplicate.

lysoPM, this activity was at least 20-fold higher than the same reaction using wild-type Sf9 membranes (Fig. 2C) but 20-fold lower than the corresponding LPAAT- β reaction using lysoPA (compare Fig. 2B, C). Together, these results suggest that lysoPA is the preferred acyl acceptor for LPAAT- β .

Acyl donor and acceptor preferences of LPAAT- α

Experiments were performed to examine the substrate preferences of the LPAAT- α activity, and the results showed many similarities with LPAAT- β . For example, like LPAAT- β , LPAAT- α activity was able to transfer the

fatty acyl groups of all acyl-CoAs tested to *sn*-1-18:1 lysoPA (Fig. 1B), and LPAAT- α activity used lysoPA exclusively as the acyl acceptor over other lysophospholipids, glycerol-3-phosphate, 1-monoolein, and diolein (Fig. 2A and data not shown). In contrast, LPAAT- α activity was unable to transfer oleoyl groups from 18:1-CoA to *sn*-1-18:1 lysoPM (Fig. 2C).

Intracellular formation of PM from exogenous lysoPM

When *sn*-1-18:1 lysoPM and 14 C-labeled oleate were added to PC-3 or DU145 cells, a radioactive band was formed that comigrated with authentic 18:1-18:1 PM by

TLC (Fig. 3A and data not shown). This band was absolutely dependent on the presence of the lysoPM and was confirmed using two separate solvent systems (see Experimental Procedures). PM formation was proportional with the concentration of *sn*-1-18:1 lysoPM and time (Fig. 3B, C). Given that the LPAAT activity assay in Sf9 membranes suggested that lysoPM was an acyl acceptor in the LPAAT- β but not the LPAAT- α -catalyzed reaction, we next examined whether LPAAT- β played a role in forming PM in these cells. Experiments were performed in which PC-3 wild-type cells or populations transduced with LPAAT- β , LPAAT- α , GFP, or vector only were incubated with *sn*-1-18:1 lysoPM and ^{14}C -labeled 18:1, and ^{14}C -labeled PM formation was measured. Although lysates from cells transduced with either LPAAT- α or LPAAT- β showed a 5- to 25-fold increase in total LPAAT enzymatic activity compared with wild-type or vector control cells (Fig. 3D), only LPAAT- β -transduced cells showed an increase (3- to 5-fold) in ^{14}C -labeled PM formation (Fig. 3E). In PC-3/ β 1, a PC-3 clonal cell line transduced with LPAAT- β , PM formation was proportional with lysoPM concentration and time (Fig. 3B, C). Similar results were observed in separate experiments that used DU145 wild-type cells and DU145 cells transduced with LPAAT- β , LPAAT- α , alkaline phosphatase, or vector only (data not shown). In addition, in similar experiments in which *sn*-1-16:0 lysoPM was added to PC-3 or PC-3/ β 1 cells, PM formation was also observed (data not shown).

Cell Therapeutics, Inc. has developed triazine-based compounds that inhibit LPAAT- β biochemical activity at sub-200 nM IC_{50} concentrations but do not inhibit the activities of LPAAT- α , γ , δ , or ϵ , phospholipase D₁ (PLD₁), or PLD₂ at 40 μM (Table 2) (20, 28). When LPAAT activity was assayed in Sf9 membranes containing recombinant LPAAT- β , CT-32228 inhibited LPAAT- β activity when either lysoPA or lysoPM was used as the acyl acceptor (Fig. 2D), with similar IC_{50} values. Furthermore, when LPAAT activity was measured in PC-3 or DU145 cell lysates using lysoPA as the acyl acceptor (see Experimental Procedures), CT-32228 at saturating (10 μM) concentrations inhibited $58 \pm 6\%$ ($n = 4$) of the total endogenous LPAAT activity in PC-3 lysates and $66 \pm 3\%$ ($n = 7$) of the total endogenous LPAAT activity in DU145 lysates. As these lysates could potentially contain LPAAT- β , LPAAT- α , and perhaps other LPAAT activities, and the assay, which uses 18:1-CoA and *sn*-1-18:1 lysoPA as substrates, will not necessarily distinguish between these activities, we interpret these data as indicating that LPAAT- β accounts for $\sim 60\%$ of the total endogenous LPAAT activity in these cells. Similar results were reported by Hideshima et al. (25).

Several experiments were then performed to test the effect of LPAAT- β inhibitors on the incorporation of ^{14}C -labeled 18:1 into lysoPM in cells. Figure 3E shows the effect of CT-32228 on PM formation in PC-3 cells. The results indicate that CT-32228 inhibited $\sim 50\%$ of the PM formation in wild-type, control-transduced, and LPAAT- α -transduced cells and 60–75% of the PM formation in LPAAT- β -transduced cells. The effect of the inhibitors on

PM formation appeared to be saturable, as separate experiments showed that the IC_{50} in PC-3/ β 1 cells for PM formation by CT-32228 was 321, 353, and 1,940 nM (the reason for the discrepancy in the IC_{50} values is unknown) and the IC_{50} for PM formation by CT-32501 was 460 ± 180 nM ($n = 5$) (Fig. 4). At saturating concentrations, CT-32501 inhibited as much as 95% and CT-32228 inhibited 60–75% of the PM formation in PC-3/ β 1 cells. One explanation for these differences in maximal inhibition might be the distribution of the inhibitors in the cell. It is important to note that CT-32212, a structurally similar compound to the LPAAT- β inhibitors but a weak inhibitor (Table 1), does not affect PM formation (Fig. 4). In similar experiments using DU145 cells transduced with LPAAT- β , the IC_{50} for PM formation by CT-32228 was 900 ± 400 nM ($n = 3$), whereas CT-32212 did not inhibit PM formation at 20 μM (data not shown).

As the LPAAT activity assay in Sf9 membranes suggested that LPAAT- β appears to prefer the transfer of 18:1 over 20:4 acyl groups by at least 18-fold (Table 1), experiments were performed to test whether this specificity was also observed in cells. PC-3 wild-type or PC-3/ β 1 cells were incubated with *sn*-1-18:1 lysoPM and with either ^{14}C -labeled 18:1 or ^{14}C -labeled 20:4. ^{14}C -labeled 20:4 was also incorporated into PM in a time-dependent and lysoPM concentration-dependent manner (data not shown). However, unlike ^{14}C -labeled 18:1, there was no significant difference in the incorporation of ^{14}C -labeled 20:4 into PM between the two cell lines (Fig. 5). Furthermore, CT-32501 inhibited the incorporation of ^{14}C -labeled 18:1 into lysoPM but did not inhibit the incorporation of ^{14}C -labeled 20:4 into lysoPM (Fig. 5). These data provide evidence, from a second assay method, for the specificity of LPAAT- β toward the transfer of 18:1 acyl groups over 20:4 acyl groups.

Although the structure of lysoPM suggests that it has amphipathic properties, it does not appear to damage the cells. Incorporation of ^{14}C -labeled 18:1 into lysoPM and other lipids was linear to at least 100 μM lysoPM and for at least 60 min of incubation time in PC-3 and DU145 cells. If lysoPM was disrupting cellular membranes, ATP would likely be released from the cell, preventing the acyl-CoA synthetase-catalyzed conversion of the exogenously added ^{14}C -labeled 18:1 into 18:1-CoA, a necessary step before the acylation of lysophospholipids. Second, incubation of DU145 or MCF7 cells with 50 μM lysoPM for 1–3 days did not affect their proliferation rates (data not shown). Third, as with lysoPM, 50 μM lysoPC can be added to cells without any noticeable deleterious effects on the membrane integrity of the cells (29).

It is important to note that similar experiments in which we tried to label cellular lipids with exogenously added ^{33}P -labeled lysoPA did not result in any incorporation of ^{33}P label into the lipid fraction (data not shown). This may be attributable in part to the inability of lysoPA to traverse the plasma membrane and/or to the action of one or more lipid phosphate phosphohydrolases that convert the lysoPA to inorganic phosphate and monoacylglycerol (30). In a separate matter, a significant amount of ^{14}C -labeled




Fig. 3. Acylation of *sn*-1-18:1 lysoPM with ^{14}C -labeled 18:1 in PC-3 cells: effects of time, lysoPM concentration, and total LPAAT activity. **A:** Separate wells containing wild-type (wt) PC-3 cells or PC-3 cells transduced with LPAAT- β (β 1), LPAAT- α (α 1), or green fluorescent protein (GFP) cDNAs were incubated with ^{14}C -labeled 18:1 (220,000 cpm) and the indicated concentrations of *sn*-1-18:1 lysoPM and CT-32228, as described in Experimental Procedures. After lipid extraction, TLC, and exposure of the plate to a phosphor screen (see Experimental Procedures), the screen was scanned with a Storm PhosphorImager in phosphor mode. The migration of lipid standards is indicated: NL, neutral lipids; PE, phosphatidylethanolamine; PC, phosphatidylcholine; PI/PS, phosphatidylinositol/phosphatidylserine; ori, origin. **B:** Effect of lysoPM concentration on PM formation. PC-3 (squares) and PC-3/ β 1 (circles) cells were incubated (40 min, 37°C) in the presence of $0.2\ \mu\text{Ci}$ of ^{14}C -labeled 18:1 and the indicated concentrations of *sn*-1-18:1 lysoPM. Samples were processed as described for A, and ^{14}C -labeled PM was quantitated and normalized to the total radioactivity detected in the lane. Results are representative of two experiments performed in triplicate. Error bars in all panels indicate SD. **C:** Rate of PM formation. PC-3 (squares) and PC-3/ β 1 (circles) cells were incubated in the presence of $50\ \mu\text{M}$ *sn*-1-18:1 lysoPM and $0.2\ \mu\text{Ci}$ of ^{14}C -labeled 18:1 for the indicated times at 37°C . Samples were processed and radioactivity quantitated as described for B, except that the data for this panel are expressed in cpm PM, because PM formation expressed as the percentage of the lane did not change over time. Results are representative of two experiments performed in triplicate. **D:** Total endogenous LPAAT activity in PC-3 wild-type (WT) cells and PC-3 cells transduced with retroviral vectors containing no transcript (X1 and X2), GFP (GFP1, GFP2, GFP3), LPAAT- α (α 1, α 2, α 3), or LPAAT- β (β 1, β 2, β 3) cDNA was measured with the LPAAT activity assay in cell lysates (see Experimental Procedures). For reference, total LPAAT activity in the lysates from wild-type cells was 18 ± 5 ($n = 8$) nmol PA formed/min/mg protein. **E:** Effect of the overexpression of LPAAT- β or LPAAT- α on ^{14}C -labeled PM formation in PC-3 cells. PC-3 wild-type cells or cells transduced as described for D were incubated separately (40 min, 37°C) in the presence of $0.4\ \mu\text{Ci}$ of ^{14}C -labeled 18:1 and $50\ \mu\text{M}$ *sn*-1-18:1 lysoPM (black bars). In some cases, $10\ \mu\text{M}$ of the LPAAT- β inhibitor CT-32228 was added to the cells 12 min before the addition of the ^{14}C -labeled 18:1 and the lysoPM (white bars). Approximately 4% of the total radioactivity was incorporated into the lipid fraction for each of the clones. Results are representative of two experiments performed in duplicate. Note that β 1 described in A, D, and E refers to the PC-3/ β 1 clonal cell line.

TABLE 2. LPAAT- β inhibitors

CT No.	Structure	LPAAT- β IC ₅₀	LPAAT- α IC ₅₀	Reference
		<i>nM</i>		
32228		60	>40,000	28
32501		27	>40,000	19
32212		36,000	>40,000	20

LPAAT activity was measured in Sf9 membranes containing recombinant LPAAT- β or LPAAT- α , as described previously (20).

20:4 is incorporated into PM (Fig. 5). Although LPAAT- β does not appear to be involved, other acyltransferase activities that transfer 20:4 acyl groups to various lysophospholipids (3) may also acylate lysoPM.

DISCUSSION

This report describes the enzymatic properties of LPAAT- β and LPAAT- α and introduces a novel assay to estimate intracellular LPAAT- β activity. We designed the LPAAT activity assay in Sf9 membranes under strict guidelines in an attempt to ensure that the assay did not mask the properties of the enzyme in question. First, the assay




Fig. 4. Inhibition of the acylation of lysoPM in PC-3/β1 cells by LPAAT- β inhibitors. ¹⁴C-labeled PM formation was determined by separately incubating PC-3/β1 cells with the indicated concentrations of CT-32228 (squares), CT-32501 (circles), or CT-32212 (triangles), 0.2 μCi of ¹⁴C-labeled 18:1, and 50 μM *sn*-1-18:1 lysoPM, as described in Experimental Procedures. Results are representative of three to five experiments performed in triplicate. Error bars indicate SD.




Fig. 5. ¹⁴C-labeled 18:1 or ¹⁴C-labeled 20:4 acylation of lysoPM in PC-3 and PC-3/β1 cells. ¹⁴C-labeled PM formation was determined by separately incubating PC-3 cells (black bars) or PC-3/β1 cells (white bars) with the indicated concentrations of CT-32501, 50 μM *sn*-1-18:1 lysoPM, and either 0.2 μCi of ¹⁴C-labeled 18:1 or 0.2 μCi of ¹⁴C-labeled 20:4, as described in Experimental Procedures. Results are representative of three experiments performed in triplicate. Error bars indicate SD.

used membranes from Sf9 cells that overexpressed either human LPAAT- β or human LPAAT- α . Although not their natural membrane, each enzyme was separately expressed with high activity such that other acyltransferase activities were not detected. Second, BSA, often included in assays with acyl-CoA-metabolizing enzymes, was not included because it has submicromolar affinities for long-chain acyl-CoA (31–33) and lysoPA (34) and would most likely influence the substrates' availability to the LPAAT. Indeed, inclusion of BSA in the membrane assay did not significantly affect the V_{max} of the reaction but did increase the apparent K_m of the substrates (data not shown). Third, the substrate concentrations were kept below reported critical micellar concentration values (35–38), as changes in the physical structure of the substrates during the assay would make interpretation of the data nearly impossible. With these guidelines established, it is interesting that LPAAT- α and LPAAT- β showed subtle differences in their preference for different acyl-CoAs: 14:0-CoA, 16:0-CoA, and 18:2-CoA were most active in the LPAAT- α reaction, whereas 18:1-CoA was most active in the LPAAT- β reaction (Fig. 1B). Furthermore, only LPAAT- β was able to use lysoPM as a substrate (Fig. 2C). Although greater confidence in the biochemical characterization of these isozymes will have to await their purification and insertion into well-defined lipid vesicles, it seems reasonable to assume that the membrane assay displays, at least in part, the properties of each LPAAT and is not a limitation of the system.

To confirm the results observed in the LPAAT activity assay in Sf9 membranes, a second, cell-based assay was used in which LPAAT- β activity was measured by acylation of lysoPM. LPAAT- β appears to acylate lysoPM in cells based on the following observations. 1) Using the membrane assay, lysoPM was shown to be a substrate for LPAAT- β but not LPAAT- α (Fig. 2C). 2) Addition of lysoPM and ¹⁴C-labeled 18:1 to PC-3 or DU145 cells resulted in the formation of a ¹⁴C-labeled lipid that comigrated with

authentic PM by TLC in two separate solvent systems; the formation of this band is absolutely dependent on the presence of lysoPM (Fig. 3A). 3) Cells transduced with LPAAT- β (5- to 20-fold higher total LPAAT activity as measured in cell lysates) showed greater incorporation of label into PM than the corresponding wild-type cells or cells transduced with LPAAT- α , GFP, or vector only (Fig. 3D, E and data not shown). 4) Addition to cells of isoform-specific, small molecule LPAAT- β inhibitors decreased the incorporation of ^{14}C -labeled 18:1 into PM (Figs. 3–5). 5) Treatment of DU145 cells, either wild type or containing recombinant LPAAT- β , with LPAAT- β small, interfering RNA resulted in a knockdown of LPAAT- β activity, as measured in cell lysates, and a decrease in the incorporation of ^{14}C -labeled 18:1 into PM, as measured in cells (M. Coon, personal communication).


It is curious that there is an apparent discrepancy between the observed lysoPM acylation rates in the membrane assay and the whole cell assay. In the membrane assay, lysoPM is acylated by LPAAT- β at a much lower rate than lysoPA (compare Fig. 2B, C), but in the whole cell assay, PM formation accounts for as much as 5% and 25% of the total radioactivity incorporated into the lipid fraction of PC-3 and PC-3/ β 1 cells, respectively (Fig. 3E). There are several possible explanations. First, the membrane assay suggests that both substrates are active at similar concentrations, suggesting similar affinities for LPAAT- β . Second, the whole cell assay has the advantage of measuring LPAAT- β activity within an active membrane, perhaps within a unique microenvironment that is not duplicated in the membrane assay. Third, the whole cell assay may take into account proteins that may affect LPAAT- β activity. For example, the cellular acyl-CoA pool appears to be controlled by acyl-CoA binding proteins and fatty acid binding proteins, and these soluble, 10–15 kDa proteins can affect the activity of enzymes in which acyl-CoA is a substrate, product, or inhibitor (39–41), including LPAAT activities (42–44). Some or all of these factors may account for the robust, LPAAT- β -dependent PM signal in the whole cell assay.

We have tried to show an effect on lipid metabolism in cells that overexpress LPAAT- β or with the use of the LPAAT- β isoform-specific inhibitors. For example, we incubated ECV304, DU145, MCF7, and NIH3T3 cells with ^{14}C -labeled 18:1 or with ^{14}C -labeled glycerol, or we incubated DU145 cells with ^{32}P -labeled phosphate. Surprisingly, we did not detect any change in the pattern of incorporation of the radioactive tracer into PA or the major glycerolipids or any change in the total mass of each lipid between wild-type cells and the LPAAT- β -overexpressing populations (data not shown). Similar observations have been noted for overexpressed PLD isozymes (45) and certain *E. coli* lipid-metabolizing enzymes (46). Furthermore, the only significant changes caused by the addition of LPAAT- β inhibitors was on the accumulation of lysoPA in 3T3-L1-derived adipocytes incubated with ^{14}C -labeled 18:1, or in DU145 cells incubated with propranolol, a lipid phosphate phosphohydrolase inhibitor, plus ^{32}P -labeled phosphate (data not shown). There may be several pos-

sible explanations for why a clear PM signal is observed in the whole cell assay but a change in PA metabolism or mass is not observed in the assays mentioned above. Exogenously added 18:1, glycerol, and phosphate must first be converted to acyl-CoA and/or lysoPA. These substrates can then be used by LPAAT- β , but also by other enzymes. In addition, 18:1, glycerol, and phosphate can each be incorporated into lipids by other pathways. These factors may dilute the signal we are trying to observe from LPAAT- β . On the other hand, the fate of the lysoPM may be more limited, as LPAAT- β and perhaps a few other unidentified acyltransferases are able to acylate it, and the resulting PM may not be metabolized as rapidly as PA.

It would appear that the major product of the LPAAT- β -catalyzed reaction is PA of a restricted molecular species, based on the following observations. First, data from the LPAAT activity assay in Sf9 membranes suggest that lysoPA but not other lysophospholipids, glycerol-3-phosphate, 1-monooleoylglycerol, or dioleoylglycerol could be used as an acyl acceptor (Fig. 2A and data not shown). Second, JC201 cells, an *E. coli* strain with a temperature-sensitive mutation within its sole LPAAT gene, can be rescued at a nonpermissive temperature by expression of LPAAT- β (8). Third, LPAAT- β appears to be specific for the transfer of 16:0, 18:0, and 18:1 acyl groups over 12:0, 14:0, and 20 carbon acyl groups (Table 1). Fourth, the high selectivity of LPAAT- β for the transfer of 18:1 groups over 20:4 groups was also observed in the whole cell assay (Fig. 5).

Questions still remain concerning the fate of the PA produced by LPAAT- β . With the cloning of six LPAAT genes, questions arise regarding the role of each isoform's cellular functions (47). Based on two separate assays in this study, the fact that 20:4 groups are not transferred if 18:1 groups are present strongly suggests that LPAAT- β is not closely involved in synthesizing glycerolipids esterified with 20:4 that are fated for the eicosinoid pathways. On the other hand, LPAAT- β may function within a lipid synthetic pathway. It is widely accepted that de novo-synthesized glycerolipids contain primarily monoenoic and dienoic acyl groups at the *sn*-2 position (48). Our data suggest that LPAAT- β can contribute to this molecular species profile. Consistent with this, LPAAT- β is highly expressed in adipocytes, and Lu et al. (47) have suggested that mouse LPAAT- β expression is regulated by peroxisome proliferator-activated receptor α , a transcription factor that regulates the expression of lipid-metabolizing enzymes. Patients with a form of congenital generalized lipodystrophy, a disorder characterized by a lack of nonmechanical adipose tissue from birth, contain mutations in the LPAAT- β gene, rendering it nonfunctional (11, 13, 49). Although a simple explanation is that patients without active LPAAT- β are unable to synthesize triglyceride in adipocytes, they have functional LPAAT- α , which is also present in healthy adipocytes, produce high levels of triglyceride in the plasma compartment, and appear to have normal mechanical fat. Therefore, it is of interest that these two isoforms appear not to have entirely redundant biological functions. To make matters more complex, PA of molecular species similar to those described above may

also play a role in signaling (45, 50, 51), and evidence has been presented raising the possibility that LPAAT- β may contribute to tumor progression (20). Further study will be required to address questions regarding the cellular role of LPAAT- β . 

The authors thank Dr. Mike Gelb for his many helpful discussions. The authors thank Dr. Peter Klein's laboratory for the synthesis of CT-32228, CT-32212, and CT-32501, Dr. Anil Kumar for the synthesis of lysoPM, and Tod Martin, ManShun Lai, Dr. Jessica Freiberg, and Dr. Scott Shaffer for their excellent technical assistance.

REFERENCES

- Dircks, L., and H. S. Sul. 1999. Acyltransferases of de novo glycerophospholipid biosynthesis. *Prog. Lipid Res.* **38**: 461–479.
- Athenstaedt, K., and G. Daum. 1999. Phosphatidic acid, a key intermediate in lipid metabolism. *Eur. J. Biochem.* **266**: 1–16.
- Yamashita, A., T. Sugiura, and K. Waku. 1997. Acyltransferases and transacylases involved in fatty acid remodeling of phospholipids and metabolism of bioactive lipids in mammalian cells. *J. Biochem. (Tokyo)*. **122**: 1–16.
- Leung, D. W. 2001. The structure and functions of human lysophosphatidic acid acyltransferases. *Front. Biosci.* **6**: D944–D953.
- Lewin, T. M., P. Wang, and R. A. Coleman. 1999. Analysis of amino acid motifs diagnostic for the sn-glycerol-3-phosphate acyltransferase reaction. *Biochemistry*. **38**: 5764–5771.
- Haque, W., A. Garg, and A. K. Agarwal. 2005. Enzymatic activity of naturally occurring 1-acylglycerol-3-phosphate-O-acyltransferase 2 mutants associated with congenital generalized lipodystrophy. *Biochem. Biophys. Res. Commun.* **327**: 446–453.
- Dircks, L. K., J. Ke, and H. S. Sul. 1999. A conserved seven amino acid stretch important for murine mitochondrial glycerol-3-phosphate acyltransferase activity. Significance of arginine 318 in catalysis. *J. Biol. Chem.* **274**: 34728–34734.
- West, J., C. K. Tompkins, N. Balantac, E. Nudelman, B. Meengs, T. White, S. Bursten, J. Coleman, A. Kumar, J. W. Singer, et al. 1997. Cloning and expression of two human lysophosphatidic acid acyltransferase cDNAs that enhance cytokine-induced signaling responses in cells. *DNA Cell Biol.* **16**: 691–701.
- Stamps, A. C., M. A. Elmore, M. E. Hill, K. Kelly, A. A. Makda, and M. J. Finnen. 1997. A human cDNA sequence with homology to non-mammalian lysophosphatidic acid acyltransferases. *Biochem. J.* **326**: 455–461.
- Eberhardt, C., P. W. Gray, and L. W. Tjoelker. 1997. Human lysophosphatidic acid acyltransferase. cDNA cloning, expression, and localization to chromosome 9q34.3. *J. Biol. Chem.* **272**: 20299–20305.
- Agarwal, A. K., E. Arioglu, S. De Almeida, N. Akkoc, S. I. Taylor, A. M. Bowcock, R. I. Barnes, and A. Garg. 2002. AGPAT2 is mutated in congenital generalized lipodystrophy linked to chromosome 9q34. *Nat. Genet.* **31**: 21–23.
- Garg, A. 2000. Lipodystrophies. *Am. J. Med.* **108**: 143–152.
- Magre, J., M. Delepine, and E. Khallouf. 2001. Identification of the gene altered in Berardinelli-Seip congenital lipodystrophy on chromosome 11q13. *Nat. Genet.* **28**: 365–370.
- Kornberg, A., and W. E. Pricer, Jr. 1953. Enzymatic esterification of alpha-glycerophosphate by long chain fatty acids. *J. Biol. Chem.* **204**: 345–357.
- Yamashita, A., N. Kawagishi, T. Miyashita, T. Nagatsuka, T. Sugiura, K. Kume, T. Shimizu, and K. Waku. 2001. ATP-independent fatty acyl-coenzyme A synthesis from phospholipid: coenzyme A-dependent transacylation activity toward lysophosphatidic acid catalyzed by acyl-coenzyme A:lysophosphatidic acid acyltransferase. *J. Biol. Chem.* **276**: 26745–26752.
- Aguado, B., and R. D. Campbell. 1998. Characterization of a human lysophosphatidic acid acyltransferase that is encoded by a gene located in the class III region of the human major histocompatibility complex. *J. Biol. Chem.* **273**: 4096–4105.
- Hollenback, D., and J. A. Glomset. 1998. On the relation between a stearyl-specific transacylase from bovine testis membranes and a copurifying acyltransferase. *Biochemistry*. **37**: 363–376.
- Bhatt, R., B. Gong, F. Hong, S. A. Jenkins, J. P. Klein, A. M. Kumar, and J. Tulinsky. 2003. PCT Int. Appl. WO 03/037346.
- Bhatt, R., B. Gong, F. Hong, S. A. Jenkins, J. P. Klein, C. T. Kohm, and J. Tulinsky. 2004. US Patent Appl. 20040204386.
- Bonham, L., D. W. Leung, T. White, D. Hollenback, P. Klein, J. Tulinsky, M. Coon, P. de Vries, and J. W. Singer. 2003. Lysophosphatidic acid acyltransferase-beta: a novel target for induction of tumour cell apoptosis. *Expert Opin. Ther. Targets*. **7**: 643–661.
- Jain, M. K., and M. H. Gelb. 1991. Phospholipase A2-catalyzed hydrolysis of vesicles: uses of interfacial catalysis in the scooting mode. *Methods Enzymol.* **197**: 112–125.
- Miller, A. D., and G. J. Rosman. 1989. Improved retroviral vectors for gene transfer and expression. *Biotechniques*. **7**: 980–986, 989.
- Sauer, B. 1987. Functional expression of the cre-lox site-specific recombination system in the yeast *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **7**: 2087–2096.
- Miller, A. D., D. G. Miller, J. V. Garcia, and C. M. Lynch. 1993. Use of retroviral vectors for gene transfer and expression. *Methods Enzymol.* **217**: 581–599.
- Hideshima, T., D. Chauhan, T. Hayashi, K. Podar, M. Akiyama, C. Mitsiades, N. M. Itsiades, B. Gong, L. Bonham, P. de Vries, et al. 2003. Antitumor activity of lysophosphatidic acid acyltransferase-beta inhibitors, a novel class of agents, in multiple myeloma. *Cancer Res.* **63**: 8428–8436.
- White, T., S. Bursten, D. Federighi, R. A. Lewis, and E. Nudelman. 1998. High-resolution separation and quantification of neutral lipid and phospholipid species in mammalian cells and sera by multi-one-dimensional thin-layer chromatography. *Anal. Biochem.* **258**: 109–117.
- Ghomashchi, F., B. Z. Yu, O. Berg, M. K. Jain, and M. H. Gelb. 1991. Interfacial catalysis by phospholipase A2: substrate specificity in vesicles. *Biochemistry*. **30**: 7318–7329.
- Coon, M., A. Ball, J. Pound, S. Ap, D. Hollenback, T. White, J. Tulinsky, L. Bonham, D. K. Morrison, R. Finney, et al. 2003. Inhibition of lysophosphatidic acid acyltransferase beta disrupts proliferative and survival signals in normal cells and induces apoptosis of tumor cells. *Mol. Cancer Ther.* **2**: 1067–1078.
- Baburina, I., and S. Jackowski. 1999. Cellular responses to excess phospholipid. *J. Biol. Chem.* **274**: 9400–9408.
- Brindley, D. N., and D. W. Waggoner. 1998. Mammalian lipid phosphate phosphohydrolases. *J. Biol. Chem.* **273**: 24281–24284.
- Bartlett, K., P. Bartlett, N. Bartlett, and H. S. Sherratt. 1985. Kinetics of enzymes requiring long-chain acyl-CoA esters as substrates: effects of substrate binding to albumin. *Biochem. J.* **229**: 559–560.
- Richards, E. W., M. W. Hamm, J. E. Fletcher, and D. A. Otto. 1990. The binding of palmitoyl-CoA to bovine serum albumin. *Biochim. Biophys. Acta.* **1044**: 361–367.
- Boylan, J. G., and J. A. Hamilton. 1992. Interactions of acyl-coenzyme A with phosphatidylcholine bilayers and serum albumin. *Biochemistry*. **31**: 557–567.
- Goetzl, E. J., H. Lee, T. Azuma, T. P. Stossel, C. W. Turck, and J. S. Karliner. 2000. Gelsolin binding and cellular presentation of lysophosphatidic acid. *J. Biol. Chem.* **275**: 14573–14578.
- Das, A. K., and A. K. Hajra. 1992. Critical micellar concentrations of palmitoyl dehydroxyacetone phosphate and 1-palmitoyl-rac-glycerol 3-phosphate. *J. Biol. Chem.* **267**: 9731.
- Powell, G. L., J. R. Grothusen, J. K. Zimmerman, C. A. Evans, and W. W. Fish. 1981. A re-examination of some properties of fatty acyl-CoA micelles. *J. Biol. Chem.* **256**: 12740–12747.
- Smith, R. H., and G. L. Powell. 1986. The critical micelle concentration of some physiologically important fatty acyl-coenzyme A's as a function of chain length. *Arch. Biochem. Biophys.* **244**: 357–360.
- Constantinides, P. P., and J. M. Steim. 1988. Micellization of fatty acyl-CoA mixtures and its relevance to the fatty acyl selectivity of acyltransferases. *Arch. Biochem. Biophys.* **261**: 430–436.
- Rasmussen, J. T., J. Rosendal, and J. Knudsen. 1993. Interaction of acyl-CoA binding protein (ACBP) on processes for which acyl-CoA is a substrate, product or inhibitor. *Biochem. J.* **292**: 907–913.
- Fyrst, H., J. Knudsen, M. A. Schott, B. H. Lubin, and F. A. Kuypers. 1995. Detection of acyl-CoA-binding protein in human red blood cells and investigation of its role in membrane phospholipid renewal. *Biochem. J.* **306**: 793–799.
- Kerkhoff, C., M. Beuck, J. Threige-Rasmussen, F. Spener, J. Knudsen, and G. Schmitz. 1997. Acyl-CoA binding protein (ACBP) regulates

- acyl-CoA:cholesterol acyltransferase (ACAT) in human mononuclear phagocytes. *Biochim. Biophys. Acta.* **1346**: 163–172.
42. Jolly, C. A., D. C. Wilton, and F. Schroeder. 2000. Microsomal fatty acyl-CoA transacylation and hydrolysis: fatty acyl-CoA species dependent modulation by liver fatty acyl-CoA binding proteins. *Biochim. Biophys. Acta.* **1483**: 185–197.
43. Jolly, C. A., E. J. Murphy, and F. Schroeder. 1998. Differential influence of rat liver fatty acid binding protein isoforms on phospholipid fatty acid composition: phosphatidic acid biosynthesis and phospholipid fatty acid remodeling. *Biochim. Biophys. Acta.* **1390**: 258–268.
44. Jolly, C. A., T. Hubbell, W. D. Behnke, and F. Schroeder. 1997. Fatty acid binding protein: stimulation of microsomal phosphatidic acid formation. *Arch. Biochem. Biophys.* **341**: 112–121.
45. Pettitt, T. R., M. McDermott, K. M. Saqib, N. Shimwell, and M. J. Wakelam. 2001. Phospholipase D1b and D2a generate structurally identical phosphatidic acid species in mammalian cells. *Biochem. J.* **360**: 707–715.
46. Raetz, C. R., and W. Dowhan. 1990. Biosynthesis and function of phospholipids in *Escherichia coli*. *J. Biol. Chem.* **265**: 1235–1238.
47. Lu, B., Y. J. Jiang, Y. Zhou, F. Y. Xu, G. M. Hatch, and P. C. Choy. 2005. Cloning and characterization of murine 1-acyl-sn-glycerol 3-phosphate acyltransferases and their regulation by PPARalpha in murine heart. *Biochem. J.* **385**: 469–477.
48. Glomset, J. A. 1996. A branched metabolic pathway in animal cells converts 2-monoacylglycerol into *sn*-1-stearoyl-2-arachidonoyl phosphatidylinositol and other phosphoglycerides. In *Advances in Lipobiology*, Vol. 1. R. W. Gross, editor. Elsevier, St. Louis, MO. 61–100.
49. Agarwal, A. K., and A. Garg. 2003. Congenital generalized lipodystrophy: significance of triglyceride biosynthetic pathways. *Trends Endocrinol. Metab.* **14**: 214–221.
50. Hodgkin, M. N., T. R. Pettitt, A. Martin, R. H. Michell, A. J. Pemberton, and M. J. Wakelam. 1998. Diacylglycerols and phosphatidates: which molecular species are intracellular messengers? *Trends Biochem. Sci.* **23**: 200–204.
51. Pettitt, T. R., A. Martin, T. Horton, C. Liassis, J. M. Lord, and M. J. Wakelam. 1997. Diacylglycerol and phosphatidate generated by phospholipases C and D, respectively, have distinct fatty acid compositions and functions. Phospholipase D-derived diacylglycerol does not activate protein kinase C in porcine aortic endothelial cells. *J. Biol. Chem.* **272**: 17354–17359.