

## Cumulative Subject Index for 1998<sup>1</sup> Volumes 130–135

### A

- Accordion spectroscopy  
diffusion coefficient measurement by 2D NMR, Millet and Pons, **131**, 166  
3D, measurement of <sup>15</sup>N and <sup>13</sup>CO relaxation rates in poorly resolved NMR spectra, Carr, Fearing, and Palmer, **132**, 25
- N*-Acetyl aspartate  
singlet, simultaneous detection in *in vivo* lactate editing at 1.5 T using PRESS excitation, in head and neck tumors, Star-Lack *et al.*, **133**, 243  
and water, non-mono-exponential signal attenuation due to diffusion in brain tissue, Assaf and Cohen, **131**, 69
- Active-coupling-pattern tilting  
determination of homonuclear coupling constants, Koźmiński, **134**, 189
- Adamantane  
high-speed <sup>13</sup>C MAS NMR spectra, Ernst, Verhoeven, and Meier, **130**, 176
- Adiabatic decoupling  
Bloch–Siegert shift-compensated and cyclic sideband-eliminated double-adiabatic homonuclear decoupling for <sup>13</sup>C- and <sup>15</sup>N-double-labeled proteins, Zhang and Gorenstein, **132**, 81  
efficient broadband, in single transient, calibration of STUD+ parameters for, Bendall and Skinner, **134**, 331  
vector model, Skinner and Bendall, **134**, 315
- Adiabatic mixing  
in liquid state, Kupče *et al.*, **135**, 361
- Adiabatic pulses  
effect on IS, I<sub>2</sub>S, and I<sub>3</sub>S spin systems, analytical description, Zwahlen, Vincent, and Kay, **130**, 169  
fast broadband inversion by, Hwang, Zijl, and Garwood, **133**, 200  
multiple spin-echo, Zweckstetter and Holak, **133**, 134  
off-resonance rotating frame relaxation experiment for protein dynamics, Mulder *et al.*, **131**, 351  
sech/tanh, adiabatic behavior, Rosenfeld and Zur, **132**, 102
- Aerogels  
silica, pore-structure determination by <sup>129</sup>Xe NMR and imaging, Gregory, Gerald, and Botto, **131**, 327
- Affinity NMR  
isotope-filtered, Gonnella *et al.*, **131**, 336
- Al<sup>+3</sup>  
species in aqueous solution, NMR shieldings, effects of hydrolysis and oligomerization, Tossell, **135**, 203
- <sup>27</sup>Al  
in Al<sub>2</sub>O<sub>3</sub> single crystal, five-quantum coherence of  $I = \frac{5}{2}$  nuclei, Kaikkonen, Ylinen, and Punkkinen, **133**, 123  
<sup>27</sup>Al → <sup>31</sup>P INEPT experiments on (CH<sub>3</sub>)<sub>3</sub>P–AlCl<sub>3</sub> complex in zeolite NaX, Kao and Grey, **133**, 313  
chemical-shielding anisotropy, definitive example, Schurko, Wasylshen, and Phillips, **133**, 388  
spin–lattice relaxation in  $\gamma$ -irradiated ruby crystal, separation of impurity and lattice contributions and determination of charge-exchanged chromium ions, Chandoul *et al.*, **135**, 113
- Alanine  
<sup>13</sup>C, intrinsic asymmetry in multidimensional solid-state NMR correlation spectra, Caldarelli and Emsley, **129**, 233  
<sup>13</sup>C–<sup>15</sup>N multiply labeled, counting spins with SEDOR, Cull *et al.*, **133**, 352
- L-Alanine  
<sup>13</sup>C–<sup>14</sup>N REAPDOR NMR experiment optimization, Ba *et al.*, **133**, 104  
multiple-quantum MAS NMR, resolution enhancement, Schnell *et al.*, **133**, 61
- AlCl<sub>3</sub> · OPCI<sub>3</sub>  
solid-state <sup>27</sup>Al NMR: chemical-shielding anisotropy, Schurko, Wasylshen, and Phillips, **133**, 388
- Algorithms  
FI-HTLS: frequency-selective MRS data quantification with frequency prior knowledge, Dologlou, Van Huffel, and Van Ormondt, **130**, 238  
generalized, for automated analysis of complex anisotropic NMR spectra, Castiglione *et al.*, **132**, 1
- Alignment  
magnetic, paramagnetic-assisted, determination of DNA internuclear angles, Beger *et al.*, **135**, 256
- Alignment echo  
spin- $\frac{3}{2}$  <sup>9</sup>Be nuclei: detection of ultraslow motion, Tang and Wu, **133**, 155
- Alignment tensor  
asymmetric, of oriented macromolecules, magnitude determination in absence of structural information, Clore, Gronenborn, and Bax, **133**, 216
- Aliphatic carbons  
gamma, and backbone amides, resonances in deuterated proteins, unambiguous correlations, McCallum, Hitchens, and Rule, **134**, 350
- Al<sub>2</sub>O<sub>3</sub>  
single crystal, <sup>27</sup>Al: five-quantum coherence of  $I = \frac{5}{2}$  nuclei, Kaikkonen, Ylinen, and Punkkinen, **133**, 123
- Amide  
backbone, and aliphatic gamma resonances, in deuterated proteins, unambiguous correlations, McCallum, Hitchens, and Rule, **134**, 350  
deuterium isotope effects on C <sup>$\beta$</sup>  chemical shifts in <sup>13</sup>C, <sup>15</sup>N-labeled proteins, multidimensional editing for, Meissner and Sørensen, **135**, 547  
protons, cross-relaxation between, detection in protein NMR, Mutzenhardt and Bodenhausen, **132**, 159
- 4-Amino-2,2,6,6-tetramethyl-1-piperidinyloxy (4-Amino-TEMPO)  
 $\pi$  and  $\sigma$  transitions, DNP study at 1.53 mT, Salman *et al.*, **134**, 1
- Ammonia  
identification as metabolite in kidney/urine proton chemical exchange, Guivel-Scharen *et al.*, **133**, 36
- Ampicillin  
<sup>13</sup>C and <sup>15</sup>N chemical-shift anisotropy by 2D-PASS and CP/MAS NMR, Antzutkin, Lee, and Levitt, **135**, 144
- Amplitude images  
at low field, use of phase reference for field mapping with, Tomasi *et al.*, **131**, 310
- Amplitude modulation  
rotor synchronized cross-polarization with, application in <sup>13</sup>C–<sup>1</sup>H spin pair under fast MAS, Raya and Hirschinger, **133**, 341
- Analog filtering  
large solvent signals for improved dynamic range in high-resolution NMR, Redfield and Kunz, **130**, 111

<sup>1</sup> Boldface numbers indicate appropriate volume; lightface numbers indicate pagination.

- Analytical transfer functions  
polarization and coherence, for three coupled spins  $\frac{1}{2}$  under planar mixing conditions, Schedletzky, Luy, and Glaser, **130**, 27
- Anderson–Weiss model  
1–370 GHz EPR linewidths for  $K_3CrO_8$ , Cage *et al.*, **135**, 178
- Anisometric NMR spectra  
complex, generalized algorithm for automated analysis, Castiglione *et al.*, **132**, 1
- Anisotropy  
in frozen nickel ferrite ionic magnetic fluid, MR analysis, Saenger *et al.*, **134**, 180  
in  $^{31}P$ ,  $^{63,65}Cu$  indirect spin–spin coupling and  $^{31}P$  nuclear shielding tensors of linear copper(I) phosphines, Kroeker *et al.*, **135**, 208
- Antarctica  
Earth's field NMR: pulsed gradient spin echo NMR study of restricted diffusion in sea ice, Callaghan *et al.*, **133**, 148
- Antibiotics  
 $^{13}C$  and  $^{15}N$  chemical-shift anisotropy by 2D-PASS and CP/MAS NMR, Antzutkin, Lee, and Levitt, **135**, 144
- Aqueous samples  
rectangular loop–gap resonator for EPR studies, Piasecki, Froncisz, and Hubbell, **134**, 36
- Aqueous solution  
 $Be^{+2}$  and  $Al^{+3}$  species, NMR shieldings, effects of hydrolysis and oligomerization, Tossell, **135**, 203  
 $^{13}C$ -labeled proteins, H(C)CH-COSY and (H)CCH-COSY experiments, Gehring and Ekiel, **135**, 185  
measurement of diffusion of proteins, practical aspects, Tillett, Lian, and Norwood, **133**, 379
- Aqueous suspensions  
char, molecular diffusion and DNP enhancement, Odintsov *et al.*, **135**, 435
- Aromatic side chains  
assignment in  $^{13}C$ -labeled proteins, 2D NMR experiments for, Prompers *et al.*, **130**, 68
- Arthritis  
collagen-induced, in DA rat, Sandimmune Neoral effects by high-resolution 3D MRI and histology, Beckmann *et al.*, **131**, 8
- Artifacts  
convection, in gradient-enhanced NMR, compensation, Jerschow and Müller, **132**, 13  
in diffusion studies using RF field gradients, remedies, Humbert *et al.*, **134**, 245  
induced by weakly ferromagnetic metals: field mapping with amplitude images at low field, Tomasi *et al.*, **131**, 310  
in multishot echo-planar imaging, image-based reduction, Hennel, **134**, 206  
rapid-pulsing, double-quantum-filtered homonuclear spectroscopy free of: 2D-INADEQUATE experiment revisited, Bourdonneau and Ancian, **132**, 316  
solvent magnetization, in high-field NMR studies of macromolecular hydration, Sobol *et al.*, **130**, 262
- Artificial neural networks  
in automated quantification of human brain metabolites from *in vivo* single-voxel  $^1H$  NMR spectra, Kaartinen *et al.*, **134**, 176
- ASICS  
software for single-crystal NMR, Vosegaard *et al.*, **135**, 126
- Asparagine  
and Gln, triple-resonance experiment selective for, multiplicity-selective coherence transfer steps for design of, Schmieider *et al.*, **131**, 199
- Asymmetry  
intrinsic, in multidimensional solid-state NMR correlation spectra, Caldarelli and Emsley, **129**, 233
- Atomic motion  
ultraslow, detection using alignment echo of spin- $\frac{3}{2}$   $^9Be$  nuclei, Tang and Wu, **133**, 155
- Atropisomerism  
effect on  $^{57}Fe$  shieldings in superstructured hemoprotein models, Gerthanssis *et al.*, **131**, 163
- Attenuation  
non-mono-exponential, water and *N*-acetyl aspartate signals due to diffusion in brain tissue, Assaf and Cohen, **131**, 69
- Automated peak picking  
and peak integration, in macromolecular NMR spectra, using AUTOPSY, Koradi *et al.*, **135**, 288
- Automated processing  
complex anisometric NMR spectra, generalized algorithm for, Castiglione *et al.*, **132**, 1  
2D correlation spectra, Sengstschmid, Sterk, and Freeman, **131**, 315
- Automated shimming  
with normal spectrometer hardware: 3D profile edge shimming, Chilvers and Morris, **133**, 210
- AUTOPSY  
automated peak picking and peak integration in macromolecular NMR spectra, Koradi *et al.*, **135**, 288
- Average Liouvillian theory  
homogeneous master equation for heteronuclear two-spin system on basis of Cartesian product operators, Allard, Helgstrand, and Hård, **134**, 7
- Azoles  
nitrogen shielding, *ab initio* calculation, Witanowski *et al.*, **131**, 54
- B
- Bacteriochlorophyll *a*  
 $^{15}N$ -labeled, low-frequency nuclear coherence detection with simple two-pulse detection scheme in pulsed EPR, Dzuba, Borovykh, and Hoff, **133**, 286
- Bandwidth extension  
in noise spectroscopy, Yang and Zax, **135**, 267
- Bayesian time-domain parameter estimation  
Metropolis Monte Carlo implementation: coupling constant estimation from antiphase multiplets, Andreac and Prestegard, **130**, 217
- $Be^{+2}$   
species in aqueous solution, NMR shieldings, effects of hydrolysis and oligomerization, Tossell, **135**, 203
- $^9Be$   
spin- $\frac{3}{2}$  nuclei, alignment echo: detection of ultraslow motion, Tang and Wu, **133**, 155
- Benzene  
dynamic isotope effect and deuterium quadrupole coupling constant, Hardy *et al.*, **134**, 300  
molecules oriented in liquid crystals, natural abundance  $^2H$  NMR spectra, Khetrapal *et al.*, **135**, 265  
polycrystalline, vacancy diffusion in, analysis with 2D deuteron exchange NMR and stimulated echoes, Isfort, Geil, and Fujara, **130**, 45
- Bicelles  
phospholipid, dilute solutions for structural NMR work, Struppe and Vold, **135**, 541
- Bilayers (*see* Lipid bilayers)
- Binary spin–bath model  
contrast prediction using: magnetization transfer imaging of rat brain under non-steady-state conditions, Quesson *et al.*, **130**, 321
- Biomembranes  
uniformly aligned, application of magic angle-oriented sample spinning, Glaubitz and Watts, **130**, 305
- Biopolymers  
magnetization transfer in, role of specific side groups and pH, Gochberg *et al.*, **131**, 191
- BIRD  
TIG-BIRD pulse sequence element for simultaneous and independent rotations for I,  $IS^\alpha$ , and  $IS^\beta$  spin systems, Briand and Sørensen, **135**, 44

- Birdcage coil  
tuned by RF shielding, for application at 9.4 T, Dardzinski *et al.*, **131**, 32
- Blind source separation  
model-free analysis of mixtures by NMR, Nuzillard, Bourg, and Nuzillard, **133**, 358
- Bloch–Siegert shift  
compensation, in double-adiabatic homonuclear decoupling for  $^{13}\text{C}$ - and  $^{15}\text{N}$ -double-labeled proteins, Zhang and Gorenstein, **132**, 81
- BOLD response  
fMRI study of visuomotor system activation during visually guided movements, Ellermann *et al.*, **131**, 272
- Bone  
trabecular,  $R_2'$  measurement in presence of multiple spectral components using reference spectrum deconvolution, Wehrli *et al.*, **131**, 61
- Bone marrow  
 $R_2'$  measurement in presence of multiple spectral components using reference spectrum deconvolution, Wehrli *et al.*, **131**, 61
- Bovine pancreatic trypsin inhibitor  
application of model-free analysis of stretched dispersions, Halle, Jóhannesson, and Venu, **135**, 1
- Bovine serum albumin  
cross-linked, magnetic coupling between lactate and water protons, Swanson, **135**, 248
- Brain  
activation of visuomotor systems during visually guided movements, functional MRI, Ellermann *et al.*, **131**, 272  
blood volume, sequential dynamic susceptibility contrast experiments,  $T_1$  effects, Levin *et al.*, **130**, 292  
excised tissue, non-mono-exponential attenuation of water and *N*-acetyl aspartate signals due to diffusion, Assaf and Cohen, **131**, 69  
frequency-selective MRS data quantification with frequency prior knowledge, Dologlou, Van Huffel, and Van Ormondt, **130**, 238  
*in vivo*  $^1\text{H}$  NMR spectra, resolution improvements with increased magnetic field strength, Gruetter *et al.*, **135**, 260  
*in vivo* multiple spin echoes, Bifone, Payne, and Leach, **135**, 30  
Li visibility *in vivo* by  $^7\text{Li}$  NMR imaging, Komoroski, Pearce, and Newton, **133**, 98  
magnetization transfer imaging under non-steady-state conditions, Quesson *et al.*, **130**, 321  
metabolites, automated quantification by artificial neural network analysis from *in vivo* single-voxel  $^1\text{H}$  NMR spectra, Kaartinen *et al.*, **134**, 176  
multivariate image analysis with direct exponential curve resolution algorithm (DECRA), Antalek, Hornak, and Windig, **132**, 307  
 $^{31}\text{P}$  MRS spectra, precision of parameters in measuring, Golding and Golding, **135**, 380  
taurine detection *in vivo* and *in vitro*, spectral editing technique for, Hardy and Norwood, **133**, 70
- Breast ductal carcinomas  
evaluation with high-resolution MAS IHMRS, Cheng *et al.*, **135**, 194
- Broadband adiabatic decoupling  
efficient, in single transient, calibration of STUD+ parameters for, Bendall and Skinner, **134**, 331
- Broadband decoupling  
liquid crystals, sequence for, Yu and Fung, **130**, 317
- Broadband inversion  
fast, by adiabatic pulses, Hwang, Zijl, and Garwood, **133**, 200
- Brownian motions  
rotational and translational, couplings between, effect on NMR spin relaxation: Green function of rigid body diffusion, Beloborodov, Orekhov, and Arseniev, **132**, 328
- Bulk magnetic moment dynamics  
in MRI of lung with hyperpolarized noble gases, Möller *et al.*, **135**, 133
- Bulk magnetic susceptibility  
effect on solid state NMR spectra of paramagnetic compounds, Kubo, Spaniol, and Terao, **133**, 330
- C
- $^{13}\text{C}$   
adamantane, high-speed MAS NMR spectra, Ernst, Verhoeven, and Meier, **130**, 176  
alanine, intrinsic asymmetry in multidimensional solid-state NMR correlation spectra, Caldarelli and Emsley, **129**, 233  
carbon–phosphorus *J* coupling constants in RNA, by spin-echo difference constant-time HCCH–COSY, Hoogstraten and Pardi, **133**, 236  
chemical-shift anisotropy of ampicillin and penicillin-V by 2D-PASS, Antzutkin, Lee, and Levitt, **135**, 144  
chemical-shift imaging  
alternative scheme via inverse detection of protons through MILS technique, Lee *et al.*, **130**, 102  
in rotating solids by RF field gradients, Malveau *et al.*, **134**, 171  
 $^{13}\text{CH}_3\text{I}$  high-resolution NMR, control of spinning sidebands, Borer and Maple, **131**, 177  
 $^{13}\text{C}$ – $^{14}\text{N}$  REAPDOR NMR, optimization, Ba *et al.*, **133**, 104  
complete dipolar decoupling, and use in 2D double-quantum solid-state NMR for determining polymer conformations, Schmidt-Rohr, **131**, 209  
 $\alpha$ – $\alpha'$ – $^{13}\text{C}_2$ -phthalic acid, polarization transfer under RF-driven and slow-magic-angle-sample-spinning conditions, Robyr and Gan, **131**, 254  
CPMAS linewidths, in solids for rigid strongly coupled carbons under CW proton decoupling, VanderHart and Campbell, **134**, 88  
deuterated glutathione *S*-transferase, correlation of backbone amide and aliphatic gamma resonances, McCallum, Hitchens, and Rule, **134**, 350  
dimethylsulfone, direct determination of motional correlational times by 1D MAS and 2D exchange NMR, Favre, Schaefer, and Chmelka, **134**, 261  
dimyristoylphosphatidylcholine bilayers, magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305  
DNA–europium complex, internuclear angles of DNA, determination using paramagnetic-assisted magnetic alignment, Beger *et al.*, **135**, 256  
fast broadband inversion by adiabatic pulses, Hwang, Zijl, and Garwood, **133**, 200  
ferrocene,  $^{13}\text{C}$ – $^1\text{H}$  spin pair under fast MAS, application of rotor-synchronized amplitude-modulated cross polarization, Raya and Hirschinger, **133**, 341  
glycine, six-pulse PASS MAS-NMR technique solved by phasor method, Effey, Butcher, and Cappelletti, **132**, 266  
heteronuclear X-filter  $^1\text{H}$  PFG double-quantum experiment for proton resonance assignment of ligand bound to protein, Dalvit, Ramage, and Hommel, **131**, 148  
HSQC-based methyl group selection via gradients in multidimensional NMR of proteins, Diercks, Schwaiger, and Kessler, **130**, 335  
improved resolution from double constant-time evolution of 3D and 4D triple-resonance experiments, McCoy, **130**, 341  
*in vivo* NMR  
on clinical 1.5 T MR system, heteronuclear cross polarization for enhanced sensitivity, van den Bergh, van den Boogert, and Heerschap, **135**, 93  
2D localized, using modified INEPT and DEPT, Watanabe *et al.*, **134**, 214  
low-temperature NMR: high-precision  $^{13}\text{C}$  shift thermometer for 100–300 K, Quast *et al.*, **134**, 236  
methyl salicylate, reduced relaxation time: sensitivity enhancement in INADEQUATE, Mattiello and Freeman, **135**, 514  
and  $^{15}\text{N}$   
alanine and urea multiply labeled with, counting spins with SEDOR, Cull *et al.*, **133**, 352  
ubiquitin labeled with, measurement of dipolar couplings for methylene and methyl sites and use in structure determination, Ottiger *et al.*, **134**, 365

- and  $^{15}\text{N}$ , proteins labeled with  
 amide deuterium isotope effects on  $C^\beta$  chemical shifts, multidimensional editing for, Meissner and Sørensen, **135**, 547  
 double-labeled, Bloch–Siegert shift-compensated and cyclic sideband-eliminated double-adiabatic homonuclear decoupling for, Zhang and Gorenstein, **132**, 81  
 in  $\text{H}_2\text{O}$  solution, H(C)CH-COSY and (H)CCH-COSY experiments, Gehring and Ekiel, **135**, 185  
 spin relaxation, measurement of cross correlation between dipolar coupling and chemical-shift anisotropy, Ghose, Huang, and Prestegard, **135**, 487  
 NMR imaging, cyclic  $J$  cross-polarization-based, analysis of carbohydrate metabolism and transport in castor bean seedlings by, Heidenreich *et al.*, **132**, 109  
 NMR with RF field gradients, self-diffusion measurements by, Malveau *et al.*, **130**, 131  
 OCS, rovibrationally averaged nuclear shielding constants, Jackowski, **135**, 444  
 1D spectroscopy, pulse sequence DEPTQ for, Burger and Bigler, **135**, 529  
 4-*n*-pentyl-4'-cyanobiphenyl liquid crystals, application of broadband decoupling sequence SPARC-16, Yu and Fung, **130**, 317  
 polystyrene labeled with, 2D NMR experiments for structural analysis, Saito and Rinaldi, **132**, 41  
 proteins labeled with, 2D NMR for assignment of aromatic side chains, Prompers *et al.*, **130**, 68  
 proteins uniformly labeled with,  $^{13}\text{C}$ -NOESY-HSQC with split carbon evolution for increased resolution, Baur, Gemmecker, and Kessler, **132**, 191  
 protons attached to, scalar spin-spin coupling constants between, methylene specific experiments, Carlomagno *et al.*, **135**, 216  
 purine nucleotides labeled with, DNA decamer duplex with,  $^{13}\text{C}$  relaxation, Kojima *et al.*, **135**, 310  
 raffinose, heteronuclear long-range couplings by gradient-enhanced 2D multiple-bond correlation, Sheng and van Halbeek, **130**, 296  
 rapid-pulsing artifact-free double-quantum-filtered homonuclear spectroscopy: 2D-INADEQUATE experiment revisited, Bourdonneau and Ancian, **132**, 316  
 relaxation data, analysis of protein side-chain dynamics, Daragan and Mayo, **130**, 329  
 RNA uniformly labeled with, selectively  $^2\text{H}$  labeled, proton and carbon assignment, 3D C(CC)H TOCSY for, Dayie, Tolbert, and Williamson, **130**, 97  
 rotational motion of solute molecule in highly viscous liquid, Lienin, Brüschweiler, and Ernst, **131**, 184  
 self-diffusion measurements by  $^{13}\text{C}$  NMR using RF field gradients, Malveau *et al.*, **130**, 131  
 sensitivity- and gradient-enhanced hetero ( $\omega_1$ ) half-filtered TOCSY for long-range heteronuclear coupling constants, Uhrin *et al.*, **130**, 155  
 solid sample enriched with, phase-sensitive detection of 2D homonuclear correlation spectra in MAS NMR, Boender and Vega, **133**, 281  
 solute-solvent interactions of fullerenes  $\text{C}_{60}$  and  $\text{C}_{70}$  in solution: 1D steady-state intermolecular  $^{13}\text{C}$ ,  $^1\text{H}$  Overhauser effect, Troganis, Gerothanassis, and Papademitriou, **130**, 149  
 spin-lattice relaxation in 99%  $^{13}\text{C}$ -enriched diamonds, Shabanova, Schaumburg, and Sellschop, **130**, 8  
 STUD+ parameter calibration for efficient broadband adiabatic decoupling in single transient, Bendall and Skinner, **134**, 331  
 triple-resonance HCN experiments, optimization for larger RNA oligonucleotides, Sklenář *et al.*, **130**, 119  
 2D CP/MAS  $^{13}\text{C}$  isotropic chemical shift correlation established by  $^1\text{H}$  spin diffusion, Wilhelm *et al.*, **134**, 255  
 ubiquitin, dipole-CSA cross-correlated relaxation, quantitative analysis by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346  
 $^{13}\text{C}_{60}$   
 triplet state, pulsed ENDOR studies at 95 GHz, van den Berg *et al.*, **131**, 39  
 $\text{C}_{60}$   
 solute-solvent interactions in solution: 1D steady-state intermolecular  $^{13}\text{C}$ ,  $^1\text{H}$  Overhauser effect, Troganis, Gerothanassis, and Papademitriou, **130**, 149  
 $\text{C}_{70}$   
 solute-solvent interactions in solution: 1D steady-state intermolecular  $^{13}\text{C}$ ,  $^1\text{H}$  Overhauser effect, Troganis, Gerothanassis, and Papademitriou, **130**, 149  
 $^{13}\text{C}^\alpha$ - $^1\text{H}^\alpha$  dipolar tensor  
 and  $^{15}\text{N}$  chemical shift tensor, correlation in solid-state MAS NMR, peptide torsion angle determination by, Hong *et al.*, **135**, 169  
 Carbohydrates  
 metabolism and transport in castor bean seedlings, cyclic  $J$  cross-polarization imaging and spectroscopy, Heidenreich *et al.*, **132**, 109  
 $\beta$ -Carboline  
 derivatives, interaction with human serum albumin,  $^1\text{H}$  NMR studies, Veglia *et al.*, **130**, 281  
 Carbon  
 assignment in uniformly  $^{13}\text{C}$ - and selectively  $^2\text{H}$ -labeled RNA, 3D C(CC)H TOCSY for, Dayie, Tolbert, and Williamson, **130**, 97  
 gamma, aliphatic, and backbone amides, resonances in deuterated proteins, unambiguous correlations, McCallum, Hitchens, and Rule, **134**, 350  
 rigid strongly coupled, under CW proton decoupling,  $^{13}\text{C}$  CPMAS line-widths in solids for, VanderHart and Campbell, **134**, 88  
 Carbon-based paramagnetic materials  
 for use in EPR oximetry, separation and enrichment of active component, Liu *et al.*, **133**, 291  
 Carbon disulfide  
 self-diffusion measurements by  $^{13}\text{C}$  NMR using RF field gradients, Malveau *et al.*, **130**, 131  
 Carbon tetrachloride  
 self-diffusion measurements by  $^{13}\text{C}$  NMR using RF field gradients, Malveau *et al.*, **130**, 131  
 Carboplatin  
 local disposition kinetics, monitoring *in vivo* after sc injection in rat using  $^{195}\text{Pt}$  NMR, Becker *et al.*, **133**, 115  
 3-Carboxy proxyl  
 Gaussian line broadening of EPR spectrum, contributions from magnetic-field modulation and unresolved proton hyperfine structure, Bales, Peric, and Lamy-Freund, **132**, 279  
 Cardiac spectroscopy  
 localized  $^{31}\text{P}$  NMR: SLOOP, Löffler *et al.*, **134**, 287  
 Cartesian product operators  
 homogeneous master equation for heteronuclear two-spin system on basis of, Allard, Helgstrand, and Härd, **134**, 7  
 CARVE  
 excitation of complicated shapes in three dimensions, Serša and Macura, **135**, 466  
 Catalase  
 NMR spectra acquisition with PASE: paramagnetic signals enhancement, Bondon and Mouro, **134**, 154  
 Catalysis  
 analysis using parahydrogen: evolution of zero-quantum coherence in AA'X spin systems, Natterer *et al.*, **133**, 92  
 Cation binding  
 induced changes in  $^{15}\text{N}$  chemical-shift anisotropy in membrane-bound polypeptide, Tian and Cross, **135**, 535  
 $\text{C}^\beta$   
 chemical shifts, in  $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, amide deuterium isotope effects, multidimensional editing for, Meissner and Sørensen, **135**, 547

- Cesium pentadecafluorooctanoate  
 $^{19}\text{F}$ - $^1\text{H}$  HOESY study of water-surfactant contact, Raulet *et al.*, **133**, 324
- Char  
 aqueous suspensions, molecular diffusion and DNP enhancement, Odintsov *et al.*, **135**, 435
- Charge exchange  
 chromium ions, and impurity nuclear spin-lattice relaxation, in  $\gamma$ -irradiated ruby crystal, Chandoul *et al.*, **135**, 113
- Chemical exchange  
 exchange-edited gradient HCP-HSQC in sensitivity-enhanced detection of fast exchanging protons, Zangger and Armitage, **135**, 70  
 peaks, multiple-pulse mixing sequences selectively enhancing, Hwang and Shaka, **135**, 280  
 proton, between metabolites and water, detection in biological tissues, Guivel-Scharen *et al.*, **133**, 36  
 rate constants, measurement with solvent protons using radiation damping, Chen and Mao, **131**, 358
- Chemical-shielding anisotropy  
 $^{27}\text{Al}$ , definitive example, Schurko, Wasylishen, and Phillips, **133**, 388  
 dipole-CSA cross-correlated relaxation, quantitative analysis by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346  
 $^1\text{H}$ , from polycrystalline powders, using MSHOT-3 based CRAMPS, Hohwy *et al.*, **133**, 374
- Chemical shift  
 analysis of conformational effect of nitroxide spin labels on Ala-rich helical peptides, Bolin *et al.*, **131**, 248  
 backbone amide proton and nitrogen in deuterated proteins, correlation with aliphatic gamma resonances, McCallum, Hitchens, and Rule, **134**, 350  
 $^{13}\text{C}$   
 in 3D DOSY-HMQC experiment for high-resolution analysis of complex mixtures, Barjat, Morris, and Swanson, **131**, 131  
 high-precision  $^{13}\text{C}$  shift thermometer for temperature range 100–300 K, Quast *et al.*, **134**, 236  
 $\text{C}^{\beta\text{T}}$ , in  $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, amide deuterium isotope effects, multi-dimensional editing for, Meissner and Sørensen, **135**, 547  
 $^{57}\text{Fe}$ , in superstructured hemoprotein models, effects of atropisomerism and porphyrin deformation, Gerothanassis *et al.*, **131**, 163  
 hyperfine, in solution, high magnetic field consequences, Bertini, Felli, and Luchinat, **134**, 360  
 isotropic, interactions, distributions in 2D multiple-quantum MAS NMR spectra following, Bodart, **133**, 207  
 in NMR temperature measurements using paramagnetic lanthanide complex, Zuo *et al.*, **133**, 53  
 proton  
 in 3D DOSY-HMQC experiment for high-resolution analysis of complex mixtures, Barjat, Morris, and Swanson, **131**, 131  
 identification in systems with rapid chemical exchange using water stripe information in TOCSY, Nakashima, McClung, and Kotovych, **133**, 222
- Chemical-shift anisotropy  
 $^{13}\text{C}$ , ampicillin and penicillin-V, 2D-PASS and CP/MAS NMR, Antzutkin, Lee, and Levitt, **135**, 144  
 -dipolar coupling cross-correlation rates  
 estimation from laboratory-frame relaxation experiments, Ghose and Prestegard, **134**, 308  
 in spin relaxation of  $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, Ghose, Huang, and Prestegard, **135**, 487  
 interference, differential line broadening in presence of, Elbayed *et al.*, **130**, 209  
 $^{15}\text{N}$   
 ampicillin and penicillin-V, 2D-PASS and CP/MAS NMR, Antzutkin, Lee, and Levitt, **135**, 144  
 in membrane-bound polypeptide, cation binding-induced changes, Tian and Cross, **135**, 535  
 purine, in DNA dynamics study via  $^{13}\text{C}$  relaxation, Kojima *et al.*, **135**, 310
- Chemical-shift correlation  
 isotropic, 2D CP/MAS  $^{13}\text{C}$ , established by  $^1\text{H}$  spin diffusion, Wilhelm *et al.*, **134**, 255
- Chemical-shift imaging  
 $^{13}\text{C}$ , alternative scheme via inverse detection of protons through MILS technique, Lee *et al.*, **130**, 102  
 in rotating solids, by RF field gradients, Malveau *et al.*, **134**, 171
- Chemical-shift tensor  
 $^{15}\text{N}$ , and  $^{13}\text{C}^{\alpha-1}\text{H}^{\alpha}$  dipolar tensor, correlation in solid-state MAS NMR, peptide torsion angle determination by, Hong *et al.*, **135**, 169
- Chloroform  
 molecules oriented in liquid crystals, natural abundance  $^2\text{H}$  NMR spectra, Khetrapal *et al.*, **135**, 265
- Cholestane spin labels  
 shape of Hahn echo responses, influence of high orientational order, van der Struijf *et al.*, **130**, 253
- Choline  
 singlet, simultaneous detection in *in vivo* lactate editing at 1.5 T using PRESS excitation, in head and neck tumors, Star-Lack *et al.*, **133**, 243
- $(\text{CH}_3)_3\text{P}-\text{AlCl}_3$  complex  
 in zeolite NaX,  $^{27}\text{Al} \rightarrow ^{31}\text{P}$  INEPT experiments, Kao and Grey, **133**, 313
- Chromium ions  
 charge exchange, and impurity nuclear spin-lattice relaxation, in  $\gamma$ -irradiated ruby crystal, Chandoul *et al.*, **135**, 113  
 spin- $\frac{3}{2}$ , in ruby, looping transition, simulation methods for, Gaffney and Silverstone, **134**, 57
- Circuits  
 double- and triple-resonance, for high-frequency probes, Zhang *et al.*, **132**, 167
- $^{13}\text{C}$  isotope editing  
 affinity NMR, Gonnella *et al.*, **131**, 336
- $^{35}\text{Cl}$   
 $\text{NaClO}_3$ , single crystal 2D NQR spectroscopy, off-resonance effects, Ajithkumar *et al.*, **135**, 165
- $^{13}\text{C}$ -labeling  
 3D NMR pulse sequences for characterization of polymer chain end structures, Saito and Rinaldi, **130**, 135
- CLEANEX-PM  
 attenuation of cross-relaxation peaks, Hwang and Shaka, **135**, 280
- $^{13}\text{CO}$   
 relaxation rate in poorly resolved NMR spectra, 3D accordion spectroscopy Carr, Fearing, and Palmer, **132**, 25
- $^{59}\text{Co}$   
 $\text{HfFeCo}_3(\text{CO})_{11}\text{PPh}_2\text{H}$ ,  $^{31}\text{P}$  *J*-coupled to, differential line broadening in presence of quadrupolar-CSA interference, Elbayed *et al.*, **130**, 209
- Coals  
 for use in EPR oximetry, separation and enrichment of active component, Liu *et al.*, **133**, 291
- Coaxial line  
 based truncated transmission line probe, FT-EPR with, Rubinson *et al.*, **132**, 255
- Coherence  
 five-quantum,  $I = \frac{5}{2}$  nuclei:  $^{27}\text{Al}$  in  $\text{Al}_2\text{O}_3$  single crystal, Kaikkonen, Ylinen, and Punkkinen, **133**, 123  
 intermolecular zero-quantum, of multi-component spin systems in solution NMR, Ahn, Lisitza, and Warren, **133**, 266  
 nuclear, low-frequency, simple two-pulse detection scheme in pulsed EPR for, Dzuba, Borovykh, and Hoff, **133**, 286  
 zero-quantum, evolution in AA'X spin systems: analysis of catalytic processes using parahydrogen, Natterer *et al.*, **133**, 92
- Coherence selection  
 pulsed-field gradient, in 2D MAS spectroscopy of dipolar solids, Fritzhanns *et al.*, **134**, 355

- RF-gradient, in selective 1D and nonselective 2D experiments, Heikkinen, Rahkamaa, and Kilpeläinen, **133**, 183
- Coherence transfer
- homogeneous master equation for heteronuclear two-spin system on basis of Cartesian product operators, Allard, Helgstrand, and Härd, **134**, 7
  - multiplicity-selective transfer steps for design of amino acid-selective triple-resonance experiments, Schmieder *et al.*, **131**, 199
  - pathways, selection by phase cycling and pulsed-field gradients in NMR, simulation, Jerschow and Müller, **134**, 17
- Coherence transfer functions
- evolution of zero-quantum coherence in AA'X spin systems: analysis of catalytic processes using parahydrogen, Natterer *et al.*, **133**, 92
  - three coupled spins  $\frac{1}{2}$  under planar mixing conditions, Schedletzky, Luy, and Glaser, **130**, 27
- Collagen
- induced arthritis in DA rat, Sandimmune Neoral effects by high-resolution 3D MRI and histology, Beckmann *et al.*, **131**, 8
- Colloidal suspensions
- experimental determination of four-time stimulated echoes, Hinze *et al.*, **131**, 218
- Combined rotation and multiple-pulse spectroscopy (CRAMPS)
- MSHOT-3 based,  $^1\text{H}$  chemical-shielding anisotropies from polycrystalline powders, Hohwy *et al.*, **133**, 374
- Common gateway interface program
- for World Wide Web NMR spectrometer, Buszko, Buszko, and Wang, **131**, 362
- Complex motion
- associated  $^1\text{H}$  and  $^2\text{H}$  NMR relaxation in hydrogen-bonded solids: classical jumps over barrier and incoherent tunneling, Reynhardt and Latanowicz, **130**, 195
- Component-resolved spectroscopy (CORE)
- component separation in NMR imaging and multidimensional spectroscopy based on prior knowledge, Stilbs, **135**, 236
- Component separation
- in NMR imaging and multidimensional spectroscopy, through global least-squares analysis based on prior knowledge, Stilbs, **135**, 236
- Composite pulses
- excitation schemes, for MQMAS NMR of half-integer quadrupolar spins, Marinelli, Medek, and Frydman, **132**, 88
  - $90^\circ$ : distortionless NMR spectra and suppression without destruction of water magnetization, Sodano, Landon, and Ptak, **133**, 194
- Computers
- high-speed personal computer, real-time image reconstruction and display system for MRI, Haishi and Kose, **134**, 138
  - improved hardware and software for single-crystal NMR, Vosegaard *et al.*, **135**, 126
- Computer simulation (*see* Simulation)
- Conformation
- Ala-rich helical peptides, effect of nitroxide spin labels, NMR analysis, Bolin *et al.*, **131**, 248
  - information from water stripe in TOCSY experiments on systems with exchangeable protons, Nakashima, McClung, and Kotovych, **133**, 222
  - polymers, 2D double-quantum solid-state NMR for determining, complete dipolar decoupling of  $^{13}\text{C}$  for, Schmidt-Rohr, **131**, 209
- Constant-time evolution
- double constant-time, 3D and 4D triple-resonance experiments, improved resolution, McCoy, **130**, 341
- Constant-time HCCH-COSY
- spin-echo difference, measurement of carbon-phosphorus  $J$  coupling constants in RNA, Hoogstraten and Pardi, **133**, 236
- Continuous-wave EPR
- nonlinear,  $T_1$  measurement by, Cianflone, Francia, and Leporini, **131**, 86
  - time-domain EPR for high-spin iron(III)-nitroxyl distances in spin-labeled porphyrins, Rakowsky *et al.*, **131**, 97
- Contrast
- prediction using binary spin-bath model and super-Lorentzian lineshape: magnetization transfer imaging of rat brain under non-steady-state conditions, Quesson *et al.*, **130**, 321
  - $T_1$ -dependent, optimization with nonstandard flip angles in spin-echo and inversion-recovery MR imaging, Bonny *et al.*, **130**, 51
- Contrast agents
- single-electron, for oximetric imaging, EPR and DNP properties, Ardenkjær-Larsen *et al.*, **133**, 1
  - superparamagnetic particles, intravascular and intracellular hepatic relaxivities, Colet *et al.*, **134**, 199
- Convection compensation
- in gradient-enhanced NMR, Jerschow and Müller, **132**, 13
- Convection suppression
- and temperature imaging by  $^1\text{H}$  NMR, in NMR probes, Hedin and Furó, **131**, 126
- Coordination complex
- Li, formation with nitroxides in aqueous phase: magnetic relaxation dispersion of  $^7\text{Li}$ , Dinesen and Bryant, **132**, 19
- Cope rearrangement
- semibullvalene valence tautomers, high-precision  $^{13}\text{C}$  shift thermometer for temperature range 100–300 K based on, Quast *et al.*, **134**, 236
- Copper
- planar, echo decay in high-temperature superconductors, contributions of spin-lattice relaxation, Curro and Slichter, **130**, 186
- Copper(I) phosphines
- linear, anisotropy in  $^{31}\text{P}$ ,  $^{63,65}\text{Cu}$  indirect spin-spin coupling and  $^{31}\text{P}$  nuclear shielding tensors, Kroeker *et al.*, **135**, 208
- CORE
- component separation in NMR imaging and multidimensional spectroscopy based on prior knowledge, Stilbs, **135**, 236
- Correction method
- for  $k$ -space trajectory deviations in MRI, Duyn *et al.*, **132**, 150
- Corrections and additions, **132**, 330; **134**, 194
- Correlated noise
- complex patterns, effect on measurement of set of peak parameters, Golding and Golding, **135**, 380
- Correlational times
- motional, direct determination by 1D MAS and 2D exchange NMR, Favre, Schaefer, and Chmelka, **134**, 261
- Correlation spectra
- multidimensional solid-state NMR, intrinsic asymmetry, Caldarelli and Emsley, **129**, 233
  - 2D, automated processing, Sengstschmid, Sterk, and Freeman, **131**, 315
  - 2D homonuclear, in MAS NMR, phase-sensitive detection, Boender and Vega, **133**, 281
- Correlation spectroscopy (*see* COSY)
- Correlation time
- extremely long, in glassy materials, sensitivity of saturation-transfer ESR extended to, van den Dries, de Jager, and Hemminga, **131**, 241
- COSY
- active-coupling-pattern tilting experiment for determination of homonuclear coupling constants, Koźmiński, **134**, 189
  - H(C)CH-COSY and (H)CCH-COSY,  $^{13}\text{C}$ -labeled proteins in  $\text{H}_2\text{O}$  solution, Gehring and Ekiel, **135**, 185
  - multiple-pulse, simplification of  $^{19}\text{F}$  NMR spectra of liquid crystalline samples, Lesot *et al.*, **133**, 166
  - quantitative  $J$  correlation, determination of three-bond  $^1\text{H}3'-^{31}\text{P}$  couplings in nucleic acids and protein-nucleic acid complexes by, Clore *et al.*, **134**, 164
  - spin-echo difference constant-time HCCH-COSY, carbon-phosphorus  $J$  coupling constants in RNA using, Hoogstraten and Pardi, **133**, 236
  - suppression of diagonal peaks with singular value decomposition, Zhe *et al.*, **132**, 176

- Coupled spin systems  
in-phase double selective excitation using excitation sculpting, Bourg and Nuzillard, **133**, 173
- Coupling (*see also* Dipolar coupling; Spin–spin coupling)  
magnetic, protein-mediated, between lactate and water protons, Swanson, **135**, 248
- Coupling constants (*see* Spin–spin coupling)
- CP-MAS (*see* Cross polarization–magic-angle spinning)
- CRAMPS  
MSHOT-3 based,  $^1\text{H}$  chemical-shielding anisotropies from polycrystalline powders, Hohwy *et al.*, **133**, 374
- Creatine  
singlet, simultaneous detection in *in vivo* lactate editing at 1.5 T using PRESS excitation, in head and neck tumors, Star-Lack *et al.*, **133**, 243
- Cross-correlation rates  
chemical-shift anisotropy–dipolar coupling  
estimation from laboratory-frame relaxation experiments, Ghose and Prestegard, **134**, 308  
in spin relaxation of  $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, Ghose, Huang, and Prestegard, **135**, 487
- Cross peaks  
detection in 2D proton NMR spectra using excitation sculpting, Van and Shaka, **132**, 154  
IS, spin-state selective editing,  $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for, Andersson, Annala, and Otting, **133**, 364
- Cross polarization (*see also* Heteronuclear cross polarization)  
cyclic  $J$  CP imaging and spectroscopy, analysis of carbohydrate metabolism and transport in castor bean seedlings by, Heidenreich *et al.*, **132**, 109  
for  $^1\text{H}$  NMR image contrast in solids, Nakai *et al.*, **134**, 44  
rotor-synchronized amplitude-modulated, application in  $^{13}\text{C}$ – $^1\text{H}$  spin pair under fast MAS, Raya and Hirschinger, **133**, 341  
surface selective  $^1\text{H}/^{29}\text{Si}$  CP NMR by NOE enhancement from laser-polarized xenon, Pietraß, Seydoux, and Pines, **133**, 299
- Cross polarization–magic-angle spinning  
chemical-shift imaging in rotating solids by RF field gradients, Malveau *et al.*, **134**, 171  
 $^{13}\text{C}$  linewidths, in solids for rigid strongly coupled carbons under CW proton decoupling, VanderHart and Campbell, **134**, 88  
 $^{15}\text{N}$  chemical-shift anisotropy of ampicillin and penicillin-V, Antzutkin, Lee, and Levitt, **135**, 144  
signals at high spinning speeds, simulation, Ray, Ladizhansky, and Vega, **135**, 427  
 $^{119}\text{Sn}$  NMR spectra of *para*-substituted triaryltin(pentacarbonyl)manganese(I) complexes, quadrupole–dipole effects, Christendat *et al.*, **131**, 1  
2D,  $^{13}\text{C}$  isotropic chemical shift correlation, established by  $^1\text{H}$  spin diffusion, Wilhelm *et al.*, **134**, 255
- Cross relaxation  
peaks, multiple-pulse mixing sequences selectively enhancing, Hwang and Shaka, **135**, 280  
water protons and  $^{19}\text{F}$  nuclei in perfluoroalkyl chain, Raulet *et al.*, **133**, 324
- Cross-relaxation time  
 $T_{IS}$ ,  $^1\text{H}$ – $^{13}\text{C}$  dipolar interactions through, cross polarization for  $^1\text{H}$  NMR image contrast in solids, Nakai *et al.*, **134**, 44
- Cryoprotection  
effects on structure and activity of p21 *ras*: implications for ESEEM, Halkides, Farrar, and Singel, **134**, 142
- Crystals  
experimental determination of four-time stimulated echoes, Hinze *et al.*, **131**, 218
- CSA (*see* Chemical-shielding anisotropy; Chemical-shift anisotropy)
- $^{63}\text{Cu}$   
 $\text{Cu}_2\text{O}$  powder, NMR imaging using second-order quadrupole broadened resonances, Swaminathan and Suits, **132**, 274
- $^{63,65}\text{Cu}$   
anisotropy in  $^{31}\text{P}$ ,  $^{63,65}\text{Cu}$  indirect spin–spin coupling and  $^{31}\text{P}$  nuclear shielding tensors of linear copper(I) phosphines, Kroeker *et al.*, **135**, 208
- $\text{Cu}_2\text{O}$  powder  
 $^{63}\text{Cu}$  NMR imaging using second-order quadrupole broadened resonances, Swaminathan and Suits, **132**, 274
- Cutinase  
 $^{13}\text{C}$ -labeled, assignment of aromatic side chains, 2D NMR experiments for, Prompers *et al.*, **130**, 68
- Cyclic  $J$  cross-polarization  
based imaging and spectroscopy, analysis of carbohydrate metabolism and transport in plant seedlings by, Heidenreich *et al.*, **132**, 109
- cyclo*-Pro-Gly  
analysis of error propagation from NMR-derived internuclear distances into molecular structure, Džakula *et al.*, **135**, 454
- Cytochrome  $c_3$   
uniformly  $^{15}\text{N}$ -labeled, interference contribution to  $T_2$  of imido protons of coordinated imidazoles, paramagnetic inversion of sign of, Ohmura *et al.*, **131**, 367

## D

- Data processing  
blind source separation use in NMR for model-free analysis of mixtures, Nuzillard, Bourg, and Nuzillard, **133**, 358  
frequency-selective MRS data quantification with frequency prior knowledge, Dologlou, Van Huffel, and Van Ormondt, **130**, 238  
spectral parameter estimation by iterative quadratic maximum likelihood method, Zhu, Choy, and Sanctuary, **135**, 37  
2D NMR, application of filter diagonalization method, Mandelshtam, Taylor, and Shaka, **133**, 304
- Davies ENDOR  
matrix line, Astashkin and Kawamori, **135**, 406
- Decay data  
multiexponential, uniform-penalty inversion, Borgia, Brown, and Fantazzini, **132**, 65
- Decomposition  
rank-revealing orthogonal, fast removal of residual water in proton spectra, Vanhamme *et al.*, **132**, 197  
singular value, suppression of diagonal peaks with, Zhe *et al.*, **132**, 176
- Deconvolution  
reference, NMR spectra by 1D filter diagonalization method, Hu *et al.*, **134**, 76  
reference spectrum,  $R_2'$  measurement in presence of multiple spectral components, Wehrli *et al.*, **131**, 61
- Decoupling  
adiabatic (*see* Adiabatic decoupling)  
complete dipolar, of  $^{13}\text{C}$ , and use in 2D double-quantum solid-state NMR for determining polymer conformations, Schmidt-Rohr, **131**, 209  
CW proton,  $^{13}\text{C}$  CPMAS linewidths in solids for rigid strongly coupled carbons under, VanderHart and Campbell, **134**, 88  
WALTZ-16, heating effects: high-precision  $^{13}\text{C}$  shift thermometer for temperature range 100–300 K, Quast *et al.*, **134**, 236
- DECRA (*see* Direct exponential curve resolution algorithm)
- Deltorphan I analog  
gradient- and sensitivity-enhanced TOCSY, Kövér, Uhrin, and Hruby, **130**, 162
- Demagnetizing field  
solvent magnetization artifacts in high-field NMR studies of macromolecular hydration, Sobol *et al.*, **130**, 262
- Density matrix  
INEPT experiments involving quadrupolar nuclei in solids, Kao and Grey, **133**, 313
- Density operator theory  
analysis of catalytic processes with parahydrogen, Natterer *et al.*, **133**, 92

- DEPT  
modified, *in vivo* 3D localized  $^{13}\text{C}$  spectroscopy, Watanabe *et al.*, **134**, 214
- DEPTQ  
distortionless enhancement by polarization transfer including detection of quaternary nuclei, Burger and Bigler, **135**, 529
- Detection coil quality factor  
super-Q detection of transient MR signals, Suits, Garroway, and Miller, **132**, 54
- Deuteration  
effects on accuracy of HN–HN distance constraints, Briercheck and Rule, **134**, 52
- Deuteron exchange NMR  
2D, in study of vacancy diffusion in polycrystalline benzene, comparison with NMR stimulated echo, Isfort, Geil, and Fajara, **130**, 45
- Diagonal peaks  
suppression with singular value decomposition, Zhu *et al.*, **132**, 176
- Diamonds  
99%  $^{13}\text{C}$ -enriched, spin–lattice relaxation,  $^{13}\text{C}$  NMR analysis, Shabanova, Schaumburg, and Sellschop, **130**, 8
- Dianisyl-nitroxide radical  
disordered powder sample, high-frequency/high-field ESEEM study of nitrogen hyperfine and quadrupole interactions, Bloß, Möbius, and Prisner, **134**, 30
- 1,3-Dibromoadamantane  
rotational motion in polychlorotrifluoroethene,  $^{13}\text{C}$  NMR, Lienin, Brüsche-weiler, and Ernst, **131**, 184
- o*-Dichlorobenzene  
solute–solvent interactions of fullerenes  $\text{C}_{60}$  and  $\text{C}_{70}$  in solution: 1D steady-state intermolecular  $^{13}\text{C}$ ,  $^1\text{H}$  Overhauser effect study, Troganis, Gerothanassis, and Papadimitriou, **130**, 149
- Diffusion  
and DNP enhancement, in aqueous char suspensions, Odintsov *et al.*, **135**, 435  
laser-polarized noble gas, time-dependent, PFG measurements, Mair *et al.*, **135**, 478  
measurements using RF field gradient, Humbert *et al.*, **134**, 245  
modeling of self-diffusion and relaxation time NMR in multi-compartment systems, Novikov, van Dusschoten, and Van As, **135**, 522  
non-mono-exponential attenuation of water and *N*-acetyl aspartate signals in brain tissue due to, Assaf and Cohen, **131**, 69  
proteins in aqueous solution, practical aspects of measurement, Tillett, Lian, and Norwood, **133**, 379  
proton PFG diffusion and  $T_1$  relaxation in polyacrylamide gels, unified approach using volume averaging, Penke *et al.*, **132**, 240  
restricted  
in sea ice in Antarctica, study by Earth's field NMR: pulsed gradient spin echo NMR, Callaghan *et al.*, **133**, 148  
spin-echo attenuation, as discord of structure waves, Stepišnik, **131**, 339  
rigid body, Green function: effect of coupling between rotational and translational Brownian motions on NMR spin relaxation: Beloborodov, Orekhov, and Arseniev, **132**, 328  
temperature imaging by  $^1\text{H}$  NMR and suppression of convection, Hedin and Furó, **131**, 126  
3D DOSY–HMQC experiment for high-resolution analysis of complex mixtures, Barjat, Morris, and Swanson, **131**, 131
- Diffusion coefficients  
measurement by 2D NMR using accordion spectroscopy, Millet and Pons, **131**, 166
- Diffusion NMR  
and  $^{13}\text{C}$  isotope editing: isotope-filtered NMR, Gonnella *et al.*, **131**, 336  
imaging, component separation through global least-squares analysis based on prior knowledge, Stilbs, **135**, 236
- Diffusion-ordered spectroscopy (DOSY)  
experiments on polymers, multivariate curve resolution, Van Gorkom and Hancewicz, **130**, 125
- Diffusivity  
ionic charge carriers, NMR measurement, Heil and Holz, **135**, 17
- Digitization  
signal, improved resolution and signal-to-noise ratio in MRI via, Elliott *et al.*, **130**, 300
- Dimers  
magnetic exchange coupled, nuclear and electron relaxation, Bertini *et al.*, **130**, 33
- Dimethylsulfone  
motional correlational times, direct determination by 1D MAS and 2D exchange NMR, Favre, Schaefer, and Chmelka, **134**, 261
- Dimyristoylphosphatidylcholine  
uniformly aligned bilayers, magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305
- 2,2-Di(4-*tert*-octylphenyl)-1-picrylhydrazyl (DPPH)  
 $\pi$  and  $\sigma$  transitions, DNP study at 1.53 mT, Salman *et al.*, **134**, 1
- $\alpha,\alpha$ Diphenyl- $\beta$ -picrylhydrazyl  
application of radiofrequency 220-MHz FT EPR spectrometer, Alecci *et al.*, **130**, 272
- Dipolar coupling  
–chemical-shift anisotropy cross-correlation rates  
estimation from laboratory-frame relaxation experiments, Ghose and Prestegard, **134**, 308  
in spin relaxation of  $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, Ghose, Huang, and Prestegard, **135**, 487  
methylene and methyl sites in weakly oriented macromolecules, Ottiger *et al.*, **134**, 365  
residual, direct structure refinement against, in presence of rhombicity of unknown magnitude, Clore, Gronenborn, and Tjandra, **131**, 159  
simplified 2D NMR spectra, Ottiger, Delaglio, and Bax, **131**, 373
- Dipolar decoupling  
complete, of  $^{13}\text{C}$ , and use in 2D double-quantum solid-state NMR for determining polymer conformations, Schmidt-Rohr, **131**, 209
- Dipolar interactions  
 $^1\text{H}$ – $^{13}\text{C}$ , through cross relaxation time  $T_{IS}$ , cross polarization for  $^1\text{H}$  NMR image contrast in solids, Nakai *et al.*, **134**, 44  
 $^{23}\text{Na}$  in solution, determination by triple-quantum relaxation time measurements, Eliav and Navon, **130**, 63
- Dipolar solids  
2D MAS spectroscopy, pulsed-field gradient selection, Fritzhanns *et al.*, **134**, 355
- Dipolar tensor  
 $^{13}\text{C}^\alpha$ – $^1\text{H}^\alpha$ , and  $^{15}\text{N}$  chemical shift tensor, correlation in solid-state MAS NMR, peptide torsion angle determination by, Hong *et al.*, **135**, 169
- Dipole–CSA cross-correlated relaxation  
quantitative analysis by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346
- Dipole–dipole interactions  
nuclear spin systems with, parametrical instability, Buishvili, Giorgadze, and Khomeriki, **130**, 82
- Direct exponential curve resolution algorithm  
multivariate image analysis of MR images  
algorithm and model study, Windig, Hornak, and Antalek, **132**, 298  
human brain images, Antalek, Hornak, and Windig, **132**, 307
- Director fluctuations  
formalism for evaluation of order-fluctuation modes in liquid crystals from field-cycling NMR-relaxometry data, Grinberg *et al.*, **135**, 54
- Direct summation  
peak integrals obtained by, influence of noise on, Bourg and Nuzillard, **134**, 184



- Disposition kinetics  
 carboplatin, monitoring *in vivo* after sc injection in rat using  $^{195}\text{Pt}$  NMR, Becker *et al.*, **133**, 115
- Dissipative media  
 surface and gradiometer coils near conducting body: lift-off effect, Suits, Garroway, and Miller, **135**, 373
- Distance constraints  
 HN–HN, deuteration effects on accuracy, Briercheck and Rule, **134**, 52
- Distance geometry  
 analysis of error propagation from NMR-derived internuclear distances into molecular structure of *cyclo*-Pro-Gly, Džakula *et al.*, **135**, 454
- Distributed parallel processing  
 for multidimensional maximum entropy reconstruction, Li, Stern, and Hoch, **134**, 161
- Distributions  
 quadrupolar and isotropic chemical shift interactions in 2D multiple-quantum MAS NMR spectra, Bodart, **133**, 207
- DNA  
 duplex, dynamics,  $^{13}\text{C}$  relaxation study, Kojima *et al.*, **135**, 310  
 internuclear angles, determination using paramagnetic-assisted magnetic alignment, Beger *et al.*, **135**, 256  
 structural studies, interresidue quiet NOE for, Beger *et al.*, **132**, 34  
 weak labile  $^1\text{H}$  resonances of protein side chains contacting, enhancement:  $^{15}\text{N}$ -edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87
- DNP (*see* dynamic nuclear polarization)
- Dorsal stream  
 activation during visually guided movements, functional MRI, Ellermann *et al.*, **131**, 272
- DOSY  
 experiments on polymers, multivariate curve resolution, Van Gorkom and Hancewicz, **130**, 125
- DOSY-HMQC  
 3D experiment for high-resolution analysis of complex mixtures, Barjat, Morris, and Swanson, **131**, 131
- Double-adiabatic decoupling  
 homonuclear, Bloch–Siegert shift-compensated and cyclic sideband-eliminated, for  $^{13}\text{C}$ - and  $^{15}\text{N}$ -double-labeled proteins, Zhang and Gorenstein, **132**, 81
- Double pulsed-field-gradient spin echo  
 cross peak detection in 2D proton NMR spectra using excitation sculpting, Van and Shaka, **132**, 154  
 ultraclean multisite selective excitation using excitation sculpting, Parella, Sánchez-Ferrando, and Virgili, **135**, 50
- Double-quantum-filtered homonuclear spectroscopy  
 rapid-pulsing artifact-free: 2D-INADEQUATE experiment revisited, Bourdonneau and Ancian, **132**, 316
- Double-quantum filtered NMR  
 selective detection of biexponential relaxation in isotropic solutions: mixed coherence filtering NMR, Yamada *et al.*, **135**, 99
- Double-quantum NMR  
 2D solid-state, for determining polymer conformations, complete dipolar decoupling of  $^{13}\text{C}$  for, Schmidt-Rohr, **131**, 209
- Double-quantum spectroscopy  
 heteronuclear X-filter  $^1\text{H}$  PFG, for proton resonance assignment of ligand bound to protein, Dalvit, Ramage, and Hommel, **131**, 148
- Double-quantum transition  
 in high-frequency EPR study of integer spin system, van Dam *et al.*, **130**, 140
- Double resonance  
 probe, for close frequencies, Haase, Curro, and Slichter, **135**, 273
- DPPH (2,2-Di(4-*tert*-octylphenyl)-1-picrylhydrazyl)  
 $\pi$  and  $\sigma$  transitions, DNP study at 1.53 mT, Salman *et al.*, **134**, 1
- DQ-COSY  
 automated processing of 2D correlation spectra, Sengstschmid, Sterk, and Freeman, **131**, 315
- DQ/ZQ-HCCH-E.-COSY  
 methylene specific experiments for scalar spin–spin coupling constants between protons attached to  $^{13}\text{C}$ , Carlomagno *et al.*, **135**, 216
- Drosomycin  
 NOESY: composite  $90^\circ$  pulse sequence application, Sodano, Landon, and Ptak, **133**, 194
- Drying  
*in situ*, for preparing NMR samples with extremely low water content, Goetz, **135**, 14
- Dynamic nuclear polarization  
 enhancement  
 in aqueous char suspensions, Odintsov *et al.*, **135**, 435  
 in proton–electron double-resonance imaging, mobility and free radical concentration effects, de Sousa *et al.*, **135**, 118  
 $\pi$  and  $\sigma$  transitions for nitroxide spin probes at 1.53 mT, Salman *et al.*, **134**, 1  
 properties, of single-electron contrast agents for oximetric imaging, Ardenkjær-Larsen *et al.*, **133**, 1
- Dynamic range  
 improved, in high-resolution NMR, analog filtering of large solvent signals for, Redfield and Kunz, **130**, 111
- Dynamics  
 bulk magnetic moment, in MRI of lung with hyperpolarized noble gases, Möller *et al.*, **135**, 133  
 DNA duplex,  $^{13}\text{C}$  relaxation study, Kojima *et al.*, **135**, 310  
 protein backbone: quantitative analysis of dipole–CSA cross-correlated relaxation by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346  
 protein side-chain, analysis from  $^{13}\text{C}$  NMR relaxation data, Daragan and Mayo, **130**, 329
- Dynamic susceptibility contrast  
 sequential,  $T_1$  effects, Levin *et al.*, **130**, 292
- E
- Earth's field NMR  
 in Antarctica: pulsed gradient spin echo NMR study of restricted diffusion in sea ice, Callaghan *et al.*, **133**, 148
- Echo (*see* Spin echo)
- Echo decay  
 planar Cu in high-temperature superconductors, contributions of spin–lattice relaxation, Curro and Slichter, **130**, 186
- Echo-planar imaging  
 multishot, image-based reduction of artifacts, Hennel, **134**, 206  
 radial EPI, description and experimental results, Silva *et al.*, **135**, 242
- Echo-planar imaging relaxometry  
 measurement of viscosity of model meal, Marciani *et al.*, **135**, 82
- Echo response  
 Hahn pulse sequence, shape of, influence of high orientational order, van der Struijf *et al.*, **130**, 253  
 nitroxide labels in proteins in lipid bilayers: use of phase memory time  $T_2$  for quantitation of rotational motions, van der Struijf and Levine, **130**, 244
- E.COSY  
 DQ/ZQ-HCCH-E.-COSY, methylene specific experiments for scalar spin–spin coupling constants between protons attached to  $^{13}\text{C}$ , Carlomagno *et al.*, **135**, 216
- Eddy currents  
 localized compensation using quantitative field mapping, Terpstra, Andersen, and Gruetter, **131**, 139
- Editing  
 DEPTQ: distortionless enhancement by polarization transfer including detection of quaternary nuclei, Burger and Bigler, **135**, 529

- in vivo* lactate editing with simultaneous detection of upfield singlets at 1.5 T using PRESS excitation, tumor applications, Star-Lack *et al.*, **133**, 243
- multidimensional, for amide deuterium isotope effects on  $C^\beta$  chemical shifts in  $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, Meissner and Sørensen, **135**, 547
- $^{15}\text{N}$ -edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87
- spin-state selective, of IS cross peaks,  $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for, Andersson, Annala, and Otting, **133**, 364
- taurine detection *in vitro* and *in vivo*, Hardy and Norwood, **133**, 70
- Editorial
- new Associate Editor and Editorial Board members, **130**, v
- Effective gradients
- in porous media due to susceptibility differences, Hürlimann, **131**, 232
- EIS
- direct determination of motional correlational times, Favre, Schaefer, and Chmelka, **134**, 261
- Electrical mobility
- ionic charge carriers, NMR measurement, Heil and Holz, **135**, 17
- Electrical transport
- in disordered medium, NMR studies, Heil and Holz, **135**, 17
- Electron-nuclear double resonance
- pulsed (*see* Pulsed ENDOR)
- Electron paramagnetic resonance (*see* EPR; Pulsed EPR)
- Electron relaxation
- and nuclear relaxation, in magnetic exchange coupled dimers, Bertini *et al.*, **130**, 33
- Electron spin echo envelope modulation (*see* ESEEM)
- Electron spin resonance (*see* EPR)
- Electrophoretic NMR
- with PFG NMR, analysis of electrical transport in disordered medium, Heil and Holz, **135**, 17
- ENDOR
- pulsed (*see* Pulsed ENDOR)
- EPR
- absorption, first-harmonic out-of-phase signals,  $T_1$  measurements, Livshits, Páli, and Marsh, **134**, 113
- aqueous samples, rectangular loop-gap resonator for, Piasecki, Froncisz, and Hubbell, **134**, 36
- continuous-wave nonlinear ESR spectroscopies,  $T_1$  measurement by, Cianflone, Francia, and Leporini, **131**, 86
- FT-EPR with nonresonant probe: use of truncated coaxial line, Rubinson *et al.*, **132**, 255
- 1–370 GHz linewidths for  $\text{K}_3\text{CrO}_8$ : test of Anderson–Weiss model, Cage *et al.*, **135**, 178
- 220-MHz RF Fourier transform EPR spectrometer, Alecci *et al.*, **130**, 272
- properties, of single-electron contrast agents for oximetric imaging, Ardenkjær-Larsen *et al.*, **133**, 1
- proxyl spin probe, Gaussian line broadening, contributions from magnetic-field modulation and unresolved proton hyperfine structure, Bales, Peric, and Lamy-Freund, **132**, 279
- pyridinium tetrafluoroborate, ferroelectric phase transitions, radicals as probes, Krzymiński, Bielewicz-Mordalska, and Kowalczyk, **135**, 76
- slow motional spectra, rapid computation of rotational correlation time, neural network approach, Martinez and Millhauser, **134**, 124
- spectra, computer simulation employing homotopy, Gates *et al.*, **135**, 104
- EPR imaging
- spectral-spatial imaging algorithm, implementation by iterative Fourier reconstruction, Placidi *et al.*, **134**, 280
- EPR oximetry
- carbon-based paramagnetic materials for use in, separation and enrichment of active component, Liu *et al.*, **133**, 291
- Error propagation
- from NMR-derived internuclear distances into molecular structure of cyclo-Pro-Gly, Džakula *et al.*, **135**, 454
- Errors
- noise-induced: precision of parameters in measuring resonance spectra, Golding and Golding, **135**, 380
- Erythrocytes
- suspensions,  $^{23}\text{Na}$  multiple-quantum-filtered NMR spectra, contribution of extracellular sodium, Knubovets, Shinar, and Navon, **131**, 92
- ESEEM
- high-frequency/high-field, study of nitrogen hyperfine and quadrupole interactions on disordered powder sample, Bloß, Möbius, and Prisner, **134**, 30
- implications of effects of cryoprotection on p21 *ras* structure and activity, Halkides, Farrar, and Singel, **134**, 142
- multifrequency powder spectra, determination of nitrogen quadrupole and hyperfine tensors in oxovanadium complex by simultaneous fitting of, Reijerse, Tyryshkin, and Dikanov, **131**, 295
- 1D and 2D, sensitivity enhancement by matched microwave pulses, Jeschke, Rakhmatullin, and Schweiger, **131**, 261
- optically detected, on photoexcited triplet state in zero magnetic field, Weis, Möbius, and Prisner, **131**, 17
- spectra, efficient simulation using Gamma, Shane, Liesum, and Schweiger, **134**, 72
- 2D nuclear-Zeeman-resolved experiment, Willer *et al.*, **133**, 46
- Evolution strategy
- optimization, for selective pulses in NMR, Lunati *et al.*, **134**, 223
- Exchange-edited gradient HCP-HSQC
- sensitivity-enhanced detection of fast exchanging protons, Zangger and Armitage, **135**, 70
- Exchange NMR
- 2D, and 1D MAS, direct determination of motional correlational times by, Favre, Schaefer, and Chmelka, **134**, 261
- Excitation
- composite pulse schemes, for MQMAS NMR of half-integer quadrupolar spins, Marinelli, Medek, and Frydman, **132**, 88
- PRESS, *in vivo* lactate editing with simultaneous detection of upfield singlets at 1.5 T with, applications to brain and head and neck tumors, Star-Lack *et al.*, **133**, 243
- Excitation sculpting
- cross peak detection in 2D proton NMR spectra, Van and Shaka, **132**, 154
- in-phase double selective excitation of coupled spin systems, Bourg and Nuzillard, **133**, 173
- RF-gradient, in selective 1D and nonselective 2D experiments, Heikkinen, Rahkamaa, and Kilpeläinen, **133**, 183
- ultraclean multisite selective excitation, Parella, Sánchez-Ferrando, and Virgili, **135**, 50
- Extended Karplus equation
- vicinal fluorine-proton coupling constants: individual substituent effects, Fabián, Guilleme, and Díez, **133**, 255
- F
- $^{19}\text{F}$
- cesium pentadecafluorooctanoate:  $^{19}\text{F}$ - $^1\text{H}$  HOESY study of water-surfactant contact, Raulet *et al.*, **133**, 324
- liquid crystalline samples, NMR, simplification of spectra by multiple-pulse COSY, Lesot *et al.*, **133**, 166
- perfluorocarbon FC40, MRI in inhomogeneous fields by SPARE sequence, McIntyre, Hennel, and Morris, **130**, 58
- Fast imaging
- real-time image reconstruction and display system for MRI using high-speed personal computer, Haishi and Kose, **134**, 138

- <sup>57</sup>Fe  
shieldings, in superstructured hemoprotein models, effects of atropisomerism and porphyrin deformation, Geroathanassis *et al.*, **131**, 163
- Ferrocene  
<sup>13</sup>C–<sup>1</sup>H spin pair under fast MAS, application of rotor-synchronized amplitude-modulated cross polarization, Raya and Hirschinger, **133**, 341
- Field-cycling relaxometry  
data, formalism for evaluation of order-fluctuation modes in liquid crystals, Grinberg *et al.*, **135**, 54
- Field-gradient NMR  
spatial resolution, demonstrating, Feiweir *et al.*, **131**, 203
- Field mapping  
with amplitude images at low field, use of phase reference for, Tomasi *et al.*, **131**, 310  
automated shimming with normal spectrometer hardware using: Chilvers and Morris, **133**, 210
- Fields  
eddy current, quantitative mapping, localized eddy current compensation, Terpstra, Andersen, and Gruetter, **131**, 139
- Filter diagonalization method  
ID, reference deconvolution, phase correction, and line listing of NMR spectra by, Hu *et al.*, **134**, 76  
1D and 2D NMR spectra, Mandelstam, Taylor, and Shaka, **133**, 304
- Filtered back projection  
in iterative applications, Fourier reconstruction as alternative: Fourier spectral spatial EPR imaging, Placidi *et al.*, **134**, 280
- Filtering  
frequency-selective MRS data quantification with frequency prior knowledge, Dologlou, Van Huffel, and Van Ormondt, **130**, 238
- Flavodoxin  
detection of <sup>15</sup>N–<sup>15</sup>N *J*-couplings, Löhner and Rüterjans, **132**, 130
- Flip angles  
arbitrary, simultaneous and independent rotations with, for I, IS<sup>α</sup>, and IS<sup>β</sup> spin systems, Briand and Sørensen, **135**, 44  
effects in STEAM and PRESS—optimized vs sinc RF pulses, Ryner, Ke, and Thomas, **131**, 118  
nonstandard, in optimization of signal intensity and *T*<sub>1</sub>-dependent contrast in spin-echo and inversion-recovery MR imaging, Bonny *et al.*, **130**, 51
- Floquet theory  
spectral simulation using: proton MAS NMR at high magnetic fields and high spinning frequencies, Ray *et al.*, **135**, 418  
time propagation technique for MAS NMR simulation derived from, application to quadrupolar nuclei, Charpentier, Fermon, and Virlet, **132**, 181
- Flowing spins  
steady-state saturation, pulse sequences for, Marshall, **133**, 13
- Fluids  
flowing, steady-state saturation of spins, pulse sequences for, Marshall, **133**, 13  
in sedimentary rock, *in situ* NMR analysis, de Swiet *et al.*, **133**, 385
- Fluorine  
vicinal F–H coupling constants: individual substituent effects, Fabián, Guilleme, and Díez, **133**, 255
- Fluoroethane derivatives  
vicinal fluorine–proton coupling constants: individual substituent effects, Fabián, Guilleme, and Díez, **133**, 255
- Fourier reconstruction  
as alternative to filtered back projection in iterative applications: Fourier spectral spatial EPR imaging, Placidi *et al.*, **134**, 280
- Free radicals  
concentration effects in proton–electron double-resonance imaging, de Sousa *et al.*, **135**, 118
- Frequency  
frequency-selective MRS data quantification with frequency prior knowledge, Dologlou, Van Huffel, and Van Ormondt, **130**, 238  
Frequency prior knowledge  
frequency-selective MRS data quantification with, Dologlou, Van Huffel, and Van Ormondt, **130**, 238
- Frequency separation  
precision in estimating, Jupp, Harris, and Aliev, **135**, 23
- FT-EPR  
with nonresonant probe: use of truncated coaxial line, Rubinson *et al.*, **132**, 255
- Fullerenes  
C<sub>60</sub> and C<sub>70</sub> solute–solvent interactions in solution: 1D steady-state intermolecular <sup>13</sup>C, <sup>1</sup>H Overhauser effect study, Troganis, Geroathanassis, and Papademitriou, **130**, 149
- G
- Gamma  
efficient simulation of ESEEM spectra, Shane, Liesum, and Schweiger, **134**, 72
- Gamma carbon  
aliphatic, and backbone amides, resonances in deuterated proteins, unambiguous correlations, McCallum, Hitchens, and Rule, **134**, 350
- Gas diffusion  
time-dependent, PFG measurements, Mair *et al.*, **135**, 478
- Gases  
hyperpolarized, low-frequency NMR polarimeter for, Saam and Conradi, **134**, 67
- Gastrointestinal tract  
viscosity of model meal, measurement with echo-planar imaging relaxometry, Marciani *et al.*, **135**, 82
- Gaussian approximation  
in analysis of *T*<sub>1</sub> contributions to planar Cu echo decay in high-temperature superconductors, Curro and Slichter, **130**, 186
- Gaussian line broadening  
proxyl spin probe EPR spectrum, contributions from magnetic-field modulation and unresolved proton hyperfine structure, Bales, Peric, and Lamy-Freund, **132**, 279
- Gaussian spherical quadrature  
computation of orientational averages in solid-state NMR by, Edén and Levitt, **132**, 220
- Gel permeation chromatography-NMR  
experiments on polymers, multivariate curve resolution, Van Gorkom and Hancewicz, **130**, 125
- Glassy materials  
extremely slow molecular mobility, sensitivity of saturation-transfer ESR extended to, van den Dries, de Jager, and Hemminga, **131**, 241  
ultraslow motion in metallic glass, detection using alignment echo of spin-<sup>3</sup>/<sub>2</sub> <sup>9</sup>Be nuclei, Tang and Wu, **133**, 155
- Global correlation time  
protein, improved estimation from <sup>15</sup>N relaxation measurements, Yao, Hinds, and Norton, **131**, 347
- Glutamine  
and Asn, triple-resonance experiment selective for, multiplicity-selective coherence transfer steps for design of, Schmieler *et al.*, **131**, 199
- Glutathione *S*-transferase  
deuterated, correlation of backbone amide and aliphatic gamma resonances, McCallum, Hitchens, and Rule, **134**, 350
- Glycerol  
cryoprotection, effects on structure and activity of p21 *ras*: implications for ESEEM, Halkides, Farrar, and Singel, **134**, 142  
saturation-transfer ESR spectra of TEMPOL spin label, van den Dries, de Jager, and Hemminga, **131**, 241  
solutions, dipolar interactions of <sup>23</sup>Na, determination by triple-quantum relaxation time measurements, Eliav and Navon, **130**, 63

- Glycine  
<sup>13</sup>C, six-pulse PASS MAS-NMR technique solved by phasor method, Effey, Butcher, and Cappellotti, **132**, 266  
<sup>13</sup>C-<sup>14</sup>N REAPDOR NMR experiment optimization, Ba *et al.*, **133**, 104
- GOESY  
convection compensation, Jerschow and Müller, **132**, 13
- gp41  
SIV, 44-kDa ectodomain, <sup>15</sup>N relaxation measurements in 3D NMR experiments, Caffrey *et al.*, **135**, 368
- GPC-NMR  
experiments on polymers, multivariate curve resolution, Van Gorkom and Hancewicz, **130**, 125
- Gradient coils  
multilayer, design, Bowtell and Robyr, **131**, 286
- Gradient-echo sequences  
artifacts induced by weakly ferromagnetic metals: field mapping with amplitude images at low field, Tomasi *et al.*, **131**, 310
- Gradient-enhanced SIMBA  
ID determination of long-range heteronuclear coupling constants by, Vogt and Benesi, **132**, 214
- Gradients (*see also* Pulsed-field gradients)  
effective, in porous media due to susceptibility differences, Hürlimann, **131**, 232  
exchange-edited gradient HCP-HSQC, sensitivity-enhanced detection of fast exchanging protons, Zangger and Armitage, **135**, 70  
HSQC-based methyl group selection via, in multidimensional NMR of proteins, Diercks, Schwaiger, and Kessler, **130**, 335  
use with multiple-pulse mixing sequences enhancing chemical exchange or cross-relaxation peaks in high-resolution spectra, Hwang and Shaka, **135**, 280
- Gradiometer coils  
and surface coils, near conducting body: lift-off effect, Suits, Garroway, and Miller, **135**, 373
- Gramicidin A  
membrane-bound, <sup>15</sup>N chemical-shift anisotropy, cation binding-induced changes, Tian and Cross, **135**, 535
- Green function  
rigid body diffusion: effect of coupling between rotational and translational Brownian motions on NMR spin relaxation: Beloborodov, Orekhov, and Arseniev, **132**, 328
- GROESY  
convection compensation, Jerschow and Müller, **132**, 13
- Guanidine  
dilute solution, measurement of chemical exchange rate constants with solvent protons using radiation damping, Chen and Mao, **131**, 358
- Gyrase B  
*Staphylococcus aureus*,  $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for spin-state selective editing of IS cross peaks, Andersson, Annala, and Otting, **133**, 364
- H
- <sup>2</sup>H  
amide, effects on C <sup>$\beta$</sup>  chemical shifts in <sup>13</sup>C, <sup>15</sup>N-labeled proteins, multidimensional editing for, Meissner and Sørensen, **135**, 547  
glutathione S-transferase, correlation of backbone amide and aliphatic gamma resonances, McCallum, Hitchens, and Rule, **134**, 350  
naphthazarin B, hydrogen-banded, <sup>1</sup>H and <sup>2</sup>H NMR relaxation due to complex motion: classical jumps over barrier and incoherent tunneling, Reynhardt and Latanowicz, **130**, 195  
natural abundance spectra of molecules oriented in liquid crystals, Khetrpal *et al.*, **135**, 265  
nematic phospholipid bicellar solutions, structural study, Struppe and Vold, **135**, 541  
quadrupole coupling constant, and dynamic isotope effect, determination in liquids, Hardy *et al.*, **134**, 300  
selectively labeled RNA, uniformly <sup>13</sup>C-labeled, proton and carbon assignment, 3D C(CC)H TOCSY for, Dayie, Tolbert, and Williamson, **130**, 97  
time domain analysis of ultraslow reorientations in supercooled liquids, Geil, Fujara, and Sillescu, **130**, 18  
vacancy diffusion study in polycrystalline benzene by 2D deuteron exchange NMR and stimulated echoes: Isfort, Geil, and Fujara, **130**, 45
- Hardware  
double resonance probe for close frequencies, Haase, Curro, and Slichter, **135**, 273  
single-crystal NMR, Vosegaard *et al.*, **135**, 126
- HCN experiments  
triple-resonance, optimization for larger RNA oligonucleotides, Sklenář *et al.*, **130**, 119
- <sup>3</sup>He  
hyperpolarized, MRI of lung with, signal dynamics, Möller *et al.*, **135**, 133
- Heat shock proteins  
small, selective NMR experiments: QUIET-NOESY, Esposito *et al.*, **132**, 204
- Helical peptides  
Ala-rich, conformational effect of nitroxide spin labels on, NMR analysis, Bolin *et al.*, **131**, 248
- Heme proteins  
models, superstructured, effects of atropisomerism and porphyrin deformation on <sup>57</sup>Fe shieldings, Gerotheranassis *et al.*, **131**, 163  
resonance assignment in active site, <sup>1</sup>H-<sup>13</sup>C-<sup>13</sup>C-edited <sup>1</sup>H NMR experiment for, Qiu, Rivera, and Stark, **130**, 76
- Heterogeneity  
2D CP/MAS <sup>13</sup>C isotropic chemical shift correlation established by <sup>1</sup>H spin diffusion, Wilhelm *et al.*, **134**, 255
- Heteronuclear correlation  
multiple-bond, gradient-enhanced 2D, heteronuclear long-range couplings by, Sheng and van Halbeek, **130**, 296  
multiple-quantum (*see* HMQC)  
single-quantum (*see* HSQC)
- Heteronuclear coupling constants (*see* Spin-spin coupling, heteronuclear)
- Heteronuclear cross polarization  
enhanced sensitivity of *in vivo* <sup>13</sup>C MR spectroscopy on clinical 1.5 T MR system, van den Bergh, van den Boogert, and Heerschap, **135**, 93  
exchange-edited gradient HCP-HSQC in sensitivity-enhanced detection of fast exchanging protons, Zangger and Armitage, **135**, 70
- Heteronuclear NMR  
solid-state, high-frequency probes in, double- and triple-resonance circuits for, Zhang *et al.*, **132**, 167  
2D, experiments for assignment of aromatic side chains in <sup>13</sup>C-labeled proteins, Prompers *et al.*, **130**, 68  
3D, correlation of backbone amide and aliphatic gamma resonances in deuterated proteins, McCallum, Hitchens, and Rule, **134**, 350
- Heteronuclear Overhauser effect spectroscopy  
<sup>19</sup>F-<sup>1</sup>H, water-surfactant contact studied by, Raulet *et al.*, **133**, 324
- Heteronuclear scalar interaction  
and proton spin diffusion, self-decoupling: high-speed <sup>13</sup>C MAS of adamantane, Ernst, Verhoeven, and Meier, **130**, 176
- Heteronuclear two-spin system  
homogeneous master equation for, on basis of Cartesian product operators, Allard, Helgstrand, and Härd, **134**, 7
- Heteronuclear X filter  
<sup>1</sup>H PFG double-quantum, for proton resonance assignment of ligand bound to protein, Dalvit, Ramage, and Hommel, **131**, 148
- HFeCo<sub>3</sub>(CO)<sub>11</sub>PPh<sub>2</sub>H  
<sup>31</sup>P J-coupled to <sup>59</sup>Co nucleus, differential line broadening in presence of quadrupolar-CSA interference, Elbayed *et al.*, **130**, 209
- High-energy phosphates  
cardiac localized <sup>31</sup>P spectroscopy, Löffler *et al.*, **134**, 287

- High-field EPR  
ESEEM, study of nitrogen hyperfine and quadrupole interactions on disordered powder sample, Bloëß, Möbius, and Prisner, **134**, 30
- High-frequency EPR  
ESEEM, study of nitrogen hyperfine and quadrupole interactions on disordered powder sample, Bloëß, Möbius, and Prisner, **134**, 30  
integer spin systems: unusual behavior of double-quantum line, van Dam *et al.*, **130**, 140
- High magnetic field  
proton MAS NMR: spectral simulations using Floquet theory, Ray *et al.*, **135**, 418
- High pressure  
high-resolution high-pressure NMR probe for experiments at 500 MHz, Ballard *et al.*, **133**, 190
- High-resolution MRI  
3D, for Sandimmune Neoral effects on collagen arthritis in DA rat, Beckmann *et al.*, **131**, 8
- High-resolution NMR  
analog filtering of large solvent signals for improved dynamic range, Redfield and Kunz, **130**, 111  
control of spinning sidebands, Borer and Maple, **131**, 177  
multiple-pulse mixing sequences selectively enhancing chemical exchange or cross-relaxation peaks, Hwang and Shaka, **135**, 280  
spectra, multilinear interactions, Corio, **134**, 131  
transition-band signals, Szántay, **135**, 334
- HMQC  
alternative scheme for  $^{13}\text{C}$  chemical-shift imaging via inverse detection of protons through MILS technique, Lee *et al.*, **130**, 102  
-DOSY 3D, for high-resolution analysis of complex mixtures, Barjat, Morris, and Swanson, **131**, 131  
gradient- and sensitivity-enhanced, Zhu, Kong, and Sze, **135**, 232
- HN  
HN-HN distance constraints, accuracy, deuteration effects, Briercheck and Rule, **134**, 52
- HN(CACB)CG  
correlation of backbone amide and aliphatic gamma resonances in deuterated proteins, McCallum, Hitchens, and Rule, **134**, 350
- HN(COCACB)CG  
correlation of backbone amide and aliphatic gamma resonances in deuterated proteins, McCallum, Hitchens, and Rule, **134**, 350
- HOESY  
 $^{19}\text{F}$ - $^1\text{H}$ , water-surfactant contact studied by, Raulet *et al.*, **133**, 324
- Homeodomain  
vnd/NK-2, -DNA complex, weak labile  $^1\text{H}$  resonances of side chains, enhancement:  $^{15}\text{N}$ -edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87
- Homogeneous master equation  
for heteronuclear two-spin system, on basis of Cartesian product operators, Allard, Helgstrand, and Härd, **134**, 7
- Homonuclear correlation spectra  
2D, in MAS NMR, phase-sensitive detection, Boender and Vega, **133**, 281
- Homotopy  
computer simulation of MR spectra employing, Gates *et al.*, **135**, 104
- $^1\text{H}$ - $^{31}\text{P}$  couplings  
three-bond, in nucleic acids and protein-nucleic acid complexes, determination by quantitative  $J$  correlation spectroscopy, Clore *et al.*, **134**, 164
- HSQC  
 $^{13}\text{C}$ -NOESY-HSQC with split carbon evolution for increased resolution with uniformly labeled proteins, Baur, Gemmecker, and Kessler, **132**, 191  
exchange-edited gradient HCP-HSQC, sensitivity-enhanced detection of fast exchanging protons, Zangger and Armitage, **135**, 70  
 $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for spin-state selective editing of IS cross peaks, Andersson, Annala, and Otting, **133**, 364  
methyl group selection based on, via gradients in multidimensional NMR of proteins, Diercks, Schwaiger, and Kessler, **130**, 335  
TROSY experiment, sensitivity enhancement, Czisch and Boelens, **134**, 158
- Human serum albumin  
interaction of  $\beta$ -carboline derivatives with,  $^1\text{H}$  NMR studies, Veglia *et al.*, **130**, 281
- Hydration  
macromolecular, solvent magnetization artifacts in high-field NMR studies, Sobol *et al.*, **130**, 262
- Hydrogen bond dynamics  
 $^1\text{H}$  and  $^2\text{H}$  NMR relaxation in hydrogen-bonded solids due to complex motion, Reynhardt and Latanowicz, **130**, 195
- Hydrogen bonding effects  
on nitrogen NMR shielding of isomeric tetrazoles, Witanowski *et al.*, **131**, 54
- Hydrolysis  
effect on NMR shieldings of  $\text{Be}^{+2}$  and  $\text{Al}^{+3}$  species in aqueous solution, Tossell, **135**, 203
- 4-Hydroxy-TEMPO (*see* 4-Hydroxy-2,2,6,6-tetramethyl-1-piperidinyloxy)
- 4-Hydroxy-2,2,6,6-tetramethyl-1-piperidinyloxy  
 $\pi$  and  $\sigma$  transitions, DNP study at 1.53 mT, Salman *et al.*, **134**, 1  
saturation-transfer ESR spectra in glycerol, van den Dries, de Jager, and Hemminga, **131**, 241
- Hyperfine shifts  
in solution, high magnetic field consequences, Bertini, Felli, and Luchinat, **134**, 360
- Hyperfine structure  
proton, unresolved, contribution to Gaussian line broadening of proxyl spin probe EPR spectrum, Bales, Peric, and Lamy-Freund, **132**, 279
- Hyperfine tensor  
and quadrupole tensor,  $^{14}\text{N}$ , determination in oxovanadium complex by simultaneous fitting of multifrequency ESEEM powder spectra, Reijerse, Tyryshkin, and Dikanov, **131**, 295
- Hyperpolarized gases  
low-frequency NMR polarimeter for, Saam and Conradi, **134**, 67
- Hyperpolarized noble gases  
MRI of lung with, signal dynamics, Möller *et al.*, **135**, 133
- HYSOCORE  
matched three-pulse, sensitivity enhancement by, Jeschke, Rakhmatullin, and Schweiger, **131**, 261
- I
- Image analysis  
multivariate (*see* Multivariate image analysis)
- Image contrast  
 $^1\text{H}$  NMR, in solids, cross polarization for, Nakai *et al.*, **134**, 44
- Image reconstruction  
and display system, real-time, for MRI, using high-speed personal computer, Haishi and Kose, **134**, 138
- Imaging  
cyclic  $J$  cross-polarization, analysis of carbohydrate metabolism and transport in castor bean seedlings by, Heidenreich *et al.*, **132**, 109  
excitation of complicated shapes in three dimensions, Serša and Macura, **135**, 466  
Fourier spectral spatial EPR, implementation, Placidi *et al.*, **134**, 280  
 $^7\text{Li}$  NMR, Li visibility in rat brain and muscle *in vivo* by, Komoroski, Pearce, and Newton, **133**, 98  
oximetric, single-electron contrast agents for, EPR and DNP properties, Ardenkjær-Larsen *et al.*, **133**, 1  
rotating-frame NQR, mapping molecular orientation in solids by, Casanova, Robert, and Pusiol, **133**, 129  
stray field, by magnetic field sweep, Mallett, Halse, and Strange, **132**, 172  
using second-order quadrupole broadened resonances, Swaminathan and Suits, **132**, 274

- Imidazoles  
 coordinated, in uniformly  $^{15}\text{N}$ -labeled cytochrome  $c_3$ , imido protons, paramagnetic inversion of sign of interference contribution to  $T_2$ , Ohmura *et al.*, **131**, 367
- Imido group  
 protons, of coordinated imidazoles in uniformly  $^{15}\text{N}$ -labeled cytochrome  $c_3$ , interference contribution to  $T_2$ , paramagnetic inversion of sign of, Ohmura *et al.*, **131**, 367
- Impurity relaxation  
 nuclear spin–lattice relaxation, suppression, and charge exchange of chromium ions, in  $\gamma$ -irradiated ruby crystal, Chandoul *et al.*, **135**, 113
- INADEQUATE  
 2D experiment revisited: rapid-pulsing artifact-free double-quantum-filtered homonuclear spectroscopy, Bourdonneau and Ancian, **132**, 316  
 sensitivity improvement, Mattiello and Freeman, **135**, 514
- India ink  
 for use in EPR oximetry, separation and enrichment of active component, Liu *et al.*, **133**, 291
- INEPT  
 modified, *in vivo* 3D localized  $^{13}\text{C}$  spectroscopy, Watanabe *et al.*, **134**, 214  
 quadrupolar nuclei in solids, Kao and Grey, **133**, 313
- INEPT–INADEQUATE  
 $^1\text{H}$ – $^{13}\text{C}$ – $^{13}\text{C}$ -edited  $^1\text{H}$  NMR experiment for resonance assignment in active site of heme proteins, Qiu, Rivera, and Stark, **130**, 76
- Inhomogeneous fields  
 robust MR imaging method: SPARE, McIntyre, Hennel, and Morris, **130**, 58
- In situ* drying  
 for preparing NMR samples with extremely low water content, Goetz, **135**, 14
- In situ* NMR  
 analysis of fluids in sedimentary rock, de Swiet *et al.*, **133**, 385
- Integer spin systems  
 application of high-frequency EPR, unusual behavior of double-quantum line, van Dam *et al.*, **130**, 140
- Integration  
 by direct summation, influence of noise on peak integrals, Bourg and Nuzillard, **134**, 184
- Interference  
 quadrupolar–CSA, differential line broadening in presence of, Elbayed *et al.*, **130**, 209  
 in  $T_2$  of imido protons of coordinated imidazoles in uniformly  $^{15}\text{N}$ -labeled cytochrome  $c_3$ , paramagnetic inversion of sign of, Ohmura *et al.*, **131**, 367
- Interference effects  
 between dipolar coupling and chemical-shift anisotropy in spin relaxation of  $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, extracting, Ghose, Huang, and Prestegard, **135**, 487
- Interleaving  
 in image-based reduction of artifacts in multishot echo-planar imaging, Hennel, **134**, 206
- Internet technology  
 in magnetic resonance: common gateway interface program for World Wide Web NMR spectrometer, Buszko, Buszko, and Wang, **131**, 362
- Internuclear angles  
 DNA, determination using paramagnetic-assisted magnetic alignment, Berger *et al.*, **135**, 256
- Internuclear distance  
 NMR-derived, analysis of error propagation from, into molecular structure of *cyclo*-Pro-Gly, Džakula *et al.*, **135**, 454
- Interspin distances  
 iron(III)–nitroxyl, high-spin, determination in spin-labeled porphyrins by time-domain EPR, Rakowsky *et al.*, **131**, 97
- Inversion  
 fast broadband, by adiabatic pulses, Hwang, Zijl, and Garwood, **133**, 200  
 REDOR, d’Espinose de la Caillerie and Fretigny, **133**, 273  
 uniform-penalty, of multiexponential decay data, Borgia, Brown, and Fantazzini, **132**, 65
- Inversion recovery  
 sequential, with RIDE: simultaneous suppression of two solvent signals in  $^{17}\text{O}$  NMR, Schulte, **134**, 168
- Inversion–recovery imaging  
 signal intensity and  $T_1$ -dependent contrast, optimization with nonstandard flip angles, Bonny *et al.*, **130**, 51
- In vivo* spectroscopy  
 $^1\text{H}$  NMR spectra, resolution improvement with increased magnetic field strength, Gruetter *et al.*, **135**, 260  
 lactate editing at 1.5 T using PRESS excitation, with simultaneous detection of upfield singlets, applications to brain and head and neck tumors, Star-Lack *et al.*, **133**, 243  
 large samples, submicrosecond resonator and receiver system for pulsed MR, Alecci *et al.*, **132**, 162  
 observation of multiple spin echoes, Bifone, Payne, and Leach, **135**, 30  
 potential applicability of protein-mediated magnetic coupling between lactate and water protons, Swanson, **135**, 248  
 spectral editing technique for *in vitro* and *in vivo* detection of taurine, Hardy and Norwood, **133**, 70  
 2D localized  $^{13}\text{C}$  MRS using modified INEPT and DEPT, Watanabe *et al.*, **134**, 214
- Iron(III)  
 –nitroxyl distances, high-spin, determination in spin-labeled porphyrins by time-domain EPR, Rakowsky *et al.*, **131**, 97
- Iron ion  
 spin- $\frac{5}{2}$ , in transferrin oxalate, looping transition, simulation methods for, Gaffney and Silverstone, **134**, 57
- Isotope editing  
 $^1\text{H}$ – $^{13}\text{C}$ – $^{13}\text{C}$ -edited  $^1\text{H}$  NMR experiment for resonance assignment in active site of heme proteins, Qiu, Rivera, and Stark, **130**, 76
- Isotope effect  
 dynamic, and deuterium quadrupole coupling constant, determination in liquids, Hardy *et al.*, **134**, 300
- Isotope-filtered affinity NMR  
 $^{13}\text{C}/^{15}\text{N}$ -labeled stromelysin catalytic domain, Gonnella *et al.*, **131**, 336
- Isotropic solutions  
 biexponential relaxation, selective detection by mixed coherence filtering NMR, Yamada *et al.*, **135**, 99
- IS spin systems  
 IS, I $_2$ S, and I $_3$ S, effects of adiabatic pulses, analytical description, Zwahlen, Vincent, and Kay, **130**, 169
- Iterative applications  
 Fourier reconstruction as alternative to filtered back projection: Fourier spectral spatial EPR imaging, Placidi *et al.*, **134**, 280  
 lineshape fitting of MAS NMR spectra, for analysis of homonuclear  $J$  coupling in isolated spin pairs, Dusold, Milius, and Sebald, **135**, 500
- Iterative quadratic maximum likelihood  
 spectral parameter estimation, Zhu, Choy, and Sanctuary, **135**, 37
- J
- $J$ -coupling (*see* Spin–spin coupling)
- Jump and return  
 impact of offset-dependent saturation, Hsu and Gregory, **131**, 46
- Jump and return echo  
 impact of offset-dependent saturation, Hsu and Gregory, **131**, 46
- Jumps over a barrier  
 and incoherent tunneling:  $^1\text{H}$  and  $^2\text{H}$  NMR relaxation in hydrogen-bonded solids due to complex motion, Reynhardt and Latanowicz, **130**, 195

## K

- $K_3CrO_8$   
1–370 GHz EPR linewidths: test of Anderson–Weiss model, Cage *et al.*, **135**, 178
- Kidney  
detection of proton chemical exchange between metabolites and water, Guivel-Scharen *et al.*, **133**, 36
- k*-space trajectory  
deviations in MRI, simple correction method for, Duyn *et al.*, **132**, 150  
excitation of complicated shapes in three dimensions, Serša and Macura, **135**, 466

## L

- Laboratory-frame relaxation  
in estimation of chemical-shift anisotropy–dipolar coupling cross-correlation rates, Ghose and Prestegard, **134**, 308
- Lactate  
and water, protons, protein-mediated magnetic coupling, Swanson, **135**, 248
- Lactate editing  
*in vivo*, with simultaneous detection of upfield singlets at 1.5 T using PRESS excitation, applications to brain and head and neck tumors, Star-Lack *et al.*, **133**, 243
- Lanthanide complex  
paramagnetic, NMR temperature measurement, Zuo *et al.*, **133**, 53
- Large signals  
solvent, analog filtering for improved dynamic range in high-resolution NMR, Redfield and Kunz, **130**, 111
- Laser-polarized  $^{129}\text{Xe}$   
surface NMR under MAS conditions, Brunner *et al.*, **130**, 145
- Least-squares analysis  
global, based on prior knowledge, component separation in NMR imaging and multidimensional spectroscopy through, Stilbs, **135**, 236
- Leg  
human calf, heteronuclear cross polarization for enhanced sensitivity of *in vivo*  $^{13}\text{C}$  MR spectroscopy on clinical 1.5 T MR system, van den Bergh, van den Boogert, and Heerschap, **135**, 93
- $^7\text{Li}$   
magnetic relaxation dispersion: complex formation with nitroxides in aqueous phase, Dinesen and Bryant, **132**, 19  
NMR analysis of lithium visibility in rat brain and muscle *in vivo*, Komoroski, Pearce, and Newton, **133**, 98
- Lift-off effect  
surface and gradiometer coils near conducting body, Suits, Garroway, and Miller, **135**, 373
- Linear prediction  
spectral parameter estimation by, comparison with iterative quadratic maximum likelihood and total least squares methods, Zhu, Choy, and Sanctuary, **135**, 37
- Line broadening  
differential, in presence of quadrupolar–CSA interference, Elbayed *et al.*, **130**, 209  
Gaussian, of proxyl spin probe EPR spectrum, contributions from magnetic-field modulation and unresolved proton hyperfine structure, Bales, Peric, and Lamy-Freund, **132**, 279
- Line list  
NMR spectra, generation by 1D filter diagonalization method, Hu *et al.*, **134**, 76
- LINEPT  
*in vivo* 3D localized  $^{13}\text{C}$  spectroscopy, Watanabe *et al.*, **134**, 214
- Lineshape  
implications of transition-band signals in high-resolution NMR, Szántay, **135**, 334  
super-Lorentzian, contrast prediction using: magnetization transfer imaging of rat brain under non-steady-state conditions, Quesson *et al.*, **130**, 321
- Lineshape fitting  
iterative, of MAS NMR spectra, for analysis of homonuclear *J* coupling in isolated spin pairs, Dusold, Milius, and Sebald, **135**, 500
- Lipid  
singlets, simultaneous detection in *in vivo* lactate editing at 1.5 T using PRESS excitation, in head and neck tumors, Star-Lack *et al.*, **133**, 243
- Lipid bilayers  
cholestane spin labels, influence of high orientational order on shape of Hahn echo responses, van der Struijf *et al.*, **130**, 253  
rotational motion of proteins, use of phase memory time  $T_2$  for quantitative characterization, van der Struijf and Levine, **130**, 244  
uniformly aligned, dimyristoylphosphatidylcholine, magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305
- Lipid membrane  
magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305
- Lipid transfer protein  
nonspecific, of wheat, NOESY: composite  $90^\circ$  pulse application, Sodano, Landon, and Ptak, **133**, 194
- Liquid crystals  
asymmetric alignment tensor of oriented macromolecules, determination in absence of structural information, Clore, Gronenborn, and Bax, **133**, 216  
broadband decoupling sequence for, Yu and Fung, **130**, 317  
cholestane spin labels in, influence of high orientational order on shape of Hahn echo responses, van der Struijf *et al.*, **130**, 253  
 $^{19}\text{F}$  NMR spectra, simplification by multiple-pulse COSY, Lesot *et al.*, **133**, 166  
molecules oriented in, natural abundance  $^2\text{H}$  NMR spectra, Khetrapal *et al.*, **135**, 265  
nematic phospholipid bicellar solutions,  $^2\text{H}$  NMR, Struppe and Vold, **135**, 541  
order-fluctuation modes, formalism for evaluation from field-cycling NMR-relaxometry data, Grinberg *et al.*, **135**, 54  
symmetry and phase-selected NMR spectra, Carravetta *et al.*, **135**, 298
- Liquids  
dynamic isotope effect and deuterium quadrupole coupling constant, Hardy *et al.*, **134**, 300  
experimental determination of four-time stimulated echoes, Hinze *et al.*, **131**, 218  
highly viscous, rotational motion of solute molecule,  $^{13}\text{C}$  NMR, Lienin, Brüschweiler, and Ernst, **131**, 184  
supercooled, ultraslow reorientations,  $^2\text{H}$  NMR time domain analysis, Geil, Fajara, and Silleseu, **130**, 18
- Liquid state  
adiabatic mixing in, Kupče *et al.*, **135**, 361
- Lithium  
visibility in rat brain and muscle *in vivo* by  $^7\text{Li}$  NMR imaging, Komoroski, Pearce, and Newton, **133**, 98
- Lithium phthalocyanine  
application of RF 220-MHz FT EPR spectrometer, Alecci *et al.*, **130**, 272
- Liver  
isolated perfused, intravascular and intracellular hepatic relaxivities of superparamagnetic particles, Colet *et al.*, **134**, 199
- $\text{Ln}(\text{C}_2\text{D}_5\text{SO}_4)_3 \cdot 8\text{H}_2\text{O}$  ( $\text{Ln} = \text{Pr, Yb}$ )  
paramagnetic, solid-state NMR, effect of bulk magnetic susceptibility, Kubo, Spaniol, and Terao, **133**, 330
- Localization  
cardiac  $^{31}\text{P}$  MRS in humans, Löffler *et al.*, **134**, 287  
*in vivo* 3D localized  $^{13}\text{C}$  MRS using modified INEPT and DEPT, Watanabe *et al.*, **134**, 214

## LODEPT

*in vivo* 3D localized  $^{13}\text{C}$  spectroscopy, Watanabe *et al.*, **134**, 214

## Logic gates

quantum, and NMR pulse sequences, Jones, Hansen, and Mosca, **135**, 353  
Longitudinal relaxation (*see* Spin–lattice relaxation)

## Loop–gap resonator

rectangular, for EPR studies of aqueous samples, Piasecki, Froncisz, and Hubbell, **134**, 36

## Looping transitions

simulation methods for, Gaffney and Silverstone, **134**, 57

## Lung

guinea pig, MRI with hyperpolarized noble gases, signal dynamics, Möller *et al.*, **135**, 133

## Lysozyme

diffusion in aqueous solution, practical aspects of measurement, Tillett, Lian, and Norwood, **133**, 379

## M

## Magic angle-oriented sample spinning

uniformly aligned biomembrane samples, Glaubitz and Watts, **130**, 305

## Magic-angle spinning

CP-MAS (*see* Cross polarization–magic-angle spinning)

fast,  $^{13}\text{C}$ – $^1\text{H}$  spin pair under, application of rotor-synchronized amplitude-modulated cross polarization, Raya and Hirschinger, **133**, 341

high-resolution, IHMRS evaluation of breast ductal carcinomas, Cheng *et al.*, **135**, 194

high-speed, adamantane:  $^{13}\text{C}$  MAS NMR spectra, Ernst, Verhoeven, and Meier, **130**, 176

*in situ* NMR analysis of fluids in sedimentary rock, de Swiet *et al.*, **133**, 385

iterative spectral lineshape fitting, for analysis of homonuclear  $J$  coupling in isolated spin pairs, Dusold, Milius, and Sebald, **135**, 500

magic angle-oriented sample spinning approach for biomembrane studies, Glaubitz and Watts, **130**, 305

multidimensional solid-state NMR correlation spectra, intrinsic asymmetry, Caldarelli and Emsley, **129**, 233

## multiple-quantum

effect of RF power and spinning speed, Amoureux *et al.*, **131**, 170

half-integer quadrupolar spins, composite pulse excitation schemes for, Marinelli, Medek, and Frydman, **132**, 88

resolution enhancement, Schnell *et al.*, **133**, 61

$^{15}\text{N}$  chemical shift and  $^{13}\text{C}$ – $^1\text{H}$  dipolar tensor correlation, peptide torsion angle determination by, Hong *et al.*, **135**, 169

1D, and 2D exchange NMR, direct determination of motional correlational times by, Favre, Schaefer, and Chmelka, **134**, 261

proton, at high magnetic fields and high spinning frequencies: spectral simulations using Floquet theory, Ray *et al.*, **135**, 418

QCPMG–MAS NMR of half-integer quadrupolar nuclei, Larsen *et al.*, **131**, 144

REDOR signal and inversion analysis, d’Espinose de la Caillerie and Fréty, **133**, 273

simulation, time propagation technique for, application to quadrupolar nuclei, Charpentier, Fermon, and Virlet, **132**, 181

six-pulse PASS MAS-NMR technique solved by phasor method, Effey, Butcher, and Cappelletti, **132**, 266

S-MAS, polarization transfer under, comparison with RF-driven conditions, Robyr and Gan, **131**, 254

surface NMR using laser-polarized  $^{129}\text{Xe}$ , Brunner *et al.*, **130**, 145

2D, of dipolar solids, pulsed-field gradient selection, Fritzhanns *et al.*, **134**, 355

2D homonuclear correlation spectra, phase-sensitive detection, Boender and Vega, **133**, 281

## Magic-sandwich echo

in complete dipolar decoupling of  $^{13}\text{C}$  and 2D double-quantum solid-state NMR for determining polymer conformations, Schmidt-Rohr, **131**, 209

## Magnetic alignment

paramagnetic-assisted, determination of DNA internuclear angles, Beger *et al.*, **135**, 256

## Magnetic anisotropy

in frozen nickel ferrite ionic magnetic fluid, MR analysis, Saenger *et al.*, **134**, 180

## Magnetic coupling

protein-mediated, between lactate and water protons, Swanson, **135**, 248

## Magnetic exchange coupling

nuclear and electron relaxation in magnetic exchange coupled dimers, Bertini *et al.*, **130**, 33

Magnetic field (*see also* Gradients; Pulsed-field gradients)

high, effects on NMR hyperfine shifts in solution, Bertini, Felli, and Luchinat, **134**, 360

increased field strength, resolution improvements in *in vivo*  $^1\text{H}$  NMR spectra by, Gruetter *et al.*, **135**, 260

inhomogeneous, robust MR imaging method: SPARE, McIntyre, Hennel, and Morris, **130**, 58

Magnetic-field gradient (*see* Gradients; Pulsed-field gradients)

## Magnetic-field modulation

role in Gaussian line broadening of proxyl spin probe EPR spectrum, Bales, Peric, and Lamy-Freund, **132**, 279

## Magnetic fluid

frozen ionic,  $\text{NiFe}_2\text{O}_4$ , MR anisotropy analysis, Saenger *et al.*, **134**, 180

## Magnetic relaxation dispersion

$^7\text{Li}$ : complex formation with nitroxides in aqueous phase, Dinesen and Bryant, **132**, 19

## Magnetic resonance imaging

birdcage coil tuned by RF shielding for application at 9.4 T, Dardzinski *et al.*, **131**, 32

component separation through global least-squares analysis based on prior knowledge, Stilbs, **135**, 236

detection of proton chemical exchange between metabolites and water in biological tissues, Guivel-Scharen *et al.*, **133**, 36

field mapping with amplitude images at low field, use of phase reference for, Tomasi *et al.*, **131**, 310

fMRI, study of activation of visuomotor systems during visually guided movements, Ellermann *et al.*, **131**, 272

$^1\text{H}$ , quantitative, and MRS microscopy, of individual V79 lung tumor spheroids, Minard, Guo, and Wind, **133**, 368

high-resolution 3D, for Sandimmune Neoral effects on collagen arthritis in DA rat, Beckmann *et al.*, **131**, 8

improved resolution and signal-to-noise ratio by enhanced signal digitization, Elliott *et al.*, **130**, 300

in inhomogeneous fields, single-point acquisition method SPARE for, McIntyre, Hennel, and Morris, **130**, 58

$k$ -space trajectory deviations, simple correction method for, Duyn *et al.*, **132**, 150

lung, with hyperpolarized noble gases, signal dynamics, Möller *et al.*, **135**, 133

multiecho, with suboptimal spoiler gradients, Does and Snyder, **131**, 25

multivariate image analysis with direct exponential curve resolution algorithm (DECRA)

algorithm and model study, Windig, Hornak, and Antalek, **132**, 298

human brain images, Antalek, Hornak, and Windig, **132**, 307

oximetry with, single-electron contrast agents for, EPR and DNP properties, Ardenkjær-Larsen *et al.*, **133**, 1

proton–electron double-resonance imaging: mobility and free radical concentration effects, de Sousa *et al.*, **135**, 118

real-time image reconstruction and display system using high-speed personal computer, Haishi and Kose, **134**, 138

$R_2'$  measurement in presence of multiple spectral components using reference spectrum deconvolution, Wehrli *et al.*, **131**, 61

sequential dynamic susceptibility contrast,  $T_1$  effects, Levin *et al.*, **130**, 292



- spin-echo and inversion-recovery MR, signal intensity and  $T_1$ -dependent contrast, optimization with nonstandard flip angles, Bonny *et al.*, **130**, 51  
with  $^{129}\text{Xe}$ , in pore-structure determination of silica aerogels, Gregory, Gerald, and Botto, **131**, 327
- Magnetic resonance imaging contrast agents  
superparamagnetic particles, intravascular and intracellular hepatic relaxivities, Colet *et al.*, **134**, 199
- Magnetic susceptibility  
differences, effective gradients in porous media due to, Hürlimann, **131**, 232  
mismatches, solenoidal microcoil design for assessment, Subramanian, Lam, and Webb, **133**, 227  
 $T_1$  effects in sequential dynamic susceptibility contrast experiments, Levin *et al.*, **130**, 292
- Magnetization  
water, undisturbed, composite  $90^\circ$  pulse sequence providing, Sodano, Landon, and Ptak, **133**, 194
- Magnetization transfer  
model, general solution, Roell, Dreher, and Leibfritz, **132**, 96  
in polymers, role of specific side groups and pH, Gochberg *et al.*, **131**, 191  
protein-mediated magnetic coupling between lactate and water protons, Swanson, **135**, 248
- Magnetization transfer imaging  
rat brain under non-steady-state conditions, Quesson *et al.*, **130**, 321
- D-Mannopyranose  
 $^{17}\text{O}$  NMR, simultaneous suppression of two solvent signals: sequential inversion recovery with RIDE, Schulte, **134**, 168
- Mannose transporter  
IIAMan domain, multidimensional NMR, HSQC-based methyl group selection via gradients, Diercks, Schwaiger, and Kessler, **130**, 335
- Mapping  
quantitative, eddy current fields, localized eddy current compensation, Terpstra, Andersen, and Gruetter, **131**, 139
- Mass-limited samples  
RF microcoil design for NMR, Subramanian, Lam, and Webb, **133**, 227
- Matched pulses  
microwave, sensitivity enhancement by, in 1D and 2D ESEEM spectroscopy, Jeschke, Rakhmatullin, and Schweiger, **131**, 261
- Matrix line  
in pulsed ENDOR, Astashkin and Kawamori, **135**, 406
- Maximum entropy reconstruction  
multidimensional, distributed parallel processing for, Li, Stern, and Hoch, **134**, 161
- Maximum likelihood  
analysis of error propagation from NMR-derived internuclear distances into molecular structure of *cyclo*-Pro-Gly, Džakula *et al.*, **135**, 454
- Meetings and Announcements, **130**, 153, 352; **131**, 176, 379; **132**, 179, 331; **133**, 241, 395; **134**, 195, 370; **135**, 271, 551
- Membrane protein  
magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305  
 $^{15}\text{N}$  chemical-shift anisotropy, cation binding-induced changes, Tian and Cross, **135**, 535
- Membranes  
oriented, magic angle-oriented sample spinning for, Glaubitz and Watts, **130**, 305
- Metabolism  
carbohydrate in plants, cyclic  $J$  cross-polarization imaging and spectroscopy for analysis, Heidenreich *et al.*, **132**, 109
- Metabolite imaging of lines in a spectrum (MILS)  
inverse detection of protons through, alternative scheme for  $^{13}\text{C}$  chemical-shift imaging via, Lee *et al.*, **130**, 102
- Metabolites  
human brain, automated quantification by artificial neural network analysis from *in vivo* single-voxel  $^1\text{H}$  NMR spectra, Kaartinen *et al.*, **134**, 176  
proton chemical exchange with water, detection in biological tissues, Guivel-Scharen *et al.*, **133**, 36  
V79 lung tumor: quantitative  $^1\text{H}$  MRI and MRS microscopy, Minard, Guo, and Wind, **133**, 368
- Metallic glass  
ultraslow motion in, detection using alignment echo of spin- $\frac{3}{2}$   $^9\text{Be}$  nuclei, Tang and Wu, **133**, 155
- Methylene  
carbons, in linear polyethylene,  $^{13}\text{C}$  CPMAS linewidths under CW proton decoupling, VanderHart and Campbell, **134**, 88  
specific experiments, for scalar spin-spin coupling constants between protons attached to  $^{13}\text{C}$ , Carlomagno *et al.*, **135**, 216
- Methyl  $\alpha$ -D-glucopyranoside  
cryoprotection, effects on structure and activity of p21 *ras*: implications for ESEEM, Halkides, Farrar, and Singel, **134**, 142
- Methyl groups  
HSQC-based selection via gradients in multidimensional NMR of proteins, Diercks, Schwaiger, and Kessler, **130**, 335
- 1-Methyl-1,2,3,4-tetrazole  
nitrogen NMR shielding, solvent polarity and hydrogen bonding effects, Witanowski *et al.*, **131**, 54
- 1-Methyl-1,2,3,5-tetrazole  
nitrogen NMR shielding, solvent polarity and hydrogen bonding effects, Witanowski *et al.*, **131**, 54
- Metropolis Monte Carlo algorithm  
implementation of Bayesian time-domain parameter estimation, application to coupling constant estimation from antiphase multiplets, Andrec and Prestegard, **130**, 217
- Micelles  
water-surfactant contact studied by  $^{19}\text{F}$ - $^1\text{H}$  HOESY, Raulet *et al.*, **133**, 324
- Microcoils  
RF, design for NMR of mass-limited samples, Subramanian, Lam, and Webb, **133**, 227
- Microscopy  
NMR (*see* NMR microscopy)
- Microwaves  
matched pulses, sensitivity enhancement by, in 1D and 2D ESEEM spectroscopy, Jeschke, Rakhmatullin, and Schweiger, **131**, 261
- MILS  
inverse detection of protons through, alternative scheme for  $^{13}\text{C}$  chemical-shift imaging via, Lee *et al.*, **130**, 102
- Mims ENDOR  
matrix line, Astashkin and Kawamori, **135**, 406
- Mixed coherence filtering NMR  
selective detection of biexponential relaxation in isotropic solutions, Yamada *et al.*, **135**, 99
- Mixing  
adiabatic, in liquid state, Kupče *et al.*, **135**, 361
- Mixture analysis  
high-resolution, complex mixtures, 3D DOSY-HMQC experiment for, Barjat, Morris, and Swanson, **131**, 131  
model-free, by NMR using blind source separation, Nuzillard, Bourg, and Nuzillard, **133**, 358
- $^{55}\text{Mn}$   
quadrupole-dipole effects in solid-state CP-MAS  $^{119}\text{Tm}$  NMR spectra of *para*-substituted triaryltin(pentacarbonyl)manganese(I) complexes, Christendat *et al.*, **131**, 1
- Mobile universal surface explorer (MOUSE)  
surface NMR analysis of polymer materials, Guthausen *et al.*, **130**, 1

- Model-free analysis  
 in DNA dynamics study via  $^{13}\text{C}$  relaxation, Kojima *et al.*, **135**, 310  
 stretched relaxation dispersions, Halle, Jóhannesson, and Venu, **135**, 1
- Mode relaxation rates  
 formalism for evaluation of order-fluctuation modes in liquid crystals from field-cycling NMR-relaxometry data, Grinberg *et al.*, **135**, 54
- Molecular alignment  
 weakly oriented macromolecules, measurement of dipolar couplings for methylene and methyl sites, Ottiger *et al.*, **134**, 365
- Molecular diffusion (*see* Diffusion)
- Molecular dynamics  
 pyridinium tetrafluoroborate: radicals as probes of ferroelectric phase transitions in EPR studies, Krzyminiewski, Bielewicz-Mordalska, and Kowalczyk, **135**, 76  
 vacancy diffusion in polycrystalline benzene, analysis with 2D deuteron exchange NMR and stimulated echoes, Isfort, Geil, and Fujara, **130**, 45
- Molecular mobility  
 effects in proton–electron double-resonance imaging, de Sousa *et al.*, **135**, 118  
 extremely slow, in glassy materials, sensitivity of saturation-transfer ESR extended to, van den Dries, de Jager, and Hemminga, **131**, 241
- Molecular motion  
 effect on spin label relaxation time determination by progressive saturation EPR, Livshits, Páli, and Marsh, **133**, 79
- Molecular reorientation  
 ultraslow, in supercooled liquids,  $^2\text{H}$  NMR time domain analysis, Geil, Fujara, and Sillescu, **130**, 18
- Monte Carlo algorithm  
 implementation of Bayesian time-domain parameter estimation, application to coupling constant estimation from antiphase multiplets, Andrec and Prestegard, **130**, 217
- Motion  
 rotational, solute molecule in highly viscous liquid,  $^{13}\text{C}$  NMR, Lienin, Brüschweiler, and Ernst, **131**, 184
- Motional correlational times  
 direct determination by 1D MAS and 2D exchange NMR, Favre, Schaefer, and Chmelka, **134**, 261
- Motional dynamics  
 protein side-chain, analysis from  $^{13}\text{C}$  NMR relaxation data, Daragan and Mayo, **130**, 329
- Motor task  
 visually guided, activation of visuomotor systems during, functional MRI, Ellermann *et al.*, **131**, 272
- MOUSE (Mobile universal surface explorer)  
 surface NMR analysis of polymer materials, Guthausen *et al.*, **130**, 1
- MP-ROESY  
 enhancement of cross-relaxation peaks, Hwang and Shaka, **135**, 280
- MQMAS NMR (*see* Magic-angle spinning, multiple-quantum)
- MSHOT-3  
 based CRAMPS,  $^1\text{H}$  chemical-shielding anisotropies from polycrystalline powders, Hohwy *et al.*, **133**, 374
- Multi-compartment systems  
 self-diffusion and relaxation time NMR, modeling, Novikov, van Dusschoten, and Van As, **135**, 522
- Multidimensional NMR  
 component separation through global least-squares analysis based on prior knowledge, Stilbs, **135**, 236  
 proteins, HSQC-based methyl group selection via gradients, Diercks, Schwaiger, and Kessler, **130**, 335
- Multiecho imaging  
 with suboptimal spoiler gradients, Does and Snyder, **131**, 25
- Multiexponential decay  
 data, uniform-penalty inversion, Borgia, Brown, and Fantazzini, **132**, 65  
 selective detection of biexponential relaxation in isotropic solutions by mixed coherence filtering NMR, Yamada *et al.*, **135**, 99
- Multilayer perceptron  
 determining rotational correlation times from slow motional ESR spectra with, comparison with radial basis network, Martinez and Millhauser, **134**, 124
- Multilinear interactions  
 in high-resolution spectra, Corio, **134**, 131
- Multiple-pulse mixing sequences  
 selectively enhancing chemical exchange or cross-relaxation peaks in high-resolution NMR spectra, Hwang and Shaka, **135**, 280
- Multiple-pulse ROESY (MP-ROESY)  
 enhancement of cross-relaxation peaks, Hwang and Shaka, **135**, 280
- Multiple-pulse sequence  
 COSY, simplification of  $^{19}\text{F}$  NMR spectra of liquid crystalline samples, Lesot *et al.*, **133**, 166
- Multiple-quantum-filtered NMR  
 $^{23}\text{Na}$  spectra of erythrocyte suspensions, contribution of extracellular sodium, Knubovets, Shinar, and Navon, **131**, 92
- Multiple quantum filtering  
 simplification of NMR spectra of liquid crystalline samples, Carravetta *et al.*, **135**, 298
- Multiple-quantum MAS NMR spectroscopy  
 half-integer quadrupolar spins, composite pulse excitation schemes for, Marinelli, Medek, and Frydman, **132**, 88  
 resolution enhancement, Schnell *et al.*, **133**, 61  
 spin- $\frac{3}{2}$  quadrupolar spin systems, using shaped pulses, Ding and McDowell, **135**, 61  
 2D, distributions of quadrupolar and isotropic chemical shift interactions, Bodart, **133**, 207
- Multiple spin echoes  
*in vivo* observation, Bifone, Payne, and Leach, **135**, 30
- Multiplets  
 antiphase, coupling constant estimation from, application of Metropolis Monte Carlo implementation of Bayesian time-domain parameter estimation, Andrec and Prestegard, **130**, 217
- Multivariate curve resolution  
 analysis of DOSY and GPC-NMR experiments on polymers by, Van Gorkom and Hancewicz, **130**, 125
- Multivariate image analysis  
 MR images, with direct exponential curve resolution algorithm (DECRA) algorithm and model study, Windig, Hornak, and Antalek, **132**, 298  
 human brain images, Antalek, Hornak, and Windig, **132**, 307
- Muscle  
 Li visibility *in vivo* by  $^7\text{Li}$  NMR imaging, Komoroski, Pearce, and Newton (rat), **133**, 98
- Myocardial hypertrophy  
 phosphomonoester signal detection in proton-decoupled  $^{31}\text{P}$  NMR spectra, Jung *et al.*, **133**, 232
- Myoglobin  
 NMR spectra acquisition with PASE: paramagnetic signals enhancement, Bondon and Mouro, **134**, 154

## N

 $^{14}\text{N}$ 

- $^{13}\text{C}$ - $^{14}\text{N}$  REAPDOR NMR, optimization, Ba *et al.*, **133**, 104  
 disordered powder sample of dianisyl-nitroxide radical, high-frequency/high-field ESEEM study of nitrogen hyperfine and quadrupole interactions, Bloeb, Möbius, and Prisner, **134**, 30  
 $\text{NH}_4\text{ClO}_4$ , solid echo signal after pulse sequence  $(\theta_1)_x - \tau - (\theta_2)_y - t$ , Punkkinen *et al.*, **130**, 287  
 nitrogen quadrupole and hyperfine tensors, determination in oxovanadium complex by simultaneous fitting of multifrequency ESEEM powder spectra, Reijerse, Tyryshkin, and Dikanov, **131**, 295

- unshielded NQR measurements using noise-immune coil, Suits, Garroway, and Miller, **131**, 154
- <sup>15</sup>N  
 bacteriochlorophyll *a* labeled with, low-frequency nuclear coherence detection with simple two-pulse detection scheme in pulsed EPR, Dzuba, Borovykh, and Hoff, **133**, 286
- and <sup>13</sup>C  
 alanine and urea multiply labeled with, counting spins with SEDOR, Cull *et al.*, **133**, 352  
 nucleotides labeled with, in DNA duplex dynamics study via <sup>13</sup>C relaxation, Kojima *et al.*, **135**, 310  
 ubiquitin labeled with, measurement of dipolar couplings for methylene and methyl sites and use in structure determination, Ottiger *et al.*, **134**, 365
- and <sup>13</sup>C, proteins labeled with  
 amide deuterium isotope effects on C<sup>β</sup> chemical shifts, multidimensional editing for, Meissner and Sørensen, **135**, 547  
 double-labeled, Bloch–Siegert shift-compensated and cyclic sideband-eliminated double-adiabatic homonuclear decoupling for, Zhang and Gorenstein, **132**, 81  
 in H<sub>2</sub>O solution, H(C)CH-COSY and (H)CCH-COSY experiments, Gehring and Ekiel, **135**, 185  
 spin relaxation, measurement of cross correlation between dipolar coupling and chemical-shift anisotropy, Ghose, Huang, and Prestegard, **135**, 487
- chemical-shift anisotropy  
 ampicillin and penicillin-V by CP/MAS NMR, Antzutkin, Lee, and Levitt, **135**, 144  
 in membrane-bound polypeptide, cation binding-induced changes, Tian and Cross, **135**, 535
- chemical shift tensor, and <sup>13</sup>C<sup>α</sup>–<sup>1</sup>H<sup>α</sup> dipolar tensor, correlation in solid-state MAS NMR, peptide torsion angle determination by, Hong *et al.*, **135**, 169
- cytochrome *c*<sub>3</sub> uniformly labeled with, interference contribution to *T*<sub>2</sub> of imido protons of coordinated imidazoles in, paramagnetic inversion of sign of, Ohmura *et al.*, **131**, 367
- deuterated glutathione *S*-transferase, correlation of backbone amide and aliphatic gamma resonances, McCallum, Hitchens, and Rule, **134**, 350
- edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87
- gyrase B, α/β-HSQC-α/β experiment for spin-state selective editing of IS cross peaks, Andersson, Annila, and Otting, **133**, 364
- heteronuclear X-filter <sup>1</sup>H PFG double-quantum experiment for proton resonance assignment of ligand bound to protein, Dalvit, Ramage, and Hommel, **131**, 148
- improved resolution from double constant-time evolution of 3D and 4D triple-resonance experiments, McCoy, **130**, 341
- <sup>15</sup>N–<sup>15</sup>N *J*-couplings in proteins, detection, Löhner and Rüterjans, **132**, 130
- <sup>15</sup>N relaxation  
 improved estimation of protein rotational correlation times, Yao, Hinds, and Norton, **131**, 347  
 large proteins, 3D NMR experiments: 44-kDa ectodomain of SIV gp41, Caffrey *et al.*, **135**, 368  
 rate in poorly resolved NMR spectra, 3D accordion spectroscopy measurement, Carr, Fearing, and Palmer, **132**, 25
- off-resonance rotating frame relaxation experiment for analysis of macromolecular dynamics using adiabatic rotations, Mulder *et al.*, **131**, 351
- peptides, 1D solid-state NMR method for nuclear spin interaction tensors associated with peptide bond, Lee and Ramamoorthy, **133**, 204
- proteins enriched in, NMR, detection of cross-relaxation between amide protons, Mutzenhardt and Bodenhausen, **132**, 159
- rod domain of F-actin cross-linking protein, *T*<sub>2</sub> errors, removal with adiabatic multiple spin-echo pulse sequence, Zweckstetter and Holak, **133**, 134
- solvent magnetization artifacts in high-field NMR studies of macromolecular hydration, Sobol *et al.*, **130**, 262
- triple-resonance HCN experiments, optimization for larger RNA oligonucleotides, Sklenář *et al.*, **130**, 119
- TROSY NMR, sensitivity enhancement, Czisch and Boelens, **134**, 158
- ubiquitin, dipole–CSA cross-correlated relaxation, quantitative analysis by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346
- Na<sup>+</sup>  
 binding, induced changes in <sup>15</sup>N CSA in membrane-bound polypeptide, Tian and Cross, **135**, 535
- <sup>23</sup>Na  
 dipolar interaction in solution, determination by triple-quantum relaxation time measurements, Eliav and Navon, **130**, 63  
 multiple-quantum-filtered NMR, erythrocyte suspensions, contribution of extracellular sodium, Knubovets, Shinar, and Navon, **131**, 92  
 Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub>, effect of RF power and spinning speed on MQMAS NMR, Amoureux *et al.*, **131**, 170  
 spin  $\frac{3}{2}$  nuclei of incompletely disordered systems, temporal analysis of NMR signals, Woessner and Bansal, **133**, 21
- NaClO<sub>3</sub>  
<sup>35</sup>Cl, single crystal 2D NQR spectroscopy, off-resonance effects, Ajithkumar *et al.*, **135**, 165
- Nanovolume NMR  
 RF microcoil design for, Subramanian, Lam, and Webb, **133**, 227
- Naphthazarin B  
 hydrogen-bonded, <sup>1</sup>H and <sup>2</sup>H NMR relaxation due to complex motion: classical jumps over barrier and incoherent tunneling, Reynhardt and Latanowicz, **130**, 195
- Natural abundance <sup>2</sup>H NMR spectra  
 molecules oriented in liquid crystals, Khetrpal *et al.*, **135**, 265
- NaX  
 (CH<sub>3</sub>)<sub>3</sub>P–AlCl<sub>3</sub> complex in, <sup>27</sup>Al → <sup>31</sup>P INEPT experiments, Kao and Grey, **133**, 313
- Neural networks  
 artificial, in automated quantification of human brain metabolites from *in vivo* single-voxel <sup>1</sup>H NMR spectra, Kaartinen *et al.*, **134**, 176  
 in rapid computation of rotational correlation times from slow motional ESR spectra, Martinez and Millhauser, **134**, 124
- NH<sub>4</sub>ClO<sub>4</sub>  
<sup>14</sup>N NMR echo after pulse sequence (θ<sub>1</sub>)<sub>x</sub>–τ–(θ<sub>2</sub>)<sub>y</sub>–t, Punkkinen *et al.*, **130**, 287
- Nickel ferrite  
 frozen ionic magnetic fluid, MR anisotropy analysis, Saenger *et al.*, **134**, 180
- Ni(II)  
 integer spin system in host Zn-tris(ethylenediamine) dinitrate, high-frequency EPR study: unusual behavior of double-quantum line, van Dam *et al.*, **130**, 140
- Nitrogen hyperfine coupling  
 and quadrupole interactions, on disordered powder sample, high-frequency/high-field ESEEM study, Bloef, Möbius, and Prisner, **134**, 30
- Nitrogen NMR shielding  
 isomeric tetrazoles, solvent polarity and hydrogen bonding effects, and *ab initio* calculation of nitrogen shielding of azole systems, Witanowski *et al.*, **131**, 54
- Nitroxide spin labels  
 3-carboxy proxyl, Gaussian line broadening of EPR spectrum, contributions from magnetic-field modulation and unresolved proton hyperfine structure, Bales, Peric, and Lamy-Freund, **132**, 279  
 complex formation with Li in aqueous phase: magnetic relaxation dispersion of <sup>7</sup>Li, Dinesen and Bryant, **132**, 19  
 conformational effect on Ala-rich helical peptides, NMR analysis, Bolin *et al.*, **131**, 248  
 π and σ transitions, DNP study at 1.53 mT, Salman *et al.*, **134**, 1

- proteins in lipid bilayers with, rotational motions, use of phase memory time  $T_2$  for quantitative characterization, van der Struijf and Levine, **130**, 244
- relaxation time determination by progressive saturation EPR, molecular motion and Zeeman modulation effects, Livshits, Páli, and Marsh, **133**, 79
- slow motional spectra, rapid computation of rotational correlation time from, neural network approach, Martinez and Millhauser, **134**, 124
- spin–lattice relaxation measurements using first-harmonic out-of-phase absorption EPR signals, Livshits, Páli, and Marsh, **134**, 113
- Nitroxyl  
–iron(III) distances, high-spin, determination in spin-labeled porphyrins by time-domain EPR, Rakowsky *et al.*, **131**, 97
- NMRD (Nuclear magnetic relaxation dispersion)  
model-free analysis of stretched dispersions, Halle, Jóhannesson, and Venu, **135**, 1
- NMR microscopy  
multilayer gradient coil design for, Bowtell and Robyr, **131**, 286  
and quantitative  $^1\text{H}$  MRI, of individual V79 lung tumor spheroids, Minard, Guo, and Wind, **133**, 368
- NMR polarimeter  
low-frequency, for hyperpolarized gases, Saam and Conradi, **134**, 67
- NMR probes  
 $^{13}\text{C}$  CP/MAS: chemical-shift imaging in rotating solids by RF field gradients, Malveau *et al.*, **134**, 171  
design: high-resolution high-pressure NMR probe for experiments at 500 MHz, Ballard *et al.*, **133**, 190  
double resonance, for close frequencies, Haase, Curro, and Slichter, **135**, 273  
high-frequency, double- and triple-resonance circuits for, Zhang *et al.*, **132**, 167  
high-pressure high-resolution, for experiments at 500 MHz, Ballard *et al.*, **133**, 190  
pulsed-gradient, for solid-state studies, Zhang and Cory, **132**, 144  
temperature imaging by  $^1\text{H}$  NMR and suppression of convection, Hedin and Furó, **131**, 126
- NMR samples  
*in situ* drying: preparation of samples with extremely low water content, Goetz, **135**, 14
- NMR second moment  
calculation, effective spatial averaging for, Goc, **132**, 78
- NMR spectra  
computer simulation employing homotopy, Gates *et al.*, **135**, 104  
macromolecular, automated peak picking and peak integration with use of AUTOPSY, Koradi *et al.*, **135**, 288  
precision of parameters in measuring, mathematical assessment, Golding and Golding, **135**, 380  
reference deconvolution, phase correction, and line listing by 1D filter diagonalization method, Hu *et al.*, **134**, 76
- NMR spectrometer  
using Earth's field, pulsed gradient spin echo NMR study of restricted diffusion in sea ice, Callaghan *et al.*, **133**, 148  
World Wide Web-operated, common gateway interface program for, Buszko, Buszko, and Wang, **131**, 362
- NMR thermometry  
high-precision  $^{13}\text{C}$  shift, for temperature range 100–300 K, Quast *et al.*, **134**, 236  
using paramagnetic lanthanide complex, Zuo *et al.*, **133**, 53
- Noble gases  
hyperpolarized, signal dynamics in MRI of lung with, Möller *et al.*, **135**, 133  
laser-polarized, time-dependent diffusion, PFG measurements, Mair *et al.*, **135**, 478
- NOE  
quiet, interresidue, for DNA structural studies, Beger *et al.*, **132**, 34  
spin polarization-induced, in surface NMR using laser-polarized  $^{129}\text{Xe}$  under MAS conditions, Brunner *et al.*, **130**, 145
- NOESY  
analysis of conformational effect of nitroxide spin labels on Ala-rich helical peptides, Bolin *et al.*, **131**, 248  
composite  $90^\circ$  pulse sequence appended to, Sodano, Landon, and Ptak, **133**, 194  
cross peak detection in 2D proton NMR spectra using excitation sculpting, Van and Shaka, **132**, 154  
1D steady-state intermolecular  $^{13}\text{C}$ ,  $^1\text{H}$ : evidence of solute–solvent interactions of  $\text{C}_{60}$  and  $\text{C}_{70}$  in solution, Troganis, Gerotheranassis, and Papademitriou, **130**, 149  
selective 1D and nonselective 2D, coherence selection and excitation sculpting using RF-gradients, Heikkinen, Rahkamaa, and Kilpeläinen, **133**, 183
- NOESY-HMQC  
 $^{15}\text{N}$ -edited 3D, with water flipback, Gruschus and Ferretti, **135**, 87
- NOESY-HSQC  
 $^{13}\text{C}$ -NOESY-HSQC with split carbon evolution for increased resolution with uniformly labeled proteins, Baur, Gemmecker, and Kessler, **132**, 191
- Noise  
influence on peak integrals obtained by direct summation, Bourg and Nuzillard, **134**, 184
- Noise pickup  
noise-immune coil for unshielded magnetic resonance measurements, Suits, Garroway, and Miller, **131**, 154
- Noise spectroscopy  
bandwidth extension, Yang and Zax, **135**, 267
- Non-Brownian particles  
determination of structure of suspensions, PFG NMR technique with applications to sphere packings, Talini, Leblond, and Feuillebois, **132**, 287
- Nonlinear EPR  
first-harmonic out-of-phase absorption EPR signals,  $T_1$  measurements, Livshits, Páli, and Marsh, **134**, 113
- Nonlinear stimulated echo  
in presence of inequivalent spins, Ardelean *et al.*, **132**, 138
- Nonresonant probe  
FT-EPR with: use of truncated coaxial line, Rubinson *et al.*, **132**, 255
- NQR (*see* Nuclear quadrupole resonance)
- Nuclear coherence  
low-frequency, simple two-pulse detection scheme in pulsed EPR for, Dzuba, Borovykh, and Hoff, **133**, 286
- Nuclear magnetic relaxation dispersion  
model-free analysis of stretched dispersions, Halle, Jóhannesson, and Venu, **135**, 1
- Nuclear Overhauser effect (*see* NOE)
- Nuclear Overhauser effect spectroscopy (*see* NOESY)
- Nuclear quadrupole resonance  
double resonance probe for close frequencies, Haase, Curro, and Slichter, **135**, 273  
rotating-frame, mapping molecular orientation in solids by, Casanova, Robert, and Pusiol, **133**, 129  
spin-echo mapping spectroscopy applied to, Bussandri and Zuriaga, **131**, 224  
surface and gradiometer coils near conducting body: lift-off effect, Suits, Garroway, and Miller, **135**, 373  
2D, using single crystal, off-resonance effects, Ajithkumar *et al.*, **135**, 165  
unshielded MR measurements, noise-immune coil for, Suits, Garroway, and Miller, **131**, 154
- Nuclear relaxation  
and electron relaxation, in magnetic exchange coupled dimers, Bertini *et al.*, **130**, 33

- Nuclear shielding constants  
rovibrationally averaged, in OCS, Jackowski, **135**, 444
- Nuclear shielding tensor  
 $^{31}\text{P}$ , and  $^{31}\text{P}$ ,  $^{63,65}\text{Cu}$  indirect spin–spin coupling tensor, of linear copper(I) phosphines, anisotropy, Kroeker *et al.*, **135**, 208
- Nuclear spin-echo Fourier-transform mapping spectroscopy  
applied to NQR, Bussandri and Zuriaga, **131**, 224
- Nuclear spin systems  
with dipole–dipole interactions, parametrical instability, Buishvili, Giorgadze, and Khomeriki, **130**, 82
- Nuclear-Zeeman-resolved ESEEM  
2D, Willer *et al.*, **133**, 46
- Nuclei (*see also* Quadrupolar nuclei)  
 $I = \frac{5}{2}$ , five-quantum coherence:  $^{27}\text{Al}$  in  $\text{Al}_2\text{O}_3$  single crystal, Kaikkonen, Ylinen, and Punkkinen, **133**, 123  
short  $T_2^*$ , relaxation time mapping with single-point imaging methods, Beyea *et al.*, **135**, 156  
spin- $\frac{3}{2}$   
 $^9\text{Be}$ , alignment echo: detection of ultraslow motion, Tang and Wu, **133**, 155  
in incompletely disordered systems, temporal characteristics of NMR signals, Woessner and Bansal, **133**, 21
- Nucleic acids  
and protein–nucleic acid complexes, three-bond  $^1\text{H}3'-^{31}\text{P}$  couplings, determination by quantitative  $J$  correlation spectroscopy, Clore *et al.*, **134**, 164
- Numerical modeling  
self-diffusion and relaxation time NMR in multi-compartment systems, Novikov, van Dusschoten, and Van As, **135**, 522
- Nutation  
nutration-frequency correlated EPR spectroscopy: PEANUT, Stoll *et al.*, **130**, 86
- O
- $^{17}\text{O}$   
D-mannopyranose, simultaneous suppression of two solvent signals: sequential inversion recovery with RIDE, Schulte, **134**, 168  
OCS, rovibrationally averaged nuclear shielding constants, Jackowski, **135**, 444
- OCS  
rovibrationally averaged nuclear shielding constants, Jackowski, **135**, 444
- ODESSA  
direct determination of motional correlational times, Favre, Schaefer, and Chmelka, **134**, 261
- Off-resonance effects  
suppression with adiabatic multiple spin-echo pulse sequence, Zweckstetter and Holak, **133**, 134  
in 2D NQR spectroscopy using single crystal, Ajithkumar *et al.*, **135**, 165
- Off-resonance relaxation  
rotating frame, experiment for analysis of macromolecular dynamics using adiabatic rotations, Mulder *et al.*, **131**, 351
- Oil  
and water, in sedimentary rock, *in situ* NMR analysis, de Swiet *et al.*, **133**, 385
- Oligomerization  
effect on NMR shieldings of  $\text{Be}^{+2}$  and  $\text{Al}^{+3}$  species in aqueous solution, Tossell, **135**, 203
- Oligonucleotides  
RNA, large, optimization of triple-resonance HCN experiments for, Sklenář *et al.*, **130**, 119
- One-dimensional exchange spectroscopy by sideband alternation (ODESSA)  
direct determination of motional correlational times, Favre, Schaefer, and Chmelka, **134**, 261  
time-reverse, direct determination of motional correlational times, Favre, Schaefer, and Chmelka, **134**, 261
- One-dimensional spectroscopy  
selective and nonselective, coherence selection and excitation sculpting using RF-gradients, Heikkinen, Rahkamaa, and Kilpeläinen, **133**, 183  
solid-state NMR analysis of nuclear spin interaction tensors associated with peptide bond, Lee and Ramamoorthy, **133**, 204
- Optically detected magnetic resonance  
electron spin echo envelope modulation, on photoexcited triplet state in zero magnetic field, Weis, Möbius, and Prisner, **131**, 17
- Optimization  
 $^{13}\text{C}$ – $^{14}\text{N}$  REAPDOR NMR, Ba *et al.*, **133**, 104  
evolution strategy, for selective pulses in NMR, Lunati *et al.*, **134**, 223  
with nonstandard flip angles, signal intensity and  $T_1$ -dependent contrast in spin-echo and inversion-recovery MR imaging, Bonny *et al.*, **130**, 51  
triple-resonance HCN experiments for larger RNA oligonucleotides, Sklenář *et al.*, **130**, 119
- Order  
incompletely disordered systems, temporal characteristics of NMR signals from spin  $\frac{3}{2}$  nuclei, Woessner and Bansal, **133**, 21
- Orientation  
in solids, mapping by rotating-frame NQR imaging, Casanova, Robert, and Pusiol, **133**, 129
- Orientalional constraints  
derived from hydrated powder samples by 2D PISEMA, Tian, Song, and Cross, **135**, 227
- Orientalional order  
high, influence on shape of echo response from Hahn pulse sequence, van der Struijf *et al.*, **130**, 253
- Orientalional symmetry  
computation of orientational averages in solid-state NMR by Gaussian spherical quadrature, Edén and Levitt, **132**, 220
- Oriented membranes  
magic angle-oriented sample spinning for, Glaubitz and Watts, **130**, 305
- Out-of-phase EPR  
first-harmonic absorption EPR signals,  $T_1$  measurements, Livshits, Páli, and Marsh, **134**, 113
- Overhauser enhancement  
in proton–electron double-resonance imaging: mobility and free radical concentration effects, de Sousa *et al.*, **135**, 118
- Overhauser magnetic resonance imaging  
oximetry with, single-electron contrast agents for, EPR and DNP properties, Ardenkjær-Larsen *et al.*, **133**, 1
- Oximetric imaging  
single-electron contrast agents for, EPR and DNP properties, Ardenkjær-Larsen *et al.*, **133**, 1
- Oximetry  
EPR, carbon-based paramagnetic materials for use in, separation and enrichment of active component, Liu *et al.*, **133**, 291
- Oxovanadium complex  
determination of nitrogen quadrupole and hyperfine tensors by simultaneous fitting of multifrequency ESEEM powder spectra, Reijerse, Tyryshkin, and Dikanov, **131**, 295
- Oxygen  
dissolved, sensitivity improvement in INADEQUATE by, Mattiello and Freeman, **135**, 514
- P
- $^{31}\text{P}$   
 $^{27}\text{Al} \rightarrow ^{31}\text{P}$  INEPT experiments on  $(\text{CH}_3)_3\text{P}-\text{AlCl}_3$  complex in zeolite NaX, Kao and Grey, **133**, 313  
anisotropy in  $^{31}\text{P}$ ,  $^{63,65}\text{Cu}$  indirect spin–spin coupling and  $^{31}\text{P}$  nuclear shielding tensors of linear copper(I) phosphines, Kroeker *et al.*, **135**, 208

- carbon-phosphorus  $J$  coupling constants in RNA, by spin-echo difference constant-time HCC $\text{H}$ -COSY, Hoogstraten and Pardi, **133**, 236
- cardiac NMR in humans, improved sensitivity and localization, Löffler *et al.*, **134**, 287
- dimyristoylphosphatidylcholine bilayers, magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305
- frequency-selective MRS data quantification with frequency prior knowledge, Dologlou, Van Huffel, and Van Ormondt, **130**, 238
- and  $^2\text{H}$ , NMR of nematic phospholipid bicellar solutions, Struppe and Vold, **135**, 541
- $J$ -coupled to  $^{59}\text{Co}$  nucleus in  $\text{HFeCo}_3(\text{CO})_{11}\text{PPh}_2\text{H}$ , differential line broadening in presence of quadrupolar-CSA interference, Elbayed *et al.*, **130**, 209
- MRS spectra from rat brain, precision of parameters in measuring, Golding and Golding, **135**, 380
- $\text{NaH}_2\text{PO}_4$ : solvent magnetization artifacts in high-field NMR studies of macromolecular hydration, Sobol *et al.*, **130**, 262
- $\text{Na}_4\text{P}_2\text{O}_7 \cdot 10\text{H}_2\text{O}$  and  $\text{Cd}(\text{NO}_3)_2 \cdot 2\text{PPh}_3$ : iterative lineshape fitting of MAS NMR spectra, Dusold, Milius, and Sebald, **135**, 500
- paramagnetic lanthanide complex: NMR temperature measurements, Zuo *et al.*, **133**, 53
- proton-decoupled NMR, detection of phosphomonoester signals in myocardial hypertrophy, Jung *et al.*, **133**, 232
- three-bond  $^1\text{H}^3\text{-}^{31}\text{P}$  couplings in nucleic acids and protein-nucleic acid complexes, determination by quantitative  $J$  correlation spectroscopy, Clore *et al.*, **134**, 164
- Packings  
spheres, application of pulsed-field gradient NMR technique for determination of structure of suspensions of non-Brownian particles, Talini, Leblond, and Feuillebois, **132**, 287
- Parahydrogen  
analysis of catalytic processes using: evolution of zero-quantum coherence in AA'X spin systems, Natterer *et al.*, **133**, 92
- Parallel processing  
distributed, for multidimensional maximum entropy reconstruction, Li, Stern, and Hoch, **134**, 161
- Paramagnetic compounds  
solid-state NMR, effect of bulk magnetic susceptibility, Kubo, Spaniol, and Terao, **133**, 330
- Paramagnetic inversion  
of sign of interference contribution of  $T_2$  of imido protons of coordinated imidazoles in uniformly  $^{15}\text{N}$ -labeled cytochrome  $c_3$ , Ohmura *et al.*, **131**, 367
- Paramagnetic ions  
assisted DNA magnetic alignment, determination of DNA internuclear angles using, **135**, 256
- Paramagnetic materials  
carbon-based, for use in EPR oximetry, separation and enrichment of active component, Liu *et al.*, **133**, 291
- Paramagnetic proteins  
NMR study using PASE: paramagnetic signals enhancement, Bondon and Mouro, **134**, 154
- Paramagnetic signals enhancement  
method for NMR study of paramagnetic proteins: PASE, Bondon and Mouro, **134**, 154
- Paramagnetic thermometric substance  
lanthanide complex, Zuo *et al.*, **133**, 53
- Parametrical resonance  
instability, in nuclear spin systems with dipole-dipole interactions, Buishvili, Giorgadze, and Khomeriki, **130**, 82
- PASE  
method for NMR study of paramagnetic proteins: paramagnetic signals enhancement, Bondon and Mouro, **134**, 154
- PASS  
six-pulse PASS MAS-NMR technique solved by phasor method, Effey, Butcher, and Cappelletti, **132**, 266
- 2D-PASS, analysis of  $^{13}\text{C}$  and  $^{15}\text{N}$  chemical-shift anisotropy of ampicillin and penicillin-V, Antzutkin, Lee, and Levitt, **135**, 144
- Paw  
high-resolution 3D MRI and histology: Sandimmune Neoral effects on collagen arthritis in DA rat, Beckmann *et al.*, **131**, 8
- Peak integration  
automated, in macromolecular NMR spectra, using AUTOPSY, Koradi *et al.*, **135**, 288
- Peak picking  
automated, in macromolecular NMR spectra, using AUTOPSY, Koradi *et al.*, **135**, 288
- Peaks  
diagonal, suppression with singular value decomposition, Zhu *et al.*, **132**, 176
- Peak separation  
precision in estimating, Jupp, Harris, and Aliev, **135**, 23
- PEANUT  
nutration-frequency correlated EPR spectroscopy, Stoll *et al.*, **130**, 86
- Penicillin-V  
 $^{13}\text{C}$  and  $^{15}\text{N}$  chemical-shift anisotropy by 2D-PASS and CP/MAS NMR, Antzutkin, Lee, and Levitt, **135**, 144
- 4-*n*-Pentyl-4'-cyanobiphenyl  
liquid crystalline, application of broadband decoupling sequence SPARC-16, Yu and Fung, **130**, 317
- Peptide bond  
nuclear spin interaction tensors associated with, 1D solid-state NMR method for, Lee and Ramamoorthy, **133**, 204
- Peptides  
Ala-rich helical, conformational effect of nitroxide spin labels on, NMR analysis, Bolin *et al.*, **131**, 248
- membrane, magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305
- torsion angle determination by  $^{15}\text{N}$  chemical shift and  $^{13}\text{C}^{\alpha}\text{-}^1\text{H}^{\alpha}$  dipolar tensor correlation in solid-state MAS NMR, Hong *et al.*, **135**, 169
- Perdeuterated protein  
backbone amide and aliphatic gamma resonances, unambiguous correlations, McCallum, Hitchens, and Rule, **134**, 350
- Perfluorocarbon FC40  
 $^{19}\text{F}$  MRI in inhomogeneous fields by SPARE sequence, McIntyre, Hennel, and Morris, **130**, 58
- PFG (*see* Pulsed-field gradients)
- pH  
role in magnetization transfer in polymers, Gochberg *et al.*, **131**, 191
- Pharmacokinetics  
isolated perfused liver: intravascular and intracellular relaxivities of superparamagnetic particles, Colet *et al.*, **134**, 199
- Phase  
arbitrary, simultaneous and independent rotations with, for I,  $\text{IS}^{\alpha}$ , and  $\text{IS}^{\beta}$  spin systems, Briand and Sørensen, **135**, 44
- and symmetry, simplification of NMR spectra of liquid crystalline samples by, Carravetta *et al.*, **135**, 298
- Phase adjustment of spinning sidebands (*see* PASS)
- Phase correction  
NMR spectra by 1D filter diagonalization method, Hu *et al.*, **134**, 76
- Phase cycling  
coherence transfer pathway selection in NMR by, simulation, Jerschow and Müller, **134**, 17
- Phase memory time  
 $T_2$ , use for quantitative characterization of rotational motions of proteins in lipid bilayer systems, van der Struijf and Levine, **130**, 244

- Phase-modulated CLEAN chemical exchange (CLEANEX-PM)  
attenuation of cross-relaxation peaks, Hwang and Shaka, **135**, 280
- Phase reference  
use for field mapping with amplitude images at low field, Tomasi *et al.*, **131**, 310
- Phase-sensitive detection  
2D homonuclear correlation spectra in MAS NMR, Boender and Vega, **133**, 281
- Phase transitions  
ferroelectric, in EPR studies of pyridinium tetrafluoroborate, radicals as probes, Krzyminiewski, Bielewicz-Mordalska, and Kowalczyk, **135**, 76
- Phasor method  
six-pulse PASS MAS-NMR technique solved by, Effey, Butcher, and Capelletti, **132**, 266
- Phospholipase A<sub>2</sub>  
multidimensional NMR, HSQC-based methyl group selection via gradients, Diercks, Schwaiger, and Kessler, **130**, 335
- Phospholipid  
bicelles, dilute solutions for structural NMR work, Struppe and Vold, **135**, 541
- Phosphomonoesters  
signal detection in proton-decoupled <sup>31</sup>P NMR spectra in myocardial hypertrophy, Jung *et al.*, **133**, 232
- Photoexcited triplet state  
acridine-*d*<sub>9</sub> in fluorene-*h*<sub>10</sub> matrix, in zero magnetic field, optically detected ESEEM, Weis, Möbius, and Prisner, **131**, 17
- $\alpha$ - $\alpha^3$ C<sub>2</sub>-Phthalic acid  
polarization transfer under RF-driven and slow-magic-angle-sample-spinning conditions, Robyr and Gan, **131**, 254
- PISEMA  
2D, orientational constraints derived from hydrated powder samples by, Tian, Song, and Cross, **135**, 227
- Planar mixing  
analytical polarization and coherence transfer functions for three coupled spins  $\frac{1}{2}$ , Schedletzy, Luy, and Glaser, **130**, 27
- Point resolved spectroscopy (*see* PRESS)
- Polarimeter  
low-frequency NMR, for hyperpolarized gases, Saam and Conradi, **134**, 67
- Polarization inversion spin exchange at magic angle (PISEMA)  
2D, orientational constraints derived from hydrated powder samples by, Tian, Song, and Cross, **135**, 227
- Polarization transfer  
analytical transfer functions for three coupled spins  $\frac{1}{2}$  under planar mixing conditions, Schedletzy, Luy, and Glaser, **130**, 27  
distortionless enhancement by, including detection of quaternary nuclei: DEPTQ, Burger and Bigler, **135**, 529  
RF-driven and slow-magic-angle-sample-spinning techniques, comparison, Robyr and Gan, **131**, 254
- Polarizing field  
large, high magnetic field gradient PGSE NMR in presence of, Callaghan, Komlos, and Nyden, **133**, 177
- Polyacrylamide gels  
proton diffusion and *T*<sub>1</sub> relaxation, unified approach using volume averaging, Penke *et al.*, **132**, 240
- Polychlorotrifluoroethylene  
rotational motion of 1,3-dibromoadamantane in, <sup>13</sup>C NMR, Lienin, Brüsche-weiler, and Ernst, **131**, 184
- Polycrystalline sample  
powders, <sup>1</sup>H chemical-shielding anisotropies using MSHOT-3 based CRAMPS, Hohwy *et al.*, **133**, 374  
solid-state NMR, effect of bulk magnetic susceptibility, Kubo, Spaniol, and Terao, **133**, 330
- Polyethylene  
linear, rigid strongly coupled carbons, <sup>13</sup>C CPMAS linewidths under CW proton decoupling, VanderHart and Campbell, **134**, 88
- Polymers  
chain end structures, 3D pulse sequences for, Saito and Rinaldi, **130**, 135  
conformation by 2D double-quantum solid-state NMR, complete dipolar decoupling of <sup>13</sup>C for, Schmidt-Rohr, **131**, 209  
DOSY and GPC-NMR experiments, multivariate curve resolution, Van Gorkom and Hancewicz, **130**, 125  
magnetization transfer in, role of specific side groups and pH, Gochberg *et al.*, **131**, 191  
surface NMR analysis via MOUSE, Guthausen *et al.*, **130**, 1
- Polysaccharide  
model meal, viscosity measurement with echo-planar imaging relaxometry, Marciani *et al.*, **135**, 82
- Polystyrene  
chain end structures, 3D pulse sequences for characterization, Saito and Rinaldi, **130**, 135  
structure, 3D NMR experiments for characterization, Saito and Rinaldi, **132**, 41
- Pore structure  
silica aerogels, determination by <sup>129</sup>Xe NMR and imaging, Gregory, Gerald, and Botto, **131**, 327
- Porous media  
effective gradients due to susceptibility differences, Hürlimann, **131**, 232  
electrical transport in, NMR studies, Heil and Holz, **135**, 17  
sedimentary rock, fluids contained, *in situ* NMR analysis, de Swiet *et al.*, **133**, 385  
silica aerogels, pore-structure determination by <sup>129</sup>Xe NMR and imaging, Gregory, Gerald, and Botto, **131**, 327  
time-dependent gas diffusion, PFG measurements, Mair *et al.*, **135**, 478
- Porphyry deformation  
effect on <sup>57</sup>Fe shieldings in superstructured hemoprotein models, Gerothernassis *et al.*, **131**, 163
- Porphyryns  
spin-labeled, high-spin iron(III)-nitroxyl distances by time-domain EPR, Rakowsky *et al.*, **131**, 97
- Powders  
Cu<sub>2</sub>O, <sup>63</sup>Cu NMR imaging using second-order quadrupole broadened resonances, Swaminathan and Suits, **132**, 274  
disordered, high-frequency/high-field ESEEM study of nitrogen hyperfine and quadrupole interactions, Bloß, Möbius, and Prisner, **134**, 30  
hydrated, orientational constraints derived by 2D PISEMA, Tian, Song, and Cross, **135**, 227  
polycrystalline, <sup>1</sup>H chemical-shielding anisotropies using MSHOT-3 based CRAMPS, Hohwy *et al.*, **133**, 374
- Powder spectra  
<sup>2</sup>H, noise spectroscopy application to, bandwidth extension, Yang and Zax, **135**, 267  
multifrequency ESEEM, determination of nitrogen quadrupole and hyperfine tensors in oxovanadium complex by simultaneous fitting of, Reijerse, Tyryshkin, and Dikanov, **131**, 295
- p21 *ras*  
structure and activity, effects of cryoprotection: implications for ESEEM, Halkides, Farrar, and Singel, **134**, 142
- Pr(C<sub>2</sub>D<sub>5</sub>SO<sub>4</sub>)<sub>3</sub> · 8H<sub>2</sub>O  
paramagnetic, solid-state NMR, effect of bulk magnetic susceptibility, Kubo, Spaniol, and Terao, **133**, 330
- Precision  
in estimating frequency separation between spectral lines, Jupp, Harris, and Aliev, **135**, 23

## PRESS

excitation, *in vivo* lactate editing with simultaneous detection of upfield singlets at 1.5 T with, applications to brain and head and neck tumors, Star-Lack *et al.*, **133**, 243  
and STEAM, flip angle effects—optimized vs sinc RF pulses, Ryner, Ke, and Thomas, **131**, 118

Probes (*see also* NMR probes)

nonresonant FT-EPR, use of truncated coaxial line, Rubinson *et al.*, **132**, 255  
Progressive saturation EPR

spin label relaxation time determination: molecular motion and Zeeman modulation effects, Livshits, Páli, and Marsh, **133**, 79

## Protein binding

isotope-filtered NMR for, Gonnella *et al.*, **131**, 336

## Protein/DNA complex

weak labile  $^1\text{H}$  resonances of side chains, enhancement:  $^{15}\text{N}$ -edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87

## Protein dynamics

analysis using adiabatic rotations, off-resonance rotating frame relaxation experiment for, Mulder *et al.*, **131**, 351

backbone: quantitative analysis of dipole-CSA cross-correlated relaxation by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346

improved estimation of rotational correlation times from  $^{15}\text{N}$  relaxation measurements, Yao, Hinds, and Norton, **131**, 347

## Protein–ligand interactions

heteronuclear X-filter  $^1\text{H}$  PFG double-quantum experiment for proton resonance assignment of ligand bound to protein, Dalvit, Ramage, and Hommel, **131**, 148

## Protein–nucleic acid complexes

three-bond  $^1\text{H}3'-^{31}\text{P}$  couplings, determination by quantitative  $J$  correlation spectroscopy, Clore *et al.*, **134**, 164

## Proteins

accuracy of HN–HN distance constraints, deuteration effects, Briercheck and Rule, **134**, 52

automated peak picking and peak integration in NMR using AUTOPSY, Koradi *et al.*, **135**, 288

 $^{13}\text{C}$ -labeled

assignment of aromatic side chains, 2D NMR experiments for, Prompers *et al.*, **130**, 68

$^{13}\text{C}$ -NOESY–HSQC with split carbon evolution for increased resolution of, Baur, Gemmecker, and Kessler, **132**, 191

in  $\text{H}_2\text{O}$  solution, H(C)CH-COSY and (H)CCH-COSY experiments, Gehring and Ekiel, **135**, 185

 $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled

amide deuterium isotope effects on  $\text{C}^\beta$  chemical shifts, multidimensional editing for, Meissner and Sørensen, **135**, 547

Bloch–Siegert shift-compensated and cyclic sideband-eliminated double-adiabatic homonuclear decoupling for, Zhang and Gorenstein, **132**, 81  
spin relaxation, measurement of cross correlation between dipolar coupling and chemical-shift anisotropy, Ghose, Huang, and Prestegard, **135**, 487

diffusion in aqueous solution, practical aspects of measurement, Tillett, Lian, and Norwood, **133**, 379

hydration, solvent magnetization artifacts in high-field NMR studies, Sobol *et al.*, **130**, 262

large, 3D NMR experiments for  $^{15}\text{N}$  relaxation data, Caffrey *et al.*, **135**, 368

magnetic coupling between lactate and water protons mediated by, Swanson, **135**, 248

## membrane

magic angle-oriented sample spinning approach, Glaubitz and Watts, **130**, 305

$^{15}\text{N}$  chemical-shift anisotropy, cation binding-induced changes, Tian and Cross, **135**, 535

multidimensional NMR, HSQC-based methyl group selection via gradients, Diercks, Schwaiger, and Kessler, **130**, 335

$^{15}\text{N}$ – $^{15}\text{N}$   $J$ -couplings, detection, Lühr and Rüterjans, **132**, 130

oriented, asymmetric alignment tensor, determination in absence of structural information, Clore, Gronenborn, and Bax, **133**, 216

rotational motions in lipid bilayer systems, use of phase memory time  $T_2$  for quantitative characterization, van der Struijf and Levine, **130**, 244  
side chains

contacting DNA, enhancement of weak labile  $^1\text{H}$  resonances:  $^{15}\text{N}$ -edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87  
dynamics, analysis from  $^{13}\text{C}$  NMR relaxation data, Daragan and Mayo, **130**, 329

spectral window in NMR revealing cross-relaxation between amide protons, Mutzenhardt and Bodenhausen, **132**, 159

torsion angle determination by  $^{15}\text{N}$  chemical shift and  $^{13}\text{C}^\alpha$ – $^1\text{H}^\alpha$  dipolar tensor correlation in solid-state MAS NMR, Hong *et al.*, **135**, 169

weakly oriented, measurement of dipolar couplings for methylene and methyl sites and use in structure determination, Ottiger *et al.*, **134**, 365

## Proton–electron double-resonance imaging

mobility and free radical concentration effects, de Sousa *et al.*, **135**, 118

## Proton hyperfine structure

unresolved, contribution to Gaussian line broadening of proxyl spin probe EPR spectrum, Bales, Peric, and Lamy-Freund, **132**, 279

## Proton NMR

high-resolution MAS, evaluation of breast ductal carcinomas, Cheng *et al.*, **135**, 194

MAS, at high magnetic fields and high spinning frequencies: spectral simulations using Floquet theory, Ray *et al.*, **135**, 418

## Protons

amide, cross-relaxation between, detection in protein NMR, Mutzenhardt and Bodenhausen, **132**, 159

assignment in uniformly  $^{13}\text{C}$ - and selectively  $^2\text{H}$ -labeled RNA, 3D C(CC)H TOCSY for, Dayie, Tolbert, and Williamson, **130**, 97

attached to  $^{13}\text{C}$ , scalar spin–spin coupling constants between, methylene specific experiments, Carlomagno *et al.*, **135**, 216

exchangeable, TOCSY experiments on systems with, information from water stripe, Nakashima, McClung, and Kotovych, **133**, 222

fast exchanging, sensitivity-enhanced detection by exchange-edited gradient HCP-HSQC, Zangger and Armitage, **135**, 70

imido, of coordinated imidazoles in uniformly  $^{15}\text{N}$ -labeled cytochrome  $c_3$ , interference contribution to  $T_2$  of, paramagnetic inversion of sign of, Ohmura *et al.*, **131**, 367

inverse detection through MILS technique, alternative scheme for  $^{13}\text{C}$  chemical-shift imaging via, Lee *et al.*, **130**, 102

lactate and water, protein-mediated magnetic coupling, Swanson, **135**, 248  
solvent, measurement of chemical exchange rate constants with, using radiation damping, Chen and Mao, **131**, 358

spectra, residual water, fast removal, **132**, 197

vicinal F–H coupling constants: individual substituent effects, Fabián, Guillemé, and Díez, **133**, 255

 $^{195}\text{Pt}$  NMR

monitoring of local disposition kinetics of carboplatin *in vivo* after sc injection in rat, Becker *et al.*, **133**, 115

## Pulsed ENDOR

matrix line, Astashkin and Kawamori, **135**, 406

studies at 95 GHz of triplet state of  $^{13}\text{C}_{60}$ , van den Berg *et al.*, **131**, 39

## Pulsed EPR

large sample, submicrosecond resonator and receiver for, Alecci *et al.*, **132**, 162

matched three-pulse ESEEM and matched HYSORE, sensitivity enhancement by, Jeschke, Rakhmatullin, and Schweiger, **131**, 261

nutational-frequency correlated EPR spectroscopy: PEANUT, Stoll *et al.*, **130**, 86

simple two-pulse detection scheme for low-frequency nuclear coherences, Dzuba, Borovykh, and Hoff, **133**, 286

2D nuclear-Zeeman-resolved ESEEM spectroscopy, Willer *et al.*, **133**, 46



- Pulsed-field gradients  
 coherence selection in 2D MAS spectroscopy of dipolar solids, Fritzhanns *et al.*, **134**, 355  
 coherence transfer pathway selection, simulation, Jerschow and Müller, **134**, 17  
 diffusion coefficient measurement by 2D NMR, with accordion spectroscopy, Millet and Pons, **131**, 166  
 diffusion measurements, and proton  $T_1$  relaxation, in polyacrylamide gels, unified approach using volume averaging, Penke *et al.*, **132**, 240  
 with electrophoretic NMR, analysis of electrical transport in disordered medium, Heil and Holz, **135**, 17  
 enhanced hetero ( $\omega_1$ ) half-filtered TOCSY, sensitivity-enhanced, for long-range heteronuclear coupling constants, Uhrín *et al.*, **130**, 155  
 enhanced HMQC, Zhu, Kong, and Sze, **135**, 232  
 enhanced NMR, convection compensation, Jerschow and Müller, **132**, 13  
 enhanced TOCSY, sensitivity-enhanced, Kövér, Uhrín, and Hruby, **130**, 162  
 heteronuclear X-filter  $^1\text{H}$  PFG double-quantum experiment for proton resonance assignment of ligand bound to protein, Dalvit, Ramage, and Hommel, **131**, 148  
 large, PGSE NMR in presence of large polarizing field requiring, Callaghan, Komlosch, and Nyden, **133**, 177  
 NMR probes, for solid-state studies, Zhang and Cory, **132**, 144  
 non-Brownian particle suspension structure determination, with applications to sphere packings, Talini, Leblond, and Feuillebois, **132**, 287  
 self-diffusion and relaxation time modeling in multi-compartment systems, Novikov, van Dusschoten, and Van As, **135**, 522  
 time-dependent gas diffusion measurements, Mair *et al.*, **135**, 478
- Pulsed gradient spin echo  
 diffusion of proteins in aqueous solution measured with, practical aspects, Tillet, Lian, and Norwood, **133**, 379  
 Earth's field NMR in Antarctica: study of restricted diffusion in sea ice, Callaghan *et al.*, **133**, 148  
 high magnetic field gradient NMR in presence of large polarizing field, Callaghan, Komlosch, and Nyden, **133**, 177  
 multilayer gradient coil design for, Bowtell and Robyr, **131**, 286  
 time-dependent gas diffusion measurements, Mair *et al.*, **135**, 478
- Pulsed NMR  
 with large samples, submicrosecond resonator and receiver for, Alecci *et al.*, **132**, 162  
 low-frequency spectrometer for monitoring spin polarization of hyperpolarized gases, Saam and Conradi, **134**, 67
- Pulse imperfections  
 effects due to, removal with adiabatic multiple spin-echo pulse sequence, Zweckstetter and Holak, **133**, 134
- Pulses  
 adiabatic (*see* Adiabatic pulses)  
 matched microwave, sensitivity enhancement by, in 1D and 2D ESEEM spectroscopy, Jeschke, Rakhmatullin, and Schweiger, **131**, 261  
 shaped, multiple-quantum MAS NMR of spin- $\frac{3}{2}$  quadrupolar spin systems, Ding and McDowell, **135**, 61
- Pulse sequences  
 adiabatic multiple spin-echo, Zweckstetter and Holak, **133**, 134  
 binomial solvent suppression, offset-dependent partial saturation, Hsu and Gregory, **131**, 46  
 composite 90°: distortionless NMR spectra and suppression without destruction of water magnetization, Sodano, Landon, and Ptak, **133**, 194  
 for correlation of backbone amide and aliphatic gamma resonances in deuterated proteins, McCallum, Hitchens, and Rule, **134**, 350  
 Hahn, shape of echo response from, influence of high orientational order, van der Struijf *et al.*, **130**, 253  
 NMR, quantum logic gates and, Jones, Hansen, and Mosca, **135**, 353  
 for steady-state saturation of flowing spins, Marshall, **133**, 13  
 $(\theta_1)_x - \tau - (\theta_2)_y - t$ ,  $^{14}\text{N}$  NMR echo in  $\text{NH}_4\text{ClO}_4$  after, Punkkinen *et al.*, **130**, 287
- WATERGATE, improved, for solvent suppression in NMR, Liu *et al.*, **132**, 125
- Purine nucleotides  
 $^{13}\text{C}$ -labeled, DNA decamer duplex with,  $^{13}\text{C}$  relaxation, Kojima *et al.*, **135**, 310
- Pyridinium tetrafluoroborate  
 EPR, ferroelectric phase transitions in, radicals as probes, Krzyminiowski, Bielewicz-Mordalska, and Kowalczyk, **135**, 76
- Q
- QCPMG  
 –MAS NMR, half-integer quadrupolar nuclei, Larsen *et al.*, **131**, 144
- Quadrupolar interactions  
 distributions in 2D multiple-quantum MAS NMR spectra following, Bodart, **133**, 207  
 and nitrogen hyperfine coupling, on disordered powder sample, high-frequency/high-field ESEEM study, Bloeb, Möbius, and Prisner, **134**, 30
- Quadrupolar nuclei  
 application of time propagation technique for MAS NMR simulation, Charpentier, Fermon, and Virlet, **132**, 181  
 half-integer  
 MQMAS NMR, composite pulse excitation schemes for, Marinelli, Medek, and Frydman, **132**, 88  
 QCPMG–MAS NMR, Larsen *et al.*, **131**, 144  
 in solids, INEPT experiments involving, Kao and Grey, **133**, 313  
 spin  $\frac{3}{2}$ , of incompletely disordered systems, temporal characteristics of NMR signals, Woessner and Bansal, **133**, 21
- Quadrupolar relaxation  
 and CSA relaxation, differential line broadening in presence of, Elbayed *et al.*, **130**, 209
- Quadrupole broadened resonances  
 second-order, NMR imaging, Swaminathan and Suits, **132**, 274
- Quadrupole coupling constant  
 deuterium, and dynamic isotope effect, determination in liquids, Hardy *et al.*, **134**, 300
- Quadrupole–dipole effects  
 in solid-state CP–MAS  $^{119}\text{Sn}$  NMR spectra of *para*-substituted triaryltin-(pentacarbonyl)manganese(I) complexes, Christendat *et al.*, **131**, 1
- Quadrupole splitting  
 NMR spectra of incompletely disordered systems: temporal characteristics of spin  $\frac{3}{2}$  nuclei, Woessner and Bansal, **133**, 21
- Quadrupole tensor  
 and hyperfine tensor,  $^{14}\text{N}$ , determination in oxovanadium complex by simultaneous fitting of multifrequency ESEEM powder spectra, Reijerse, Tyryshkin, and Dikanov, **131**, 295
- Quality factor  
 super-Q detection of transient MR signals, Suits, Garroway, and Miller, **132**, 54
- Quantitative  $J$  correlation spectroscopy  
 determination of three-bond  $^1\text{H}3' - ^{31}\text{P}$  couplings in nucleic acids and protein–nucleic acid complexes by, Clore *et al.*, **134**, 164
- Quantum computer  
 quantum logic gates and NMR pulse sequences, Jones, Hansen, and Mosca, **135**, 353
- Quaternary nuclei  
 detection, and distortionless enhancement by polarization transfer: DEPTQ, Burger and Bigler, **135**, 529
- Qubit  
 NMR pulse sequences and quantum logic gates, Jones, Hansen, and Mosca, **135**, 353
- QUIET–BIRD–NOESY  
 detection of cross-relaxation between amide protons in protein NMR, Mutzenhardt and Bodenhausen, **132**, 159

- Quiet NOE  
interresidue, for DNA structural studies, Beger *et al.*, **132**, 34
- QUIET-NOESY  
selective NMR experiments on macromolecules, Esposito *et al.*, **132**, 204
- R
- $R_2'$   
measurement in presence of multiple spectral components using reference spectrum deconvolution, Wehrli *et al.*, **131**, 61
- Radial basis network  
determining rotational correlation times from slow motional ESR spectra with, comparison with multilayer perceptron, Martinez and Millhauser, **134**, 124
- Radial echo-planar imaging  
description and experimental results, Silva *et al.*, **135**, 242
- Radiation damping  
composite 90° pulse sequence use: suppression without destruction of water magnetization, Sodano, Landon, and Ptak, **133**, 194  
measurement of chemical exchange rate constants with solvent protons, Chen and Mao, **131**, 358  
solvent magnetization artifacts in high-field NMR studies of macromolecular hydration, Sobol *et al.*, **130**, 262
- Radicals  
as probes of ferroelectric phase transitions in EPR studies of pyridinium tetrafluoroborate, Krzyminiwski, Bielewicz-Mordalska, and Kowalczyk, **135**, 76
- Radiofrequency-driven conditions  
polarization-transfer under, comparison with slow-magic-angle-sample-spinning conditions, Robyr and Gan, **131**, 254
- Radiofrequency field gradients  
chemical-shift imaging in rotating solids by, Malveau *et al.*, **134**, 171  
coherence selection and excitation sculpting, in selective 1D and nonselective 2D experiments, Heikkinen, Rahkamaa, and Kilpeläinen, **133**, 183  
diffusion measurements, Humbert *et al.*, **134**, 245  
self-diffusion measurements by  $^{13}\text{C}$  NMR, Malveau *et al.*, **130**, 131
- Radiofrequency field strength  
and spinning speed, effect on MQMAS NMR, Amoureux *et al.*, **131**, 170
- Radiofrequency pulses  
generation, Shinnar-Le Roux *B* polynomial ripples for, generalized estimate for, Raddi and Klose, **132**, 260  
optimized vs sinc, flip angle effects in STEAM and PRESS, Ryner, Ke, and Thomas, **131**, 118  
PRESS excitation: *in vivo* lactate editing with simultaneous detection of uncoupled moieties at 1.5 T, with applications to head and neck tumors, Star-Lack *et al.*, **133**, 243
- Radiofrequency shielding  
birdcage coil tuned by, for application at 9.4 T, Dardzinski *et al.*, **131**, 32
- Raffinose  
heteronuclear long-range couplings, measurement by gradient-enhanced 2D multiple-bond correlation, Sheng and van Halbeek, **130**, 296
- ras*  
p21 *ras* structure and activity, effects of cryoprotection: implications for ESEEM, Halkides, Farrar, and Singel, **134**, 142
- $^{87}\text{Rb}$   
agar powder, selective detection of biexponential relaxation in isotropic solutions by mixed coherence filtering NMR, Yamada *et al.*, **135**, 99  
 $\text{Rb}_2\text{SO}_4$ , QCPMG-MAS NMR of half-integer quadrupolar nuclei, Larsen *et al.*, **131**, 144
- Real-time MRI  
image reconstruction and display system using high-speed personal computer, Haishi and Kose, **134**, 138
- REAPDOR  
optimization of  $^{13}\text{C}$ - $^{14}\text{N}$  REAPDOR NMR, Ba *et al.*, **133**, 104
- Receiver  
and resonator, submicrosecond system for pulsed MR with large samples, Alecci *et al.*, **132**, 162
- Reciprocity  
generalized, Insko *et al.*, **131**, 111
- REDOR  
signal and inversion, d'Espinose de la Caillerie and Fretigny, **133**, 273
- Reduction factors  
simplification of  $^{19}\text{F}$  NMR spectra of liquid crystalline samples by multiple-pulse COSY, Lesot *et al.*, **133**, 166
- Reference deconvolution  
NMR spectra by 1D filter diagonalization method, Hu *et al.*, **134**, 76
- Reference spectrum deconvolution  
 $R_2'$  measurement in presence of multiple spectral components, Wehrli *et al.*, **131**, 61
- Refinement  
structure, against residual dipolar couplings, in presence of rhombicity of unknown magnitude, Clore, Gronenborn, and Tjandra, **131**, 159
- Relaxation (*see also* Spin-lattice relaxation; Spin-spin relaxation)  
biexponential, selective detection in isotropic solutions by mixed coherence filtering NMR, Yamada *et al.*, **135**, 99  
 $^{13}\text{C}$ ,  $^{15}\text{N}$ -labeled proteins, measurement of cross correlation between dipolar coupling and chemical-shift anisotropy, Ghose, Huang, and Prestegard, **135**, 487  
 $^{13}\text{C}$  NMR  
analysis of protein side-chain dynamics, Daragan and Mayo, **130**, 329  
DNA duplex dynamics via, Kojima *et al.*, **135**, 310  
dipole-CSA cross-correlated, quantitative analysis by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346  
effect of coupling between rotational and translational Brownian motions: Green function of rigid body diffusion, Beloborodov, Orekhov, and Arseniev, **132**, 328  
electron and nuclear, in magnetic exchange coupled dimers, Bertini *et al.*, **130**, 33  
laboratory-frame, in estimation of chemical-shift anisotropy-dipolar coupling cross-correlation rates, Ghose and Prestegard, **134**, 308  
magnetization transfer in polymers, role of specific side groups and pH, Gochberg *et al.*, **131**, 191  
 $^{15}\text{N}$   
improved estimation of protein rotational correlation times, Yao, Hinds, and Norton, **131**, 347  
large proteins, 3D NMR experiments: 44-kDa ectodomain of SIV gp41, Caffrey *et al.*, **135**, 368  
rate in poorly resolved NMR spectra, 3D accordion spectroscopy measurement, Carr, Fearing, and Palmer, **132**, 25  
off-resonance rotating frame, experiment for analysis of macromolecular dynamics using adiabatic rotations, Mulder *et al.*, **131**, 351  
quadrupolar-CSA, differential line broadening in presence of, Elbayed *et al.*, **130**, 209
- Relaxation dispersions  
stretched, model-free analysis, Halle, Jóhannesson, and Venu, **135**, 1
- Relaxation rate  
echo-planar imaging measurement of viscosity of model meal, Marciari *et al.*, **135**, 82
- Relaxation time  
determination by progressive saturation EPR: effects of molecular motion and Zeeman modulation for spin labels, Livshits, Páli, and Marsh, **133**, 79  
 $T_1$ , reduction with dissolved oxygen: sensitivity improvement in INADEQUATE, Mattiello and Freeman, **135**, 514  
triple-quantum, determination of dipolar interactions of  $^{23}\text{Na}$  in solution by, Eliav and Navon, **130**, 63
- Relaxation time mapping  
short  $T_2^*$  nuclei using single-point imaging methods, Beyea *et al.*, **135**, 156

- Relaxation time NMR  
and self-diffusion, modeling in multi-compartment systems, Novikov, van Dusschoten, and Van As, **135**, 522
- Relaxometry  
echo-planar imaging, measurement of viscosity of model meal, Marciani *et al.*, **135**, 82  
field-cycling, formalism for evaluation of order-fluctuation modes in liquid crystals, Grinberg *et al.*, **135**, 54  
hepatic intravascular and intracellular, of superparamagnetic particles, Colet *et al.*, **134**, 199
- Resolution enhancement  
from double constant-time evolution of 3D and 4D triple-resonance experiments, McCoy, **130**, 341  
in *in vivo*  $^1\text{H}$  NMR spectra, with increased magnetic field strength, Gruetter *et al.*, **135**, 260  
in MRI via enhanced signal digitization, Elliott *et al.*, **130**, 300  
in multiple-quantum MAS NMR, Schnell *et al.*, **133**, 61
- Resonance assignment  
active site of heme proteins,  $^1\text{H}$ - $^{13}\text{C}$ - $^{13}\text{C}$ -edited  $^1\text{H}$  NMR experiment for, Qiu, Rivera, and Stark, **130**, 76  
aromatic side chains in  $^{13}\text{C}$ -labeled proteins, 2D NMR experiments for, Prompers *et al.*, **130**, 68  
ligand bound to protein, heteronuclear X-filter  $^1\text{H}$  PFG double-quantum experiment for, Dalvit, Ramage, and Hommel, **131**, 148  
protons and carbons in uniformly  $^{13}\text{C}$ - and selectively  $^2\text{H}$ -labeled RNA by 3D C(CC)H TOCSY, Dayie, Tolbert, and Williamson, **130**, 97
- Resonance offset  
dependent partial saturation, in binomial solvent suppression sequences, Hsu and Gregory, **131**, 46
- Resonances  
second-order quadrupole broadened, NMR imaging, Swaminathan and Suits, **132**, 274
- Resonator  
and receiver, submicrosecond system for pulsed MR with large samples, Alecci *et al.*, **132**, 162  
220-MHz RF Fourier transform EPR spectrometer, Alecci *et al.*, **130**, 272
- Rhombicity  
unknown magnitude, direct structure refinement against residual dipolar couplings in presence of, Clore, Gronenborn, and Tjandra, **131**, 159
- Ribozyme  
lead-dependent, carbon-phosphorus  $J$  coupling constants, by spin-echo difference constant-time HCCH-COSY, Hoogstraten and Pardi, **133**, 236
- RIDE  
sequential inversion recovery with: simultaneous suppression of two solvent signals in  $^{17}\text{O}$  NMR, Schulte, **134**, 168
- Ripples  
Shinnar-Le Roux  $B$  polynomial, for RF pulse generation, generalized estimate for, Raddi and Klose, **132**, 260
- RNA  
carbon-phosphorus  $J$  coupling constants, by spin-echo difference constant-time HCCH-COSY, Hoogstraten and Pardi, **133**, 236  
larger oligonucleotides, optimization of triple-resonance HCN experiments for, Sklenář *et al.*, **130**, 119  
uniformly  $^{13}\text{C}$ - and selectively  $^2\text{H}$ -labeled, proton and carbon assignment, 3D C(CC)H TOCSY for, Dayie, Tolbert, and Williamson, **130**, 97
- Rock  
sedimentary, fluids contained, *in situ* NMR analysis, de Swiet *et al.*, **133**, 385
- Rotating-frame NQR imaging  
mapping molecular orientation in solids by, Casanova, Robert, and Pusiol, **133**, 129
- Rotation  
proteins in lipid bilayer systems, use of phase memory time  $T_2$  for quantitative characterization, van der Struijff and Levine, **130**, 244
- Rotational correlation time  
rapid computation from slow motional ESR spectra, neural network approach, Martinez and Millhauser, **134**, 124
- Rotational-echo double resonance (REDOR)  
signal and inversion, d'Espinose de la Caillerie and Fretigny, **133**, 273
- Rotational motion  
solute molecule in highly viscous liquid,  $^{13}\text{C}$  NMR, Lienin, Brünschweiler, and Ernst, **131**, 184
- Rotor synchronization  
amplitude-modulated cross-polarization with, application in  $^{13}\text{C}$ - $^1\text{H}$  spin pair under fast MAS, Raya and Hirschinger, **133**, 341
- Rovibrational effects  
rovibrationally averaged nuclear shielding constants in OCS, Jackowski, **135**, 444
- Ruby  
 $\gamma$ -irradiated crystal, impurity nuclear spin-lattice relaxation suppression and charge exchange of chromium ions, Chandoul *et al.*, **135**, 113  
spin- $\frac{3}{2}$   $\text{Cr}^{3+}$  in, looping transition, simulation methods for, Gaffney and Silverstone, **134**, 57

## S

 $^{33}\text{S}$ 

OCS, rovibrationally averaged nuclear shielding constants, Jackowski, **135**, 444

## Saccharides

long-range heteronuclear coupling constants, 1D determination by gradient-enhanced SIMBA, Vogt and Benesi, **132**, 214

## Sandimmune Neoral

effects on collagen arthritis in DA rat: high-resolution 3D MRI and histology, Beckmann *et al.*, **131**, 8

## Saturation

offset-dependent partial, in binomial solvent suppression sequences, Hsu and Gregory, **131**, 46

progressive saturation EPR determination of spin label relaxation times: molecular motion and Zeeman modulation effects, Livshits, Páli, and Marsh, **133**, 79

steady-state, of flowing spins, pulse sequences for, Marshall, **133**, 13

## Saturation transfer

detection of proton chemical exchange between metabolites and water in biological tissues, Guivel-Scharen *et al.*, **133**, 36

## Saturation-transfer ESR

sensitivity, extended to extremely slow mobility in glassy materials, van den Dries, de Jager, and Hemminga, **131**, 241

## Sea ice

restricted diffusion in, pulsed gradient spin echo NMR study: Earth's field NMR, Callaghan *et al.*, **133**, 148

## Secant pulse

STUD+ parameter calibration for efficient broadband adiabatic decoupling in single transient, Bendall and Skinner, **134**, 331

## sech/tanh pulse

STUD+ parameter calibration for efficient broadband adiabatic decoupling in single transient, Bendall and Skinner, **134**, 331

## Second moment calculation

effective spatial averaging for, Goc, **132**, 78

## Sedimentary rock

fluids in, *in situ* NMR analysis, de Swiet *et al.*, **133**, 385

## SEDOR

counting spins with, Cull *et al.*, **133**, 352

## Selective excitation

complicated shapes in three dimensions, Serša and Macura, **135**, 466

in-phase double, of coupled spin systems, using excitation sculpting, Bourg and Nuzillard, **133**, 173

ultraclean multisite, using excitation sculpting, Parella, Sánchez-Ferrando, and Virgili, **135**, 50

- Selective NMR  
 on macromolecules: QUIET–NOESY, Esposito *et al.*, **132**, 204
- Selective-pulse design  
 evolution strategy optimization for, Lunati *et al.*, **134**, 223
- Selective rotations  
 simultaneous and independent, with arbitrary flip angles and phases, for I, IS<sup>α</sup>, and IS<sup>β</sup> spin systems, Briand and Sørensen, **135**, 44
- Self-decoupling  
 heteronuclear scalar interaction and proton spin diffusion: high-speed <sup>13</sup>C MAS of adamantane, Ernst, Verhoeven, and Meier, **130**, 176
- Self-diffusion  
<sup>13</sup>C NMR measurements using RF field gradients, Malveau *et al.*, **130**, 131  
 and electric mobility, in disordered medium, NMR measurement, Heil and Holz, **135**, 17  
 and relaxation time NMR, modeling in multi-compartment systems, Novikov, van Dusschoten, and Van As, **135**, 522
- Semibullvalenes  
 valence tautomers, high-precision <sup>13</sup>C shift thermometer for temperature range 100–300 K based on, Quast *et al.*, **134**, 236
- Sensitivity enhancement  
 cardiac <sup>31</sup>P MRS in humans, Löffler *et al.*, **134**, 287  
 fast exchanging proton detection by exchange-edited gradient HCP-HSQC, Zangger and Armitage, **135**, 70  
 gradient-enhanced hetero ( $\omega_1$ ) half-filtered TOCSY for long-range heteronuclear coupling constants, Uhrín *et al.*, **130**, 155  
 gradient-enhanced TOCSY, Kövér, Uhrín, and Hruby, **130**, 162  
 HMQC, Zhu, Kong, and Sze, **135**, 232  
 INADEQUATE, Mattiello and Freeman, **135**, 514  
*in vivo* <sup>13</sup>C MR spectroscopy on clinical 1.5 T MR system, heteronuclear cross polarization for, van den Bergh, van den Boogert, and Heerschap, **135**, 93  
 by matched microwave pulses, in 1D and 2D ESEEM spectroscopy, Jeschke, Rakhmatullin, and Schweiger, **131**, 261  
 TROSY, Czisch and Boelens, **134**, 158
- Separated-local field experiments  
 ID solid-state NMR method for nuclear spin interaction tensors associated with peptide bond, Lee and Ramamoorthy, **133**, 204
- Serine proteinase PB92  
 TROSY, sensitivity enhancement, Czisch and Boelens, **134**, 158
- Shaped pulses  
 multiple-quantum MAS NMR of spin- $\frac{3}{2}$  quadrupolar spin systems, Ding and McDowell, **135**, 61
- Shaping  
 pulses in multiple-quantum MAS NMR, effect in spin- $\frac{3}{2}$  quadrupolar nuclear spin systems, Ding and McDowell, **135**, 61
- Shearing  
 distributions of quadrupolar and isotropic chemical shift interactions in 2D multiple-quantum MAS NMR spectra following, Bodart, **133**, 207
- Shielding constants  
 rovibrationally averaged, in OCS, Jackowski, **135**, 444
- Shieldings  
<sup>57</sup>Fe, in superstructured hemoprotein models, effects of atropisomerism and porphyrin deformation, Gerotheranassis *et al.*, **131**, 163  
 NMR, Be<sup>+2</sup> and Al<sup>+3</sup> species in aqueous solution, effects of hydrolysis and oligomerization, Tossell, **135**, 203  
 rovibrationally averaged nuclear shielding constants in OCS, Jackowski, **135**, 444
- Shimming  
 automated, with normal spectrometer hardware: 3D profile edge shimming, Chilvers and Morris, **133**, 210
- Shinnar–Le Roux algorithm  
 SLR *B* polynomial ripples for RF pulse generation, generalized estimate for, Raddi and Klose, **132**, 260
- <sup>29</sup>Si  
 surface selective <sup>1</sup>H/<sup>29</sup>Si CP NMR by NOE enhancement from laser-polarized xenon, Pietraß, Seydoux, and Pines, **133**, 299
- Sidebands  
 cyclic, elimination in double-adiabatic homonuclear decoupling for <sup>13</sup>C- and <sup>15</sup>N-double-labeled proteins, Zhang and Gorenstein, **132**, 81
- Side chains  
 protein  
 contacting DNA, enhancement of weak labile <sup>1</sup>H resonances: <sup>15</sup>N-edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87  
 motional dynamics, analysis from <sup>13</sup>C NMR relaxation data, Daragan and Mayo, **130**, 329
- Side groups  
 specific, role in magnetization transfer in polymers, Gochberg *et al.*, **131**, 191
- Signal  
 NMR, generalized reciprocity and, Insko *et al.*, **131**, 111  
 transient MR, super-Q detection, Suits, Garroway, and Miller, **132**, 54
- Signal intensity  
 optimization with nonstandard flip angles in spin-echo and inversion-recovery MR imaging, Bonny *et al.*, **130**, 51
- Signal-to-noise ratio  
 MRI, improvement via enhanced signal digitization, Elliott *et al.*, **130**, 300
- Signal processing (*see* Data processing)
- Silica aerogels  
 pore-structure determination by <sup>129</sup>Xe NMR and imaging, Gregory, Gerald, and Botto, **131**, 327
- SIMBA  
 gradient-enhanced, 1D determination of long-range heteronuclear coupling constants by, Vogt and Benesi, **132**, 214
- Simulated annealing  
 direct structure refinement against residual dipolar couplings in presence of rhombicity of unknown magnitude, Clore, Gronenborn, and Tjandra, **131**, 159
- Simulation  
 coherence transfer pathway selection by phase cycling and pulsed-field gradients in NMR, Jerschow and Müller, **134**, 17  
 CP-MAS signals at high spinning speeds, Ray, Ladizhansky, and Vega, **135**, 427  
 ESEEM spectra using Gamma, Shane, Liesum, and Schweiger, **134**, 72  
 looping transitions, methods for, Gaffney and Silverstone, **134**, 57  
 MAS NMR, time propagation technique for, application to quadrupolar nuclei, Charpentier, Fermon, and Virlet, **132**, 181  
 MR spectra, employing homotopy, Gates *et al.*, **135**, 104  
 NMR, homogeneous master equation for heteronuclear two-spin system on basis of Cartesian product operators, Allard, Helgstrand, and Hård, **134**, 7  
 spectral, using Floquet theory: proton MAS NMR at high magnetic fields and high spinning frequencies, Ray *et al.*, **135**, 418  
 spectral lineshape: iterative lineshape fitting of MAS NMR spectra, Dusold, Milius, and Sebald, **135**, 500
- Single-crystal NMR spectroscopy  
 improved hardware and software for, Vosegaard *et al.*, **135**, 126
- Single-crystal 2D NQR spectroscopy  
 off-resonance effects, Ajithkumar *et al.*, **135**, 165
- Single-point acquisition with relaxation enhancement (SPARE)  
 robust method for MRI in inhomogeneous fields, McIntyre, Hennel, and Morris, **130**, 58
- Single-point imaging  
 relaxation time mapping of short *T*<sub>2</sub><sup>\*</sup> nuclei, Beyea *et al.*, **135**, 156
- Singular value decomposition  
 suppression of diagonal peaks with, Zhe *et al.*, **132**, 176
- SLOOP  
 localized <sup>31</sup>P NMR of human heart, Löffler *et al.*, **134**, 287

- Slow-magic-angle-sample-spinning conditions  
polarization transfer under, comparison with RF-driven conditions, Robyr and Gan, **131**, 254
- Slow motional spectra  
rapid computation of rotational correlation time from, neural network approach, Martinez and Millhauser, **134**, 124
- Small loop antenna  
surface and gradiometer coils near conducting body: lift-off effect, Suits, Garroway, and Miller, **135**, 373
- Smoothing penalty  
uniform-penalty inversion of multiexponential decay data, Borgia, Brown, and Fantazzini, **132**, 65
- Sodium  
cation, binding-induced changes in  $^{15}\text{N}$  CSA in membrane-bound polypeptide, Tian and Cross, **135**, 535  
extracellular, contribution to  $^{23}\text{Na}$  multiple-quantum-filtered NMR spectra of erythrocyte suspensions, Knubovets, Shinar, and Navon, **131**, 92
- Software  
single-crystal NMR, Vosegaard *et al.*, **135**, 126
- Solenoids  
microcoils, design for NMR of mass-limited samples, Subramanian, Lam, and Webb, **133**, 227
- Solids  
hydrogen-bonded,  $^1\text{H}$  and  $^2\text{H}$  NMR relaxation due to complex motion: classical jumps over barrier and incoherent tunneling, Reynhardt and Latanowicz, **130**, 195  
mapping molecular orientation by rotating-frame NQR imaging, Casanova, Robert, and Pusiol, **133**, 129  
quadrupolar nuclei in, INEPT experiments involving, Kao and Grey, **133**, 313  
stray field imaging by magnetic field sweep, Mallett, Halse, and Strange, **132**, 172  
surface NMR using laser-polarized  $^{129}\text{Xe}$  under MAS conditions, Brunner *et al.*, **130**, 145
- Solid-state imaging  
 $^1\text{H}$  NMR image contrast, cross polarization for, Nakai *et al.*, **134**, 44
- Solid-state NMR  
 $^{13}\text{C}$  and  $^{15}\text{N}$ , analysis of chemical-shift anisotropy of ampicillin and penicillin-V by 2D-PASS and CP/MAS NMR, Antzutkin, Lee, and Levitt, **135**, 144  
counting spins with SEDOR, Cull *et al.*, **133**, 352  
definitive example of  $^{27}\text{Al}$  chemical-shielding anisotropy, Schurko, Wasylshen, and Phillips, **133**, 388  
 $^2\text{H}$ , noise spectroscopy application to, bandwidth extension, Yang and Zax, **135**, 267  
1D, nuclear spin interaction tensors associated with peptide bond, Lee and Ramamoorthy, **133**, 204  
orientational averages, computation by Gaussian spherical quadrature, Edén and Levitt, **132**, 220  
paramagnetic compounds, effect of bulk magnetic susceptibility, Kubo, Spaniol, and Terao, **133**, 330  
pulsed-gradient NMR probes for studies, Zhang and Cory, **132**, 144  
 $^{29}\text{Si}$ : surface selective  $^1\text{H}/^{29}\text{Si}$  CP NMR by NOE enhancement from laser-polarized xenon, Pietraß, Seydoux, and Pines, **133**, 299  
spin-echo mapping spectroscopy applied to NQR, Bussandri and Zuriaga, **131**, 224
- Solute molecule  
rotational motion in highly viscous liquid,  $^{13}\text{C}$  NMR, Lienin, Brüscheweiler, and Ernst, **131**, 184
- Solute-solvent interactions  
fullerenes  $\text{C}_{60}$  and  $\text{C}_{70}$  in solution: 1D steady-state intermolecular  $^{13}\text{C}$ ,  $^1\text{H}$  Overhauser effect study, Troganis, Gerothanassis, and Papademitriou, **130**, 149
- Solution NMR  
direct structure refinement against residual dipolar couplings in presence of rhombicity of unknown magnitude, Clore, Gronenborn, and Tjandra, **131**, 159  
hyperfine shifts, high magnetic field consequences, Bertini, Felli, and Luchinat, **134**, 360  
multi-component spin systems, intermolecular zero-quantum coherences, Ahn, Lisitza, and Warren, **133**, 266  
structure determination, use of magnitude of asymmetric alignment tensor of oriented macromolecules, Clore, Gronenborn, and Bax, **133**, 216
- Solutions  
dilute bicellar, for structural NMR work, Struppe and Vold, **135**, 541  
glycerol, dipolar interactions of  $^{23}\text{Na}$ , determination by triple-quantum relaxation time measurements, Eliav and Navon, **130**, 63
- Solvent  
large signals, analog filtering for improved dynamic range in high-resolution NMR, Redfield and Kunz, **130**, 111  
protons, measurement of chemical exchange rate constants with, using radiation damping, Chen and Mao, **131**, 358
- Solvent effects  
solvent magnetization artifacts in high-field NMR studies of macromolecular hydration, Sobol *et al.*, **130**, 262
- Solvent polarity effects  
on nitrogen NMR shielding of isomeric tetrazoles, Witanowski *et al.*, **131**, 54
- Solvent suppression  
binomial sequences for, offset-dependent partial saturation, Hsu and Gregory, **131**, 46  
H(C)CH-COSY and (H)CCH-COSY for  $^{13}\text{C}$ -labeled proteins in  $\text{H}_2\text{O}$  solution, Gehring and Ekiel, **135**, 185  
in NMR, improved WATERGATE pulse sequences for, Liu *et al.*, **132**, 125  
simultaneous, two solvent signals in  $^{17}\text{O}$  NMR: sequential inversion recovery with RIDE, Schulte, **134**, 168
- SPARC-16  
broadband decoupling sequence for liquid crystals, Yu and Fung, **130**, 317
- SPARE  
robust method for MRI in inhomogeneous fields, McIntyre, Hennel, and Morris, **130**, 58
- Spatial averaging  
effective, for NMR second moment calculation, Goc, **132**, 78
- Spatial correlation  
2D CP/MAS  $^{13}\text{C}$  isotropic chemical shift correlation established by  $^1\text{H}$  spin diffusion, Wilhelm *et al.*, **134**, 255
- Spatial resolution  
field-gradient NMR, demonstrating, Feiweir *et al.*, **131**, 203
- Spectral analysis  
symmetry and phase-selected NMR spectra of liquid crystalline samples, Carravetta *et al.*, **135**, 298
- Spectral density mapping  
in DNA dynamics study via  $^{13}\text{C}$  relaxation, Kojima *et al.*, **135**, 310
- Spectral editing (*see* Editing)
- Spectral lines  
precision in estimating frequency separation between, Jupp, Harris, and Aliev, **135**, 23
- Spectral localization with optimal pointspread function (SLOOP)  
localized  $^{31}\text{P}$  NMR of human heart, Löffler *et al.*, **134**, 287
- Spectral parameter estimation  
by iterative quadratic maximum likelihood method, Zhu, Choy, and Sanctuary, **135**, 37
- Spectral simulation  
using Floquet theory: proton MAS NMR at high magnetic fields and high spinning frequencies, Ray *et al.*, **135**, 418
- Spectral spatial EPR imaging  
implementation by iterative Fourier reconstruction, Placidi *et al.*, **134**, 280

- Spectrometer (*see also* NMR spectrometer)  
 radiofrequency Fourier transform EPR, 220-MHz, Alecci *et al.*, **130**, 272
- Spectrum analysis  
 distributed parallel processing for multidimensional maximum entropy reconstruction, Li, Stern, and Hoch, **134**, 161
- Sphere packings  
 application of pulsed-field gradient NMR technique for determination of structure of suspensions of non-Brownian particles, Talini, Leblond, and Feuillebois, **132**, 287
- Spin diffusion  
 interresidue quiet NOE for suppression, for DNA structural studies, Beger *et al.*, **132**, 34  
 proton  
 and heteronuclear scalar interaction, self-decoupling: high-speed  $^{13}\text{C}$  MAS of adamantane, Ernst, Verhoeven, and Meier, **130**, 176  
 2D CP/MAS  $^{13}\text{C}$  isotropic chemical shift correlation established by, Wilhelm *et al.*, **134**, 255
- Spin echo  
 adiabatic multiple spin-echo pulse sequence, Zweckstetter and Holak, **133**, 134  
*in vivo* observation of multiple spin echoes, Bifone, Payne, and Leach, **135**, 30  
 $^{14}\text{N}$  NMR, in  $\text{NH}_4\text{ClO}_4$  after pulse sequence  $(\theta_1)_x - \tau - (\theta_2)_y - t$ , Punkkinen *et al.*, **130**, 287  
 pulsed gradient (*see* Pulsed gradient spin echo)
- Spin-echo attenuation  
 of restricted diffusion, as discord of structure waves, Stepišnik, **131**, 339
- Spin-echo difference constant-time HCCH-COSY  
 measurement of carbon-phosphorus  $J$  coupling constants in RNA, Hoogstraten and Pardi, **133**, 236
- Spin echo double resonance (SEDOR)  
 counting spins with, Cull *et al.*, **133**, 352
- Spin-echo imaging  
 signal intensity and  $T_1$ -dependent contrast, optimization with nonstandard flip angles, Bonny *et al.*, **130**, 51
- Spin-echo mapping spectroscopy  
 applied to NQR, Bussandri and Zuriaga, **131**, 224
- Spin labels  
 nitroxide (*see* Nitroxide spin labels)  
 TEMPOL, saturation-transfer ESR spectra in glycerol, van den Dries, de Jager, and Hemminga, **131**, 241
- Spin-lattice relaxation  
 $^{27}\text{Al}$  in  $\gamma$ -irradiated ruby crystal, separation of impurity and lattice contributions and determination of charge-exchanged chromium ions, Chandoul *et al.*, **135**, 113  
 dependent contrast, optimization with nonstandard flip angles in spin-echo and inversion-recovery MR imaging, Bonny *et al.*, **130**, 51  
 effects in sequential dynamic susceptibility contrast experiments, Levin *et al.*, **130**, 292  
 $^1\text{H}$  and  $^2\text{H}$ , in hydrogen-bonded solids, due to complex motion: classical jumps over barrier and incoherent tunneling, Reynhardt and Latanowicz, **130**, 195  
 $^7\text{Li}$ , magnetic relaxation dispersion technique: complex formation with nitroxides in aqueous phase, Dinesen and Bryant, **132**, 19  
 measurement  
 continuous-wave nonlinear ESR spectroscopies, Cianflone, Francia, and Leporini, **131**, 86  
 first-harmonic out-of-phase absorption EPR signals, Livshits, Páli, and Marsh, **134**, 113  
 in multivariate image analysis with direct exponential curve resolution algorithm  
 algorithm and model study, Windig, Hornak, and Antalek, **132**, 298  
 human brain images, Antalek, Hornak, and Windig, **132**, 307  
 $^{15}\text{N}$  and  $^{13}\text{CO}$  in poorly resolved NMR spectra, 3D accordion spectroscopy measurement, Carr, Fearing, and Palmer, **132**, 25  
 in 99%  $^{13}\text{C}$ -enriched diamonds,  $^{13}\text{C}$  NMR analysis, Shabanova, Schaumburg, and Sellschop, **130**, 8  
 order-fluctuation modes in liquid crystals, formalism for evaluation from field-cycling NMR-relaxometry data, Grinberg *et al.*, **135**, 54  
 and proton diffusion, in polyacrylamide gels, unified approach using volume averaging, Penke *et al.*, **132**, 240  
 role in echo decay of planar Cu in high-temperature superconductors, Curro and Slichter, **130**, 186  
 short, mapping of nuclei using single-point imaging methods, Beyea *et al.*, **135**, 156  
 spin labels, determination by progressive saturation EPR, molecular motion and Zeeman modulation effects, Livshits, Páli, and Marsh, **133**, 79
- Spin-locking  
 3D NMR pulse sequences for characterization of polymer chain end structures, Saito and Rinaldi, **130**, 135
- Spinning frequency  
 high, proton MAS NMR: spectral simulations using Floquet theory, Ray *et al.*, **135**, 418
- Spinning sidebands  
 control in high-resolution NMR, Borer and Maple, **131**, 177
- Spinning speed  
 high, simulation of CP-MAS signals at, Ray, Ladizhansky, and Vega, **135**, 427  
 and RF power, effect on MQMAS NMR, Amoureux *et al.*, **131**, 170
- Spin pairs  
 $^{13}\text{C}$ - $^1\text{H}$ , under fast MAS, application of rotor-synchronized amplitude-modulated cross polarization, Raya and Hirschinger, **133**, 341  
 isolated, homonuclear  $J$  coupling, iterative lineshape fitting of MAS NMR spectra for analysis, Dusold, Milius, and Sebald, **135**, 500
- Spin polarization  
 hyperpolarized gases, low-frequency NMR polarimeter for, Saam and Conrad, **134**, 67
- Spin polarization-induced NOE  
 in surface NMR using laser-polarized  $^{129}\text{Xe}$  under MAS conditions, Brunner *et al.*, **130**, 145  
 surface selective  $^1\text{H}/^{29}\text{Si}$  CP NMR by NOE enhancement from laser-polarized xenon, Pietraß, Seydoux, and Pines, **133**, 299
- Spins  
 $\frac{1}{2}$  coupled, under planar mixing conditions, analytical polarization and coherence transfer functions for, Schedletzky, Luy, and Glaser, **130**, 27  
 counting with SEDOR, Cull *et al.*, **133**, 352  
 flowing fluid, pulse sequences for steady-state saturation, Marshall, **133**, 13  
 half-integer quadrupolar, MQMAS NMR, composite pulse excitation schemes for, Marinelli, Medek, and Frydman, **132**, 88  
 $I = \frac{5}{2}$ , five-quantum coherence of nuclei with:  $^{27}\text{Al}$  in  $\text{Al}_2\text{O}_3$  single crystal, Kaikkonen, Ylinen, and Punkkinen, **133**, 123  
 inequivalent, nonlinear stimulated echo in presence of, Ardelean *et al.*, **132**, 138  
 $\frac{3}{2}$  nuclei of incompletely disordered systems with, temporal characteristics of NMR signals, Woessner and Bansal, **133**, 21
- Spin-spin coupling  
 $^{27}\text{Al} \rightarrow ^{31}\text{P}$  INEPT experiments on  $(\text{CH}_3)_3\text{P}-\text{AlCl}_3$  complex in zeolite NaX, Kao and Grey, **133**, 313  
 carbon-phosphorus, by spin-echo difference constant-time HCCH-COSY, Hoogstraten and Pardi, **133**, 236  
 deuterium quadrupole coupling constant in liquids, and dynamic isotope effect, Hardy *et al.*, **134**, 300  
 estimation from antiphase multiplets: Metropolis Monte Carlo implementation of Bayesian time-domain parameter estimation, Andrec and Prestegard, **130**, 217

- heteronuclear, simultaneous and independent rotations with arbitrary flip angles and phases for I, IS<sup>α</sup>, and IS<sup>β</sup> spin systems, Briand and Sørensen, **135**, 44
- heteronuclear long-range  
 gradient-enhanced 2D multiple-bond correlation experiment for, Sheng and van Halbeek, **130**, 296  
 1D determination by gradient-enhanced SIMBA, Vogt and Benesi, **132**, 214  
 sensitivity- and gradient-enhanced hetero ( $\omega_1$ ) half-filtered TOCSY for, Uhrin *et al.*, **130**, 155
- homonuclear  
 active-coupling-pattern tilting experiment for, Koźmiński, **134**, 189  
 in isolated spin pairs, iterative lineshape fitting of MAS NMR spectra for, Dusold, Milius, and Sebald, **135**, 500  
 $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for spin-state selective editing of IS cross peaks, Andersson, Annila, and Otting, **133**, 364  
<sup>3</sup>J<sub>NH $\alpha$ , in analysis of conformational effect of nitroxide spin labels on Ala-rich helical peptides, Bolin *et al.*, **131**, 248  
 measurement from simplified 2D NMR spectra, Ottiger, Delaglio, and Bax, **131**, 373  
 nitrogen–nitrogen, in proteins, detection, Löhr and Rüterjans, **132**, 130  
<sup>31</sup>P, <sup>63,65</sup>Cu indirect, and <sup>31</sup>P nuclear shielding tensor, of linear copper(I) phosphines, anisotropy, Kroeker *et al.*, **135**, 208  
 scalar, between protons attached to <sup>13</sup>C, methylene specific experiments, Carlomagno *et al.*, **135**, 216  
 three-bond <sup>1</sup>H3'-<sup>31</sup>P couplings in nucleic acids and protein–nucleic acid complexes: quantitative J correlation spectroscopy, Clore *et al.*, **134**, 164  
 vicinal fluorine–proton coupling constants: individual substituent effects, Fabián, Guilleme, and Díez, **133**, 255</sub>
- Spin–spin relaxation  
 echo-planar imaging measurement of viscosity of model meal, Marciani *et al.*, **135**, 82  
 imido protons of coordinated imidazoles in uniformly <sup>15</sup>N-labeled cytochrome *c*<sub>3</sub>, paramagnetic inversion of sign of interference contribution to, Ohmura *et al.*, **131**, 367  
 measurement with multiecho imaging with suboptimal spoiler gradients, Does and Snyder, **131**, 25  
 in multivariate image analysis with direct exponential curve resolution algorithm  
 algorithm and model study, Windig, Hornak, and Antalek, **132**, 298  
 human brain images, Antalek, Hornak, and Windig, **132**, 307  
 natural-abundance crystalline carbons of linear polyethylene under CW proton decoupling conditions and MAS, VanderHart and Campbell, **134**, 88  
 removal of systematic errors with adiabatic multiple spin-echo pulse sequence, Zweckstetter and Holak, **133**, 134  
 R<sub>2</sub>' measurement in presence of multiple spectral components using reference spectrum deconvolution, Wehrl *et al.*, **131**, 61  
 short, mapping of nuclei using single-point imaging methods, Beyea *et al.*, **135**, 156  
 spin  $\frac{3}{2}$  nuclei of incompletely disordered systems, Woessner and Bansal, **133**, 21
- Spin-state selective editing  
 IS cross peaks,  $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for, Andersson, Annila, and Otting, **133**, 364
- Spin systems  
 AA'X, evolution of zero-quantum coherence: analysis of catalytic processes using parahydrogen, Natterer *et al.*, **133**, 92  
 coupled, in-phase double selective excitation using excitation sculpting, Bourg and Nuzillard, **133**, 173  
 I, IS<sup>α</sup>, and IS<sup>β</sup>, simultaneous and independent rotations with arbitrary flip angles and phases for, Briand and Sørensen, **135**, 44  
 IS, I<sub>2</sub>S, and I<sub>3</sub>S, effects of adiabatic pulses, analytical description, Zwaalen, Vincent, and Kay, **130**, 169  
 multi-component, in solution NMR, intermolecular zero-quantum coherences, Ahn, Lisitza, and Warren, **133**, 266  
 spin- $\frac{3}{2}$  quadrupolar, multiple-quantum MAS NMR using shaped pulses, Ding and McDowell, **135**, 61
- Split evolution time  
<sup>13</sup>C-NOESY-HSQC with, for increased resolution with uniformly labeled proteins, Baur, Gemmecker, and Kessler, **132**, 191
- Split-loop resonator  
 noise-immune coil for unshielded magnetic resonance measurements, Suits, Garroway, and Miller, **131**, 154
- Spoiler gradients  
 suboptimal, MRI with, Does and Snyder, **131**, 25
- Staphylococcus aureus*  
 gyrase B,  $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for spin-state selective editing of IS cross peaks, Andersson, Annila, and Otting, **133**, 364
- Steady-state saturation  
 of flowing spins, pulse sequences for, Marshall, **133**, 13
- STEAM  
 and PRESS, flip angle effects—optimized vs sinc RF pulses, Ryner, Ke, and Thomas, **131**, 118
- Stimulated echo  
 four-time, experimental determination in liquids, colloidal suspensions, and crystals, Hinze *et al.*, **131**, 218  
<sup>2</sup>H NMR, time domain analysis of ultraslow reorientations in supercooled liquids, Geil, Fujara, and Sillescu, **130**, 18  
 nonlinear, in presence of inequivalent spins, Ardelean *et al.*, **132**, 138  
 polycrystalline benzene, comparison with 2D deuteron exchange NMR: molecular dynamics of vacancy diffusion, Isfort, Geil, and Fujara, **130**, 45
- Stimulated echo acquisition mode (STEAM)  
 and PRESS, flip angle effects—optimized vs sinc RF pulses, Ryner, Ke, and Thomas, **131**, 118
- Stray field imaging  
 by magnetic field sweep, Mallett, Halse, and Strange, **132**, 172
- Structure  
 analysis of error propagation from NMR-derived internuclear distances into molecular structure of *cyclo*-Pro-Gly, Džakula *et al.*, **135**, 454  
 direct refinement against residual dipolar couplings in presence of rhombicity of unknown magnitude, Clore, Gronenborn, and Tjandra, **131**, 159  
 DNA, interresidue quiet NOE for, Beger *et al.*, **132**, 34  
 information, determination of magnitude of asymmetric alignment tensor of oriented macromolecules in absence of, Clore, Gronenborn, and Bax, **133**, 216  
 orientational constraints derived from hydrated powder samples by 2D PISEMA, Tian, Song, and Cross, **135**, 227  
 polymer, 3D NMR experiments, Saito and Rinaldi, **132**, 41  
 suspensions of non-Brownian particles, PFG NMR technique with applications to sphere packings, Talini, Leblond, and Feuillebois, **132**, 287  
 weakly oriented macromolecules, using measured dipolar couplings for methylene and methyl sites, Ottiger *et al.*, **134**, 365
- Structure waves  
 discord, spin-echo attenuation of restricted diffusion as, Stepišnik, **131**, 339
- STUD+  
 calibration of parameters, for efficient broadband adiabatic decoupling in single transient, Bendall and Skinner, **134**, 331
- Subcutaneous injection  
 carboplatin, disposition kinetics monitoring using <sup>195</sup>Pt NMR following, Becker *et al.*, **133**, 115
- Substituent effects  
 on vicinal fluorine–proton coupling constants, Fabián, Guilleme, and Díez, **133**, 255

- Sucrose  
long-range heteronuclear coupling constants, 1D determination by gradient-enhanced SIMBA, Vogt and Benesi, **132**, 214
- Superconductors  
high-temperature, echo decay of planar Cu, contributions of spin-lattice relaxation, Curro and Slichter, **130**, 186
- Superparamagnetic particles  
intravascular and intracellular hepatic relaxivities, Colet *et al.*, **134**, 199
- Super-Q detection  
transient MR signals, Suits, Garroway, and Miller, **132**, 54
- Surface coils  
and gradiometer coils, near conducting body: lift-off effect, Suits, Garroway, and Miller, **135**, 373  
noise-immune, for unshielded magnetic resonance measurements, Suits, Garroway, and Miller, **131**, 154
- Surface groups  
specific, role in magnetization transfer in polymers, Gochberg *et al.*, **131**, 191
- Surface liquid interactions  
molecular diffusion and DNP enhancement in aqueous char suspensions, Odintsov *et al.*, **135**, 435
- Surface NMR  
analysis of polymer materials via MOUSE, Guthausen *et al.*, **130**, 1  
using laser-polarized  $^{129}\text{Xe}$  under MAS conditions, Brunner *et al.*, **130**, 145
- Surfactant  
-water contact,  $^{19}\text{F}$ - $^1\text{H}$  HOESY study, Raulet *et al.*, **133**, 324
- Susceptibility (see Magnetic susceptibility)
- Suspensions  
aqueous, char, molecular diffusion and DNP enhancement, Odintsov *et al.*, **135**, 435  
non-Brownian particles, structure determination, PFG NMR technique with applications to sphere packings, Talini, Leblond, and Feuillebois, **132**, 287
- Sweep coil  
additional, stray field imaging, Mallett, Halse, and Strange, **132**, 172
- Symmetry  
and phase, simplification of NMR spectra of liquid crystalline samples by, Carravetta *et al.*, **135**, 298
- Symposium  
The Wallace Brey Symposium, November 2, 1997, **134**, 197
- T
- $T_1$  (see Spin-lattice relaxation)  
 $T_2$  (see Spin-spin relaxation)  
 $T_2^*$   
short, nuclei with, relaxation time mapping with single-point imaging methods, Beyea *et al.*, **135**, 156
- Tangential frequency sweep  
adiabatic pulses with, fast broadband inversion by, Hwang, Zijl, and Garwood, **133**, 200
- Taurine  
detection *in vitro* and *in vivo*, spectral editing technique for, Hardy and Norwood, **133**, 70
- Temperature dependence  
rovibrationally averaged nuclear shielding constants in OCS, Jackowski, **135**, 444
- Temperature imaging  
by  $^1\text{H}$  NMR, and suppression of convection, in NMR probes, Hedin and Furó, **131**, 126
- Temperature measurement  
high-precision  $^{13}\text{C}$  shift NMR, for range 100–300 K, Quast *et al.*, **134**, 236  
NMR, using paramagnetic lanthanide complex, Zuo *et al.*, **133**, 53
- TEMPOL (see 4-Hydroxy-2,2,6,6-tetramethyl-1-piperidinyloxy)
- Temporal analysis  
NMR signals from spin  $\frac{3}{2}$  nuclei of incompletely disordered systems, Woessner and Bansal, **133**, 21
- Tensors  
asymmetric alignment, of oriented macromolecules, magnitude determination in absence of structural information, Clore, Gronenborn, and Bax, **133**, 216  
 $^{15}\text{N}$  chemical shift and  $^{13}\text{C}$ - $^1\text{H}$  dipolar, correlation in solid-state MAS NMR, peptide torsion angle determination by, Hong *et al.*, **135**, 169  
nitrogen quadrupole and hyperfine, determination in oxovanadium complex by simultaneous fitting of multifrequency ESEEM powder spectra, Reijerse, Tyryshkin, and Dikanov, **131**, 295  
nuclear spin interaction, associated with peptide bond, 1D solid-state NMR method for, Lee and Ramamoorthy, **133**, 204  
 $^{31}\text{P}$ ,  $^{63,65}\text{Cu}$  indirect spin-spin coupling and  $^{31}\text{P}$  nuclear shielding, of linear copper(I) phosphines, anisotropy, Kroeker *et al.*, **135**, 208
- ortho*-Terphenyl  
supercooled melt, ultraslow reorientations,  $^2\text{H}$  NMR time domain analysis, Geil, Fujara, and Sillescu, **130**, 18
- Tetrazoles  
isomeric, nitrogen NMR shielding, solvent polarity and hydrogen bonding effects, Witanowski *et al.*, **131**, 54
- Three-dimensional NMR  
accordion spectroscopy measurement of  $^{15}\text{N}$  and  $^{13}\text{CO}$  relaxation rates in poorly resolved NMR spectra, Carr, Fearing, and Palmer, **132**, 25  
heteronuclear, correlation of backbone amide and aliphatic gamma resonances in deuterated proteins, McCallum, Hitchens, and Rule, **134**, 350  
*in vivo* localized  $^{13}\text{C}$ , using modified INEPT and DEPT, Watanabe *et al.*, **134**, 214  
 $^{15}\text{N}$ -edited NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87  
for  $^{15}\text{N}$  relaxation of large proteins: 44-kDa ectodomain of SIV gp41, Caffrey *et al.*, **135**, 368  
polymer structure characterization, Saito and Rinaldi, **132**, 41  
pulse sequences, for characterization of polymer chain end structures, Saito and Rinaldi, **130**, 135
- Three-dimensional NMR imaging  
excitation of complicated shapes, Serša and Macura, **135**, 466  
high-resolution, for Sandimmune Neoral effects on collagen arthritis in DA rat, Beckmann *et al.*, **131**, 8
- Thulium 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetrakis(methylene phosphonate) complex  
NMR temperature measurements, Zuo *et al.*, **133**, 53
- TIG-BIRD  
pulse sequence element for simultaneous and independent rotations for I,  $\text{IS}^\alpha$ , and  $\text{IS}^\beta$  spin systems, Briand and Sørensen, **135**, 44
- Time domain analysis  
 $^2\text{H}$  NMR, ultraslow reorientations in supercooled liquids, Geil, Fujara, and Sillescu, **130**, 18  
NMR signals from spin  $\frac{3}{2}$  nuclei of incompletely disordered systems, Woessner and Bansal, **133**, 21
- Time-domain EPR  
high-spin iron(III)-nitroxyl distances in spin-labeled porphyrins, Rakowsky *et al.*, **131**, 97
- Time-domain parameter estimation  
Bayesian, Metropolis Monte Carlo implementation, application to coupling constant estimation from antiphase multiplets, Andrec and Prestegard, **130**, 217
- Time propagation technique  
for MAS NMR simulation, application to quadrupolar nuclei, Charpentier, Fermon, and Virlet, **132**, 181
- Time-reverse ODESSA  
direct determination of motional correlational times, Favre, Schaefer, and Chmelka, **134**, 261



- <sup>119</sup>Tin  
*para*-substituted triaryltin(pentacarbonyl)manganese(I) complexes, solid-state CP–MAS NMR spectra, quadrupole–dipole effects, Christendat *et al.*, **131**, 1
- TmDOTP<sup>5-</sup> complex  
 NMR temperature measurements, Zuo *et al.*, **133**, 53
- TOCSY  
 adiabatic mixing in liquid state, Kupče *et al.*, **135**, 361  
 cross peak detection in 2D proton NMR spectra using excitation sculpting, Van and Shaka, **132**, 154  
 selective 1D and nonselective 2D, coherence selection and excitation sculpting using RF-gradients, Heikkinen, Rahkamaa, and Kilpeläinen, **133**, 183  
 sensitivity- and gradient-enhanced, Kövér, Uhrín, and Hruby, **130**, 162  
 sensitivity- and gradient-enhanced hetero ( $\omega_1$ ) half-filtered, for long-range heteronuclear coupling constants, Uhrín *et al.*, **130**, 155  
 systems with exchangeable protons, information from water stripe, Nakashima, McClung, and Kotovych, **133**, 222  
 2D C(CC)H, assignment of protons and carbons in uniformly <sup>13</sup>C- and selectively <sup>2</sup>H-labeled RNA, Dayie, Tolbert, and Williamson, **130**, 97
- Torsion angle  
 peptide, determination by <sup>15</sup>N chemical shift and <sup>13</sup>C $\alpha$ –<sup>1</sup>H $\alpha$  dipolar tensor correlation in solid-state MAS NMR, Hong *et al.*, **135**, 169
- Total correlation spectroscopy (*see* TOCSY)
- Total least squares  
 spectral parameter estimation by, comparison with iterative quadratic maximum likelihood and linear prediction methods, Zhu, Choy, and Sanctuary, **135**, 37
- TQF NMR  
 relaxation time measurements by, in determination of dipolar interactions of <sup>23</sup>Na in solution, Eliav and Navon, **130**, 63
- Transferrin oxalate  
 spin- $\frac{5}{2}$  Fe<sup>3+</sup> in, looping transition, simulation methods for, Gaffney and Silverstone, **134**, 57
- Transient decay  
 magnetization transfer contrast of rat brain, Quesson *et al.*, **130**, 321
- Transient MR signals  
 super-Q detection, Suits, Garroway, and Miller, **132**, 54
- Transition-band signals  
 in high-resolution NMR, Szántay, **135**, 334
- Transport  
 carbohydrate in plants, cyclic *J* cross-polarization imaging and spectroscopy for analysis, Heidenreich *et al.*, **132**, 109
- Transverse relaxation (*see* Spin–spin relaxation)
- Triaryltin(pentacarbonyl)manganese(I) complexes  
*para*-substituted, solid-state CP–MAS <sup>119</sup>Tin NMR spectra, quadrupole–dipole effects, Christendat *et al.*, **131**, 1
- Triphenylmethyl  
 single-electron contrast agents for oximetric imaging based on, EPR and DNP properties, Ardenkjær-Larsen *et al.*, **133**, 1
- Triple-quantum-filtered (TQF) NMR  
 relaxation time measurements: dipolar interactions of <sup>23</sup>Na in solution, Eliav and Navon, **130**, 63
- Triple resonance  
 amino acid-selective experiments, multiplicity-selective coherence transfer steps for design, Schmieder *et al.*, **131**, 199  
 3D and 4D experiments, improved resolution from double constant-time evolution, McCoy, **130**, 341  
 HCN, optimization for application to larger RNA oligonucleotides, Sklenář *et al.*, **130**, 119
- Triplet spectra  
 in high-frequency EPR study of integer spin system, unusual behavior of double-quantum line, van Dam *et al.*, **130**, 140
- Triplet state  
<sup>13</sup>C<sub>60</sub>, pulsed ENDOR studies at 95 GHz, van den Berg *et al.*, **131**, 39  
 photoexcited, acridine-*d*<sub>9</sub> in fluorene-*h*<sub>10</sub> matrix in zero magnetic field, optically detected ESEEM, Weis, Möbius, and Prisner, **131**, 17
- Trisaccharide  
 gradient- and sensitivity-enhanced TOCSY, Kövér, Uhrín, and Hruby, **130**, 162
- TROSY  
 $\alpha/\beta$ -HSQC- $\alpha/\beta$  experiment for spin-state selective editing of IS cross peaks, Andersson, Annala, and Otting, **133**, 364  
 sensitivity enhancement, Czisch and Boelens, **134**, 158
- Truncated transmission line probe  
 FT-EPR with, Rubinson *et al.*, **132**, 255
- TSETSE-2  
 in-phase double selective excitation of coupled spin systems using excitation sculpting, Bourg and Nuzillard, **133**, 173
- Tumors  
 brain and head and neck, applications of *in vivo* lactate editing with simultaneous detection of upfield singlets at 1.5 T using PRESS excitation, Star-Lack *et al.*, **133**, 243  
 breast ductal carcinomas, evaluation with high-resolution MAS IHMRS, Cheng *et al.*, **135**, 194
- Tumor spheroids  
 V79, quantitative <sup>1</sup>H MRI and MRS microscopy, Minard, Guo, and Wind, **133**, 368
- Tunneling  
 incoherent: and classical jumps over barrier: <sup>1</sup>H and <sup>2</sup>H NMR relaxation in hydrogen-bonded solids due to complex motion, Reynhardt and Lantanowicz, **130**, 195
- Two-dimensional NMR  
 assignment of aromatic side chains in <sup>13</sup>C-labeled proteins, Prompers *et al.*, **130**, 68  
 chemical-shift imaging in rotating solids by RF field gradients, Malveau *et al.*, **134**, 171  
 CP/MAS <sup>13</sup>C isotropic chemical shift correlation established by <sup>1</sup>H spin diffusion, Wilhelm *et al.*, **134**, 255  
 deuteron exchange, in study of vacancy diffusion in polycrystalline benzene, comparison with NMR stimulated echo, Isfort, Geil, and Fujara, **130**, 45  
 diffusion coefficient measurement by, using accordion spectroscopy, Millet and Pons, **131**, 166  
 exchange, and 1D MAS, direct determination of motional correlational times by, Favre, Schaefer, and Chmelka, **134**, 261  
 filter diagonalization method applied to spectra, Mandelshtam, Taylor, and Shaka, **133**, 304  
*J* and dipolar coupling measurement from simplified spectra, Ottiger, Delaglio, and Bax, **131**, 373  
 MAS, of dipolar solids, pulsed-field gradient selection, Fritzhanns *et al.*, **134**, 355  
 model-free analysis of mixtures using blind source separation, Nuzillard, Bourg, and Nuzillard, **133**, 358  
 multiple-quantum MAS, distributions of quadrupolar and isotropic chemical shift interactions, Bodart, **133**, 207  
 NQR, using single crystal, off-resonance effects, Ajithkumar *et al.*, **135**, 165  
 PISEMA, orientational constraints derived from hydrated powder samples by, Tian, Song, and Cross, **135**, 227  
 proton, cross peak detection in spectra using excitation sculpting, Van and Shaka, **132**, 154  
 selective and nonselective, coherence selection and excitation sculpting using RF-gradients, Heikkinen, Rahkamaa, and Kilpeläinen, **133**, 183  
 sensitivity- and gradient-enhanced TOCSY, Kövér, Uhrín, and Hruby, **130**, 162
- Two-dimensional spectroscopy  
 nuclear-Zeeman-resolved ESEEM spectroscopy, Willer *et al.*, **133**, 46

- mutation-frequency correlated EPR spectroscopy: PEANUT, Stoll *et al.*, **130**, 86
- U
- Ubiquitin  
<sup>13</sup>C/<sup>15</sup>N-labeled, measurement of dipolar couplings for methylene and methyl sites and use in structure determination, Ottiger *et al.*, **134**, 365  
dipole–CSA cross-correlated relaxation of <sup>13</sup>C and <sup>15</sup>N nuclei, quantitative analysis by ZQ/DQ spectroscopy, Brutscher *et al.*, **130**, 346
- Urea  
<sup>13</sup>C–<sup>15</sup>N multiply labeled, counting spins with SEDOR, Cull *et al.*, **133**, 352  
identification as metabolite in kidney/urine proton chemical exchange, Guivel-Scharen *et al.*, **133**, 36
- Urine  
detection of proton chemical exchange between metabolites and water, Guivel-Scharen *et al.*, **133**, 36
- V
- Vacancy diffusion  
in polycrystalline benzene, analysis with 2D deuteron exchange NMR and stimulated echoes, Isfort, Geil, and Fujara, **130**, 45
- Vector model  
adiabatic decoupling, Skinner and Bendall, **134**, 315
- Vicinal coupling constants  
fluorine–proton: individual substituent effects, Fabián, Guillemé, and Díez, **133**, 255
- Viscosity  
dilute nematic bicellar solutions, temperature effects, Struppe and Vold, **135**, 541  
model meal, measurement with echo-planar imaging relaxometry, Marciani *et al.*, **135**, 82
- Visuomotor systems  
activation during visually guided movements, functional MRI, Ellermann *et al.*, **131**, 272
- Volume averaging  
in analysis of proton diffusion and *T*<sub>1</sub> relaxation in polyacrylamide gels, Penke *et al.*, **132**, 240
- Volume selective excitation  
complicated shapes in three dimensions, Serša and Macura, **135**, 466
- W
- Water  
and *N*-acetyl aspartate, non-mono-exponential signal attenuation due to diffusion in brain tissue, Assaf and Cohen, **131**, 69  
and lactate, protons, protein-mediated magnetic coupling, Swanson, **135**, 248  
and metabolites, proton chemical exchange between, detection in biological tissues, Guivel-Scharen *et al.*, **133**, 36  
and oil, in sedimentary rock, *in situ* NMR analysis, de Swiet *et al.*, **133**, 385  
protons, measurement of chemical exchange rate constants with, using radiation damping, Chen and Mao, **131**, 358  
residual, in proton spectra, fast removal, Vanhamme *et al.*, **132**, 197  
–surfactant contact, <sup>19</sup>F–<sup>1</sup>H HOESY study, Raullet *et al.*, **133**, 324
- Water flipback  
<sup>15</sup>N-edited 3D NOESY-HMQC with, Gruschus and Ferretti, **135**, 87
- WATERGATE  
improved pulse sequences for solvent suppression in NMR, Liu *et al.*, **132**, 125  
in <sup>15</sup>N-edited 3D NOESY-HMQC with water flipback, Gruschus and Ferretti, **135**, 87
- Water–macromolecule interactions  
in aqueous model polymers, analysis using magnetization transfer, effect of pH and specific side groups, Gochberg *et al.*, **131**, 191
- Water magnetization  
undisturbed, composite 90° pulse sequence providing, Sodano, Landon, and Ptak, **133**, 194
- Water stripe  
information, in TOCSY experiments on systems with exchangeable protons, Nakashima, McClung, and Kotovych, **133**, 222
- Wideline NMR  
bandwidth extension in noise spectroscopy, Yang and Zax, **135**, 267
- World Wide Web  
NMR spectrometer operated by, common gateway interface program for, Buszko, Buszko, and Wang, **131**, 362
- X
- <sup>129</sup>Xe  
hyperpolarized, MRI of lung with, signal dynamics, Möller *et al.*, **135**, 133  
laser-polarized, surface NMR under MAS conditions, Brunner *et al.*, **130**, 145  
NMR and imaging, pore-structure determination of silica aerogels by, Gregory, Gerald, and Botto, **131**, 327
- Xenon  
laser-polarized, surface selective <sup>1</sup>H/<sup>29</sup>Si CP NMR by NOE enhancement, Pietraß, Seydoux, and Pines, **133**, 299  
time-dependent diffusion, PFG measurements, Mair *et al.*, **135**, 478
- Y
- Yb(C<sub>2</sub>D<sub>5</sub>SO<sub>4</sub>)<sub>3</sub> · 8H<sub>2</sub>O  
paramagnetic, solid-state NMR, effect of bulk magnetic susceptibility, Kubo, Spaniol, and Terao, **133**, 330
- Z
- Zeeman modulation  
effect on spin label relaxation time determination by progressive saturation EPR, Livshits, Páli, and Marsh, **133**, 79
- Zeolites  
NaX, (CH<sub>3</sub>)<sub>3</sub>P–AlCl<sub>3</sub> complex in, <sup>27</sup>Al → <sup>31</sup>P INEPT experiments, Kao and Grey, **133**, 313
- Zero-filling  
influence on peak integrals obtained by direct summation, Bourg and Nuzillard, **134**, 184
- Zero magnetic field  
photoexcited triplet state, optically detected ESEEM, Weis, Möbius, and Prisner, **131**, 17
- Zero-quantum coherence  
in AA'X spin systems, evolution: analysis of catalytic processes using parahydrogen, Natterer *et al.*, **133**, 92  
intermolecular, of multi-component spin systems in solution NMR, Ahn, Lisitza, and Warren, **133**, 266
- Zero-quantum dephasing  
and enhancement of chemical exchange or cross-relaxation, multiple-pulse mixing sequences for, Hwang and Shaka, **135**, 280
- Zero-quantum/double-quantum (ZQ/DQ) NMR  
quantitative analysis of dipole–CSA cross-correlated relaxation by, Brutscher *et al.*, **130**, 346
- Zn-tris(ethylenediamine) dinitrate  
integer spin system Ni(II), high-frequency EPR study: unusual behavior of double-quantum line, van Dam *et al.*, **130**, 140