

Cumulative Subject Index for 2001¹ Volumes 148–153

A

- Abelson kinase
interdomain orientation, estimation using rotational diffusion tensor, Ghose, Fushman, and Cowburn, **149**, 204
- Absolute-value display
in solid-state NMR imaging, symmetric echo acquisition for, Matsui and Saito, **149**, 103
- Acetylsalicylic acid
quadrupolar echo spectra of damped CD₃ quantum rotor, iterative lineshape analysis, Szymański *et al.*, **148**, 277
- Adaptive filter bank approach
nonparametric NMR spectroscopy method based on, Stoica and Sundin, **152**, 57
- Adiabatic dipolar recoupling
in solid-state NMR: DREAM scheme, Verel, Ernst, and Meier, **150**, 81
- Adiabatic full-passage pulses
water suppression schemes WASHCODE, Starčuk *et al.*, **152**, 168
- Adiabatic pulses
principles and method for creation of, review, Garwood and DelaBarre, **153**, 155
- Adiabatic TOCSY MAS
in liquids, Kupče, Keifer, and Delepierre, **148**, 115
- Advances in Magnetic Resonance
designing adiabatic pulses, Garwood and DelaBarre, **153**, 155
formal theory of spin–lattice relaxation, Goldman, **149**, 160
high-resolution ¹H NMR spectroscopy in solid state, Schnell and Spiess, **151**, 153
- ²⁷Al
AlPO-41, high-resolution NMR of quadrupolar nuclei using mixed MQ coherences, Jerschow, Logan, and Pines, **149**, 268
- L-Alanine
 γ -irradiated, longitudinal relaxation of stable radical in, pulse EPR study, Rakvin *et al.*, **152**, 149
- Alpha helix-forming peptide
demonstration of approach to visualizing spectral densities and deriving correlation times, Idiyatullin, Daragan, and Mayo, **152**, 132
- γ -Aminobutyric acid
detection by longitudinal scalar order difference editing, de Graaf and Rothman, **152**, 124
- Amplitude-modulated cw LOD EPR
experimental setup for, Granwehr, Forrer, and Schweiger, **151**, 78
- Anabaena variabilis*
plastocyanin, R_1 measurement using signal-eliminating relaxation filter, Hansen and Led, **151**, 339
- Analytical transfer functions
planar mixing, for two coupled spin-1 nuclei, Luy and Glaser, **153**, 210
three spins $\frac{1}{2}$: superposition of scalar and residual dipolar couplings, Luy and Glaser, **148**, 169
- Angular flow
in toroid cavity probes, Trautner *et al.*, **151**, 284
- Anisotropic diffusion tensor
measurement in unoriented sample by exploiting chemical shift anisotropy: ¹⁹F PGSE NMR with homonuclear decoupling, Dvinskikh and Furó, **148**, 73
- Anisotropic magnetic susceptibility
effect on cross correlation between dipole–dipole interaction and Curie spin relaxation, Bertini *et al.*, **152**, 103
- Anisotropies
chemical shielding, large, double-quantum-filtered rotational resonance MAS NMR in presence of, Bechmann, Helluy, and Sebald, **152**, 14
- Announcements
16th European Experimental NMR Conference, Prague, Czech Republic, June 2002, **153**, 277
The Laukien Prize to be presented at 43rd Experimental Nuclear Magnetic Resonance Conference, Pacific Grove, California, April 2002, **152**, 313
The Russell Varian Prize, **153**, 277
- Antamanide
¹³C-labeled, fast MAS total through-bond correlation spectroscopy, Hardy, Verel, and Meier, **148**, 459
- Aromatic residues
and MUSIC: amino acid type-selective ¹H–¹⁵N correlations, Schubert, Oschkinat, and Schmieder, **153**, 186
- Arrays
coil, with reactive cross-talk at 1.5 T, differential shading artifact in images from, spatial dependence, Tropp and Schirmer, **151**, 146
- Artifact
differential shading, in images from coil arrays with reactive cross-talk at 1.5 T, spatial dependence, Tropp and Schirmer, **151**, 146
- Automated NMR spectra assignment
scorpion neurotoxin using NOAH/DIAMOD, Xu *et al.*, **148**, 35
- Automatic coupling
resonator with, for minimizing sample motion noise in *in vivo* EPR spectroscopy, He *et al.*, **149**, 218
- Automatic coupling control
loop–gap resonator, by variable capacitor attached coupling coil, for EPR measurements at 650 MHz, Yokoyama *et al.*, **149**, 29
- Automatic tuning
resonator with, for minimizing sample motion noise in *in vivo* EPR spectroscopy, He *et al.*, **149**, 218

B

- ¹⁰B
NMR: effects of higher-rank multipoles on relaxation of $I=3$ spin system, Nosel *et al.*, **150**, 178
- B_0 homogeneity
high, magnet design with, for fast-field-cycling NMR applications, Lips *et al.*, **149**, 22
- B_1 inhomogeneity
role in magnetization loss during TOCSY experiments under MAS, Piotto *et al.*, **149**, 114
- Backbone assignment
sequential HNCACB and CBCANH protein pulse sequences for, Meissner and Sørensen, **151**, 328

¹Boldface numbers indicate volume; lightface numbers indicate pagination.

- sequential HNCA protein NMR pulse sequence for, Meissner and Sørensen, **150**, 100
- Backbone structure
 polypeptide, dihedral angles, determination in solid-state NMR by DQ ^{13}C CSA measurements, Blanco and Tycko, **149**, 131
- Background gradients
 macroscopic, effects on high-field PGSE NMR diffusion measurements, Price *et al.*, **150**, 49
- Bacteriophages
 filamentous, *fd*, cholesteric phase of, characterization for molecular alignment, Barrientos, Louis, and Gronenborn, **149**, 154
- Base protons
 and sugar protons, through-bond correlation in unlabeled nucleic acids, Phan, **153**, 223
- p*-Benzosemiquinone
 electron-spin-polarization pathways, time-resolved EPR study, Jäger and Norris, **150**, 26
- Bicelles
 partially oriented, NMR spectra, simultaneous determination of orientational and order parameter distributions, Sternin *et al.*, **149**, 110
- Bilayers
 partially oriented, NMR spectra, simultaneous determination of orientational and order parameter distributions, Sternin *et al.*, **149**, 110
 phospholipid, magnetically aligned, EPR spectroscopy at 94 GHz, Mangels *et al.*, **151**, 253
- Biological systems
 mesoscopic magnetic field inhomogeneities, theory of induced FID NMR signal dephasing, Sukstanskii and Yablonskiy, **151**, 107
- Biological tissues
 taurine detection by ^{33}S NMR, Musio and Sciacovelli, **153**, 259
- Biomolecular probe
 potential of xenon: effects of xenon-protein interactions on ^{129}Xe shifts, Rubin *et al.*, **152**, 79
- Bitter magnets
 NMR phase noise in, Sigmund *et al.*, **148**, 309, *erratum*, **151**, 347
- Blocking temperature
 core, horse spleen ferritin, Wajnberg *et al.*, **153**, 69
- Blood
 rheology, by NMR, Han *et al.*, **152**, 87
- Blood vessels
 mesoscopic magnetic field inhomogeneities, theory of induced FID NMR signal dephasing, Sukstanskii and Yablonskiy, **151**, 107
- Bond-shift rearrangement
 in solid $\text{Li}_3\text{P}_7(\text{monoglyme})_3$, ^{31}P MAS NMR study, Jäger *et al.*, **153**, 227
- Bone
 imaging, using long echo-train summation, and high-field gradient ^{31}P solid-state NMR, Gillies, Newling, and Randall, **151**, 235
- Bone marrow
 mesoscopic magnetic field inhomogeneities, theory of induced FID NMR signal dephasing, Sukstanskii and Yablonskiy, **151**, 107
- Bovine serum albumin
 in solution, water molecules strongly bound to, characterization by triple-quantum-filtered ^{17}O NMR, Lehoux, Krzysztyniak, and Bagnuet, **148**, 11
- BP-LED experiments
 variable-temperature, convection in, diffusion-broadened velocity spectra of, Mao and Kohlmann, **150**, 35
- Brain
 γ -aminobutyric acid detection by longitudinal scalar order difference editing, de Graaf and Rothman, **152**, 124
 localized ^{31}P MR spectra, quantitation based on reciprocity principle, Kreis *et al.*, **149**, 245
 localized proton spectroscopy, optimization of residual water signal removal by HLSVD, Cabanes *et al.*, **150**, 116
 mammalian, ^{13}C - ^{15}N coupling in, *in vivo* observation, Kanamori and Ross, **153**, 193
 two-dimensional homonuclear correlation spectroscopy, two-voxel localization sequence for, Delmas *et al.*, **149**, 119
- Broadband decoupling
 two-dimensional experiment separating decoupling sidebands from main peaks, Kupče and Freeman, **151**, 142
- Broadband excitation
 stochastic microwave, high-field/high-frequency Fourier transform EPR using, Fuhs, Prisner, and Möbius, **149**, 67
- Broadband inversion pulses
 improved performance in NMR in liquids, Smith, Hu, and Shaka, **151**, 269
- ## C
- ^{13}C
 adamantane, ramped-speed cross-polarization MAS NMR, Samoson, Tuherm, and Past, **149**, 264
 antamanide labeled with, fast MAS total through-bond correlation spectroscopy, Hardy, Verel, and Meier, **148**, 459
 ^{13}C - ^{13}C coupling effects on ^{13}C PM5-REDOR: peptide torsion angle measurements, Sack *et al.*, **148**, 104
 ^{13}C chemical shift and ^{13}C - ^{14}N dipolar coupling tensors by ^{13}C rotary resonance solid-state NMR, Odgaard *et al.*, **148**, 298
 ^{13}C chemical shifts in RNA nucleosides, calculation, Rossi and Harbison, **151**, 1
 chemical-shift tensor measurement, dipolar dephasing method for slow MAT, Hu *et al.*, **152**, 7
 ^{13}C , ^2H -enriched carbohydrates, weakly aligned, residual ^2H quadrupolar couplings in, Kjellberg and Homans, **151**, 90
 ^{13}C - ^{15}N coupling in mammalian brain, *in vivo* observation, Kanamori and Ross, **153**, 193
 ^{13}C , ^{15}N -glycine, MAS NMR, double-quantum excitation by pulse-assisted rotational resonance, Karlsson *et al.*, **148**, 238
 ^{13}C , ^{15}N -labeled proteins
 backbone assignment, sequential HNCA NMR pulse sequence for, Meissner and Sørensen, **150**, 100
 measurement of $^1J_{\text{NC}\alpha}$ and $^2J_{\text{NC}\alpha}$ from simplified 2D $^{13}\text{C}\alpha$ -coupled ^{15}N SE-HSQC spectrum, Heikkinen, Permi, and Kilpeläinen, **148**, 53
 correlation of $^{15}\text{N}(i)$ with interresidue $^{13}\text{C}^\beta(i-1)$ of side chains, sequential HNCACB and CBCANH pulse sequences for, Meissner and Sørensen, **151**, 328
 dipolar-coupled ^{13}C spin pairs of enriched poly(ethylene), DQ-DQ-MAS exchange NMR, Schnell, Watts, and Spiess, **149**, 90
 double-quantum CSA measurements, for peptide backbone dihedral angles in solid-state NMR, Blanco and Tycko, **149**, 131
- MAS NMR
 double-quantum-filtered rotational resonance MAS NMR in presence of large chemical shielding anisotropies, Bechmann, Helluy, and Sebald, **152**, 14
 under moderately fast spinning conditions, spectral editing in, De Vita and Frydman, **148**, 327
 medium- and long-distance ^1H - ^{13}C heteronuclear correlation NMR in solids, Yao, Schmidt-Rohr, and Hong, **149**, 139
 NMR chemical shifts, dependence on conformations of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
 relaxation rates in trehalose: separating structure and dynamics in CSA/DD cross-correlated relaxation, Kövér and Batta, **150**, 137
 solid-state ^1H - ^{13}C heteronuclear correlation in the limit of fast MAS, recoupled polarization-transfer methods for, Saalwächter, Graf, and Spiess, **148**, 398
 spin diffusion, in organic solids, slow down by fast MAS: CODEX NMR, Reichert, Bonagamba, and Schmidt-Rohr, **151**, 129
 spin-state-selective excitation in selective 1D inverse NMR experiments, Parella and Belloc, **148**, 78

- trehalose, analysis of conformation of glycosidic linkage by double-quantum solid-state NMR, Ravindranathan *et al.*, **151**, 136
- two-dimensional dynamic-director ^{13}C NMR of liquid crystals, McElheny, Zhou, and Frydman, **148**, 436
- uniformly labeled L-alanine, dipolar and *J*-encoded DQ MAS spectra under rotational resonance, Filip *et al.*, **150**, 184
- Calbindin D_{9k}
cerium(III)-substituted, self-orientation in magnetic field of, solution structure calculation through, Bertini *et al.*, **148**, 23
- α -Calcium formate
deuterons in, precision measurement of quadrupole coupling and chemical shift tensors, Schmitt *et al.*, **151**, 65
- Calf muscle
human, *in vivo* ^{31}P echo-planar spectroscopic imaging, Wilhelm and Bachert, **149**, 126
- Canonical correlation
 ^{13}C chemical shift determination of conformations of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
- Capillary NMR flow probe
 ^1H NMR characterization of product from single solid-phase resin beads using, Lacey *et al.*, **153**, 215
- Carbohydrates
conformation of glycosidic linkage in disaccharide, double-quantum solid-state NMR, Ravindranathan *et al.*, **151**, 136
weakly aligned, residual ^2H quadrupolar couplings in, Kjellberg and Homans, **151**, 90
- Carbon-based sensors
mechanism of oxygen response in, Atsarkin *et al.*, **149**, 85
- Carbonyl chemical shift
use to relieve degeneracies in triple-resonance assignment experiments, Sayers and Torchia, **153**, 246
- Carr–Purcell–Meiboom–Gill
analysis of diffusion effects in stray field NMR using, Hürlimann, **148**, 367
CP/QCPMG strategy for NMR of half-integer quadrupolar nuclei in dilute environments, Lipton, Sears, and Ellis, **151**, 48
- Carr–Purcell sequences
with composite pulses, Hürlimann, **152**, 109
optimized homonuclear, Kramer *et al.*, **149**, 58
- Cartilage
diffusion of paramagnetically labeled proteins: enhancement of 1D NMR imaging, Foy and Blake, **148**, 126
- Cavities
in proteins in solution, probing by ^{129}Xe NMR, Locci *et al.*, **150**, 167
- Cavity insert
ferroelectric, enhanced EPR sensitivity from, Nesselov, Surek, and Thomas, **153**, 7
- CBCACO(N)H
triple-resonance assignment experiments, use of carbonyl chemical shift to relieve degeneracies in, Sayers and Torchia, **153**, 246
- CBCANH
sequential, protein NMR pulse sequence, Meissner and Sørensen, **151**, 328
- CD₃ quantum rotor
damped, quadrupolar echo spectra of, iterative lineshape analysis, Szymański *et al.*, **148**, 277
- Cellular research
integrated confocal and magnetic resonance microscope for, Wind *et al.*, **147**, 371, *erratum*, **151**, 347
- Cerium(III)
substituted calbindin D_{9k}, self-orientation in magnetic field of, solution structure calculation through, Bertini *et al.*, **148**, 23
- Chars
oxygen-sensitive synthesized, mechanism of oxygen response in, Atsarkin *et al.*, **149**, 85
- Chemical exchange
effects, neglecting when correcting *in vivo* ^{31}P MRS data for partial saturation, Ouwerkerk and Bottomley, **148**, 425; comment and reply, **149**, 251, 282; *erratum*, **151**, 347
H(N)-water proton, improved measurement of ^{15}N -{ ^1H } NOEs in presence of, Idiyatullin, Daragan, and Mayo, **153**, 138
- Chemically induced dynamic electron polarization
time-resolved EPR study of electron-spin-polarization pathways of *p*-benzosemiquinone, Jäger and Norris, **150**, 26
- Chemical shielding anisotropies
large, double-quantum-filtered rotational resonance MAS NMR in presence of, Bechmann, Helluy, and Sebald, **152**, 14
- Chemical-shift anisotropy
CSA/DD cross-correlated relaxation, separating structure and dynamics, Kövér and Batta, **150**, 137
double-quantum ^{13}C , for polypeptide backbone dihedral angles in solid-state NMR, Blanco and Tycko, **149**, 131
- Chemical-shift correlations
two-dimensional dynamic-director ^{13}C NMR of liquid crystals, McElheny, Zhou, and Frydman, **148**, 436
- Chemical-shift editing
and *J*-editing, proton MRI of ^{13}C distribution by, Casieri *et al.*, **153**, 117
- Chemical shifts
 ^{13}C
and ^{13}C - ^{14}N dipolar coupling tensors, ^{13}C rotary resonance solid-state NMR, Odgaard *et al.*, **148**, 298
in RNA nucleosides, calculation, Rossi and Harbison, **151**, 1
 ^{13}C NMR, and conformations of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
oxygen-17, in methyl formate, solvent effects, de Kowalewski *et al.*, **148**, 1
 ^{129}Xe , in aqueous solution, effects of nonspecific xenon–protein interactions, Rubin *et al.*, **152**, 79
- Chemical-shift tensor
 ^{13}C , measurement, dipolar dephasing method for slow MAT, Hu *et al.*, **152**, 7
deuterons in α -calcium formate, precision measurement, Schmitt *et al.*, **151**, 65
- Cholesteric phase
filamentous bacteriophage *fd*, characterization for molecular alignment, Barrientos, Louis, and Gronenborn, **149**, 154
- Chromatographic NMR
technique for measuring angular flow in toroid cavity probes for use in, Trautner *et al.*, **151**, 284
- Chymotrypsin inhibitor 2
clean-TROSY-adapted implementation of sequential HNCACB and CBCANH pulse sequences, Meissner and Sørensen, **151**, 328
- CIDEP (*see* Chemically induced dynamic electron polarization)
- ^{13}C indirect NMR imaging
proton MRI of ^{13}C distribution by *J* and chemical-shift editing, Casieri *et al.*, **153**, 117
- Circularly polarized RF magnetic fields
for spin-1 NQR, Miller, Suits, and Garroway, **151**, 228
- Circular polarization
excitation and detection, ^{14}N NQR signals, Lee, Robert, and Lathrop, **148**, 355
- ^{13}C isotopomers
 T_2^* measurement of protons in, 2D pulse sequence for, Lambert, Haiber, and Herzog, **150**, 175
- ^{35}Cl
in solid-state proton imaging detected by quadrupole resonance, Perlo *et al.*, **150**, 132
- Clay
kaolin, iron(III)-doped, NMR relaxation and diffusion measurements on, Valckenborg, Pel, and Kopinga, **151**, 291

- Co²⁺
salts, relaxation enhancement of spin labels in membranes by, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- CODEX NMR
fast MAS, slow down of ¹³C spin diffusion in organic solids, Reichert, Bonagamba, and Schmidt-Rohr, **151**, 129
- Coherences
fast MAS and multiple-quantum, review, Schnell and Spiess, **151**, 153
- Coherence selection
by gradients, optimization, in sensitivity enhancement of 2D experiment for long-range coupling constants, Cicero, Barbato, and Bazzo, **148**, 209
- Coherence transfer
analytical planar mixing transfer functions for two coupled spin-1 nuclei, Luy and Glaser, **153**, 210
gradient-selected DQ, localized spectral editing with, effects of slice-selective excitation/refocusing, Lei and Dunn, **150**, 17
- Coherent spontaneous emission
and quantum origins of free induction decay signal and spin noise, Hoult and Ginsberg, **148**, 182
- Coil arrays
with reactive cross-talk at 1.5 T, differential shading artifact in images from, spatial dependence, Tropp and Schirmer, **151**, 146
- COIN TACS
tailored correlation spectroscopy approach, Carlomagno, Prash, and Glaser, **149**, 52
- Composite pulses
Carr–Purcell sequences with, Hürlimann, **152**, 109
resonance offset tailored, Cummins and Jones, **148**, 338
- Composite-pulse sequence
for spurious signal suppression, determination of quadrupole parameters with, Millot and Man, **150**, 10
- Computer simulation
NMR experiments, complete Hermitian operator basis set for any spin quantum number, Allard and Härd, **153**, 15
spin-echo spatial distribution in case of restricted diffusion, Duh, Mohorič, and Stepišnik, **148**, 257
- Conducting solids
exchange-narrowed regime of ESR in, site-selective electron nuclear double resonance in, Denninger *et al.*, **148**, 248
- Confocal scanning fluorescent microscopy
integrated with magnetic resonance microscopy, for cellular research, Wind *et al.*, **147**, 371, *erratum*, **151**, 347
- Conformation
fluoroalcohol-induced, of peptides and proteins, intermolecular ¹H{¹⁹F} NOEs in studies of, Martinez and Gerig, **152**, 269
glycosidic linkage in disaccharide, double-quantum solid-state NMR, Ravindranathan *et al.*, **151**, 136
RNA nucleosides, and calculation of ¹³C chemical shifts, Rossi and Harbison, **151**, 1
RNA nucleosides and nucleotides, dependence of ¹³C NMR chemical shifts on, Ebrahimi *et al.*, **150**, 1
- Constant-time COSY
cross-peak nulling in, measurement of homonuclear proton couplings based on, Wu and Bax, **151**, 242
- Constraining fitting parameters
modeling magnetization transfer using, Ceckler, Maneval, and Melkowitz, **151**, 9
- Continuous wavelet transform
water modeled signal removal and data quantification in localized MRS using time-scale postacquisition, Serrai *et al.*, **149**, 45
- Continuous-wave MRI
short *T*₂ materials, Davies *et al.*, **148**, 289
- Contrast indicator
MRI-based D₂O/H₂O-contrast method for water flow and distribution in heterogeneous systems, Ilvonen *et al.*, **149**, 36
- Convection
in variable-temperature BP-LED experiments, diffusion-broadened velocity spectra, Mao and Kohlmann, **150**, 35
- Convection compensated electrophoretic NMR
detection of electrophoretic motion in presence of bulk convective flows, He and Wei, **150**, 126
- Convection effects
in self-diffusion NMR measurements, minimization with sample rotation, Esturau *et al.*, **153**, 48
- Convolution-based fitting
high-spatial-resolution multi-site EPR oximetry using, Grinberg, Smirnov, and Swartz, **152**, 247
- Copper
Cu²⁺ salts, relaxation enhancement of spin labels in membranes by, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
pseudo-Jahn–Teller low-symmetry Cu(II) complexes in diaqua(L-aspartato) Zn(II)·H₂O crystals, electron spin relaxation in, Hoffmann *et al.*, **153**, 92
- Correlation
spin displacements, 2D PFG sequences VEXSY and SERPENT for, Blümich *et al.*, **152**, 162
- Correlation spectra
¹⁵N, ¹H, in Asn and Glu side chains, NMR experiments for measuring dipolar couplings from, Permi, **153**, 267
- Correlation spectroscopy
¹H–¹H MAS, and distance measurements, in deuterated peptide, Reif *et al.*, **151**, 320
- Correlation time distributions
motional, derivation, and visualizing spectral density functions, Idiyatullin, Daragan, and Mayo, **152**, 132
- COSY
COIN TACS, Carlomagno, Prash, and Glaser, **149**, 52
constant-time, cross-peak nulling in, measurement of homonuclear proton couplings based on, Wu and Bax, **151**, 242
and HMQC, multiple-coil probe design for acquisition of spectra from multiple samples, Zhang, Sweedler, and Webb, **153**, 254
¹H–¹⁵N, in solid state under fast MAS, inverse detection and heteronuclear editing in, Schnell *et al.*, **150**, 57
in vivo 2D homonuclear, two-voxel localization sequence for, Delmas *et al.*, **149**, 119
two-dimensional regular, spectra, measurement of homonuclear proton couplings from, Delaglio, Wu, and Bax, **149**, 276
- Coupling
measurement with multifrequency pulsed EPR/ENDOR probe, Avdievich and Gerfen, **153**, 178
- Coupling coil
variable capacitor attached, automatic coupling control of loop–gap resonator by, for EPR measurements at 650 MHz, Yokoyama *et al.*, **149**, 29
- Coupling constants (*see J* coupling)
- Creatine
skeletal muscle, magnetic coupling of methyl protons in, role of exchangeable NH protons, Kruijskamp and Nicolay, **149**, 8
- Cross-correlated relaxation
characterization of hydrogen bond lengths in Watson–Crick base pairs, Riek, **149**, 149
CSA/DD, separating structure and dynamics in, Kövér and Batta, **150**, 137
effects in measurement of homonuclear proton couplings based on cross-peak nulling in CT-COSY, Wu and Bax, **151**, 242
¹⁹F, study of fluorinated ligand–receptor interactions, Peng, **153**, 32
- Cross correlation
between dipole–dipole interaction and Curie spin relaxation, effect of anisotropic magnetic susceptibility, Bertini *et al.*, **152**, 103
- Cross-peak nulling
in CT-COSY, measurement of homonuclear proton couplings based on, Wu and Bax, **151**, 242

- Cross peaks
NOESY, automatic assignment, and protein structure determination of scorpion neurotoxin, using NOAH/DIAMOD, Xu *et al.*, **148**, 35
- Cross polarization
CP/QCPMG strategy for NMR of half-integer quadrupolar nuclei in dilute environments, Lipton, Sears, and Ellis, **151**, 48
multiple periods, indirect NMR detection in solids with, Khitritin and Fung, **152**, 185
ramped-speed CP MAS NMR, Samoson, Tuherm, and Past, **149**, 264
- Cross-talk
reactive, at 1.5 T, differential shading artifact in images from coil arrays with, spatial dependence, Tropp and Schirmer, **151**, 146
- Cu²⁺ Jahn–Teller effect
low-symmetry Cu(II) complexes in diaqua(L-aspartato)Zn(II) · H₂O crystals, electron spin relaxation in, Hoffmann *et al.*, **153**, 92
- Cu(H₂O)₆
vibronic complexes in Tutton salt crystals at low temperatures, electron spin echo of, dephasing relaxation, Hoffmann *et al.*, **153**, 56
- Curie spin relaxation
and dipole–dipole interaction, cross correlation between, effect of anisotropic magnetic susceptibility, Bertini *et al.*, **152**, 103
- Cycling sidebands
decoupling-generated, separation from main peaks, 2D experiment, Kupče and Freeman, **151**, 142
- Cylindrical mixing conditions
analytical transfer functions for three spins $\frac{1}{2}$: superposition of scalar and residual dipolar couplings, Luy and Glaser, **148**, 169
- D**
- Data quantification
and water modeled signal removal, in localized MRS using time-scale postacquisition, Serrai *et al.*, **149**, 45
- Dead Ringer–DNA complex
application of NMR experiment for analyzing sugar puckering in unlabeled DNA, Iwahara, Wojciak, and Clubb, **153**, 262
- Deconvolution
and measurement of spin–spin splittings by modified *J* doubling in frequency domain, Garza-García, Ponzanelli-Velázquez, and del Río-Portilla, **148**, 214
potential of noise reduction in quadrupolar echo spectra at short echo times, Furó and Hedin, **152**, 214
- Decoupling
sidebands generated in, separation from main peaks with 2D experiment, Kupče and Freeman, **151**, 142
- Delayed-focus pulses
optimized using simulated annealing, Shen, **149**, 234
- Dephasing
FID NMR signal, induced by magnetic field inhomogeneities, theory, Sukstanskii and Yablonskiy, **151**, 107
- Dephasing relaxation
electron spin echo of vibronic Cu(H₂O)₆ complexes in Tutton salt crystals at low temperatures, Hoffmann *et al.*, **153**, 56
- Deuterated peptide
¹H–¹H MAS correlation spectroscopy and distance measurements in, Reif *et al.*, **151**, 320
- Deuterium oxide
MRI-based D₂O/H₂O-contrast method to study water flow and distribution in heterogeneous systems, Ilvonen *et al.*, **149**, 36
- Deuterons
in α -calcium formate, quadrupole coupling and chemical shift tensors, precision measurement, Schmitt *et al.*, **151**, 65
- Deutsch–Jozsa quantum algorithm
and logic gates, implementation by 2D NMR using spin- and transition-selective pulses, Mahesh *et al.*, **148**, 95
- DIAMOD
automatic assignment of NOESY cross peaks and determination of protein structure of scorpion neurotoxin with NOAH/DIAMOD, Xu *et al.*, **148**, 35
- Diaqua(L-aspartato)Zn(II) · H₂O crystals
pseudo-Jahn–Teller low-symmetry Cu(II) complexes in, electron spin relaxation in, Hoffmann *et al.*, **153**, 92
- Difftrain
approach to true spectroscopic single-scan diffusion measurement, Stamps *et al.*, **151**, 28
- Diffusion
effects in stray field NMR, Hürlimann, **148**, 367
measurements
accurate, of translational diffusion coefficients, Damberg, Jarvet, and Gräslund, **148**, 343
high-field PGSE NMR, macroscopic background gradient and radiation damping effects, Price *et al.*, **150**, 49
by ²H stimulated-echo-type PGSE NMR, application to liquid crystal, Dvinskikh *et al.*, **153**, 83
improved spin-echo-edited NMR, Otto and Larive, **153**, 273
on iron(III)-doped kaolin clay, Valckenborg, Pel, and Kopinga, **151**, 291
NMR, static fringe field of superconducting magnets used in, modeling, Jeglič *et al.*, **150**, 39
true spectroscopic single-scan: Difftrain approach, Stamps *et al.*, **151**, 28
using nutation spin echoes, Scharfenecker Ardelean, and Kimmich, **148**, 363
microscopic imaging: STE PFG TSE pulse sequence, Scheenen *et al.*, **151**, 94
one-dimensional DOSY, Loening, Keeler, and Morris, **153**, 103
paramagnetically labeled proteins in cartilage: enhancement of 1D NMR imaging, Foy and Blake, **148**, 126
system undergoing, velocity autocorrelation functions by multiple data acquisition in PFG NMR in, Stapf, **152**, 308
- Diffusion-broadened velocity spectra
convection in variable-temperature BP-LED experiments, Mao and Kohlmann, **150**, 35
- Diffusion coefficients
use of sample rotation for minimizing convection effects in self-diffusion NMR measurements, Esturau *et al.*, **153**, 48
- Diffusion tensor magnetic resonance imaging
eigenvalues and eigenvectors in, analytical computation, Hasan *et al.*, **152**, 41
- Diffusivities
genuine, determination in heterogeneous media with stimulated echo PFG NMR, Vasenkov *et al.*, **149**, 228
- Dilute environments
half-integer quadrupolar nuclei in, NMR observation strategy, Lipton, Sears, and Ellis, **151**, 48
- Dimensionality evolution periods
reduced, triple-resonance solid-state NMR experiments with, Astrof, Lyon, and Griffin, **152**, 303
- Dipolar correlation
¹H–¹³C, in solids under very fast MAS, recoupled polarization-transfer methods for, Saalwächter, Graf, and Spiess, **148**, 398
- Dipolar-coupled spin pairs
as probes for slow molecular dynamics: DQ-DQ-MAS exchange NMR spectroscopy, Schnell, Watts, and Spiess, **149**, 90
- Dipolar coupling (*see also* Residual dipolar coupling)
¹³C–¹³C, effects on carbon-observed deuterium-dephased PM5-REDOR: peptide torsion angle measurements, Sack *et al.*, **148**, 104

- ^{13}C - ^{14}N dipolar coupling and ^{13}C chemical shift tensors determined by ^{13}C rotary resonance solid-state NMR, Odgaard *et al.*, **148**, 298
- data sets, maximum likelihood method for determining D_a^{PO} and R for, Warren and Moore, **149**, 271
- double-quantum filtered ^1H NMR quantification of lactate in muscle, Asllani *et al.*, **152**, 195
- ^1H - ^{13}C dipolar correlation in solids under very fast MAS, recoupled polarization-transfer methods for, Saalwächter, Graf, and Spiess, **148**, 398
- scalar and residual, superposition: analytical transfer functions for three spins $\frac{1}{2}$, Luy and Glaser, **148**, 169
- Dipolar dephasing
method, for slow magic angle turning experiment, Hu *et al.*, **152**, 7
- Dipolar fields
intermolecular MQ coherence pumping efficiency in solution NMR, Zhang *et al.*, **148**, 200
- long-range, imaging in structured liquid-state samples, Bowtell, Gutteridge, and Ramanathan, **150**, 147
- Dipolar mixing sequences
optimized homonuclear Carr–Purcell-type, Kramer *et al.*, **149**, 58
- Dipolar recoupling
adiabatic, in solid-state NMR: DREAM scheme, Verel, Ernst, and Meier, **150**, 81
- heteronuclear, under MAS in SFAM and REDOR, effect of RF inhomogeneity, Nishimura, Fu, and Cross, **152**, 227
- methods, measurement of interfluorine distances in solids with, Gilchrist *et al.*, **152**, 1
- Dipolar recoupling enhanced by amplitude modulation (*see* DREAM scheme)
- Dipole–dipole interaction
CSA/DD cross-correlated relaxation, separating structure and dynamics, Kövér and Batta, **150**, 137
- and Curie spin relaxation, cross correlation between, effect of anisotropic magnetic susceptibility, Bertini *et al.*, **152**, 103
- Disaccharide
glycosidic linkage in, conformation, double-quantum solid-state NMR, Ravindranathan *et al.*, **151**, 136
- Displacements
spin, correlating, 2D PFG sequences VEXSY and SERPENT for, Blümich *et al.*, **152**, 162
- Distance measurements
 ^{19}F - ^{19}F , in solids, Gilchrist *et al.*, **152**, 1
- ^1H - ^1H , in deuterated peptide, Reif *et al.*, **151**, 320
- DNA
homonuclear proton couplings in, measurement based on cross-peak nulling in CT-COSY, Wu and Bax, **151**, 242
- unlabeled, sugar puckering in, NMR experiment for analyzing, Iwahara, Wojciak, and Clubb, **153**, 262
- Watson–Crick base pairs, characterization of hydrogen bond lengths by cross-correlated relaxation, Riek, **149**, 149
- Domain orientation
multidomain proteins, estimation using rotational diffusion tensor determination from relaxation data, Ghose, Fushman, and Cowburn, **149**, 204
- DOSY
one-dimensional, qualitative spectrum based on single 1D NMR experiment, Loening, Keeler, and Morris, **153**, 103
- Double pulsed-gradient spin-echo NMR
with stimulated echoes, and phase cycle scheme, Khrapitchev and Callaghan, **152**, 259
- Double-quantum buildup and decay curves
for crosslinked natural rubber samples, Wiesmath *et al.*, **149**, 258
- Double-quantum coherence
double-quantum excitation in MAS NMR by pulse-assisted rotational resonance, Karlsson *et al.*, **148**, 238
- Double-quantum coherence transfer
gradient-selected, localized spectral editing with, effects of slice-selective excitation/refocusing, Lei and Dunn, **150**, 17
- Double-quantum excitation
in NMR of spinning solids, by pulse-assisted rotational resonance, Karlsson *et al.*, **148**, 238
- Double-quantum filtered NMR
 ^1H , quantification of lactate in muscle, Asllani *et al.*, **152**, 195
- signals, in inhomogeneous magnetic fields, Wiesmath *et al.*, **149**, 258
- Double-quantum filtered spectroscopy
 ^{13}C CSA measurements for peptide backbone dihedral angles in solid-state NMR, Blanco and Tycko, **149**, 131
- Double-quantum filtration
effects of slice-selective excitation/refocusing, Lei and Dunn, **150**, 17
- under rotational-resonance MAS, in presence of large chemical shielding anisotropies, Bechmann, Helluy, and Sebald, **152**, 14
- Double-quantum MAS
DQ–DQ–MAS exchange NMR spectroscopy, Schnell, Watts, and Spiess, **149**, 90
- Double-quantum NMR
 ^1H - ^1H , in solid state under fast MAS, inverse detection and heteronuclear editing in, Schnell *et al.*, **150**, 57
- dipolar and J -encoded DQ MAS spectra under rotational resonance, Filip *et al.*, **150**, 184
- solid-state, analysis of conformation of glycosidic linkage in disaccharide, Ravindranathan *et al.*, **151**, 136
- Double resonance
enhancements of ^{31}P in solid-state NMR in high-field gradients, Gillies, Newling, and Randall, **151**, 235
- Doubly rotating frame
spin–lattice relaxation in, rotating-frame analog of spin-locking and, Tabuchi and Hatanaka, **148**, 121
- DREAM scheme
adiabatic dipolar recoupling in solid-state NMR, Verel, Ernst, and Meier, **150**, 81
- Droplet size
in oil-in-water emulsions, measurement with NMR transverse relaxation, Marciani *et al.*, **153**, 1
- Dy^{3+}
salts, relaxation enhancement of spin labels in membranes by, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Dynamic NMR microscopy
gas-phase Poiseuille flow, Kaiser *et al.*, **149**, 144
- Dynamics
and structure, separating in CSA/DD cross-correlated relaxation, Kövér and Batta, **150**, 137
- E
- Echo acquisition
multiple echoes, in PFG NMR experiments, determination of velocity autocorrelation functions by, Stapf, **152**, 308
- symmetric, for absolute-value display in solid-state NMR imaging, Matsui and Saito, **149**, 103
- Echo Fourier transformation
symmetric echo acquisition for absolute-value display in solid-state imaging, Matsui and Saito, **149**, 103
- Echo-planar imaging microscopy
one-shot velocimetry using, Han and Callaghan, **148**, 349
- Echo-planar spectroscopic imaging
in vivo ^{31}P , of human calf muscle, Wilhelm and Bachert, **149**, 126
- E.COSY-type spectra
improved measurement of $^3J(\text{H}_i^\alpha, \text{N}_{i+1})$ coupling constants in H_2O -dissolved proteins, Löhner *et al.*, **153**, 75

- S^3E -E.COSY methods for measurement of ^{19}F -associated scalar and dipolar coupling constants, Luy, Barchi, and Marino, **152**, 179
- Editorial
incorporation of *Advances in Magnetic Resonance*, **149**, 159
- Eigenvalues
in diffusion tensor MRI, analytical computation, Hasan *et al.*, **152**, 41
- Eigenvectors
in diffusion tensor MRI, analytical computation, Hasan *et al.*, **152**, 41
- Electrically detected magnetic resonance
resonance-field dependence, effects of exchange interaction, Fukui *et al.*, **149**, 13
of silicon wafer, imaging, Sato *et al.*, **153**, 113
- Electron-nuclear double resonance (*see* ENDOR)
- Electron spin echo
electron spin relaxation in pseudo-Jahn–Teller low-symmetry Cu(II) complexes in diaqua(*l*-aspartato)Zn(II) · H₂O crystals, Hoffmann *et al.*, **153**, 92
vibronic Cu(H₂O)₆ complexes in Tutton salt crystals at low temperatures, dephasing relaxation of, Hoffmann *et al.*, **153**, 56
- Electron-spin-polarization pathways
p-benzosemiquinone, time-resolved EPR study, Jäger and Norris, **150**, 26
- Electron spin relaxation
in pseudo-Jahn–Teller low-symmetry Cu(II) complexes in diaqua(*l*-aspartato) Zn(II) · H₂O crystals, Hoffmann *et al.*, **153**, 92
- Electron spin relaxation time
measurements using RF-LODESR, and application in oximetry, Pangiotelis, Nicholson, and Hutchison, **149**, 74
triarylmethyl radicals in fluid solution, Yong *et al.*, **152**, 156
- Electron spin resonance (*see* EPR)
- Electron-Zeeman resolved EPR
principles and theoretical description, Eichel and Schweiger, **152**, 276
- Electrophoretic NMR
convection compensated, for detection of electrophoretic motion in presence of bulk convective flows, He and Wei, **150**, 126
- Emulsification
fat, measurement with NMR transverse relaxation, Marciani *et al.*, **153**, 1
- Encoding methods
for hyperpolarized noble gas MRI, signal-to-noise ratio comparison, Zhao *et al.*, **148**, 314
- Encoding selection
phase cycle scheme for double PGSE NMR with stimulated echoes, Khrapitchev and Callaghan, **152**, 259
- ENDOR
Q-band pulse EPR/ENDOR spectrometer, Gromov *et al.*, **149**, 196
single-crystal ^{57}Fe Q-band ENDOR study of 4 iron–4 sulfur cluster in reduced [4Fe–4S]¹⁺ state, Moriaud *et al.*, **153**, 238
site-selective, in exchange-narrowed regime of ESR in conducting solids, Denninger *et al.*, **148**, 248
spectra recorded at high magnetic fields and low temperature, effect of spin relaxation, Epel *et al.*, **148**, 388
- EPR (*see also specific types*)
in conducting solids, exchange-narrowed regime of, site-selective electron nuclear double resonance in, Denninger *et al.*, **148**, 248
enhanced sensitivity from ferroelectric cavity insert, Nesselov, Surek, and Thomas, **153**, 7
horse spleen ferritin, temperature dependence of spectra, Wajnberg *et al.*, **153**, 69
magnetically aligned phospholipid bilayers at 94 GHz, Mangels *et al.*, **151**, 253
mechanism of oxygen response in carbon-based sensors, Atsarkin *et al.*, **149**, 85
Q-band pulse EPR/ENDOR spectrometer, and implementation of advanced 1D and 2D pulse EPR methodology, Gromov *et al.*, **149**, 196
topical *in vivo*, bridged loop–gap S-band surface resonator for, Petryakov *et al.*, **151**, 124
- EPR imaging
in vivo, of distribution and metabolism of nitroxide radicals in human skin, He *et al.*, **148**, 155
topical *in vivo*, bridged loop–gap S-band surface resonator for, Petryakov *et al.*, **151**, 124
- EPR instrumentation
multifrequency pulsed EPR/ENDOR probe, Avdievich and Gerfen, **153**, 178
- EPR oximetry
high-spatial-resolution multi-site, Grinberg, Smirnov, and Swartz, **152**, 247
- EPR relaxation enhancement
spin labels in membranes by paramagnetic ion salts, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Erythrocytes
mesoscopic magnetic field inhomogeneities, theory of induced FID NMR signal dephasing, Sukstanskii and Yablonskiy, **151**, 107
- Escherichia coli*
heterologous protein in, selective labeling for NMR studies, Almeida *et al.*, **148**, 142
- ESEEM
demonstration of Q-band pulse EPR/ENDOR spectrometer, Gromov *et al.*, **149**, 196
properties of HYSORE spin echo signal, Astashkin and Raitsimring, **148**, 379
- ESR (*see* EPR)
- Exchange
role of exchangeable NH protons in creatine for magnetic coupling of creatine methyl protons in muscle, Kruskamp and Nicolay, **149**, 8
- Exchange interaction
and resonance-field dependence in electrically detected magnetic resonance, Fukui *et al.*, **149**, 13
- Exchange spectroscopy
DQ-DQ-MAS exchange NMR spectroscopy, Schnell, Watts, and Spiess, **149**, 90
- Excitation
broadband stochastic microwave, high-field/high-frequency Fourier transform EPR using, Fuhs, Prisner, and Möbius, **149**, 67
spin-state-selective, in selective 1D inverse NMR experiments, Parella and Belloc, **148**, 78
- Excitation bandwidth
large, achievement in high-resolution spectra at high field, NMR instrumentation for, Luchinat *et al.*, **150**, 161
- Excitation and detection
circular polarization, in ^{14}N NQR, Lee, Robert, and Lathrop, **148**, 355
- Excitation/refocusing
slice-selective, effects in localized spectral editing with gradient-selected DQ coherence transfer, Lei and Dunn, **150**, 17
- Excitation sculpting
Hadamard, and application in NOESY1D experiments, Krishnamurthy, **153**, 144
improved resolution using selectively shifted pulses, Krishnamurthy, **153**, 124
- Exocyclic angle
from ^{13}C chemical shift data of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
- Exorcycle
in noise reduction in quadrupolar echo spectra at short echo times, Furó and Hedin, **152**, 214
- Extended Fourier transform
for 2D spectral estimation, Armstrong and Mandelshtam, **153**, 22

F

 ^{19}F

- associated scalar and dipolar coupling constants, S^3E -E.COSY methods for measurement of, Luy, Barchi, and Marino, **152**, 179

- cross-correlated relaxation measurements for fluorinated ligand–receptor interactions, Peng, **153**, 32
- intermolecular $^1\text{H}\{^{19}\text{F}\}$ NOEs in fluoroalcohol-induced conformations of peptides and proteins, Martinez and Gerig, **152**, 269
- measurement of interfluorine distances in solids, Gilchrist *et al.*, **152**, 1
- PFG spin echo NMR with homonuclear decoupling, anisotropic diffusion tensor measurement using CSA, Dvinskikh and Furó, **148**, 73
- Faraday induction
- near-field, and quantum origins of free induction decay signal and spin noise, Hoult and Ginsberg, **148**, 182
- Fast-field-cycling applications
- magnet design with high B_0 homogeneity for, Lips *et al.*, **149**, 22
- Fast magic-angle spinning
- high-resolution ^1H NMR, review, Schnell and Spiess, **151**, 153
 - ^1H – ^{15}N correlation and ^1H – ^1H DQ NMR under, inverse detection and heteronuclear editing in, Schnell *et al.*, **150**, 57
 - in the limit of, recoupled polarization-transfer methods for solid-state ^1H – ^{13}C heteronuclear correlation in, Saalwächter, Graf, and Spiess, **148**, 398
 - slow down of ^{13}C spin diffusion in organic solids by, CODEX NMR study, Reichert, Bonagamba, and Schmidt-Rohr, **151**, 129
 - total through-bond correlation spectroscopy, Hardy, Verel, and Meier, **148**, 459
- Fast rotating-frame experiments
- improving resolution in, Casanova, Robert, and Pusiol, **151**, 85
- Fast rotation
- ramped-speed cross-polarization MAS NMR, Samoson, Tuhern, and Past, **149**, 264
- Fat emulsification
- measurement with NMR transverse relaxation, Marciani *et al.*, **153**, 1
- Fe(III)
- doped kaolin clay, NMR relaxation and diffusion measurements on, Valckenborg, Pel, and Kopinga, **151**, 291
- ^{57}Fe
- single-crystal ^{57}Fe Q-band ENDOR study of 4 iron–4 sulfur cluster in reduced $[\text{4Fe–4S}]^{1+}$ state, Moriaud *et al.*, **153**, 238
- Fe^{3+}
- salts, relaxation enhancement of spin labels in membranes by, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Ferritin
- horse spleen, ferromagnetic resonance: core blocking and surface ordering temperatures, Wajnberg *et al.*, **153**, 69
- Ferroelectric cavity insert
- enhanced EPR sensitivity from, Nesmelov, Surek, and Thomas, **153**, 7
- Ferromagnetic resonance
- horse spleen ferritin, core blocking and surface ordering temperatures, Wajnberg *et al.*, **153**, 69
- FID (*see* Free induction decay)
- Field-cycling NMR
- fast-field-cycling applications, magnet design with high B_0 homogeneity for, Lips *et al.*, **149**, 22
- Field mapping
- high-precision mapping of magnetic field using harmonic function mean value property, Li and Leigh, **148**, 442
- Filtering
- approach to solvent peak suppression in MRS, critical review, Coron *et al.*, **152**, 26
- Fitting
- convolution-based, high-spatial-resolution multi-site EPR oximetry using, Grinberg, Smirnov, and Swartz, **152**, 247
 - nonlinear metabolic changes, and time averaging, Simond, Bendahan, and Cozzone, **149**, 1
- Fitting parameters
- constraining, modeling magnetization transfer using, Ceckler, Maneval, and Melkowitz, **151**, 9
- Flavodoxin
- water-dissolved, improved measurement of $^3J(\text{H}_i^\alpha, \text{N}_{i+1})$ coupling constants in, Löhr *et al.*, **153**, 75
- Flow
- angular, in toroid cavity probes, Trautner *et al.*, **151**, 284
 - blood, NMR study, Han *et al.*, **152**, 87
 - bulk convective, convection compensated electrophoretic NMR in presence of, He and Wei, **150**, 126
 - slow, microscopic imaging: STE PFG TSE pulse sequence, Scheenen *et al.*, **151**, 94
 - system undergoing, velocity autocorrelation functions by multiple data acquisition in PFG NMR in, Stapf, **152**, 308
- Fluid solution
- triarylmethyl radicals in, electron spin relaxation, Yong *et al.*, **152**, 156
- Fluorine
- fluorinated ligand–receptor interactions, cross-correlated ^{19}F relaxation measurements for study of, Peng, **153**, 32
- Fluoroalcohol
- induced conformations of peptides/proteins, intermolecular $^1\text{H}\{^{19}\text{F}\}$ NOEs in studies of, Martinez and Gerig, **152**, 269
- Fourier transform
- extended FT for 2D spectral estimation, Armstrong and Mandelshtam, **153**, 22
- Fourier-transform EPR
- demonstration of Q-band pulse EPR/ENDOR spectrometer, Gromov *et al.*, **149**, 196
 - high-field/high-frequency, using broadband stochastic microwave excitation, Fuhs, Prisner, and Möbius, **149**, 67
- Free induction decay
- FID NMR signal dephasing induced by mesoscopic magnetic field inhomogeneities in biological systems, theory of, Sukstanskii and Yablonskiy, **151**, 107
 - quantum origins of, Hoult and Ginsberg, **148**, 182
- Free radicals
- nitroxide, distribution and metabolism in human skin, *in vivo* EPR imaging, He *et al.*, **148**, 155
- Frequency domain
- modified J doubling in, measurement of spin–spin splittings by, deconvolution and, Garza-García, Ponzanelli-Velázquez, and del Río-Portilla, **148**, 214
- Frequency shift
- and angular flow in toroid cavity probes, Trautner *et al.*, **151**, 284
- Frequency sweep
- review of adiabatic principles and methods to create adiabatic pulses, Garwood and DelaBarre, **153**, 155
- G**
- ^{69}Ga
- β - Ga_2O_3 , electron nuclear double resonance in exchange-narrowed regime of ESR, Denninger *et al.*, **148**, 248
- ^{71}Ga
- β - Ga_2O_3 , electron nuclear double resonance in exchange-narrowed regime of ESR, Denninger *et al.*, **148**, 248
- Gallium oxide
- ESR, electron nuclear double resonance in exchange-narrowed regime, Denninger *et al.*, **148**, 248
- Gas-phase NMR microscopy
- dynamic, xenon gas undergoing Poiseuille flow, Kaiser *et al.*, **149**, 144
- Gelatin capsule
- T_1 relaxation time of laser-polarized ^{129}Xe gas in, size dependency and wall effect, Fujiwara *et al.*, **150**, 156
- Glass bulb
- T_1 relaxation time of laser-polarized ^{129}Xe gas in, size dependency and wall effect, Fujiwara *et al.*, **150**, 156

- Gliosarcoma
9L, subcutaneously implanted, TQ-filtered ^{23}Na NMR in rat in presence of TmDOTP $^{5-}$, Winter and Bansal, **152**, 70
- Glucan
osmoregulated periplasmic, *Ralstonia solanacearum*, *in vivo* detection by HR MAS NMR, Wieruszkeski *et al.*, **151**, 118
- Glucose metabolites imaging
proton MRI of ^{13}C distribution by *J* and chemical-shift editing, Casieri *et al.*, **153**, 117
- Glutamine
mammalian brain, ^{13}C - ^{15}N coupling, *in vivo* observation, Kanamori and Ross, **153**, 193
- Glycine
 ^{13}C , ^{15}N -labeled, MAS NMR, double-quantum excitation by pulse-assisted rotational resonance, Karlsson *et al.*, **148**, 238
- Glycosidic angle
from ^{13}C chemical shift data of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
- Glycosidic linkage
in disaccharide, conformation of, double-quantum solid-state NMR, Ravindranathan *et al.*, **151**, 136
- Gradient calibration
accurate measurement of translational diffusion coefficients, Damberg, Jarvet, and Gräslund, **148**, 343
- Gradient modified spin echo
improved spin-echo-edited NMR diffusion measurements, Otto and Larive, **153**, 273
- Gradient pulses
and radiofrequency pulses, rotor synchronization in high-resolution MAS NMR, Wieruszkeski *et al.*, **152**, 95
- Gradients
high-field, ^{31}P solid-state NMR in, Gillies, Newling, and Randall, **151**, 235
- H
- ^1H
crosslinked natural rubber, dipolar and *J*-encoded DQ MAS spectra under rotational resonance, Filip *et al.*, **150**, 184
 ^1H - ^1H MAS correlation spectroscopy and distance measurements in deuterated peptide, Reif *et al.*, **151**, 320
 ^1H - ^{15}N correlation and ^1H - ^1H DQ NMR under fast MAS, inverse detection and heteronuclear editing in, Schnell *et al.*, **150**, 57
 ^1H spectroscopy without solvent suppression, Clayton *et al.*, **153**, 203
improved measurement of ^{15}N - $\{^1\text{H}\}$ NOEs in presence of H(N)-water proton chemical exchange, Idiyatullin, Daragan, and Mayo, **153**, 138
indirect detection, sensitivity-enhanced static ^{15}N NMR of solids by, Hong and Yamaguchi, **150**, 43
measurement of homonuclear proton couplings from regular 2D COSY spectra, Delaglio, Wu, and Bax, **149**, 276
medium- and long-distance ^1H - ^{13}C heteronuclear correlation NMR in solids, Yao, Schmidt-Rohr, and Hong, **149**, 139
MRI, of ^{13}C distribution, by *J* and chemical-shift editing, Casieri *et al.*, **153**, 117
natural abundance ^1H NMR, indirect detection in solids with multiple cross-polarization periods, Khitrin and Fung, **152**, 185
solid-state ^1H - ^{13}C heteronuclear correlation in the limit of fast MAS, recoupled polarization-transfer methods for, Saalwächter, Graf, and Spiess, **148**, 398
- ^2H
dephased REDOR, carbon-observed, effects of nondilute spin pairs: peptide torsion angle measurements, Sack *et al.*, **148**, 104
indirect NMR detection in solids with multiple cross-polarization periods, Khitrin and Fung, **152**, 185
residual ^2H quadrupolar couplings in weakly aligned carbohydrates, Kjellberg and Homans, **151**, 90
stimulated-echo-type ^2H PGSE NMR, diffusion measurement in liquid crystal, Dvinskikh *et al.*, **153**, 83
- Hadamard spectroscopy
in excitation sculpting approach, Krishnamurthy, **153**, 144
- Harmonic function
mean value property, high-precision mapping of magnetic field using, Li and Leigh, **148**, 442
- Hartmann-Hahn transfer
homonuclear
COIN TACS Y experiments, Carlomagno, Prasad, and Glaser, **149**, 52
optimized homonuclear Carr-Purcell-type dipolar mixing sequences, Kramer *et al.*, **149**, 58
- H(CA)CON
triple-resonance assignment experiments, use of carbonyl chemical shift to relieve degeneracies in, Sayers and Torchia, **153**, 246
- Heisenberg spin exchange
and relaxation enhancement of spin labels in membranes by paramagnetic ion salts, Livshits, Dzikovski, and Marsh, **148**, 221
- Hermitian operator basis set
complete, for any spin quantum number, Allard and Härd, **153**, 15
- Heterogeneous media
genuine diffusivities in, determination with stimulated echo PFG NMR, Vasenkov *et al.*, **149**, 228
- Heterogeneous system
water flow and distribution in, MRI-based $\text{D}_2\text{O}/\text{H}_2\text{O}$ -contrast method for study of, Ilvonen *et al.*, **149**, 36
- Heterologous protein
in *Escherichia coli*, selective labeling for NMR studies, Almeida *et al.*, **148**, 142
- Heteronuclear correlation
single-quantum (*see* HSQC)
solid-state ^1H - ^{13}C , in the limit of fast MAS, recoupled polarization-transfer methods for, Saalwächter, Graf, and Spiess, **148**, 398
through-bond HSQC in solid-state NMR, and comparison to other through-bond and through-space experiments, Lesage and Emsley, **148**, 449
- Heteronuclear correlation NMR
medium- and long-distance ^1H - ^{13}C , in solids, Yao, Schmidt-Rohr, and Hong, **149**, 139
through-bond, of sugar and base protons in unlabeled nucleic acids, Phan, **153**, 223
- Heteronuclear editing
and inverse detection, in ^1H - ^{15}N correlation and ^1H - ^1H DQ NMR in solid state under fast MAS, Schnell *et al.*, **150**, 57
- Heteronuclear local field spectroscopy
HCCH-2Q-HLF, analysis of conformation of glycosidic linkage in disaccharide, Ravindranathan *et al.*, **151**, 136
- Heteronuclear multiple-bond correlation
sensitivity enhancement of 2D experiment for long-range coupling constants, Cicero, Barbato, and Bazzo, **148**, 209
- Heteronuclear NMR
spectra, probe design for acquisition from multiple spectra, Zhang, Sweedler, and Webb, **153**, 254
through-bond correlation of sugar and base protons in unlabeled nucleic acids, Phan, **153**, 223
- Heteronuclear shift correlation
simultaneously detected, through multiple and single bonds: HMSC, Burger, Schorn, and Bigler, **148**, 88
two-dimensional pulse sequence for T_2^* measurement of protons in ^{13}C isotopomers, Lambert, Haiber, and Herzog, **150**, 175
- Hexahydro-1,3,5-trinitro-1,3,5-triazine
circular polarization excitation and detection of ^{14}N NQR signals, Lee, Robert, and Lathrop, **148**, 355
- Higher-rank multipoles
effects on relaxation of $I = 3$ spin system, Nosel *et al.*, **150**, 178

- High field
and low temperature, ENDOR spectra recorded at, effect of spin relaxation, Epel *et al.*, **148**, 388
NMR instrumentation to achieve excitation of large bandwidths in high-resolution spectra at, Luchinat *et al.*, **150**, 161
one-micrometer-resolution NMR microscopy, Lee *et al.*, **150**, 207
- High-field EPR
Fourier-transform EPR at 3.4 T/95 GHz using broadband stochastic microwave excitation, Fuhs, Prisner, and Möbius, **149**, 67
magnetically aligned phospholipid bilayers at 94 GHz, Mangels *et al.*, **151**, 253
- High-field gradients
³¹P solid-state NMR in, Gillies, Newling, and Randall, **151**, 235
- High-field NMR
in liquids, improved broadband inversion performance for, Smith, Hu, and Shaka, **151**, 269
- High-frequency EPR
Fourier-transform EPR at 3.4 T/95 GHz using broadband stochastic microwave excitation, Fuhs, Prisner, and Möbius, **149**, 67
- High gradient field
one-micrometer-resolution NMR microscopy, Lee *et al.*, **150**, 207
- High-intensity pulsed field gradients
ultra-high-intensity, for NMR spectroscopy, generation and application, Galvosas *et al.*, **151**, 260
- High magnetic field
one-micrometer-resolution NMR microscopy, Lee *et al.*, **150**, 207
- High-resolution MAS NMR
in vivo detection of cyclic osmoregulated periplasmic glucan in *Ralstonia solanacearum*, Wieruszski *et al.*, **151**, 118
rotor synchronization of RF and gradient pulses in, Wieruszski *et al.*, **152**, 95
- High-resolution NMR
analytical solution to Lipari–Szabo model, with reduced spectral density approximation, Renner, Moroder, and Holak, **151**, 32
at high field, excitation of large bandwidths in spectra, NMR instrumentation for, Luchinat *et al.*, **150**, 161
- HLSVD method
removal of residual water signal by, optimization, Cabanes *et al.*, **150**, 116
- HMQC
and COSY, multiple-coil probe design for acquisition of spectra from multiple samples, Zhang, Sweedler, and Webb, **153**, 254
- HMSC
simultaneously detected heteronuclear shift correlation through multiple and single bonds, Burger, Schorn, and Bigler, **148**, 88
- HNCA
sequential, NMR pulse sequence for protein backbone assignment, Meissner and Sørensen, **150**, 100
- HNCACB
sequential, protein NMR pulse sequence, Meissner and Sørensen, **151**, 328
- HOHAHA
COIN TACSU experiments, Carlomagno, Prasch, and Glaser, **149**, 52
- Homogeneous master equation
complete Hermitian operator basis set for any spin quantum number, Allard and Härd, **153**, 15
- Homonuclear couplings
limits of measurements for larger proteins, extension: *J*-modulated TROSY experiment, Kövér and Batta, **151**, 60
proton, measurement based on cross-peak nulling in CT-COSY, Wu and Bax, **151**, 242
- Homonuclear dipolar decoupling
¹⁹F PGSE NMR with, measurement of anisotropic diffusion tensor measurement using CSA, Dvinskikh and Fűrő, **148**, 73
- Homonuclear NMR spectra
probe design for acquisition from multiple spectra, Zhang, Sweedler, and Webb, **153**, 254
- HSQC
detection of natural abundance ¹⁵N through ³¹P using pulsed-field gradients, Carbajo and López-Ortiz, **148**, 165
¹H–¹⁵N, in solid state under fast MAS, inverse detection and heteronuclear editing in, Schnell *et al.*, **150**, 57
improved resolution using selectively shifted pulses, Krishnamurthy, **153**, 124
MUSIC-based experiments, amino acid type-selective ¹H–¹⁵N correlations aromatic residues, Schubert, Oschkinat, and Schmieder, **153**, 186
MUSIC combined with selective pulses and tuned delays, Schubert, Oschkinat, and Schmieder, **148**, 61
¹⁵N, linewidth-resolved: 3D method to measure ¹⁵N *T*₁ and *T*₂ from linewidths, Heikkinen and Kilpeläinen, **151**, 314
NMR experiments for measuring dipolar couplings in asparagine and glutamine side chains, Permi, **153**, 267
simplified 2D ¹³C^α-coupled ¹⁵N SE–HSQC spectrum, measurement of ¹J_{NC^α} and ²J_{NC^α} from, Heikkinen, Permi, and Kilpeläinen, **148**, 53
through-bond, in solid-state NMR, and comparison to other through-bond and through-space experiments, Lesage and Emsley, **148**, 449
- Hydrogen bonds
improved measurement of ³J(H_i^α, N_{i+1}) in H₂O-dissolved proteins, Löhrl *et al.*, **153**, 75
lengths, in Watson–Crick base pairs, characterization by cross-correlated relaxation, Riek, **149**, 149
- Hyperbolic secant pulses
with controlled offsets and delays: water suppression schemes WASHCODE, Starčuk *et al.*, **152**, 168
- Hyperpolarized noble gas MRI
encoding methods for, signal-to-noise ratio comparison, Zhao *et al.*, **148**, 314
- HYSCORE
demonstration of Q-band pulse EPR/ENDOR spectrometer, Gromov *et al.*, **149**, 196
spin echo signal, properties of, Astashkin and Raitsimring, **148**, 379
- I
- Images
from coil arrays with reactive cross-talk at 1.5 T, differential shading artifact in, spatial dependence, Tropp and Schirmer, **151**, 146
- Imaging
bone, using long echo-train summation, and high-field gradient ³¹P solid-state NMR, Gillies, Newling, and Randall, **151**, 235
electrically detected magnetic resonance of silicon wafer, Sato *et al.*, **153**, 113
in vivo EPR
distribution and metabolism of nitroxide radicals in human skin, He *et al.*, **148**, 155
topical, bridged loop–gap S-band surface resonator for, Petryakov *et al.*, **151**, 124
solid-state NMR, symmetric echo acquisition for absolute-value display in, Matsui and Saito, **149**, 103
transverse relaxation times, by RF field gradients: solvent permeation into polymeric materials, Malveau, Grandclaude, and Canet, **150**, 214
- Indirect detection
¹H, sensitivity-enhanced static ¹⁵N NMR of solids by, Hong and Yamaguchi, **150**, 43
²H NMR, in solids, with multiple cross-polarization periods, Khitrin and Fung, **152**, 185
- Inhomogeneity
*B*₀, magnet design minimizing, for fast-field-cycling NMR applications, Lips *et al.*, **149**, 22
magnetic field, in biological systems, theory of FID NMR signal dephasing induced by, Sukstanskii and Yablonskiy, **151**, 107
RF, effect on heteronuclear dipolar recoupling under MAS in SFAM and REDOR, Nishimura, Fu, and Cross, **152**, 227

- Inhomogeneous fields
improved performance of Carr–Purcell sequences with composite pulses, Hürlimann, **152**, 109
- Inhomogeneous magnetic fields
double-quantum-filtered NMR signals in, Wiesmath *et al.*, **149**, 258
nutration spin echoes after two inhomogeneous RF pulses in, diffusion measurements using, Scharfenecker Ardelean, and Kimmich, **148**, 363
- Integrated microscope
confocal and magnetic resonance, for cellular research, Wind *et al.*, **147**, 371, *erratum*, **151**, 347
- Intermolecular dipolar fields
intermolecular MQ coherence pumping efficiency in solution NMR, Zhang *et al.*, **148**, 200
- Intermolecular multiple-quantum coherences
phase encoding of, MR imaging with, Velan, Narasimhan, and Jacobs, **152**, 189
pumping efficiency in solution NMR, experiment, Zhang *et al.*, **148**, 200
- Internal coordinates
for molecular dynamics and minimization in structure determination/refinement, Schwieters and Clore, **152**, 288
- Internal magnetic field gradients
genuine diffusivities by stimulated echo PFG NMR in heterogeneous media with, Vasenkov *et al.*, **149**, 228
- Inverse detected NMR spectra
probe design for acquisition from multiple spectra, Zhang, Sweedler, and Webb, **153**, 254
- Inverse detection
and heteronuclear editing, in ^1H – ^{15}N correlation and ^1H – ^1H DQ NMR in solid state under fast MAS, Schnell *et al.*, **150**, 57
- Inversion
improved broadband inversion performance for NMR in liquids, Smith, Hu, and Shaka, **151**, 269
- In vivo* echo-planar spectroscopic imaging
 ^{31}P , of human calf muscle, Wilhelm and Bachert, **149**, 126
- In vivo* EPR
at 650 MHz, automatic coupling control of loop–gap resonator by variable capacitor attached coupling coil for, Yokoyama *et al.*, **149**, 29
electron spin T_2 measurements using RF-LOESR and application in oximetry, Pangioteletis, Nicholson, and Hutchison, **149**, 74
minimizing sample motion noise in, resonator with automatic tuning and coupling for, He *et al.*, **149**, 218
topical, bridged loop–gap S-band surface resonator for, Petryakov *et al.*, **151**, 124
- In vivo* NMR
 γ -aminobutyric acid detection by longitudinal scalar order difference editing, de Graaf and Rothman, **152**, 124
high-resolution MAS, detection of cyclic osmoregulated periplasmic glucan in *Ralstonia solanacearum*, Wieruszkeski *et al.*, **151**, 118
 ^{31}P , correcting data for partial saturation, neglecting chemical exchange effects in, Ouwerkerk and Bottomley, **148**, 425; comment and reply, **149**, 251, 282; *erratum*, **151**, 347
triple-quantum-filtered ^{23}Na , of subcutaneous gliosarcoma in presence of TmDOTP $^{5-}$ in rat, Winter and Bansal, **152**, 70
- In vivo* spectroscopy
two-dimensional homonuclear correlation spectroscopy, two-voxel localization sequence for, Delmas *et al.*, **149**, 119
- Iron(III)
doped kaolin clay, NMR relaxation and diffusion measurements on, Valckenborg, Pel, and Kopinga, **151**, 291
- Iron–sulfur proteins
4 iron–4 sulfur cluster in $[\text{4Fe–4S}]^{1+}$ state, single-crystal ^{57}Fe Q-band ENDOR, Moriaud *et al.*, **153**, 238
- Isotope filter
NMR experiment for analyzing sugar puckering in unlabeled DNA using, Iwahara, Wojciak, and Clubb, **153**, 262
- Isotopomers
 ^{13}C , T_2^* measurement of protons in, 2D pulse sequence for, Lambert, Haiber, and Herzog, **150**, 175
- Iterative lineshape analysis
quadrupolar echo spectra of damped CD_3 quantum rotor, Szymański *et al.*, **148**, 277
- J
- Jahn–Teller effect
pseudo-Jahn–Teller low-symmetry Cu(II) complexes in diaqua(L-aspartato) Zn(II) · H_2O crystals, electron spin relaxation in, Hoffmann *et al.*, **153**, 92
- J coupling
 ^{13}C – ^{15}N , in mammalian brain, *in vivo* observation, Kanamori and Ross, **153**, 193
deconvolution and measurement of spin–spin splittings by modified J doubling in frequency domain, Garza-García, Ponzanelli-Velázquez, and del Río-Portilla, **148**, 214
dipolar and J -encoded DQ MAS spectra under rotational resonance, Filip *et al.*, **150**, 184
heteronuclear long-range, sensitivity enhancement of 2D experiment for, Cicero, Barbato, and Bazzo, **148**, 209
homonuclear
 J -modulated TROSY experiment for larger proteins, Kövér and Batta, **151**, 60
homonuclear ^1H – ^1H , measurement
based on cross-peak nulling in CT-COSY, Wu and Bax, **151**, 242
from regular 2D COSY spectra, Delaglio, Wu, and Bax, **149**, 276
 $^3J(\text{H}_i^\alpha, \text{N}_{i+1})$ in H_2O -dissolved proteins, improved measurement, Löhner *et al.*, **153**, 75
 $^1J_{\text{NC}\alpha}$ and $^2J_{\text{NC}\alpha}$, measurement from simplified 2D $^{13}\text{C}^\alpha$ -coupled ^{15}N SE–HSQC spectrum, Heikkinen, Permi, and Kilpeläinen, **148**, 53
proton–carbon coupling constants, determination: spin-state-selective excitation in selective 1D inverse NMR experiments, Parella and Belloc, **148**, 78
and quantification of lactate in muscle using double-quantum filtered ^1H NMR, Asllani *et al.*, **152**, 195
 S^3E -E.COSY methods for measurement of ^{19}F -associated scalar and dipolar coupling constants, Luy, Barchi, and Marino, **152**, 179
spectral editing in solid-state NMR using scalar multiple quantum filters, Sakellariou, Lesage, and Emsley, **151**, 40
spin–spin, small, accurate measurement in partially aligned molecules, Brutscher, **151**, 332
- J doubling
modified, in frequency domain, measurement of spin–spin splittings by, deconvolution and, Garza-García, Ponzanelli-Velázquez, and del Río-Portilla, **148**, 214
- J editing
and chemical-shift editing, proton MRI of ^{13}C distribution by, Casieri *et al.*, **153**, 117
- J modulation
NMR experiment for analyzing sugar puckering in unlabeled DNA using, Iwahara, Wojciak, and Clubb, **153**, 262
- K
- Kaolin clay
iron(III)-doped, NMR relaxation and diffusion measurements on, Valckenborg, Pel, and Kopinga, **151**, 291
- KTaO_3
crystal, ferroelectric cavity insert made from, enhanced EPR sensitivity using, Nesmelov, Surek, and Thomas, **153**, 7

L

- Labeling
selective, heterologous protein in *Escherichia coli* for NMR studies, Almeida *et al.*, **148**, 142
- Lactate
in muscle, quantification using double-quantum filtered ^1H NMR, Asllani *et al.*, **152**, 195
signal loss, and spatial dependence, in PRESS localized ^1H NMR spectroscopy, Jung, Bunse, and Lutz, **152**, 203
- Lanthanide complex
TmDOTA $^-$: sensitive probe for MR thermometry *in vivo*, Zuo, Mahmood, and Sherry, **151**, 101
- Laser-polarized noble gas
 ^{129}Xe , relaxation behavior, Fujiwara *et al.*, **150**, 156
- Ligand–receptor interactions
fluorinated ligand, cross-correlated ^{19}F relaxation measurements for study of, Peng, **153**, 32
- Linear response
delayed-focus pulses optimized using simulated annealing, Shen, **149**, 234
- Linear solvation shift relationships
solvent effects on ^{17}O chemical shifts in methyl formate, de Kowalewski *et al.*, **148**, 1
- Linewidth
-resolved ^{15}N HSQC: 3D method to measure ^{15}N T_1 and T_2 from linewidths, Heikkinen and Kilpeläinen, **151**, 314
- Lipari–Szabo model
analytical solution of, based on reduced spectral density approximation, Renner, Moroder, and Holak, **151**, 32
- $\text{Li}_3\text{P}_7(\text{monoglyme})_3$
solid, bond-shift rearrangement in, ^{31}P MAS NMR study, Jäger *et al.*, **153**, 227
- Liquid crystals
cholesteric phase of filamentous bacteriophage *fd*, characterization for molecular alignment, Barrientos, Louis, and Gronenborn, **149**, 154
dense cholesteric, DNA fragments in water, NMR of spin $I = 3/2$ sodium in, van der Maarel *et al.*, **151**, 298
diffusion coefficients by ^2H stimulated-echo-type PGSE NMR, Dvinskikh *et al.*, **153**, 83
dilute medium, weakly aligned carbohydrates in, residual ^2H quadrupolar couplings, Kjellberg and Homans, **151**, 90
two-dimensional dynamic-director ^{13}C NMR, McElheny, Zhou, and Frydman, **148**, 436
unoriented, measurement of anisotropic diffusion tensor by exploiting chemical shift anisotropy: ^{19}F PGSE NMR with homonuclear decoupling, Dvinskikh and Furó, **148**, 73
- Liquids
adiabatic TOCSY MAS in, Kupče, Keifer, and Delepierre, **148**, 115
high-field NMR in, improved broadband inversion performance for, Smith, Hu, and Shaka, **151**, 269
structured liquid-state samples, imaging the long-range dipolar field in, Bowtell, Gutteridge, and Ramanathan, **150**, 147
- Liquid-state NMR
accurate measurement of small spin–spin couplings in partially aligned molecules, Brutscher, **151**, 332
- Localization
two-voxel sequence, for *in vivo* 2D homonuclear correlation spectroscopy, Delmas *et al.*, **149**, 119
- Localized NMR spectroscopy
 ^1H , PRESS, lactate signal loss and spatial dependence in, Jung, Bunse, and Lutz, **152**, 203
water modeled signal removal and data quantification using time-scale postacquisition, Serrai *et al.*, **149**, 45
- Localized proton spectroscopy
STEAM, of human brain, optimization of residual water signal removal by HLSVD, Cabanes *et al.*, **150**, 116
- LOD EPR
improved instrumentation and new pulse schemes, Granwehr, Forrer, and Schweiger, **151**, 78
magnetometer, for wide-range measurements of low fields, Yokoyama *et al.*, **150**, 194
radiofrequency, electron spin relaxation time measurements using, and application to oximetry, Pangiotelis, Nicholson, and Hutchison, **149**, 74
- Logic gates
and Deutsch–Jozsa quantum algorithm, implementation by 2D NMR using spin- and transition-selective pulses, Mahesh *et al.*, **148**, 95
- Long echo-train summation
imaging bone using, and high-field gradient ^{31}P solid-state NMR, Gillies, Newling, and Randall, **151**, 235
- Longitudinally detected ESR (*see* LOD EPR)
- Longitudinal relaxation (*see* Spin–lattice relaxation)
- Longitudinal scalar order
difference editing, detection of γ -aminobutyric acid by, de Graaf and Rothman, **152**, 124
- Loop–gap resonator
automatic coupling control by variable capacitor attached coupling coil, for EPR measurements at 650 MHz, Yokoyama *et al.*, **149**, 29
bridged loop–gap S-band surface resonator for topical EPR spectroscopy, Petryakov *et al.*, **151**, 124
- Low fields
LODESR magnetometer for wide-range measurements of, Yokoyama *et al.*, **150**, 194
- Low-power water suppression
by hyperbolic secant pulses with controlled offsets and delays, Starčuk *et al.*, **152**, 168
- Low-temperature studies
ENDOR spectra at high magnetic fields in, effect of spin relaxation, Epel *et al.*, **148**, 388
strategy for NMR observation of half-integer quadrupolar nuclei in dilute environments, Lipton, Sears, and Ellis, **151**, 48
- Lysozyme
fluoroalcohol-induced conformations, intermolecular $^1\text{H}^{19}\text{F}$ NOEs in studies of, Martinez and Gerig, **152**, 269
in solution, water molecules strongly bound to, characterization by triple-quantum-filtered ^{17}O NMR, Lehoux, Krzystyniak, and Baguet, **148**, 11

M

- Magic-angle spinning
adiabatic TOCSY, in liquids, Kupče, Keifer, and Delepierre, **148**, 115
 ^{13}C , under moderately fast spinning conditions, spectral editing in, De Vita and Frydman, **148**, 327
CP, analysis of dependence of ^{13}C chemical shifts on conformations of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
dipolar and J -encoded DQ MAS spectra under rotational resonance, Filip *et al.*, **150**, 184
double-quantum double-quantum MAS exchange NMR spectroscopy, Schnell, Watts, and Spiess, **149**, 90
double-quantum excitation by pulse-assisted rotational resonance, Karlsson *et al.*, **148**, 238
double-quantum-filtered rotational-resonance, in presence of large chemical shielding anisotropies, Bechmann, Helluy, and Sebald, **152**, 14
fast (*see* Fast magic-angle spinning)
 ^1H – ^1H MAS correlation spectroscopy in deuterated peptide, Reif *et al.*, **151**, 320

- high-resolution
in vivo detection of cyclic osmoregulated periplasmic glucan in *Ralstonia solanacearum*, Wieruszkeski *et al.*, **151**, 118
 rotor synchronization of RF and gradient pulses in, Wieruszkeski *et al.*, **152**, 95
 magnetization loss during TOCSY experiments under, role of radial B_1 inhomogeneities, Piotto *et al.*, **149**, 114
 measurement of interfluorine distances in solids, Gilchrist *et al.*, **152**, 1
 multiple-quantum (*see* Multiple-quantum MAS NMR)
 ^{31}P , study of bond-shift rearrangement in solid $\text{Li}_3\text{P}_7(\text{monoglyme})_3$, Jäger *et al.*, **153**, 227
 radial-field sidebands in, Tekely and Goldman, **148**, 135
 ramped-speed cross polarization, Samoson, Tuherm, and Past, **149**, 264
- Magic angle turning
 slow, dipolar dephasing method for, Hu *et al.*, **152**, 7
- Magic echo
 application to liquid crystal of diffusion measurement by ^2H stimulated-echo-type PGSE NMR, Dvinskikh *et al.*, **153**, 83
 symmetric echo acquisition for absolute-value display in solid-state imaging, Matsui and Saito, **149**, 103
- Magnet design
 with high B_0 homogeneity for fast-field-cycling NMR applications, Lips *et al.*, **149**, 22
- Magnetic coupling
 creatine methyl protons in skeletal muscle, role of exchangeable NH protons, Kruiskamp and Nicolay, **149**, 8
- Magnetic field gradient
 nonuniform, one-dimensional DOSY using, Loening, Keeler, and Morris, **153**, 103
- Magnetic field inhomogeneities
 mesoscopic, in biological systems, theory of FID NMR signal dephasing induced by, Sukstanskii and Yablonskiy, **151**, 107
- Magnetic fields
 cerium(III)-substituted calbindin D_{9k} , self-orientation in, solution structure calculation through, Bertini *et al.*, **148**, 23
 circularly polarized RF, for spin-1 NQR, Miller, Suits, and Garroway, **151**, 228
 inhomogeneous
 double-quantum-filtered NMR signals in, Wiesmath *et al.*, **149**, 258
 nutation spin echoes after two inhomogeneous RF pulses in, diffusion measurements using, Scharfenecker Ardelean, and Kimmich, **148**, 363
 long-range dipolar, imaging in structured liquid-state samples, Bowtell, Guttridge, and Ramanathan, **150**, 147
 low, LODESR magnetometer for wide-range measurements of, Yokoyama *et al.*, **150**, 194
- Magnetic relaxation dispersion
 proton, measurement of molecular oxygen τ_1 in solutions, Teng *et al.*, **148**, 31
- Magnetic resonance (*see* EPR; *In vivo* EPR; NMR)
- Magnetic resonance imaging
 continuous-wave, of short T_2 materials, Davies *et al.*, **148**, 289
 diffusion tensor, eigenvalues and eigenvectors in, analytical computation, Hasan *et al.*, **152**, 41
 $\text{D}_2\text{O}/\text{H}_2\text{O}$ -contrast method based on, to study water flow and distribution in heterogeneous systems, Ilvonen *et al.*, **149**, 36
 ^1H , of ^{13}C distribution, by J and chemical-shift editing, Casieri *et al.*, **153**, 117
 hyperpolarized noble gas, encoding methods for, signal-to-noise ratio comparison, Zhao *et al.*, **148**, 314
 long-range dipolar field in structured liquid-state samples, Bowtell, Guttridge, and Ramanathan, **150**, 147
 with phase encoding of intermolecular multiple quantum coherences, Velan, Narasimhan, and Jacobs, **152**, 189
 theory of FID NMR signal dephasing induced by magnetic field inhomogeneities, Sukstanskii and Yablonskiy, **151**, 107
- Magnetic resonance microscopy
 integrated with confocal microscopy, for cellular research, Wind *et al.*, **147**, 371, *erratum*, **151**, 347
- Magnetic resonance thermometry
 sensitive probe for *in vivo* use: TmDOTA $^-$, Zuo, Mahmood, and Sherry, **151**, 101
- Magnetic susceptibility
 anisotropic, effect on cross correlation between dipole-dipole interaction and Curie spin relaxation, Bertini *et al.*, **152**, 103
 difference between olive oil and water: fat emulsification measurement by EPI T_2 relaxometry, Marciani *et al.*, **153**, 1
 iron(III)-doped kaolin clay, Valckenborg, Pel, and Kopinga, **151**, 291
- Magnetization
 iron(III)-doped kaolin clay, Valckenborg, Pel, and Kopinga, **151**, 291
 loss during TOCSY experiments under MAS, role of radial B_1 inhomogeneities, Piotto *et al.*, **149**, 114
 selective storage in strongly relaxing spin systems, Kilfoil and Callaghan, **150**, 110
- Magnetization transfer
 modeling using three-pool model and constraining fitting parameters, Ceckler, Maneval, and Melkowitz, **151**, 9
- Magnetization transfer NMR spectroscopy
 recovery of underwater resonances by (RECUR-NMR), Liu *et al.*, **153**, 133
- Magnetometer
 LODESR, for wide-range measurements of low fields, Yokoyama *et al.*, **150**, 194
- Maximum likelihood
 method for determining D_a^{pQ} and R for sets of dipolar coupling data, Warren and Moore, **149**, 271
- Mean value property
 harmonic function, high-precision mapping of magnetic field using, Li and Leigh, **148**, 442
- Membrane protein
 PISEMA powder patterns and PISA wheels, Denny *et al.*, **152**, 217
- Membranes
 model
 magnetically aligned phospholipid bilayers, EPR spectroscopy at 94 GHz, Mangels *et al.*, **151**, 253
 partially oriented, simultaneous determination of orientational and order parameter distributions from NMR spectra, Sternin *et al.*, **149**, 110
 spin labels in, relaxation enhancement by paramagnetic ion salts, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Metabolic changes
 nonlinear, fitting of, and time averaging, Simond, Bendahan, and Cozzone, **149**, 1
- Metabolism
 nitroxide radicals in human skin, *in vivo* EPR imaging, He *et al.*, **148**, 155
- Metabolite concentrations
 measurement in *in vivo* NMR: comment on Ouwerkerk and Bottomley (*J. Magn. Reson.*, **148**, pp. 425–435, 2001), Spencer, Fishbein, and Galban, and reply, **149**, 251, 282; *erratum*, **151**, 347
- Metalloenes
 paramagnetic, solid-state NMR, Heise, Köhler, and Xie, **150**, 198
- Methyl formate
 ^{17}O chemical shifts in, solvent effects, de Kowalewski *et al.*, **148**, 1
- ^{25}Mg
 NMR observation in dilute environments, Lipton, Sears, and Ellis, **151**, 48
- Micron resolution
 one-micrometer-resolution NMR microscopy, Lee *et al.*, **150**, 207
- Microscopic imaging
 slow flow and diffusion: STE PFG TSE pulse sequence, Scheenen *et al.*, **151**, 94
- Microscopy
 echo planar imaging, one-shot velocimetry using, Han and Callaghan, **148**, 349

- Microwave bridge
broadband stochastic, high-field/high-frequency Fourier transform EPR using, Fuhs, Prisner, and Möbius, **149**, 67
- Microwave-induced delayed phosphorescence
determination of relative triplet sublevel populating rates during optical pumping using ODMR, Ozarowski and Maki, **148**, 419
- Minimization
in structure determination/refinement, internal coordinates for, Schwieters and Clore, **152**, 288
- Mixture analysis
improved spin-echo-edited NMR diffusion measurements, Otto and Larive, **153**, 273
- Mn²⁺
salts, relaxation enhancement of spin labels in membranes by, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Molecular alignment
characterization of cholesteric phase of filamentous bacteriophage *fd* for, Barrientos, Louis, and Gronenborn, **149**, 154
small spin-spin couplings in partially aligned molecules, accurate measurement, Brutscher, **151**, 332
- Molecular dynamics
slow, dipolar-coupled spin pairs as probes: DQ-DQ-MAS exchange NMR spectroscopy, Schnell, Watts, and Spiess, **149**, 90
in structure determination/refinement, internal coordinates for, Schwieters and Clore, **152**, 288
- Motion
system undergoing, velocity autocorrelation functions by multiple data acquisition in PFG NMR in, Stapf, **152**, 308
- Motional correlation time distributions
derivation of, and visualizing spectral density functions, Idiyatullin, Daragan, and Mayo, **152**, 132
- Motional parameters
extraction protocol: analytical solution to Lipari-Szabo model, Renner, Moroder, and Holak, **151**, 32
- Multifrequency probe
for pulsed EPR and ENDOR spectroscopy, Avdievich and Gerfen, **153**, 178
- Multiple-coil probe
design, for acquisition of homonuclear, heteronuclear, and inverse detected NMR spectra from multiple samples, Zhang, Sweedler, and Webb, **153**, 254
- Multiple-quantum coherence
imaging the long-range dipolar field in structured liquid-state samples, Bowtell, Gutteridge, and Ramanathan, **150**, 147
intermolecular
phase encoding of, MR imaging with, Velan, Narasimhan, and Jacobs, **152**, 189
pumping efficiency in solution NMR, experiment, Zhang *et al.*, **148**, 200
mixed, high-resolution NMR of quadrupolar nuclei using, Jerschow, Logan, and Pines, **149**, 268
- Multiple-quantum-filtered spectroscopy
TQ-filtered ²³Na NMR of subcutaneous gliosarcoma in presence of TmDOTP⁵⁻ in rat, Winter and Bansal, **152**, 70
- Multiple-quantum filters
scalar, spectral editing in solid-state NMR using, Sakellariou, Lesage, and Emsley, **151**, 40
- Multiple-quantum MAS NMR
high-resolution ¹H, review, Schnell and Spiess, **151**, 153
quadrupolar nuclei, using mixed MQ coherences, Jerschow, Logan, and Pines, **149**, 268
RIACT/MQ-MAS experiments, enhanced sensitivity using RAPPT, Kwak *et al.*, **150**, 71; *erratum*, **153**, 151
- Multipoles
higher-rank, effects on relaxation of *I* = 3 spin system, Nosel *et al.*, **150**, 178
- Multi-site EPR oximetry
high-spatial-resolution, Grinberg, Smirnov, and Swartz, **152**, 247
- Multivoxel localization
two-voxel sequence, for *in vivo* 2D homonuclear correlation spectroscopy, Delmas *et al.*, **149**, 119
- Muscle
lactate in, quantification using double-quantum filtered ¹H NMR, Asllani *et al.*, **152**, 195
- Muscle fiber
EPR spectra, enhanced sensitivity using ferroelectric cavity insert, Nesmelov, Surek, and Thomas, **153**, 7
- MUSIC
amino acid type-selective ¹H-¹⁵N correlations
aromatic residues, Schubert, Oschkinat, and Schmieder, **153**, 186
MUSIC combined with selective pulses and tuned delays, Schubert, Oschkinat, and Schmieder, **148**, 61
- N
- ¹⁴N
¹³C-¹⁴N dipolar coupling and ¹³C chemical shift tensors determined by ¹³C rotary resonance solid-state NMR, Odgaard *et al.*, **148**, 298
NQR, circular polarization excitation and detection in, Lee, Robert, and Lathrop, **148**, 355
- ¹⁵N
amino acid type-selective ¹H-¹⁵N correlations
aromatic residues, Schubert, Oschkinat, and Schmieder, **153**, 186
MUSIC with selective pulses and tuned delays, Schubert, Oschkinat, and Schmieder, **148**, 61
¹³C-¹⁵N coupling in mammalian brain, *in vivo* observation, Kanamori and Ross, **153**, 193
[¹³C, ¹⁵N]-glycine, MAS NMR, double-quantum excitation by pulse-assisted rotational resonance, Karlsson *et al.*, **148**, 238
correlation of ¹⁵N(*i*) with interresidue ¹³C^{*β*}(*i* - 1) of side chains, sequential HNCACB and CBCANH pulse sequences for, Meissner and Sørensen, **151**, 328
¹H-¹⁵N correlation NMR under fast MAS, inverse detection and heteronuclear editing in, Schnell *et al.*, **150**, 57
improved measurement of ¹⁵N-¹H NOEs in presence of H(N)-water proton chemical exchange, Idiyatullin, Daragan, and Mayo, **153**, 138
linewidth-resolved ¹⁵N HSQC: 3D method to measure ¹⁵N *T*₁ and *T*₂ from linewidths, Heikkinen and Kilpeläinen, **151**, 314
natural abundance, detection through ³¹P using pulsed-field gradients, Carbajo and López-Ortiz, **148**, 165
¹⁵N, ¹³C-labeled proteins
backbone assignment, sequential HNCA NMR pulse sequence for, Meissner and Sørensen, **150**, 100
measurement of ¹J_{NC α} and ²J_{NC α} from simplified 2D ¹³C α -coupled ¹⁵N SE-HSQC spectrum, Heikkinen, Permi, and Kilpeläinen, **148**, 53
¹⁵N, ¹H correlation spectrum in Asn and Gln side chains, NMR experiments for measuring dipolar couplings, Permi, **153**, 267
¹⁵N³-¹H³ · · · ¹⁵N¹ hydrogen bond lengths in Watson-Crick base pairs, characterization by cross-correlated relaxation, Riek, **149**, 149
static NMR, of solids, sensitivity enhancement by ¹H indirect detection, Hong and Yamaguchi, **150**, 43
uniformly labeled larger proteins, homonuclear coupling measurements, *J*-modulated TROSY experiment for, Kövér and Batta, **151**, 60
uniformly labeled ubiquitin, separating structure and dynamics in CSA/DD cross-correlated relaxation, Kövér and Batta, **150**, 137
- ²³Na
triple-quantum-filtered NMR of subcutaneous gliosarcoma in presence of TmDOTP⁵⁻ in rat, Winter and Bansal, **152**, 70
- Natural abundance ¹H NMR
indirect detection in solids with multiple cross-polarization periods, Khitrin and Fung, **152**, 185

- Neurotoxin
 scorpion, automatic assignment of NOESY cross peaks and determination of protein structure with NOAH/DIAMOD, Xu *et al.*, **148**, 35
- NH₂ editing
 NMR experiments for measuring dipolar couplings in Asn and Gln side chains, Permi, **153**, 267
- Ni²⁺
 salts, relaxation enhancement of spin labels in membranes by, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Nitroxide radicals
 distribution and metabolism in human skin, *in vivo* EPR imaging, He *et al.*, **148**, 155
- NMR
¹⁰B: effects of higher-rank multipoles on relaxation of *I* = 3 spin system, Nosel *et al.*, **150**, 178
¹³C, chemical shifts, dependence on conformations of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
 double-quantum filtered ¹H, quantification of lactate in muscle, Asllani *et al.*, **152**, 195
 dynamics of spin *I* = 3/2 under spin-locking conditions in ordered environment, van der Maarel *et al.*, **151**, 298
 experiment for analyzing sugar puckering in unlabeled DNA, Iwahara, Wojciak, and Clubb, **153**, 262
 experiments for measuring dipolar couplings in asparagine and glutamine side chains, Permi, **153**, 267
¹H, detection of GABA by longitudinal scalar order difference editing, de Graaf and Rothman, **152**, 124
 improved measurement of ¹⁵N-¹H NOEs in presence of H(N)-water proton chemical exchange, Idiyatullin, Daragan, and Mayo, **153**, 138
 improved spin-echo-edited NMR diffusion measurements, Otto and Larive, **153**, 273
 in liquids, improved broadband inversion performance for, Smith, Hu, and Shaka, **151**, 269
 maximum likelihood method for determining D_a^{pq} and *R* for sets of dipolar coupling data, Warren and Moore, **149**, 271
 modeling static fringe field of superconducting magnets, Jeglič *et al.*, **150**, 39
 noise reduction in quadrupolar echo spectra at short echo times, Furó and Hedin, **152**, 214
 nonparametric, Stoica and Sundin, **152**, 57
¹⁵N static, of solids, sensitivity enhancement by ¹H indirect detection, Hong and Yamaguchi, **150**, 43
¹⁷O, triple-quantum-filtered, characterization of water molecules strongly bound to macromolecules in solution, Lehoux, Krzystyniak, and Baguet, **148**, 11
 observation of half-integer quadrupolar nuclei in dilute environments, strategy, Lipton, Sears, and Ellis, **151**, 48
 phase noise, in Bitter magnets, Sigmund *et al.*, **148**, 309, *erratum*, **151**, 347
³¹P *in vivo*
 data correction for partial saturation, neglecting chemical exchange effects in, Ouwkerk and Bottomley, **148**, 425; comment and reply, **149**, 251, 282; *erratum*, **151**, 347
 fitting of metabolic changes in muscle recorded by, time averaging and, Simond, Bendahan, and Cozzone, **149**, 1
 localized, quantitation of spectra based on reciprocity principle, Kreis *et al.*, **149**, 245
 quantum computation, resonance offset tailored composite pulses suitable for, Cummins and Jones, **148**, 338
 residual ²H quadrupolar couplings in weakly aligned carbohydrates, Kjellberg and Homans, **151**, 90
³³S, taurine detection in biological tissues by, Musio and Sciacovelli, **153**, 259
 selective 1D inverse experiments, spin-state-selective excitation in, Parella and Belloc, **148**, 78
 selective labeling of heterologous protein in *Escherichia coli* for, Almeida *et al.*, **148**, 142
 simulations of experiments, complete Hermitian operator basis set for any spin quantum number, Allard and Härd **153**, 15
 solvent peak suppression in, filtering approach, critical review, Coron *et al.*, **152**, 26
 spectra
 localized proton, of human brain, optimization of residual water signal removal by HLSVD, Cabanes *et al.*, **150**, 116
 partially oriented model membranes, simultaneous determination of orientational and order parameter distributions, Sternin *et al.*, **149**, 110
 spin- $\frac{1}{2}$ nuclei scalar coupled to two equivalent spin-1 nuclei in the limit of slow quadrupolar relaxation, Bernatowicz and Szymański, **148**, 455
 structure refinement, VMD-XPLOR visualization package for, Schwieters and Clore, **149**, 239
 two-dimensional homonuclear correlation, two-voxel localization sequence for, Delmas *et al.*, **149**, 119
¹²⁹Xe, probing proteins in solution by, Locci *et al.*, **150**, 167
 NMR instrumentation
 to achieve excitation of large bandwidths in high-resolution spectra at high field, Luchinat *et al.*, **150**, 161
 NMR microscopy
 dynamic
 gas-phase Poiseuille flow, Kaiser *et al.*, **149**, 144
 imaging of slow flow and diffusion, Scheenen *et al.*, **151**, 94
 one-micrometer-resolution, Lee *et al.*, **150**, 207
 NMR MOUSE
 detection of proton DQ-filtered signals in inhomogeneous magnetic fields, Wiesmath *et al.*, **149**, 258
 NMR probe
 to achieve excitation of large bandwidths in high-resolution spectra at high field, Luchinat *et al.*, **150**, 161
 for acquisition of homonuclear, heteronuclear, and inverse detected NMR spectra from multiple samples, Zhang, Sweedler, and Webb, **153**, 254
 arbitrary geometry, NMR reciprocity theorem for, van der Klink, **148**, 147
 NMR reciprocity theorem
 for arbitrary probe geometry, van der Klink, **148**, 147
 NMR spectrometer
 temperature sensitivity, analyzing and correcting, Bowyer, Swanson, and Morris, **152**, 234
 NOAH/DIAMOD
 automatic assignment of NOESY cross peaks and determination of protein structure of scorpion neurotoxin, Xu *et al.*, **148**, 35
 NOESY
 cross peaks, automatic assignment, and protein structure determination of scorpion neurotoxin, using NOAH/DIAMOD, Xu *et al.*, **148**, 35
 NOESY 1D
 Hadamard excitation sculpting in, Krishnamurthy, **153**, 144
 Noise reduction
 in quadrupolar echo spectra at short echo times, Furó and Hedin, **152**, 214
 sample motion noise, minimizing in *in vivo* EPR, resonator with automatic tuning and coupling for, He *et al.*, **149**, 218
 Non-Fourier-encoded MRI
 encoding methods for, signal-to-noise ratio comparison, Zhao *et al.*, **148**, 314
 Nonlinear gradients
 accounting for, practical method: measurement of translational diffusion coefficients, Damberg, Jarvet, and Gräslund, **148**, 343
 Nonparametric method
 high-resolution, in NMR spectroscopy, Stoica and Sundin, **152**, 57
¹⁵N relaxation
 ubiquitin: analytical solution to Lipari-Szabo model, Renner, Moroder, and Holak, **151**, 32
 Nuclear Overhauser effects
 intermolecular ¹H{¹⁹F}, in fluoroalcohol-induced conformations of peptides and proteins, Martinez and Gerig, **152**, 269

^{15}N -(^1H), improved measurement in presence of H(N)-water proton chemical exchange, Idiyatullin, Daragan, and Mayo, **153**, 138

Nuclear quadrupole resonance

^{14}N , circular polarization excitation and detection in, Lee, Robert, and Lathrop, **148**, 355

pure, revision of spin echoes in, Meriles, **149**, 188

spin-1, circularly polarized RF magnetic fields for, Miller, Suits, and Garroway, **151**, 228

Nuclear quadrupole resonance imaging

improving resolution in fast rotating-frame experiments, Casanova, Robert, and Pusiol, **151**, 85

Nuclei

fast-relaxing, longitudinal relaxation rates, measurement using signal-eliminating relaxation filter, Hansen and Led, **151**, 339

Nucleic acids

unlabeled, through-bond correlation of sugar and base protons in, Phan, **153**, 223

Nutation spectroscopy

improving resolution in fast rotating-frame experiments, Casanova, Robert, and Pusiol, **151**, 85

Nutation spin echoes

diffusion measurements using, Scharfenecker Ardelean, and Kimmich, **148**, 363

O

^{17}O

chemical shifts in methyl formate, solvent effects, de Kowalewski *et al.*, **148**, 1

ODMR

determination of relative triplet sublevel populating rates during optical pumping using, Ozarowski and Maki, **148**, 419

Off-resonance

resonance offset tailored composite pulses, Cummins and Jones, **148**, 338

Olive oil

emulsification, measurement with NMR transverse relaxation, Marciani *et al.*, **153**, 1

One-dimensional DOSY

qualitative spectrum, based on single 1D NMR experiment, Loening, Keeler, and Morris, **153**, 103

One-dimensional NMR

selective inverse, spin-state-selective excitation in, Parella and Belloc, **148**, 78

One-dimensional NMR imaging

enhancement, for measurement of diffusion of paramagnetically labeled proteins in cartilage, Foy and Blake, **148**, 126

Oocytes

Xenopus, integrated confocal and magnetic resonance microscopy, Wind *et al.*, **147**, 371, *erratum*, **151**, 347

Optically detected magnetic resonance

determination of relative triplet sublevel populating rates during optical pumping using, Ozarowski and Maki, **148**, 419

Optical pumping

determination of relative triplet sublevel populating rates using ODMR during, Ozarowski and Maki, **148**, 419

Ordered environment

dynamics of spin $I=3/2$ under spin-locking conditions in, van der Maarel *et al.*, **151**, 298

Ordering temperature

surface, horse spleen ferritin, Wajnberg *et al.*, **153**, 69

Order parameter

extraction: analytical solution to Lipari-Szabo model, Renner, Moroder, and Holak, **151**, 32

Order parameter distributions

and orientational distributions, simultaneous determination from NMR spectra of partially oriented model membranes, Sternin *et al.*, **149**, 110

Orientation

interdomain, multidomain proteins, estimation using rotational diffusion tensor determination from relaxation data, Ghose, Fushman, and Cowburn, **149**, 204

magnetically aligned phospholipid bilayers, EPR spectroscopy at 94 GHz, Mangels *et al.*, **151**, 253

Oriental distributions

and order parameter distributions, simultaneous determination from NMR spectra of partially oriented model membranes, Sternin *et al.*, **149**, 110

Oximetry

application of electron spin relaxation measurements using RF-LODESR, Pangiotelis, Nicholson, and Hutchison, **149**, 74

high-spatial-resolution multi-site EPR, Grinberg, Smirnov, and Swartz, **152**, 247

mechanism of oxygen response in carbon-based sensors, Atsarkin *et al.*, **149**, 85

Oxygen

molecular, spin-lattice relaxation in solutions measured by ^1H magnetic relaxation dispersion, Teng *et al.*, **148**, 31

tissue pO_2 , high-spatial-resolution multi-site EPR oximetry, Grinberg, Smirnov, and Swartz, **152**, 247

Oxygen-17

chemical shifts in methyl formate, solvent effects, de Kowalewski *et al.*, **148**, 1

Oxygen response

in carbon-based sensors, mechanism, Atsarkin *et al.*, **149**, 85

P

^{31}P

detection of natural abundance ^{15}N through, using pulsed-field gradients, Carbajo and López-Ortiz, **148**, 165

in vivo echo-planar spectroscopic imaging of human calf muscle, Wilhelm and Bachert, **149**, 126

in vivo ^{31}P MRS data, correcting for partial saturation, neglecting chemical exchange effects in, Ouwerkerk and Bottomley, **148**, 425; comment and reply, **149**, 251, 282; *erratum*, **151**, 347

localized MR spectra, quantitation based on reciprocity principle, Kreis *et al.*, **149**, 245

MAS NMR, study of bond-shift rearrangement in solid Li_3P_7 (monoglyme) $_3$, Jäger *et al.*, **153**, 227

MRS, fitting of metabolic changes in muscle recorded by, time averaging and, Simond, Bendahan, and Cozzone, **149**, 1

solid-state NMR in high-field gradients, Gillies, Newling, and Randall, **151**, 235

Parallel data acquisition

homonuclear, heteronuclear, and inverse detected NMR spectra from multiple samples, probe design for, Zhang, Sweedler, and Webb, **153**, 254

Paramagnetic center

stable radical in γ -irradiated L-alanine, electron relaxation rate, pulse EPR study, Rakvin *et al.*, **152**, 149

Paramagnetic ion salts

relaxation enhancement of spin labels in membranes by, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221

Paramagnetic metallocenes

solid-state NMR, Heise, Köhler, and Xie, **150**, 198

Paramagnetic molecules

NMR instrumentation to achieve excitation of large bandwidths in high-resolution spectra at high field, Luchinat *et al.*, **150**, 161

- Paramagnetic NMR
relaxation enhancement: test of controlling influence of ZFS rhombicity for $S = 1$, Miller, Lohr, and Sharp, **148**, 267
- Paramagnetism
solution structure calculation through self-orientation in magnetic field of Ce(III)-substituted calbindin D_{9k}, Bertini *et al.*, **148**, 23
- Peak suppression
solvent peaks in MRS, filtering approach to, critical review, Coron *et al.*, **152**, 26
- Peptides
 α -helix-forming, demonstration of approach to visualizing spectral densities and deriving correlation times, Idiyatullin, Daragan, and Mayo, **152**, 132
deuterated, ^1H - ^1H MAS correlation spectroscopy and distance measurements in, Reif *et al.*, **151**, 320
fluoroalcohol-induced conformations, intermolecular $^1\text{H}\{^{19}\text{F}\}$ NOEs in studies of, Martinez and Gerig, **152**, 269
torsion angle measurements: effects of nondilute spin pairs on carbon-observed deuterium-dephased PM5-REDOR, Sack *et al.*, **148**, 104
- Periplasm
Ralstonia solanacearum, cyclic osmoregulated glucan in, *in vivo* detection by HR MAS NMR, Wieruszkeski *et al.*, **151**, 118
- Phase
high-precision mapping of magnetic field using harmonic function mean value property, Li and Leigh, **148**, 442
- Phase cycle scheme
for double PGSE NMR with stimulated echoes, Khrapitchev and Callaghan, **152**, 259
- Phase encoding
intermolecular multiple quantum coherences, MR imaging with, Velan, Narasimhan, and Jacobs, **152**, 189
- Phase noise
NMR, in Bitter magnets, Sigmund *et al.*, **148**, 309, *erratum*, **151**, 347
- Phospholipid bilayers
magnetically aligned, EPR spectroscopy at 94 GHz, Mangels *et al.*, **151**, 253
- Photons
virtual, and quantum origins of free induction decay signal and spin noise, Hoult and Ginsberg, **148**, 182
- π -Echo planar imaging microscopy
one-shot velocimetry using, Han and Callaghan, **148**, 349
- PISA wheels
PISEMA powder patterns and, Denny *et al.*, **152**, 217
- PISEMA
powder patterns, and PISA wheels, Denny *et al.*, **152**, 217
- Planar mixing
analytical transfer functions for two coupled spin-1 nuclei under conditions of, Luy and Glaser, **153**, 210
- Plastocyanin
Anabaena, R_1 , measurement using signal-eliminating relaxation filter, Hansen and Led, **151**, 339
- Poiseuille flow
gas-phase, dynamic NMR microscopy, Kaiser *et al.*, **149**, 144
- Polarization transfer
recoupled polarization-transfer methods for solid-state ^1H - ^{13}C heteronuclear correlation in the limit of fast MAS, Saalwächter, Graf, and Spiess, **148**, 398
- Polycrystalline solids
measurement of anisotropic diffusion tensor by exploiting chemical shift anisotropy: ^{19}F PGSE NMR with homonuclear decoupling, Dvinskikh and Furó, **148**, 73
- Polymeric materials
solvent permeation into, study using T_2 images by RF field gradients, Malveau, Grandclaude, and Canet, **150**, 214
- Polypeptides
backbone dihedral angles, determination in solid-state NMR by DQ ^{13}C CSA measurements, Blanco and Tycko, **149**, 131
- Postacquisition
time-scale, water modeled signal removal and data quantification in localized MRS using, Serrai *et al.*, **149**, 45
- Potassium chloride
crystal, vibronic level of rhodium(II) hexacyanide complex in, electron spin relaxation via, Coelho Neto and Vugman, **150**, 105
- Powder patterns
PISEMA, and PISA wheels, Denny *et al.*, **152**, 217
- Powder spectra
quadrupolar echo spectra at short echo times, noise reduction in, Furó and Hedin, **152**, 214
- PRESS
localized ^1H NMR spectroscopy using, lactate signal loss and spatial dependence in, Jung, Bunse, and Lutz, **152**, 203
- Propagator
PFG, measurements, blood rheology study using, Han *et al.*, **152**, 87
- Protein-DNA complex
26-kDa Dead Ringer-DNA, application of NMR experiment for analyzing sugar puckering in unlabeled DNA, Iwahara, Wojciak, and Clubb, **153**, 262
- Protein interactions
structure characterization with convection compensated electrophoretic NMR, He and Wei, **150**, 126
with xenon: effects on ^{129}Xe chemical shifts in aqueous solution, Rubin *et al.*, **152**, 79
- Proteins
amino acid type-selective ^1H - ^{15}N correlations
aromatic residues, Schubert, Oschkinat, and Schmieder, **153**, 186
MUSIC combined with selective pulses and tuned delays, Schubert, Oschkinat, and Schmieder, **148**, 61
backbone assignment, sequential HNCA NMR pulse sequence for, Meissner and Sørensen, **150**, 100
 ^{13}C , ^{15}N -labeled, measurement of $^1J_{\text{NC}\alpha}$ and $^2J_{\text{NC}\alpha}$ from simplified 2D $^{13}\text{C}\alpha$ -coupled ^{15}N SE-HSQC spectrum, Heikkinen, Permi, and Kilpeläinen, **148**, 53
fluoroalcohol-induced conformations, intermolecular $^1\text{H}\{^{19}\text{F}\}$ NOEs in studies of, Martinez and Gerig, **152**, 269
 H_2O -dissolved, improved measurement of $^3J(\text{H}_i^{\alpha}, \text{N}_{i+1})$ coupling constants in, Löhr *et al.*, **153**, 75
heterologous, in *Escherichia coli*, selective labeling for NMR studies, Almeida *et al.*, **148**, 142
larger, homonuclear coupling measurements, J -modulated TROSY experiment for, Kövér and Batta, **151**, 60
paramagnetically labeled, diffusion in cartilage: enhancement of 1D NMR imaging, Foy and Blake, **148**, 126
in solution
probing by ^{129}Xe NMR, Locci *et al.*, **150**, 167
rotational diffusion tensor from relaxation data, combination of exact and approximate methods, Ghose, Fushman, and Cowburn, **149**, 204
water molecules strongly bound to, characterization by triple-quantum-filtered ^{17}O NMR, Lehoux, Krzystyniak, and Baguet, **148**, 11
well-folded, demonstration of approach to visualizing spectral densities and deriving correlation times, Idiyatullin, Daragan, and Mayo, **152**, 132
- Protein structure
scorpion neurotoxin, determination using NOAH/DIAMOD, Xu *et al.*, **148**, 35
- Proton-carbon coupling constants
spin-state-selective excitation in selective 1D inverse NMR experiments, Parella and Belloc, **148**, 78
- Protons
in ^{13}C isotopomers, T_2^* measurements, 2D pulse sequence for, Lambert, Haiber, and Herzog, **150**, 175
imaging solid materials by quadrupole resonance via, Perlo *et al.*, **150**, 132
methyl, in muscle creatine, magnetic coupling, role of exchangeable NH protons, Kruijskamp and Nicolay, **149**, 8

Proton solid-state NMR

high-resolution, review, Schnell and Spiess, **151**, 153

Pulse cluster

pure-phase selective excitation in fast-relaxing systems, Zangger, Oberer, and Sterk, **152**, 48

Pulsed ENDOR

multifrequency probe for, Avdievich and Gerfen, **153**, 178

Pulsed-field gradient NMR

multiple data acquisition in, determination of velocity autocorrelation functions by, Stapf, **152**, 308

pure-phase selective excitation in fast-relaxing systems, Zangger, Oberer, and Sterk, **152**, 48

stimulated spin echo, determination of genuine diffusivities in heterogeneous media, Vasenkov *et al.*, **149**, 228

ultra-high-intensity magnetic field gradient pulses for, generation and application, Galvosas *et al.*, **151**, 260

Pulsed-field gradients

detection of natural abundance ^{15}N through ^{31}P using, Carbajo and López-Ortiz, **148**, 165

optimization of coherence selection by, sensitivity enhancement of 2D experiment for long-range coupling constants, Cicero, Barbato, and Bazzo, **148**, 209

propagator measurements, blood rheology study using, Han *et al.*, **152**, 87

Pulsed-field gradient sequences

for 2D NMR of velocity exchange: VEXSY and SERPENT, Blümich *et al.*, **152**, 162

Pulsed-field-gradient spin echo NMR

^{19}F , with homonuclear decoupling, measurement of anisotropic diffusion tensor measurement using CSA, Dvinskikh and Furó, **148**, 73

^2H , stimulated-echo-type, diffusion measurement in liquid crystal, Dvinskikh *et al.*, **153**, 83

Pulsed-field-gradient stimulated echo turbo spin echo sequence

microscopic imaging of slow flow and diffusion, Scheenen *et al.*, **151**, 94

Pulsed-gradient spin-echo NMR

double, with stimulated echoes, and phase cycle scheme, Khrapitchev and Callaghan, **152**, 259

high-field, diffusion measurements by, macroscopic background gradient and radiation damping effects, Price *et al.*, **150**, 49

Pulse EPR

improved instrumentation and new pulse schemes for LOD EPR, Granwehr, Forrer, and Schweiger, **151**, 78

longitudinal relaxation of stable radical in γ -irradiated L-alanine, Rakvin *et al.*, **152**, 149

multifrequency probe for, Avdievich and Gerfen, **153**, 178

Q-band pulse EPR/ENDOR spectrometer and implementation of advanced 1D and 2D pulse EPR methodology, Gromov *et al.*, **149**, 196

Pulses

delayed-focus, optimized using simulated annealing, Shen, **149**, 234

selectively shifted, improved resolution using, Krishnamurthy, **153**, 124

spin- and transition-selective, 2D NMR using, implementing logic gates and Deutsch–Jozsa quantum algorithm by, Mahesh *et al.*, **148**, 95

Pulse scheme

two-dimensional, for residual ^2H quadrupolar couplings in weakly aligned carbohydrates, Kjellberg and Homans, **151**, 90

Pulse sequence

Difftrain: true spectroscopic single-scan diffusion measurement, Stamps *et al.*, **151**, 28

sequential HNCA, for protein backbone assignment, Meissner and Sørensen, **150**, 100

two-dimensional, for T_2^* measurement of protons in ^{13}C isotopomers, Lambert, Haiber, and Herzog, **150**, 175

Pulse-train LOD EPR

pulse scheme, Granwehr, Forrer, and Schweiger, **151**, 78

Q

Q-band ENDOR

single-crystal ^{57}Fe , study of 4 iron–4 sulfur cluster in reduced $[\text{4Fe–4S}]^{1+}$ state, Moriaud *et al.*, **153**, 238

Quadrupolar couplings

application to liquid crystal of diffusion measurement by ^2H stimulated-echo-type PGSE NMR, Dvinskikh *et al.*, **153**, 83

residual ^2H , in weakly aligned carbohydrates, Kjellberg and Homans, **151**, 90

Quadrupolar echo spectra

damped CD_3 quantum rotor, iterative lineshape analysis, Szymański *et al.*, **148**, 277

at short echo times, noise reduction in, Furó and Hedin, **152**, 214

Quadrupolar nuclei

half-integer, in dilute environments, NMR observation strategy, Lipton, Sears, and Ellis, **151**, 48

high-resolution NMR using mixed MQ coherences, Jerschow, Logan, and Pines, **149**, 268

spin-3/2, RIACT/MQ-MAS NMR experiments, sensitivity enhancement using RAPT, Kwak *et al.*, **150**, 71; *erratum*, **153**, 151

Quadrupolar relaxation

slow, limit of, NMR spectra of spin- $\frac{1}{2}$ nuclei scalar coupled to two equivalent spin-1 nuclei in, Bernatowicz and Szymański, **148**, 455

Quadrupole coupling tensors

deuterons in α -calcium formate, precision measurement, Schmitt *et al.*, **151**, 65

Quadrupole parameters

determination with composite pulse for spurious signal suppression, Millot and Man, **150**, 10

Quadrupole resonance

solid-state proton imaging detected by, Perlo *et al.*, **150**, 132

Quantum computation

implementing logic gates and Deutsch–Jozsa quantum algorithm by 2D NMR using spin- and transition-selective pulses, Mahesh *et al.*, **148**, 95

resonance offset tailored composite pulses suitable for, Cummins and Jones, **148**, 338

Quantum electrodynamics

quantum origins of free induction decay signal and spin noise, Hoult and Ginsberg, **148**, 182

Quantum rotation

damped, of CD_3 groups, quadrupolar echo spectral information, iterative lineshape analysis, Szymański *et al.*, **148**, 277

Quiet decoupling

two-dimensional experiment separating decoupling sidebands from main peaks, Kupče and Freeman, **151**, 142

R

Radial RF field

sidebands in MAS spectra due to, Tekely and Goldman, **148**, 135

Radiation damping

effect on high-field PGSE NMR diffusion measurements, Price *et al.*, **150**, 49

Radiofrequency field gradients

transverse relaxation time images by, study of solvent permeation into polymeric materials, Malveau, Grandclaude, and Canet, **150**, 214

Radiofrequency inhomogeneity

effect on heteronuclear dipolar recoupling under MAS in SFAM and REDOR, Nishimura, Fu, and Cross, **152**, 227

Radiofrequency-LOD EPR

electron spin relaxation measurements using, and application to oximetry, Pangiotelis, Nicholson, and Hutchison, **149**, 74

Radiofrequency magnetic fields

circularly polarized, for spin-1 NQR, Miller, Suits, and Garroway, **151**, 228

- Radiofrequency pulses
 assisted rotational resonance, double-quantum excitation in NMR of spinning solids by, Karlsson *et al.*, **148**, 238
 delayed-focus pulses optimized using simulated annealing, Shen, **149**, 234
 and gradient pulses, rotor synchronization in high-resolution MAS NMR, Wieruszkeski *et al.*, **152**, 95
 inhomogeneous, nutation spin echoes after, diffusion measurements using, Scharfenecker Ardelean, and Kimmich, **148**, 363
 T_2 -selective, Klose and Raddi, **148**, 47
- Ralstonia solanacearum*
 cyclic osmoregulated periplasmic glucan in, *in vivo* detection by HR MAS NMR, Wieruszkeski *et al.*, **151**, 118
- Ramped spinning speed
 cross-polarization MAS NMR using, Samoson, Tuherm, and Past, **149**, 264
- RAPT (*see* Rotor-assisted population transfer)
- ^{87}Rb
 RbClO_4 , RIACT/MQ-MAS NMR experiments, sensitivity enhancement using RAPT, Kwak *et al.*, **150**, 71; *erratum*, **153**, 151
- RDX
 circular polarization excitation and detection of ^{14}N NQR signals, Lee, Robert, and Lathrop, **148**, 355
- Reciprocity principle
 quantitation of localized ^{31}P MR spectra based on, Kreis *et al.*, **149**, 245
- Reciprocity theorem
 NMR, for arbitrary probe geometry, van der Klink, **148**, 147
- Recombination
 resonance-field dependence in electrically detected MR, effects of exchange interaction, Fukui *et al.*, **149**, 13
- Recoupled polarization transfer
 methods for solid-state ^1H - ^{13}C heteronuclear correlation in the limit of fast MAS, Saalwächter, Graf, and Spiess, **148**, 398
- Recoupling (*see also* Dipolar recoupling)
 recoupled polarization-transfer techniques for solid-state ^1H - ^{13}C heteronuclear correlation in the limit of fast MAS, Saalwächter, Graf, and Spiess, **148**, 398
- RECUR-NMR
 recovery of underwater resonances by magnetization transfer NMR spectroscopy, Liu *et al.*, **153**, 133
- REDOR
 carbon-observed deuterium-dephased PM5-REDOR, effects of nondilute spin pairs: peptide torsion angle measurements, Sack *et al.*, **148**, 104
 heteronuclear dipolar recoupling under MAS for, effect of RF inhomogeneity, Nishimura, Fu, and Cross, **152**, 227
 recoupled polarization-transfer methods for solid-state ^1H - ^{13}C heteronuclear correlation in the limit of fast MAS, Saalwächter, Graf, and Spiess, **148**, 398
- Regularized resolvent transform
 extended Fourier transform for 2D spectral estimation, Armstrong and Mandelshtam, **153**, 22
- Relaxation (*see also* Spin-lattice relaxation; Spin-spin relaxation)
 ^{13}C and ^{15}N NMR: visualizing spectral densities and deriving correlation times, Idiyatullin, Daragan, and Mayo, **152**, 132
 cross-correlated (*see* Cross-correlated relaxation)
 dephasing, of electron spin echo of vibronic $\text{Cu}(\text{H}_2\text{O})_6$ complexes in Tutton salt crystals at low temperatures, Hoffmann *et al.*, **153**, 56
 effects in stray field NMR, Hürlimann, **148**, 367
 $I = 3$ spin system, effects of higher-rank multipoles, Nosel *et al.*, **150**, 178
 measurements
 Carr-Purcell sequences with composite pulses for, Hürlimann, **152**, 109
 improved, of ^{15}N - ^1H NOEs in presence of H(N)-water proton chemical exchange, Idiyatullin, Daragan, and Mayo, **153**, 138
 iron(III)-doped kaolin clay, Valckenborg, Pel, and Kopinga, **151**, 291
 pure-phase selective excitation in fast-relaxing systems, Zangger, Oberer, and Sterk, **152**, 48
 rotational diffusion tensor determination from data, combination of exact and approximate methods, Ghose, Fushman, and Cowburn, **149**, 204
 slow quadrupolar, limit of, NMR spectra of spin- $\frac{1}{2}$ nuclei scalar coupled to two equivalent spin-1 nuclei in, Bernatowicz and Szymański, **148**, 455
- Relaxation behavior
 laser-polarized ^{129}Xe gas, Fujiwara *et al.*, **150**, 156
- Relaxation enhancement
 NMR paramagnetic: test of controlling influence of ZFS rhombicity for $S = 1$, Miller, Lohr, and Sharp, **148**, 267
 spin labels in membranes by paramagnetic ion salts, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Relaxation filter
 signal-eliminating, measurement of longitudinal relaxation rates of fast-relaxing nuclei using, Hansen and Led, **151**, 339
- Relaxation time
 electron spin, measurements using RF-LODESR, and application in oximetry, Pangiotelis, Nicholson, and Hutchison, **149**, 74
- Residual dipolar coupling
 in asparagine and glutamine side chains, NMR experiments for measuring, Permi, **153**, 267
 characterization of cholesteric phase of filamentous bacteriophage *fd* for molecular alignment, Barrientos, Louis, and Gronenborn, **149**, 154
 ^{19}F -associated, S^3E -E.COSY methods for measurement of, Luy, Barchi, and Marino, **152**, 179
 method for accurate measurement of small spin-spin in partially aligned molecules, Brutscher, **151**, 332
 optimized homonuclear Carr-Purcell-type dipolar mixing sequences for, Kramer *et al.*, **149**, 58
 and scalar dipolar coupling, superposition: analytical transfer functions for three spins $\frac{1}{2}$, Luy and Glaser, **148**, 169
- Resin beads
 single solid-phase, cleaved product from, ^1H NMR characterization using capillary NMR flow probe, Lacey *et al.*, **153**, 215
- Resonance-field dependence
 in electrically detected magnetic resonance, effects of exchange interaction, Fukui *et al.*, **149**, 13
- Resonance offset tailoring
 composite pulse sequences produced using, Cummins and Jones, **148**, 338
- Resonance overlaps
 in triple-resonance assignment experiments, use of carbonyl chemical shift to relieve, Sayers and Torchia, **153**, 246
- Resonator
 with automatic tuning and coupling, for minimizing sample motion noise in *in vivo* EPR spectroscopy, He *et al.*, **149**, 218
 multifrequency pulsed EPR/ENDOR probe, Avdievich and Gerfen, **153**, 178
- Restricted self-diffusion
 spin-echo spatial distribution in case of, computer simulation, Duh, Mohorič, and Stepšnik, **148**, 257
- Rheology
 blood, by NMR, Han *et al.*, **152**, 87
 studies, technique for measuring angular flow in toroid cavity probes for use in, Trautner *et al.*, **151**, 284
- Rhodium(II) hexacyanide complex
 in KCl crystal, vibronic level, electron spin relaxation via, Coelho Neto and Vugman, **150**, 105
- Rhombicity
 controlling influence of ZFS rhombicity for $S = 1$: EPR paramagnetic relaxation enhancement, Miller, Lohr, and Sharp, **148**, 267
- RIACT (*see* Rotationally induced adiabatic coherence transfer)
- Ribosomal proteins
 S4 , triple-resonance assignment experiments, use of carbonyl chemical shift to relieve degeneracies, Sayers and Torchia, **153**, 246
- Rifampicin
 selective labeling of heterologous protein in *Escherichia coli* with, for NMR, Almeida *et al.*, **148**, 142

- Ringing
in quadrupolar echo spectra at short echo times, reducing, Furó and Hedin, **152**, 214
- RNA nucleosides
¹³C chemical shifts in, calculation, Rossi and Harbison, **151**, 1
conformations, dependence of ¹³C NMR chemical shifts on, Ebrahimi *et al.*, **150**, 1
- RNA nucleotides
conformations, dependence of ¹³C NMR chemical shifts on, Ebrahimi *et al.*, **150**, 1
- ROESY
recovery of underwater resonances by RECUR-ROESY, Liu *et al.*, **153**, 133
- Rotary resonance solid-state NMR
¹³C chemical shift and ¹³C–¹⁴N dipolar coupling tensors determined by, Odgaard *et al.*, **148**, 298
- Rotating field
circularly polarized RF magnetic fields for spin-1 NQR, Miller, Suits, and Garroway, **151**, 228
- Rotating-frame analog of spin-locking
and spin–lattice relaxation in doubly rotating frame, Tabuchi and Hatanaka, **148**, 121
- Rotating-frame imaging
fast rotating-frame, improving resolution in, Casanova, Robert, and Pusiol, **151**, 85
- Rotational diffusion tensor
from NMR relaxation data, combination of exact and approximate methods, Ghose, Fushman, and Cowburn, **149**, 204
- Rotational echo double resonance (*see* REDOR)
- Rotationally induced adiabatic coherence transfer
RIACT/MQ-MAS NMR experiments, enhanced sensitivity using RAPT, Kwak *et al.*, **150**, 71; *erratum*, **153**, 151
- Rotational resonance
dipolar and *J*-encoded DQ MAS spectra under, Filip *et al.*, **150**, 184
double-quantum-filtered *R*² MAS NMR in presence of large chemical shielding anisotropies, Bechmann, Helluy, and Sebald, **152**, 14
pulse-assisted, double-quantum excitation in NMR of spinning solids by, Karlsson *et al.*, **148**, 238
- Rotor-assisted population transfer
enhanced sensitivity in RIACT/MQ-MAS NMR experiments using, Kwak *et al.*, **150**, 71; *erratum*, **153**, 151
- Rotor synchronization
radiofrequency and gradient pulses in high-resolution MAS NMR, Wieruszkeski *et al.*, **152**, 95
- Rubber
crosslinked natural, DQ buildup and decay curves, Wiesmath *et al.*, **149**, 258
- S
- S³E-E.COSY
methods for measurement of ¹⁹F associated scalar and dipolar coupling constants, Luy, Barchi, and Marino, **152**, 179
- S4
triple-resonance assignment experiments, use of carbonyl chemical shift to relieve degeneracies, Sayers and Torchia, **153**, 246
- Sample rotation
minimization of convection effects in self-diffusion NMR measurements, Esturau *et al.*, **153**, 48
- Saturation
partial, correcting *in vivo* ³¹P MRS data for, neglecting chemical exchange effects in, Ouwerkerk and Bottomley, **148**, 425; comment and reply, **149**, 251, 282; *erratum*, **151**, 347
- Scalar coupling
double-quantum filtered ¹H NMR quantification of lactate in muscle, Asllani *et al.*, **152**, 195
- ¹⁹F-associated, S³E-E.COSY methods for measurement of, Luy, Barchi, and Marino, **152**, 179
- heteronuclear: spectral editing in solid-state NMR using scalar MQ filters, Sakellariou, Lesage, and Emsley, **151**, 40
- NMR spectra of spin- $\frac{1}{2}$ nuclei scalar coupled to two equivalent spin-1 nuclei in the limit of slow quadrupolar relaxation, Bernatowicz and Szymafski, **148**, 455
- Selective excitation
pure-phase, in fast-relaxing systems, Zangger, Oberer, and Sterk, **152**, 48
- Selective experiments
one-dimensional inverse NMR, spin-state-selective excitation in, Parella and Belloc, **148**, 78
- Selective labeling
heterologous protein in *Escherichia coli* for NMR studies, Almeida *et al.*, **148**, 142
- Selective pulses
with MUSIC, and aromatic residues: amino acid type-selective ¹H–¹⁵N correlations, Schubert, Oschkinat, and Schmieder, **153**, 186
and tuned delays, combined with MUSIC: amino acid type-selective ¹H–¹⁵N correlations, Schubert, Oschkinat, and Schmieder, **148**, 61
- Selective storage
magnetization in strongly relaxing spin systems, Kilfoil and Callaghan, **150**, 110
- Self-correcting distance geometry
applied to scorpion neurotoxin, Xu *et al.*, **148**, 35
- Self-diffusion
restricted, spin-echo spatial distribution in case of, computer simulation, Duh, Mohorič, and Stepišnik, **148**, 257
- Self-diffusion NMR
convection effects in measurements, minimization with sample rotation, Esturau *et al.*, **153**, 48
- Self-orientation
in magnetic field of Ce(III)-substituted calbindin D_{9k}, solution structure calculations through, Bertini *et al.*, **148**, 23
- Sensitivity enhancement
¹⁵N spectra of nonspinning solids, by ¹H indirect detection, Hong and Yamaguchi, **150**, 43
in RIACT/MQ-MAS NMR experiments using RAPT, Kwak *et al.*, **150**, 71; *erratum*, **153**, 151
spin-1 NQR, with circularly polarized RF magnetic fields, Miller, Suits, and Garroway, **151**, 228
two-dimensional experiment for measurement of heteronuclear long-range coupling constants, Cicero, Barbato, and Bazzo, **148**, 209
- Sensors
carbon-based, mechanism of oxygen response in, Atsarkin *et al.*, **149**, 85
- Sequential assignment
sequential HNCACB and CBCANH protein NMR pulse sequences for, Meissner and Sørensen, **151**, 328
- Sequential CBCANH
protein NMR pulse sequence, Meissner and Sørensen, **151**, 328
- Sequential HNCA
NMR pulse sequence for protein backbone assignment, Meissner and Sørensen, **150**, 100
- Sequential HNCACB
protein NMR pulse sequence, Meissner and Sørensen, **151**, 328
- SERPENT
and VEXSY, comparison: 2D NMR of velocity exchange, Blümich *et al.*, **152**, 162
- SFAM
heteronuclear dipolar recoupling under MAS for, effect of RF inhomogeneity, Nishimura, Fu, and Cross, **152**, 227
- Shift reagent
TmDOTP⁵⁻, triple-quantum-filtered ²³Na NMR of subcutaneous gliosarcoma in rat in presence of, Winter and Bansal, **152**, 70

- Sidebands
in ^1H spectra without water suppression, Clayton *et al.*, **153**, 203
- Sideband suppression
of decoupling sidebands from main peaks, 2D experiment for, Kupče and Freeman, **151**, 142
- Signal-eliminating relaxation filter
measurement of longitudinal relaxation rates of fast-relaxing nuclei using, Hansen and Led, **151**, 339
- Signal loss
lactate, and spatial dependence, in PRESS localized ^1H NMR spectroscopy, Jung, Bunse, and Lutz, **152**, 203
- Signal-to-noise ratio
and circularly polarized RF magnetic fields for spin-1 NQR, Miller, Suits, and Garroway, **151**, 228
comparison of encoding methods for hyperpolarized noble gas MRI using, Zhao *et al.*, **148**, 314
- Signal processing
methods, and taurine detection in biological tissues by ^{33}S NMR, Musio and Sciacovelli, **153**, 259
- Signal removal
water modeled, and data quantification, in localized MRS using time-scale postacquisition, Serrai *et al.*, **149**, 45
- Signal suppression
residual water signal by HLSVD, optimization, Cabanes *et al.*, **150**, 116
spurious signal, composite pulse for, determination of quadrupole parameters with, Millot and Man, **150**, 10
- Silicon wafer
electrically detected magnetic resonance imaging, Sato *et al.*, **153**, 113
- Simulated annealing
delayed-focus pulses optimized using, Shen, **149**, 234
- Single-bead cleavage NMR
product from, ^1H NMR characterization using capillary NMR flow probe, Lacey *et al.*, **153**, 215
- Single-turn coil
LODESR magnetometer for wide-range measurements of low fields, Yokoyama *et al.*, **150**, 194
- Site-selective spin density determination
electron nuclear double resonance in exchange-narrowed regime of ESR in conducting solids, Denninger *et al.*, **148**, 248
- Size dependency
 T_1 relaxation time of laser-polarized ^{129}Xe gas in glass and gelatin bulbs, Fujiwara *et al.*, **150**, 156
- Skeletal muscle
calf muscle, *in vivo* ^{31}P echo-planar spectroscopic imaging (human), Wilhelm and Bachert, **149**, 126
fitting of metabolic changes recorded by ^{31}P MRS, time averaging and, Simond, Bendahan, and Cozzone, **149**, 1
magnetic coupling of creatine methyl protons, role of exchangeable NH protons, Kruskamp and Nicolay, **149**, 8
- Skin
distribution and metabolism of nitroxide radicals in, *in vivo* EPR imaging, He *et al.*, **148**, 155
topical EPR spectroscopy *in vivo*, bridged loop-gap S-band surface resonator for, Petryakov *et al.*, **151**, 124
- Slice selection
in localized DQ spectral editing, Lei and Dunn, **150**, 17
- Slow magic angle turning
dipolar dephasing method for, Hu *et al.*, **152**, 7
- Slow molecular dynamics
dipolar-coupled spin pairs as probes: DQ-DQ-MAS exchange NMR spectroscopy, Schnell, Watts, and Spiess, **149**, 90
- ^{33}S NMR spectroscopy
taurine detection in biological tissues by, Musio and Sciacovelli, **153**, 259
- Sodium
spin $I = 3/2$, in dense cholesteric liquid crystal in water, NMR, van der Maarel *et al.*, **151**, 298
- Solenoidal coils
for acquisition of homonuclear, heteronuclear, and inverse detected NMR spectra from multiple samples, Zhang, Sweedler, and Webb, **153**, 254
- Solid-phase synthesis
 ^1H NMR characterization of product from single solid-phase resin beads using capillary NMR flow probes, Lacey *et al.*, **153**, 215
- Solids
conducting, exchange-narrowed regime of ESR in, site-selective electron nuclear double resonance in, Denninger *et al.*, **148**, 248
continuous-wave MRI of short T_2 materials, Davies *et al.*, **148**, 289
indirect NMR detection in, with multiple cross-polarization periods, Khitritin and Fung, **152**, 185
organic, ^{13}C spin diffusion in, slow down by fast MAS: CODEX NMR, Reichert, Bonagamba, and Schmidt-Rohr, **151**, 129
sensitivity-enhanced static ^{15}N NMR by ^1H indirect detection, Hong and Yamaguchi, **150**, 43
- Solid-state imaging
proton, by quadrupole resonance, Perlo *et al.*, **150**, 132
symmetric echo acquisition for absolute-value display in, Matsui and Saito, **149**, 103
- Solid-state NMR
adiabatic dipolar recoupling: DREAM scheme, Verel, Ernst, and Meier, **150**, 81
double-quantum, analysis of conformation of glycosidic linkage in disaccharide, Ravindranathan *et al.*, **151**, 136
heteronuclear dipolar recoupling in, effect of RF inhomogeneity, Nishimura, Fu, and Cross, **152**, 227
medium- and long-distance ^1H - ^{13}C heteronuclear correlation NMR, Yao, Schmidt-Rohr, and Hong, **149**, 139
 ^{31}P , in high-field gradients, Gillies, Newling, and Randall, **151**, 235
paramagnetic metallocenes, Heise, Köhler, and Xie, **150**, 198
peptide torsion angle measurements: effects of ^{13}C homonuclear interactions on ^{13}C PM5-REDOR, Sack *et al.*, **148**, 104
PISEMA powder patterns and PISA wheels, Denny *et al.*, **152**, 217
polypeptide backbone dihedral angles in, determination by DQ ^{13}C CSA measurements, Blanco and Tycko, **149**, 131
rotary resonance, ^{13}C chemical shift and ^{13}C - ^{14}N dipolar coupling tensors determined by, Odgaard *et al.*, **148**, 298
spectral editing in, using scalar multiple quantum filters, Sakellariou, Lesage, and Emsley, **151**, 40
through-bond HSQC in, and comparison to other through-bond and through-space experiments, Lesage and Emsley, **148**, 449
triple-resonance, with reduced dimensionality evolution periods, Astrof, Lyon, and Griffin, **152**, 303
- Solid-state NMR imaging
symmetric echo acquisition for absolute-value display in, Matsui and Saito, **149**, 103
- Solution NMR
intermolecular multiple-quantum coherence pumping efficiency, experiment, Zhang *et al.*, **148**, 200
- Solutions
proteins in, probing by ^{129}Xe NMR, Locci *et al.*, **150**, 167
- Solution structure
calculations, through self-orientation in magnetic field of Ce(III)-substituted calcium-binding protein, Bertini *et al.*, **148**, 23
- Solvent effects
on ^{17}O chemical shifts in methyl formate, de Kowalewski *et al.*, **148**, 1
- Solvent peaks
suppression in MRS, filtering approach to, critical review, Coron *et al.*, **152**, 26

- Solvents
permeation into polymeric materials, study using T_2 images by RF field gradients, Malveau, Grandclaude, and Canet, **150**, 214
- Solvent suppression
 ^1H spectroscopy without, Clayton *et al.*, **153**, 203
- Spatial distribution
and lactate signal loss in PRESS localized ^1H NMR spectroscopy, Jung, Bunse, and Lutz, **152**, 203
spin-echo, in case of restricted diffusion, computer simulation, Duh, Mohorič, and Stepišnik, **148**, 257
- Spatial localization
effects of slice-selective excitation/refocusing in localized spectral editing with gradient-selected DQ coherence transfer, Lei and Dunn, **150**, 17
- Spectral assignment
automated NMR, scorpion neurotoxin using NOAH/DIAMOD, Xu *et al.*, **148**, 35
spectral editing
in ^{13}C MAS NMR under moderately fast spinning conditions, De Vita and Frydman, **148**, 327
in solid-state NMR using scalar MQ filters, Sakellariou, Lesage, and Emsley, **151**, 40
- Spectral density approximation
reduced, analytical solution to Lipari–Szabo model based on, Renner, Moroder, and Holak, **151**, 32
- Spectral density functions
visualizing, and deriving motional correlation time distributions, Idiyatullin, Daragan, and Mayo, **152**, 132
- Spectral editing
in ^{13}C MAS NMR under moderately fast spinning conditions, De Vita and Frydman, **148**, 327
detection of GABA by longitudinal scalar order difference editing, de Graaf and Rothman, **152**, 124
improved resolution using selectively shifted pulses, Krishnamurthy, **153**, 124
localized, with gradient-selected DQ coherence transfer, effects of slice-selective excitation/refocusing, Lei and Dunn, **150**, 17
in solid-state NMR, using scalar multiple quantum filters, Sakellariou, Lesage, and Emsley, **151**, 40
- Spectral estimation
two-dimensional, extended Fourier transform for, Armstrong and Mandelshtam, **153**, 22
- Spectral folding
improved resolution using selectively shifted pulses, Krishnamurthy, **153**, 124
- Spectrometer
temperature sensitivity, analyzing and correcting, Bowyer, Swanson, and Morris, **152**, 234
- Spin- $\frac{1}{2}$ nuclei
scalar coupled to two equivalent spin-1 nuclei, NMR spectra in the limit of slow quadrupolar relaxation, Bernatowicz and Szymański, **148**, 455
- Spin-1 nuclei
coupled, analytical planar mixing transfer functions for, Luy and Glaser, **153**, 210
- Spin decoherence
NMR phase noise in Bitter magnets, Sigmund *et al.*, **148**, 309, *erratum*, **151**, 347
- Spin density distribution
in solid-state NMR of paramagnetic metallocenes, Heise, Köhler, and Xie, **150**, 198
- Spin diffusion
 ^{13}C , in organic solids, slow down by fast MAS: CODEX NMR, Reichert, Bonagamba, and Schmidt-Rohr, **151**, 129
- Spin displacements
correlating, 2D PFG sequences VEXSY and SERPENT for, Blümich *et al.*, **152**, 162
- Spin dynamics
spin $I=3/2$ under spin-locking conditions in ordered environment, van der Maarel *et al.*, **151**, 298
- Spin echoes
improved spin-echo-edited NMR diffusion measurements, Otto and Larive, **153**, 273
nutating, diffusion measurements using, Scharfenecker Ardelean, and Kimmich, **148**, 363
properties of HYSORE spin echo signal, Astashkin and Raitsimring, **148**, 379
in pure nuclear quadrupole resonance, revision of, Meriles, **149**, 188
stimulated spin echo PFG NMR, determination of genuine diffusivities in heterogeneous media, Vasenkov *et al.*, **149**, 228
- Spin-echo spatial distribution
in case of restricted diffusion, computer simulation, Duh, Mohorič, and Stepišnik, **148**, 257
- Spin exchange
stochastic, novel mechanism: iterative lineshape analysis of quadrupolar echo spectra of damped CD_3 quantum rotor, Szymański *et al.*, **148**, 277
- Spin labels
in *in vivo* skin, bridged loop-gap S-band surface resonator for topical EPR spectroscopy, Petryakov *et al.*, **151**, 124
in membranes, relaxation enhancement by paramagnetic ion salts, mechanism, Livshits, Dzikovski, and Marsh, **148**, 221
- Spin–lattice relaxation
in doubly rotating frame, rotating-frame analog of spin-locking and, Tabuchi and Hatanaka, **148**, 121
effect on ENDOR spectra at high magnetic fields and low temperature, Epel *et al.*, **148**, 388
effect of neglecting chemical exchange effects when correcting *in vivo* ^{31}P MRS data for partial saturation, Ouwerkerk and Bottomley, **148**, 425; comment and reply, **149**, 251, 282; *erratum*, **151**, 347
formal theory, Advances in Magnetic Resonance review, Goldman, **149**, 160
laser-polarized ^{129}Xe gas in glass and gelatin bulbs, size dependency and wall effect, Fujiwara *et al.*, **150**, 156
and T_2 , ^{15}N , measurement from linewidths, 3D method: linewidth-resolved ^{15}N HSQC, Heikkinen and Kilpeläinen, **151**, 314
- Spin–lattice relaxation rate
fast-relaxing nuclei, measurement using signal-eliminating relaxation filter, Hansen and Led, **151**, 339
molecular oxygen in solutions, measurement by ^1H magnetic relaxation dispersion, Teng *et al.*, **148**, 31
stable radical in γ -irradiated L-alanine, pulse EPR study, Rakvin *et al.*, **152**, 149
via vibronic level of rhodium(II) hexacyanide complex in KCl crystal, Coelho Neto and Vugman, **150**, 105
- Spin-locking
dynamics of spin $I=3/2$ in ordered environment under conditions of, van der Maarel *et al.*, **151**, 298
rotating-frame analog of, and spin–lattice relaxation in doubly rotating frame, Tabuchi and Hatanaka, **148**, 121
- Spinning sidebands
radial-field, in MAS, Tekely and Goldman, **148**, 135
- Spin noise
quantum origins of, Hoult and Ginsberg, **148**, 182
- Spin pairs
dipolar-coupled, as probes for slow molecular dynamics: DQ-DQ-MAS exchange NMR, Schnell, Watts, and Spiess, **149**, 90
- Spins
 $\frac{1}{2}$, three, analytical transfer functions: superposition of scalar and residual dipolar couplings, Luy and Glaser, **148**, 169
 $I=3/2$, dynamics under spin-locking conditions in ordered environment, van der Maarel *et al.*, **151**, 298
spin-1 NQR, circularly polarized RF magnetic fields for, Miller, Suits, and Garroway, **151**, 228

- Spin–spin coupling (*see J* coupling)
- Spin–spin relaxation
- imaging by RF field gradients: solvent permeation into polymeric materials, Malveau, Grandclaude, and Canet, **150**, 214
 - measurement of fat emulsification using, Marciani *et al.*, **153**, 1
 - resistance of delayed-focus pulses optimized using simulated annealing, Shen, **149**, 234
 - selectivity, comparison between different kinds of RF pulses, Klose and Raddi, **148**, 47
 - short T_2 values, continuous-wave MRI of materials with, Davies *et al.*, **148**, 289
 - strong, selective storage of magnetization in system with, Kilfoil and Callaghan, **150**, 110
 - and T_1 , ^{15}N , measurement from linewidths, 3D method: linewidth-resolved ^{15}N HSQC, Heikkinen and Kilpeläinen, **151**, 314
- Spin–spin splittings
- measurement by modified J doubling in frequency domain, deconvolution and, Garza-García, Ponzanelli-Velázquez, and del Río-Portilla, **148**, 214
- Spin-state selective filters
- J -mismatch compensated, accurate measurement of small spin–spin couplings in partially aligned molecules, Brutscher, **151**, 332
 - NMR experiments for measuring dipolar couplings in Asn and Gln side chains, Permi, **153**, 267
- Spin systems
- $I = 3$, relaxation of, effects of higher-rank multipoles, Nosel *et al.*, **150**, 178
 - strongly relaxing, selective storage of magnetization in, Kilfoil and Callaghan, **150**, 110
- Spurious signal suppression
- composite pulse for, determination of quadrupole parameters with, Millot and Man, **150**, 10
- Stable alanine radical
- in γ -irradiated L-alanine, longitudinal relaxation, pulse EPR study, Rakvin *et al.*, **152**, 149
- Static fringe magnetic field
- superconducting magnets, modeling, Jeglič *et al.*, **150**, 39
- STEAM
- localized proton spectroscopy of human brain, optimization of residual water signal removal by HLSVD, Cabanes *et al.*, **150**, 116
- Stimulated echoes
- double PGSE NMR with, and phase cycle scheme, Khrapitchev and Callaghan, **152**, 259
 - STE PFG TSE pulse sequence, in microscopic imaging of slow flow and diffusion, Scheenen *et al.*, **151**, 94
- Stimulated spin echo pulsed-field gradient NMR
- determination of genuine diffusivities in heterogeneous media, Vasenkov *et al.*, **149**, 228
- Stochastic excitation
- broadband microwave, high-field/high-frequency Fourier transform EPR using, Fuhs, Prisner, and Möbius, **149**, 67
- Stochastic spin exchange
- novel mechanism: iterative lineshape analysis of quadrupolar echo spectra of damped CD_3 quantum rotor, Szymański *et al.*, **148**, 277
- Stray field NMR
- ^{31}P solid-state NMR in high-field gradients, Gillies, Newling, and Randall, **151**, 235
 - diffusion and relaxation effects in, Hürlimann, **148**, 367
- Structure
- determination, internal coordinates for molecular dynamics and minimization in, Schwieters and Clore, **152**, 288
 - and dynamics, separating in CSA/DD cross-correlated relaxation, Kövér and Batta, **150**, 137
 - interacting proteins, characterization with convection compensated electrophoretic NMR, He and Wei, **150**, 126
- Structure refinement
- internal coordinates for molecular dynamics and minimization in, Schwieters and Clore, **152**, 288
- NMR, VMD-XPLOR visualization package for, Schwieters and Clore, **149**, 239
- Sugar protons
- and base protons, through-bond correlation in unlabeled nucleic acids, Phan, **153**, 223
- Sugar pucker
- from ^{13}C chemical shift data of RNA nucleosides and nucleotides, Ebrahimi *et al.*, **150**, 1
 - ribose, and calculation of ^{13}C chemical shifts in RNA nucleosides, Rossi and Harbison, **151**, 1
 - in unlabeled DNA, NMR experiment for analyzing, Iwahara, Wojciak, and Clubb, **153**, 262
- Superconducting magnets
- static fringe field of, modeling, Jeglič *et al.*, **150**, 39
- Symmetric echo acquisition
- for absolute-value display in solid-state NMR imaging, Matsui and Saito, **149**, 103
- T
- T_1 (*see* Spin–lattice relaxation)
- T_2 (*see* Spin–spin relaxation)
- T_2^* measurement
- protons in ^{13}C isotopomers, 2D pulse sequence for, Lambert, Haiber, and Herzog, **150**, 175
- TACSy
- COIN TACSy, Carlomagno, Prasch, and Glaser, **149**, 52
- Tailored correlation spectroscopy (*see* TACSy)
- Taurine
- detection in biological tissues by ^{33}S NMR spectroscopy, Musio and Sciacovelli, **153**, 259
- Temperature
- effect on EPR spectra of horse spleen ferritin: core blocking and surface ordering temperatures, Wajnberg *et al.*, **153**, 69
 - variable, BP-LED experiments with, diffusion-broadened velocity spectra of convection in, Mao and Kohlmann, **150**, 35
- Temperature sensitivity
- spectrometer, analyzing and correcting, Bowyer, Swanson, and Morris, **152**, 234
- Tentagel beads
- single solid-phase, cleaved product from, ^1H NMR characterization using capillary NMR flow probe, Lacey *et al.*, **153**, 215
- 1,2,4,5-Tetramethylbenzene
- natural abundance ^2H NMR spectrum, indirect detection in solids, Khitrin and Fung, **152**, 185
- Thermometry
- magnetic resonance, sensitive probe for *in vivo* use: TmDOTA $^-$, Zuo, Mahmood, and Sherry, **151**, 101
- Three-dimensional NMR
- automatic assignment of NOESY cross peaks and determination of protein structure of scorpion neurotoxin with NOAH/DIAMOD, Xu *et al.*, **148**, 35
 - method to measure ^{15}N T_1 and T_2 from linewidths: linewidth-resolved ^{15}N HSQC, Heikkinen and Kilpeläinen, **151**, 314
- Three-pool model
- modeling magnetization transfer using, Ceckler, Maneval, and Melkowitz, **151**, 9
- Through-bond correlation
- sugar and base protons in unlabeled nucleic acids, Phan, **153**, 223
- Through-bond spectroscopy
- fast MAS total through-bond correlation spectroscopy, Hardy, Verel, and Meier, **148**, 459
 - HSQC, in solid-state NMR, and comparison to other through-bond and through-space experiments, Lesage and Emsley, **148**, 449
- Thulium 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (*see* TmDOTA $^-$)

- Thulium 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetrakis(methylene phosphonate) (*see* TmDOTP⁵⁻)
- Time averaging
and fitting of nonlinear metabolic changes, Simond, Bendahan, and Cozzone, **149**, 1
- Time index
choice, for fitting nonlinear metabolic changes: ³¹P MRS analysis of muscle energetics, Simond, Bendahan, and Cozzone, **149**, 1
- Time-resolved EPR
electron-spin-polarization pathways of *p*-benzosemiquinone, Jäger and Norris, **150**, 26
- Tissue pO₂
high-spatial-resolution multi-site EPR oximetry, Grinberg, Smirnov, and Swartz, **152**, 247
- TmDOTA⁻
sensitive probe for MR thermometry *in vivo*, Zuo, Mahmood, and Sherry, **151**, 101
- TmDOTP⁵⁻
triple-quantum-filtered ²³Na NMR of subcutaneous gliosarcoma in rat in presence of, Winter and Bansal, **152**, 70
- TOBSY
fast MAS, Hardy, Verel, and Meier, **148**, 459
- TOCSY
fast MAS total through-bond correlation spectroscopy, Hardy, Verel, and Meier, **148**, 459
under MAS conditions, loss of magnetization during, role of radial *B*₁ inhomogeneities, Piotto *et al.*, **149**, 114
recovery of underwater resonances by RECUR-TOCSY, Liu *et al.*, **153**, 133
- TOCSY MAS
adiabatic, in liquids, Kupče, Keifer, and Delepierre, **148**, 115
- Topical spectroscopy
in vivo, bridged loop-gap S-band surface resonator for, Petryakov *et al.*, **151**, 124
- Toroid cavity detector
angular flow in, Trautner *et al.*, **151**, 284
- Torsion angle
peptide, measurements: effects of nondilute spin pairs on carbon-observed deuterium-dephased PM5-REDOR, Sack *et al.*, **148**, 104
- Total correlation spectroscopy (*see* TOCSY)
- Total through-bond correlation spectroscopy (*see* TOBSY)
- Transient nutation LOD EPR
pulse scheme, Granwehr, Forrer, and Schweiger, **151**, 78
- Translational diffusion coefficients
accurate measurement, Damberg, Jarvet, and Gräslund, **148**, 343
- Transverse relaxation (*see* Spin-spin relaxation)
- Trehalose
conformation of glycosidic linkage in, double-quantum solid-state NMR, Ravindranathan *et al.*, **151**, 136
CSA/DD cross-correlated relaxation, separating structure and dynamics in, Kövér and Batta, **150**, 137
- Triarylmethyl radicals
in fluid solution, electron spin relaxation, Yong *et al.*, **152**, 156
- Trifluoroethanol
induced conformations of peptides/proteins, intermolecular ¹H¹⁹F NOEs in studies of, Martinez and Gerig, **152**, 269
- Triple-quantum-filtered NMR
²³Na, of subcutaneous gliosarcoma in presence of TmDOTP⁵⁻ in rat, Winter and Bansal, **152**, 70
- Triple-quantum-filtered spectra
¹⁷O NMR, characterization of water molecules strongly bound to macromolecules in solution, Lehoux, Krzystyniak, and Baguet, **148**, 11
- Triple resonance
amino acid type-selective ¹H-¹⁵N correlations
aromatic residues, Schubert, Oschkinat, and Schmieder, **153**, 186
MUSIC combined with selective pulses and tuned delays, Schubert, Oschkinat, and Schmieder, **148**, 61
solid-state NMR experiments, with reduced dimensionality evolution periods, Astrof, Lyon, and Griffin, **152**, 303
- Triple-resonance assignment experiments
use of carbonyl chemical shift to relieve degeneracies in, Sayers and Torchia, **153**, 246
- Triplet state sublevels
photoexcited, relative populating rates during optical pumping, determination using ODMR, Ozarowski and Maki, **148**, 419
- TROSY
accurate measurement of small spin-spin in partially aligned molecules, Brutscher, **151**, 332
clean-TROSY-adapted implementation of sequential HNCACB and CBCANH pulse sequences, Meissner and Sørensen, **151**, 328
improved measurement of ³J(H_{*i*}^α, N_{*i*+1}) coupling constants in H₂O-dissolved proteins, Löhner *et al.*, **153**, 75
J-modulated, extension of limits of homonuclear coupling measurements for larger proteins, Kövér and Batta, **151**, 60
sequential HNCA, protein backbone assignment, Meissner and Sørensen, **150**, 100
- Tuned delays
and selective pulses, combined with MUSIC: amino acid type-selective ¹H-¹⁵N correlations, Schubert, Oschkinat, and Schmieder, **148**, 61
- Tuning
automatic, resonator with, for minimizing sample motion noise in *in vivo* EPR spectroscopy, He *et al.*, **149**, 218
- Tunneling
and longitudinal relaxation of stable radical in γ -irradiated L-alanine, pulse EPR study, Rakvin *et al.*, **152**, 149
- Turbo spin echo imaging
PFG STE sequence combined with, in microscopic imaging of slow flow and diffusion, Scheenen *et al.*, **151**, 94
- Turbulent flow
rheology of blood by NMR, Han *et al.*, **152**, 87
- Tutton salts
vibronic Cu(H₂O)₆ complexes in crystals at low temperatures, dephasing relaxation of, Hoffmann *et al.*, **153**, 56
- Two-dimensional NMR
dynamic-director ¹³C NMR of liquid crystals, McElheny, Zhou, and Frydman, **148**, 436
FIREMAT and 2D-PASS, dipolar dephasing method for slow MAT, Hu *et al.*, **152**, 7
measurement of heteronuclear long-range coupling constants, sensitivity enhancement, Cicero, Barbato, and Bazzo, **148**, 209
pulse sequence for T₂^{*} measurement of protons in ¹³C isotopomers, Lambert, Haiber, and Herzog, **150**, 175
regular COSY spectra, measurement of homonuclear proton couplings from, Delaglio, Wu, and Bax, **149**, 276
using spin- and transition-selective pulses, implementing logic gates and Deutsch-Jozsa quantum algorithm by, Mahesh *et al.*, **148**, 95
velocity exchange: VEXSY and SERPENT, Blümich *et al.*, **152**, 162
- Two-dimensional processing
nonparametric NMR spectroscopy, Stoica and Sundin, **152**, 57
- Two-dimensional spectral estimation
extended Fourier transform for, Armstrong and Mandelshtam, **153**, 22
- Two-dimensional spectroscopy
in vivo homonuclear correlation, two-voxel localization sequence for, Delmas *et al.*, **149**, 119
separation of decoupling sidebands from main peaks, Kupče and Freeman, **151**, 142

- linewidth-resolved ^{15}N HSQC: 3D method to measure ^{15}N T_1 and T_2 from linewidths, Heikkinen and Kilpeläinen, **151**, 314
- ^{15}N , ^{13}C -labeled, measurement of $^1J_{\text{NC}\alpha}$ and $^2J_{\text{NC}\alpha}$ from simplified 2D $^{13}\text{C}\alpha$ -coupled ^{15}N SE-HSQC spectrum, Heikkinen, Permi, and Kilpeläinen, **148**, 53
- ^{15}N relaxation: analytical solution to Lipari-Szabo model, Renner, Moroder, and Holak, **151**, 32
- Ultra-high-intensity pulsed-field gradients
for NMR spectroscopy, generation and application, Galvosas *et al.*, **151**, 260
- Underwater resonances
recovery by magnetization transfer NMR spectroscopy (RECUR-NMR), Liu *et al.*, **153**, 133
- Unlabeled DNA
sugar puckering in, NMR experiment for analyzing, Iwahara, Wojciak, and Clubb, **153**, 262
- Unlabeled nucleic acids
through-bond correlation of sugar and base protons in, Phan, **153**, 223
- Unoriented sample
anisotropic diffusion tensor measurement using CSA: ^{19}F PGSE NMR with homonuclear decoupling, Dvinskikh and Furó, **148**, 73

V

- Varactor diode
resonator with automatic tuning and coupling for minimizing sample motion noise in *in vivo* EPR, He *et al.*, **149**, 218
- Velocimetry
one-shot, using echo planar imaging microscopy, Han and Callaghan, **148**, 349
- Velocity autocorrelation function
determination by multiple data acquisition in NMR PFG experiments, Stapf, **152**, 308
- Velocity exchange
two-dimensional NMR: VEXSY and SERPENT, Blümich *et al.*, **152**, 162
- Velocity fluctuations
double PGSE NMR with stimulated echoes for study of, and phase cycle scheme, Khrapitchev and Callaghan, **152**, 259
- Velocity imaging
two-dimensional NMR, blood rheology study using, Han *et al.*, **152**, 87
- Velocity spectra
diffusion-broadened, of convection in variable-temperature BP-LED experiments, Mao and Kohlmann, **150**, 35
- VEXSY
and SERPENT, comparison: 2D NMR of velocity exchange, Blümich *et al.*, **152**, 162
- Vibronic level
rhodium(II) hexacyanide complex in KCl crystal, electron spin relaxation via, Coelho Neto and Vugman, **150**, 105
- Virtual photons
and quantum origins of free induction decay signal and spin noise, Hoult and Ginsberg, **148**, 182
- VMD-XPLOR visualization package
for NMR structure refinement, Schwieters and Clore, **149**, 239

W

- Wall effect
on T_1 relaxation time of laser-polarized ^{129}Xe gas in glass and gelatin bulbs, Fujiwara *et al.*, **150**, 156

WASHCODE

- low-power water suppression by hyperbolic secant pulses with controlled offsets and delays, Starčuk *et al.*, **152**, 168
- Water
 ^{17}O , strongly bound to macromolecules in solution, characterization by triple-quantum-filtered NMR, Lehoux, Krzystyniak, and Baguet, **148**, 11
- Water flow
and distribution, in heterogeneous systems, MRI-based $\text{D}_2\text{O}/\text{H}_2\text{O}$ -contrast method for study of, Ilvonen *et al.*, **149**, 36
- Water signal
residual, removal by HLSVD, optimization, Cabanes *et al.*, **150**, 116
- Water suppression
data quantification without, in localized MRS using time-scale postacquisition, Serrai *et al.*, **149**, 45
- low-power, by hyperbolic secant pulses with controlled offsets and delays, Starčuk *et al.*, **152**, 168
- Watson-Crick base pairs
hydrogen bond lengths in, characterization by cross-correlated relaxation, Riek, **149**, 149
- Wavelet encoding
signal-to-noise ratio comparison of encoding methods for hyperpolarized noble gas MRI, Zhao *et al.*, **148**, 314
- Wide-range measurements
low fields, LODESR magnetometer for, Yokoyama *et al.*, **150**, 194
- Wood xylem
water flow and distribution in, analysis with MRI-based $\text{D}_2\text{O}/\text{H}_2\text{O}$ -contrast method, Ilvonen *et al.*, **149**, 36

X

- ^{129}Xe
chemical shifts in aqueous solution, effects of nonspecific xenon-protein interactions, Rubin *et al.*, **152**, 79
- laser-polarized, relaxation behavior, Fujiwara *et al.*, **150**, 156
- ^{129}Xe NMR
probing proteins in solution by, Locci *et al.*, **150**, 167
- Xenon
gas-phase Poiseuille flow, dynamic NMR microscopy, Kaiser *et al.*, **149**, 144
- Xenon-protein interactions
nonspecific, effects on ^{129}Xe chemical shifts in aqueous solution, Rubin *et al.*, **152**, 79

Z

- Zero field splitting
controlling influence of ZFS rhombicity for $S = 1$: EPR paramagnetic relaxation enhancement, Miller, Lohr, and Sharp, **148**, 267
- magnetic anisotropy from, and cross correlation between dipolar interaction and Curie spin relaxation, Bertini *et al.*, **152**, 103
- ^{67}Zn
NMR observation in dilute environments, Lipton, Sears, and Ellis, **151**, 48
- $\text{Zn}(\text{L-aspartate}) \cdot 3\text{H}_2\text{O}$
pseudo-Jahn-Teller low-symmetry Cu(II) complexes in, electron spin relaxation in, Hoffmann *et al.*, **153**, 92