



ACADEMIC
PRESS

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Journal of Magnetic Resonance 159 (2002) 255–275

JMR

Journal of
Magnetic Resonance

www.academicpress.com

Cumulative Subject Index for 2002¹ Volumes 154–159

A

- Acoustic ringing
in rapid imaging pulse trains, compensation: RIDE'n RIPT, Woelk *et al.*, **159**, 207
- Adiabatic passage
in earth's-field NMR technique, condition for, Melton and Pollak, **158**, 15
- Adiabatic pulses
linear and partial, universal equations for, Tesiram and Bendall, **156**, 26
MR tagging using tandem of two adiabatic DANTE inversion sequences, Tsekos, Garwood, and Ugurbil, **156**, 187
- Adjustable Broadband Selective Excitation with Uniform Phase
for NMR of liquids, Cano, Smith, and Shaka, **155**, 131
- Adsorption
cesium in hydrated iron oxide particle suspensions, NMR study, Gossuin *et al.*, **157**, 132
nitric oxide on H-ZSM-5, Na-ZSM-5, Na-A, EPR study, Rudolf, Böhlmann, and Pöpl, **155**, 45
- Aeration
and mixing, cell and organelle suspensions during NMR experiments, device for, Bourne and Kuchel, **159**, 158
- Affinity NMR
analysis of protein–ligand interactions with NMR diffusion measurements, Derrick, McCord, and Larive, **155**, 217
- Age
and porcine muscle water characteristics, *in vitro* low-field NMR relaxation study, Bertram *et al.*, **157**, 267
- Akaganeite
particle suspensions, NMR study of cesium adsorption, Gossuin *et al.*, **157**, 132
relaxation enhancement by, role of diffusion and exchange, Gossuin *et al.*, **158**, 36
- Algorithms
entropy minimization-based, for automatic phase correction of NMR spectra, Chen *et al.*, **158**, 164
Lanczos, improved, for blackbox MRS data quantitation, Laudadio *et al.*, **157**, 292
subspace time-domain, for automated NMR spectral normalization, Lemmerling *et al.*, **157**, 190
- Alignment
magnetic, of phage Pf1, capture and manipulation with aqueous polymer gel, Riley, Giuliani, and Augustine, **159**, 83
- Alignment tensors
relative orientation, for protein dissolved in different anisotropic media, graphical analysis, Nomura and Kainosho, **154**, 146
- Alpha-helical peptides and proteins
torsion angle Ψ in, NMR determination, Ladizhansky, Veshtort, and Griffin, **154**, 317
- Amide protons
relaxation and chemical shift in perdeuterated protein, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181
- temperature coefficients, location of hydrogen bonds in human ubiquitin using, Cierpicki *et al.*, **157**, 178
- Amide resonances
backbone, correlation with side-chain ¹³C resonances in perdeuterated proteins, Löhr and Rüterjans, **156**, 10
- Amino acids
¹³C NMR, magnesium chloride as relaxation agent for, Tian *et al.*, **159**, 137
- Amorphous matrices
large-angle jumps of small molecules in, 2D exchange NMR analysis, Medick, Vogel, and Rössler, **159**, 126
- Anisotropic media
different, relative orientation of molecular alignment tensors for protein dissolved in, graphical analysis, Nomura and Kainosho, **154**, 146
- Anisotropic diffusion
water in macroscopically oriented lipid bilayers, pulsed magnetic field gradient NMR, Wästerby, Orädd, and Lindblom, **157**, 156
- Anisotropic interaction tensors
in polypeptides, specification and visualization, and numerical simulations in biological solid-state NMR, Bak *et al.*, **154**, 28
- Anisotropic molecular motion
half-integer quadrupolar nuclei, analysis by central transition NMR spectroscopy, Kristensen and Farnan, **158**, 99
- Anisotropic relaxation
half-integer quadrupolar nuclei, analysis by central transition NMR spectroscopy, Kristensen and Farnan, **158**, 99
- Announcements
NMR Nomenclature: Nuclear Spin Properties and Conventions for Chemical Shifts—IUPAC Recommendations, **156**, 323
The Russell Varian Prize, **212**, 161
- Aqueous polymer gel
capture and manipulation of magnetically aligned phage Pf1 with, Riley, Giuliani, and Augustine, **159**, 83
- ARMA modeling
subband, approach to high-resolution NMR spectroscopy, Tomczak and Djermoune, **158**, 86
- Artifact
automated correction of unwanted phase jumps in reference signals corrupting MRSI spectra after eddy current correction, Simonetti *et al.*, **159**, 151
two-dimensional, from asynchronous decoupling, van Ingen, Vuister, and Tessari, **156**, 258
- Artificial neural network
automated, quantification of human brain metabolites from *in vivo* ¹H NMR magnitude spectra, Hiltunen *et al.*, **154**, 1
- Asymmetric populations
*R*_{1ρ} relaxation outside of fast-exchange limit, Trott and Palmer, **154**, 157
- Asynchronous decoupling
two-dimensional artifact from, van Ingen, Vuister, and Tessari, **156**, 258
- Automated correction
unwanted phase jumps in reference signals corrupting MRSI spectra after eddy current correction, Simonetti *et al.*, **159**, 151

¹Boldface numbers indicate volume; lightface numbers indicate pagination.

- Automatic frequency alignment
in multiple MR spectra via complex principal component analysis, Van Huffel *et al.*, **158**, 1
- Automatic tuning
MRI RF coil with, for multinuclear imaging of small animals at 3T, Muftuler *et al.*, **155**, 39
- Autoregressive moving-average process
subband ARMA modeling approach to high-resolution NMR spectroscopy, Tomczak and Djermoune, **158**, 86
- ## B
- B_0 homogeneity
variation with head pitch, quantification by registered 3D field mapping, Tyszka and Mamelak, **159**, 213
- Backbone
amide resonances, correlation with side-chain ^{13}C resonances in perdeuterated proteins, Löhr and Rüterjans, **156**, 10
- Backbone assignment
protein, intraresidue HNCA and COHNCA experiments for, Brutscher, **156**, 155
- Backbone carbons
in membrane-bound fusion peptides, filtered MAS observation, application of REDOR subtraction, Yang *et al.*, **159**, 101
- Bacteriophage
Pfl1, magnetically aligned, capture and manipulation with aqueous polymer gel, Riley, Giuliani, and Augustine, **159**, 83
- Benzene
in amorphous matrices, large-angle jumps of, 2D exchange NMR analysis, Medick, Vogel, and Rössler, **159**, 126
- Bicelles
transmembrane peptides in, study methods: hydrophobic mismatch and peptide sequence, Whiles *et al.*, **158**, 149
- Biomedical signal processing
improved Lanczos algorithms for blackbox MRS data quantitation, Laudadio *et al.*, **157**, 292
- Bi-radical
electron–electron interaction in, 130 GHz ESEEM induced by, Kulik, Paschenko, and Dzuba, **159**, 237
- Birdcage coil
automatic tuned MRI RF coil for multinuclear imaging of small animals at 3T, Muftuler *et al.*, **155**, 39
double end-cap, for small animal whole-body imaging, Gulsen, Muftuler, and Nalcioglu, **156**, 309
- BIRD pulse
determination of small one-bond proton-carbon residual dipolar coupling constants in ^{13}C natural abundance carbohydrates, Pham *et al.*, **157**, 200
- Bisadducts
fullerene, radical anions of, multifrequency CW-EPR study, Zoleo *et al.*, **159**, 226
- Bitter magnet
NMR phase noise, inductive shielding, Sigmund *et al.*, **159**, 190
- Brain
structure, *in vivo* magnetic resonance microscopy in unanesthetized flies, Jasanoff and Sun, **158**, 79
- Brain metabolites
automated artificial neural network quantification from *in vivo* ^1H NMR magnitude spectra, Hiltunen *et al.*, **154**, 1
quantitation errors due to saturation factor changes in MRS, optimized pulse parameters for reducing, Galbán and Spencer, **156**, 161
- Broadband HMBC
pulse sequence, optimizing delays in, Schulte-Herbrüggen *et al.*, **156**, 282
- Broadband pulse
adjustable broadband selective excitation with uniform phase, Cano, Smith, and Shaka, **155**, 131
- Broadband XLOC
pulse sequence, optimizing delays in, Schulte-Herbrüggen *et al.*, **156**, 282
- BURP
simple scheme for design of solvent-suppression pulses, Prost *et al.*, **159**, 77
- ## C
- ^{13}C
chemical shifts, prediction, using neural networks for, Meiler *et al.*, **157**, 242
 ^{13}C -labeled backbone carbons of membrane fusion peptides, filtered MAS observation with REDOR subtraction, Yang *et al.*, **159**, 101
 ^{13}C - ^{14}N dipolar coupling constants, ^{13}C FIREMAT experiments, Strohmeier, Alderman, and Grant, **155**, 263
 ^{13}C , ^{15}N -labeled peptides and proteins, uniformly labeled, *soft*-triple-resonance solid-state NMR for assignments of, Astrof and Griffin, **158**, 157
 ^{13}C , ^{15}N -labeled proteins
backbone assignment, intraresidue HNCA and COHNCA experiments for, Brutscher, **156**, 155
perdeuterated, correlation of backbone amide and side-chain ^{13}C resonances in, Löhr and Rüterjans, **156**, 10
 ^{13}C NMR spectral assignments for nematic liquid crystals by 2D chemical-shift γ -encoding NMR, Nishiyama, Kubo, and Terao, **158**, 60
CP/MAS NMR, of solids at 21.15 T, 4-mm probe for, Jakobsen *et al.*, **156**, 152
FIREMAT experiments, molecular and structural information from ^{13}C - ^{14}N systems with, Strohmeier, Alderman, and Grant, **155**, 263
 ^1H -detected ^{13}C echo planar imaging, Hudson *et al.*, **155**, 64
high-resolution 2D ^1H - ^{13}C correlation of cholesterol in model membrane, Soubias *et al.*, **158**, 143
one-bond ^{15}N - $^1\text{H}^{\text{N}}$ and ^{15}N - $^{13}\text{C}'$ and two-bond $^{13}\text{C}'$ - $^1\text{H}^{\text{N}}$ residual dipolar couplings in proteins, sensitivity-enhanced E.COSY-type HSQC for, Ding and Gronenborn, **158**, 173
relaxation study, hexamethylenetetramine: dipole-dipole coupling constant for directly bonded CH pair, Kowalewski, Effemey, and Jokisaari, **157**, 171
selective detection of ^1H NMR spectra of molecules with rare spins in liquid crystal samples, Emsley *et al.*, **154**, 303
side-chain resonances, correlation with backbone amide resonances in perdeuterated proteins, Löhr and Rüterjans, **156**, 10
spin pairs, coupled to ^{14}N , double-quantum solid-state NMR, Hughes *et al.*, **159**, 25
- ^{14}C
 ^{13}C spin pairs coupled to, double-quantum solid-state NMR, Hughes *et al.*, **159**, 25
- Canonical orientations
 g -tensor, nuclear frequencies at, determination of hyperfine tensor components from, Shubin and Dikanov, **155**, 100
- Carbohydrates
3D HSQC-HSQMBC for increasing resolution of long-range proton-carbon correlation experiments, Uhrín, **159**, 145
 ^{13}C natural abundance, partially oriented, measurement of small one-bond proton-carbon residual dipolar coupling constants, Pham *et al.*, **157**, 200
- Carboxylic acids
 ^{13}C NMR, magnesium chloride as relaxation agent for, Tian *et al.*, **159**, 137
- Carr–Purcell–Meiboom–Gill
detection, sensitivity enhancement in multiple-quantum NMR experiments with, Lim *et al.*, **157**, 160
non-CPMG fast spin echo with full signal, Le Roux, **155**, 278
sequence, categories of coherence pathways for, Song, **157**, 82

- Catheter placement
during interventional MRI, use of Overhauser enhancement to assist with, Vahala *et al.*, **157**, 298
- Cells
suspensions, aeration and mixing during NMR experiments, device for, Bourne and Kuchel, **159**, 158
uptake of MRI contrast agents, high-resolution MAS NMR, Calabi *et al.*, **156**, 222
- Central transition NMR spectroscopy
anisotropic relaxation and motion of half-integer quadrupolar nuclei, Kristensen and Farnan, **158**, 99
- Cesium
adsorption in hydrated iron oxide particle suspensions, NMR study, Gossuin *et al.*, **157**, 132
- Chemical exchange
effect on high-resolution DOSY NMR with spins in different chemical surroundings, Cabrita *et al.*, **157**, 124
effects on metabolite quantitation: optimized pulse parameters for error reduction due to saturation factor changes in MRS, Galbán and Spencer, **156**, 161
 $R_{1\rho}$ relaxation outside of fast-exchange limit, Trott and Palmer, **154**, 157
- Chemical-shift anisotropy
amplified interactions, 2D MAS solid-state NMR method to recover, Wei *et al.*, **158**, 23
relative CSA–dipolar orientation from REDOR sidebands, O'Connor and Schaefer, **154**, 46
and RF offset, effects in cross-polarization solid-state NMR, Shekar, Lee, and Ramamoorthy, **157**, 223
shielding tensor orientation from two or one rotations of single crystal, Shekar, Ramamoorthy, and Wittebort, **155**, 257
undistorted powder patterns, 2D separation in MAS NMR, Liu, Mao, and Schmidt-Rohr, **155**, 15
- Chemical-shift anisotropy recoupling
 ^1H , selective dephasing of OH and NH proton magnetization based on, Schmidt-Rohr and Mao, **157**, 210
- Chemical shifts
amide proton, in perdeuterated protein, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181
amino acid-specific, assignment strategy for proteins with known structure using, Hus, Prompers, and Brüschweiler, **157**, 119
 ^{13}C , prediction, using neural networks for, Meiler *et al.*, **157**, 242
 ^{14}N , inorganic nitrates, Marburger, Fung, and Khitritin, **154**, 205
sum and difference, 2D correlation spectroscopy, combining, enhanced spectral resolution of immobilized peptides and proteins by, Luca and Baldus, **159**, 243
- Chemical-shift tensor
 ^{13}C , FIREMAT experiments, Strohmeier, Alderman, and Grant, **155**, 263
- Chlorophyll *a*/H₂O
solid self-assembled, 3D structural model, van Rossum *et al.*, **155**, 1
- para*-Chlorotoluene
in liquid crystal, simultaneous selective detection of multiple-quantum spectra, Syvitski *et al.*, **155**, 251
- Cholesterol
in model membrane, high-resolution 2D ^1H – ^{13}C correlation, Soubias *et al.*, **158**, 143
- CH pair
directly bonded, dipole–dipole coupling constant for, ^{13}C relaxation study, Kowalewski, Effemey, and Jokisaari, **157**, 171
- ^{13}C NMR
relaxation agent, for amino acids and carboxylic acids, magnesium chloride as, Tian *et al.*, **159**, 137
spectral assignments for nematic liquid crystals by 2D chemical-shift γ -encoding NMR, Nishiyama, Kubo, and Terao, **158**, 60
subband ARMA modeling approach to high-resolution NMR, Tomczak and Djermoune, **158**, 86
- Cogwheel phase cycling
method for constructing phase cycles, Levitt, Madhu, and Hughes, **155**, 300
- Coherence pathways
for CPMG sequence, categories of, Song, **157**, 82
- Coherence transfer
in high-molecular-weight systems, evaluation and optimization, Permi and Annala, **155**, 123
pathways, cogwheel phase cycling for experiments with, Levitt, Madhu, and Hughes, **155**, 300
time-optimal coherence-order-selective, of in-phase coherence in heteronuclear *IS* spin systems, Reiss, Khaneja, and Glaser, **154**, 192
- Coherent excitation
with phase-incremented pulses, Zhang and Gorenstein, **154**, 73
- COHNCA
intraresidue, experiments for protein backbone assignment, Brutscher, **156**, 155
- Complete relaxation and conformational exchange matrix
analysis of intermolecular saturation transfer effects in reversibly forming ligand–receptor complexes, Jayalakshmi and Krishna, **155**, 106
- Conditional phase-shift gate
implementation for quantum information processing by NMR, using transition-selective pulses, Das, Mahesh, and Kumar, **159**, 46
- Conjoined rigid body/torsion angle simulated annealing
determination of relative orientation of two domains from dipolar couplings, Clore and Bewley, **154**, 329
- Constant-time multidimensional electrophoretic NMR
improved EF-time-incrementing approach, Li and He, **156**, 181
- Continuous-wave EPR
1.1-GHz, with frequency modulation method, Hirata *et al.*, **155**, 140
measurement, active resonator system operating at 700 MHz for, Sato *et al.*, **159**, 161
multifrequency, radical anions of fullerene bisadducts, Zoleo *et al.*, **159**, 226
- Contrast agents
cell uptake, high-resolution MAS NMR, Calabi *et al.*, **156**, 222
MRI, relaxation enhancement by, role of diffusion and exchange, Gossuin *et al.*, **158**, 36
- Contrast parameter
for visualization of crosslink density in rubber based on dipolar-correlation effect, Grinberg, Heidenreich, and Kuhn, **159**, 88
- Contrast solid-state imaging
spin interaction filter in, Casieri and De Luca, **159**, 63
- CORCEMA (*see* Complete relaxation and conformational exchange matrix)
- Correlation spectroscopy
2D ^{13}C – ^{13}C MAS NMR, with mixing by true ^1H spin diffusion, de Boer *et al.*, **157**, 286
chemical shift sum and difference, combining, enhanced spectral resolution of immobilized peptides and proteins by, Luca and Baldus, **159**, 243
 T_2 – T_2 spectra obtained using fast 2D Laplace inversion, Song *et al.*, **154**, 261
- Cosine modulation
 ^{15}N -COSMO-HSQC rapid determination of $^3J_{\text{HNH}_2}$ scalar couplings in ^{15}N -labeled proteins, Petit, Vincent, and Zwahlen, **156**, 313
- COSY-type mixing
and homonuclear Hartmann–Hahn mixing sequences, in presence of scalar and residual dipolar couplings, Kramer and Glaser, **155**, 83
- Coupling constants
dipole–dipole, for directly bonded CH pair, ^{13}C relaxation study, Kowalewski, Effemey, and Jokisaari, **157**, 171
one-bond carbon–proton, in ^{13}C natural abundance carbohydrates, *J*-modulated 2D HSQC, Pham *et al.*, **157**, 200

- quadrupole, ^{14}N , of inorganic nitrates, Marburger, Fung, and Khitrin, **154**, 205
- Cramer–Rao lower bound
achievable accuracy of parameter estimation for multidimensional NMR, Ober *et al.*, **157**, 1
- Cristobalite
 ^{17}O , central transition NMR spectroscopy: anisotropic relaxation and motion of half-integer quadrupolar nuclei, Kristensen and Farnan, **158**, 99
- Cross correlation
effects involving Curie spin relaxation in methyl groups, Madhu, Mandal, and Müller, **155**, 29
- Crosslink density
in rubber, visualization based on dipolar-correlation effect, contrast parameter for, Grinberg, Heidenreich, and Kuhn, **159**, 88
- Crosslinking
in natural rubber, effects on multipolar spin states, Wiesmath *et al.*, **154**, 60
- Cross polarization
from dipolar reservoir under mismatched Hartmann–Hahn condition, Venkatraman, Sinha, and Ramanathan, **157**, 137
- Cross-polarization/MAS NMR
 ^{13}C , at 21.15 T, 4-mm probe for, Jakobsen *et al.*, **156**, 152
and hyperpolarized ^{129}Xe , for spin polarization transfer to surface nuclei, Smith *et al.*, **159**, 111
- Cross-polarization solid-state NMR
chemical-shift anisotropy and offset effects in, Shekar, Lee, and Ramamoorthy, **157**, 223
- Crystal field tensors
low-symmetry, analysis: Fe^{3+} in natrolite, Vinokurov *et al.*, **155**, 57
- Curie spin relaxation
in methyl groups, cross-correlation effects involving, Madhu, Mandal, and Müller, **155**, 29
- Current magnetization
ferro-refraction, open magnet utilizing, Pulyer and Hrovat, **154**, 298
- Curve fitting
2D, comparison with trilinear SLICING for curve resolution of low-field NMR relaxation data, Pedersen, Bro, and Engelsen, **157**, 141
- Cyclic J cross polarization (CYCLCROP)
 ^1H -detected ^{13}C echo planar imaging using, Hudson *et al.*, **155**, 64
- Cycling sidebands
and two-dimensional artifact from asynchronous decoupling, van Ingen, Vuister, and Tessari, **156**, 258
- Cylinders
multi-compartment systems, modeling of self-diffusion and relaxation time NMR in, van der Weerd *et al.*, **156**, 213
- D**
- DANTE
MR tagging using tandem of two adiabatic DANTE inversion sequences, Tsekos, Garwood, and Ugurbil, **156**, 187
- Data processing
automated correction of unwanted phase jumps in reference signals corrupting MRSI spectra after eddy current correction, Simonetti *et al.*, **159**, 151
- Deconvolution
automated correction of unwanted phase jumps in reference signals corrupting MRSI spectra after eddy current correction, Simonetti *et al.*, **159**, 151
compartmental water diffusion coefficients in yeast cell suspensions, combined T_1 and D , Silva *et al.*, **156**, 52
- Decoupling
asynchronous, 2D artifact from, van Ingen, Vuister, and Tessari, **156**, 258
low-power, local field spectroscopy with, measurement of dipolar couplings in partially oriented molecules, Marjanska *et al.*, **158**, 52
- Dephasing
selective, of OH and NH proton magnetization, based on ^1H chemical-shift anisotropy recoupling, Schmidt-Rohr and Mao, **157**, 210
- Desorption
nitric oxide in H-ZSM-5, Na-ZSM-5, Na-A, EPR study, Rudolf, Böhlmann, and Pöpl, **155**, 45
- Deuterated benzene
in amorphous matrices, large-angle jumps of, 2D exchange NMR analysis, Medick, Vogel, and Rössler, **159**, 126
- Deuterium solid-state NMR
transmembrane peptides in bicelles, and hydrophobic mismatch and peptide sequence, Whiles *et al.*, **158**, 149
- 1,3-Dichloro-4-fluorobenzene
in nematic solvents, dipolar couplings in, local field NMR with low-power decoupling, Marjanska *et al.*, **158**, 52
- 1,3-Dichloro-4-fluoro-5-nitrobenzene
in nematic solvents, dipolar couplings in, local field NMR with low-power decoupling, Marjanska *et al.*, **158**, 52
- Difference spectroscopy
dual-coil probe for, Macnaughtan *et al.*, **156**, 97
- Diffusion
anisotropic, of water in macroscopically oriented lipid bilayers, pulsed magnetic field gradient NMR, Wästerby, Orädd, and Lindblom, **157**, 156
and exchange, role in relaxation enhancement by MRI contrast agents, Gossuin *et al.*, **158**, 36
measurements, NMR, analysis of protein–ligand interactions with, Derrick, McCord, and Larive, **155**, 217
modeling in multi-compartment systems with cylindrical geometry, van der Weerd *et al.*, **156**, 213
and relaxation, 2D distribution functions, quantitative measurement in grossly inhomogeneous fields, Hürlimann and Venkataraman, **157**, 31
restricted (*see* Restricted diffusion)
xenon gas diffusion NMR of porous media, narrow pulse approximation and long length scale determination in, Mair *et al.*, **156**, 202
- Diffusion coefficients
compartmental water, deconvolution in yeast cell suspensions, Silva *et al.*, **156**, 52
in mixtures, resolving: 2D- J -DOSY experiment, Lucas, Otto, and Larive, **156**, 138
- Diffusion MR
effects of restricted diffusion on MR signal formation, Sukstanskii and Yablonskiy, **157**, 92
- Diffusion-ordered spectroscopy
high-resolution, with spins in different chemical surroundings, effect of particle exchange, Cabrita *et al.*, **157**, 124
2D- J -DOSY experiment: resolving diffusion coefficients in mixtures, Lucas, Otto, and Larive, **156**, 138
- Diffusion processes
analysis using EPR imaging with modulated field gradient, Herrling, Fuchs, and Groth, **154**, 6
- Diffusion tensor imaging
intervoxel anisotropy index mapping with, application of Euclidian distance-weighted smoothing for quantitative MRI, Bonny and Renou, **159**, 183
- Diffusion tensor magnetic resonance imaging
discrete data, continuous tensor field approximation, Pajevic, Aldroubi, and Basser, **154**, 85
- 1,2-Difluorobenzene
in nematic solvents, dipolar couplings in, local field NMR with low-power decoupling, Marjanska *et al.*, **158**, 52
- Dilaurylphosphatidylcholine
and DPPC, bicelles, study methods for transmembrane peptides in, Whiles *et al.*, **158**, 149

- Dipalmitoylphosphatidylcholine
and DLPC, bicelles, study methods for transmembrane peptides in,
Whiles *et al.*, **158**, 149
- Dipolar correlation
HCCN experiment, torsion angle Ψ determination in α -helical peptides
and proteins, Ladizhansky, Veshtort, and Griffin, **154**, 317
high-resolution 2D ^1H - ^{13}C , of cholesterol in model membrane, Soubias
et al., **158**, 143
- Dipolar-correlation effect
visualization of crosslink density in rubber based on, contrast parameter
for, Grinberg, Heidenreich, and Kuhn, **159**, 88
- Dipolar-correlation spectroscopy
MAS NMR 2D high-field, 3D structural model of aggregated
chlorophyll *a*/ H_2O from, van Rossum *et al.*, **155**, 1
- Dipolar couplings (*see also* Residual dipolar couplings)
2D MAS solid-state NMR method to recover amplified heteronuclear
dipolar interactions, Wei *et al.*, **158**, 23
2D NMR correlations of liquid crystals using switched angle spinning,
Havlin, Park, and Pines, **157**, 163
covalently linked protein domains from, relative orientation by
conjoined rigid body/torsion angle simulated annealing, Clore
and Bewley, **154**, 329
electron dipole–dipole ESEEM in field-step ELDOR of nitroxide
biradicals, Kulik *et al.*, **157**, 61
high-resolution 2D ^1H - ^{13}C correlation of cholesterol in model
membrane, Soubias *et al.*, **158**, 143
measurement in partially oriented molecules by local field NMR with
low-power decoupling, Marjanska *et al.*, **158**, 52
to ^{14}N , effect on ^{13}C FIREMAT experiments, Strohmeier, Alderman,
and Grant, **155**, 263
scalar and residual, homonuclear Hartmann-Hahn and COSY-type
mixing sequences in presence of, Kramer and Glaser, **155**, 83
separation from exchange coupling, and PELDOR at S- and X-band
frequencies, Weber *et al.*, **157**, 277
- Dipolar fields
scaling laws in NMR scattering via, Brown, Sen, and Cory, **154**,
154
- Dipolar interactions
heteronuclear, amplified, 2D MAS solid-state NMR method to recover,
Wei *et al.*, **158**, 23
long-range, in heterogeneous systems: origins of dips in multiple spin
echoes, Alessandri, Capuani, and Maraviglia, **156**, 72
- Dipolar order
NMR of multipolar spin states excited in strongly inhomogeneous
magnetic fields, Wiesmath *et al.*, **154**, 60
- Dipolar recoupling
 ^{13}C - ^{13}C , measurements, sensitivity enhancement by pulsed spin lock-
ing, Petkova and Tycko, **155**, 293
double-quantum, at high magic-angle spinning rates, Kristiansen,
Mitchell, and Evans, **157**, 253
homonuclear zero-quantum, in fast MAS NMR, Brinkmann, Schmedt
auf der Gönne, and Levitt, **156**, 79
narrowband, for homonuclear distance measurements in rotating solids,
improved, Goobes and Vega, **154**, 236
proton, in resin-bound peptides under high-resolution MAS, Raya *et al.*,
157, 43
- Dipolar reservoir
cross-polarization from, under mismatched Hartmann–Hahn condition,
Venkatraman, Sinha, and Ramanathan, **157**, 137
- Dipolar tensor
relative orientation with respect to CSA orientation, REDOR sidebands,
O'Connor and Schaefer, **154**, 46
- Dipolar time evolution data
EPR, direct conversion to distance distributions, Jeschke *et al.*, **155**, 72
- Dipole–dipole coupling constant
for directly bonded CH pair, ^{13}C relaxation study, Kowalewski,
Effemey, and Jokisaari, **157**, 171
- Discrete wavelet transforms
improved solvent suppression in NMR spectra using: WAVEMAT,
Günther, Ludwig, and Rüterjans, **156**, 19
- Distance distributions
direct conversion of EPR dipolar time evolution data to, Jeschke *et al.*,
155, 72
- Distance measurements
homonuclear, in rotating solids, improved narrowband dipolar coupling
for, Goobes and Vega, **154**, 236
- Distribution functions
two-dimensional, of diffusion and relaxation, quantitative measurement
in grossly inhomogeneous fields, Hürlimann and Venkataraman,
157, 31
- Double frequency sweeps
influencing satellite transitions of half-integer quadrupolar nuclei
for enhancement of MAS spectra, Iuga and Kentgens, **158**,
65
- Double-quantum coherences
NMR of multipolar spin states excited in strongly inhomogeneous
magnetic fields, Wiesmath *et al.*, **154**, 60
- Double-quantum NMR
solid-state, ^{13}C spin pairs coupled to ^{14}N , Hughes *et al.*, **159**, 25
- Double-quantum recoupling
dipolar, at high magic-angle spinning rates, Kristiansen, Mitchell, and
Evans, **157**, 253
- Double resonance probe
4-mm, for ^{13}C CP/MAS NMR of solids at 21.15 T, Jakobsen *et al.*, **156**,
152
- Double spin–echo
method for variable temperature gradient shimming, Evans, Morris, and
Davis, **154**, 325
- Droplets
emulsion, sizing under flow using flow-compensating NMR-PFG
techniques, Johns and Gladden, **154**, 142
- Dynamics
multiple-quantum, in linear chains and rings of nuclear spins in solids at
low temperatures, Fel'dman and Maximov, **157**, 106

E

- Earth's-field NMR
condition for adiabatic passage in, Melton and Pollak, **158**, 15
- Echo planar imaging
 ^1H -detected ^{13}C , Hudson *et al.*, **155**, 64
- E.COSY-type HSQC
sensitivity-enhanced, for one-bond ^{15}N - $^1\text{H}^{\text{N}}$ and ^{15}N - $^{13}\text{C}'$ and two-bond
 $^{13}\text{C}'$ - $^1\text{H}^{\text{N}}$ residual dipolar couplings in proteins, Ding and Gronen-
born, **158**, 173
- Eddy current correction
unwanted phase jumps in reference signals corrupting MRSI spectra
after, automated correction, Simonetti *et al.*, **159**, 151
- ELDOR
field-step, of nitroxide biradicals, electron dipole–dipole ESEEM in,
Kulik *et al.*, **157**, 61
- Electron dipole–dipole interactions
in nitroxide biradicals, field-step ELDOR, Kulik *et al.*, **157**, 61
- Electron–electron interaction
in biradical, 130 GHz ESEEM induced by, Kulik, Paschenko, and
Dzuba, **159**, 237
- Electron spin–lattice relaxation
in radicals with two methyl groups in γ -irradiated polycrystalline solids,
Harbridge, Eaton, and Eaton, **159**, 195
- Electron spin relaxation times
level-averaged, outside Zeeman limit, closed-form expressions, Sharp,
154, 269
long, out-of-phase rapid-passage EPR for samples with, enhanced signal
intensities, Harbridge *et al.*, **156**, 41

- Electrophoretic NMR
 constant-time multidimensional, Li and He, **156**, 181
- Emulsion droplets
 sizing under flow using flow-compensating NMR-PFG techniques, Johns and Gladden, **154**, 142
- Enantiomeric excess
 measurements in weakly oriented chiral liquid crystal solvents using 2D ^1H selective refocusing, Farjon *et al.*, **158**, 169
- End cap
 double end-cap birdcage coil for small animal whole-body imaging, Gulsen, Muftuler, and Nalcioglu, **156**, 309
- Endohedral fullerenes
 N@C_{60} and N@C_{70} , in liquid crystal, EPR study, Jakes *et al.*, **156**, 303
- Endovascular imaging
 opposed-solenoid MRI probes for, reciprocity and sensitivity, Crottet *et al.*, **159**, 219
- Entropy minimization
 efficient algorithm for automatic phase correction of NMR spectra based on, Chen *et al.*, **158**, 164
- EPR
 adsorption/desorption behavior of NO on H-ZSM-5, Na-ZSM-5, Na-A, Rudolf, Böhlmann, and Pöpl, **155**, 45
 combined k-space q-space pulsed EPR imaging, Feintuch *et al.*, **157**, 69
 continuous-wave (*see* Continuous-wave EPR)
 dipolar time evolution data, direct conversion to distance distributions, Jeschke *et al.*, **155**, 72
 endohedral fullerenes N@C_{60} and N@C_{70} in liquid crystal, Jakes *et al.*, **156**, 303
 Fe^{3+} in natrolite: low-symmetry spin Hamiltonian and crystal field analysis, Vinokurov *et al.*, **155**, 57
 high-field, 130 GHz ESEEM induced by electron–electron interaction in biradical, Kulik, Paschenko, and Dzuba, **159**, 237
 multifrequency, CW-EPR study of radical anions of fullerene bisadducts, Zoleo *et al.*, **159**, 226
 out-of-phase rapid-passage, for samples with long electron spin relaxation times, enhanced signal intensities, Harbridge *et al.*, **156**, 41
 PELDOR at S- and X-band frequencies and separation of exchange coupling from dipolar coupling, Weber *et al.*, **157**, 277
 pulsed and continuous-wave RF, for *in vivo* free radical detection/imaging techniques, comparison, Yamada *et al.*, **154**, 287
 signal intensity, frequency dependence
 248 MHz to 1.4 GHz, Rinard *et al.*, **154**, 80
 250 MHz to 9.1 GHz, Rinard *et al.*, **156**, 113
 spectra decomposition, quantification of lateral lipid domain properties in erythrocyte ghost membranes using, Arsov, Schara, and Štrancar, **157**, 52
 spectrum of magnetic center in kaolin, temperature-induced changes, Vasyukov *et al.*, **154**, 15
 structural determination of spin label immobilization and orientation, Sale *et al.*, **156**, 104
- EPR imaging
 with modulated field gradient, kinetic measurements using, Herrling, Fuchs, and Groth, **154**, 6
- EPR signal intensity
 frequency dependence
 248 MHz to 1.4 GHz, Rinard *et al.*, **154**, 80
 250 MHz to 9.1 GHz, Rinard *et al.*, **156**, 113
- Equations
 universal, for linear and partial adiabatic pulses, Tesiram and Bendall, **156**, 26
- Erythrocyte ghosts
 lateral lipid domain properties, quantification using EPR-spectra decomposition, Arsov, Schara, and Štrancar, **157**, 52
- ESEEM
 130 GHz, induced by electron–electron interaction in biradical, Kulik, Paschenko, and Dzuba, **159**, 237
 electron dipole–dipole, in field-step ELDOR of nitroxide biradicals, Kulik *et al.*, **157**, 61
 one- and two-dimensional experiments, numerical simulations, Madi, Van Doorslaer, and Schweiger, **154**, 181
 spectra of $S > 1/2$ centers in orientationally disordered systems, Benetis, Dave, and Goldfarb, **158**, 126
- ESR (*see* EPR)
- Ethylbenzene
 920-MHz high-resolution NMR, Hashi *et al.*, **156**, 318
- Euclidian distance-weighted smoothing
 for quantitative MRI, Bonny and Renou, **159**, 183
- Exchange
 and diffusion, role in relaxation enhancement by MRI contrast agents, Gossuin *et al.*, **158**, 36
 effect of particle exchange on high-resolution DOSY NMR with spins in different chemical surroundings, Cabrera *et al.*, **157**, 124
- Exchange coupling
 separation from dipolar coupling, and PELDOR at S- and X-band frequencies, Weber *et al.*, **157**, 277
- Excitation delay
 optimizing delays in MBOB, broadband HMBC, and broadband XLOC pulse sequences, Schulte-Herbrüggen *et al.*, **156**, 282
- Excitation sculpting
 simple scheme for design of solvent-suppression pulses, Prost *et al.*, **159**, 77
- Extramyocellular lipids
 and intramyocellular lipids, signal separation in proton MR spectra by magnetic field distribution, Steidle *et al.*, **154**, 228
- F**
- (FA) $_2$ PF $_6$
 restricted diffusion in, mapping: combined k-space q-space pulsed ESR imaging, Feintuch *et al.*, **157**, 69
- Falling liquid films
 NMR velocimetry, Heine *et al.*, **154**, 311
- FAM–RIACT–FAM
 signal enhancement in triple-quantum MAS NMR of spins-3/2 in solids, Madhu and Levitt, **155**, 150
- Fast magic-angle spinning
 NMR, homonuclear zero-quantum recoupling in, Brinkmann, Schmedt auf der Günne, and Levitt, **156**, 79
- Fast spin echo
 non-CPMG, with full signal, Le Roux, **155**, 278
- Fat
 and changes in porcine muscle water characteristics during growth, *in vitro* low-field ^1H NMR, Bertram *et al.*, **157**, 267
- Feedback
 RF power: active resonator system for CW-ESR operating at 700 MHz, Sato *et al.*, **159**, 161
- Fercayl
 relaxation enhancement by, role of diffusion and exchange, Gossuin *et al.*, **158**, 36
- Ferrihydrite
 dextran-coated particle suspensions, NMR study of cesium adsorption, Gossuin *et al.*, **157**, 132
- Ferritin
 particle suspensions, NMR study of cesium adsorption, Gossuin *et al.*, **157**, 132
 relaxation enhancement by, role of diffusion and exchange, Gossuin *et al.*, **158**, 36
- Ferro-refraction current magnetization
 open magnet utilizing, **154**, 298
- Fiber
 microstructure/architecture, continuous tensor field approximation of DT-MRI data, Pajevic, Aldroubi, and Bassar, **154**, 85

- Fiber tractography
continuous tensor field approximation of DT-MRI data, Pajevic, Aldroubi, and Basser, **154**, 85
- Fibronectin
perdeuterated, amide proton relaxation and chemical shift in, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181
- Field gradients
internal, enhanced transverse relaxation in porous media due to, Dunn, **156**, 171
- Filtered MAS
observation of backbone carbons of membrane-bound fusion peptides, with REDOR subtraction, Yang *et al.*, **159**, 101
- FIREMAT
¹³C, molecular and structural information from ¹³C–¹⁴N systems, Strohmeier, Alderman, and Grant, **155**, 263
- Flow
gradient echo imaging of flowing hyperpolarized nuclei, Johansson *et al.*, **159**, 69
through porous media, PFG NMR measurements: effects of spatial correlations of magnetic field and velocity field, Lebon and Leblond, **159**, 13
sizing of emulsion droplets under, using flow-compensating NMR-PFG techniques, Johns and Gladden, **154**, 142
two-phase, spectrally resolved velocity exchange spectroscopy, Khra-pitchev *et al.*, **159**, 36
- Flow compensation
NMR-PFG techniques with, sizing of emulsion droplets under flow using, Johns and Gladden, **154**, 142
- Free induction decay
frequency dependence of EPR signal intensity, 248 MHz to 1.4 GHz, Rinard *et al.*, **154**, 80
signal behavior in presence of magnetic field gradients, theoretical framework for analysis, Sukstanskii and Yablonskiy, **157**, 92
- Free radicals
in vivo detection and imaging, pulsed and continuous-wave RF EPR techniques, comparison, Yamada *et al.*, **154**, 287
- Frequency dependence
EPR signal intensity
248 MHz to 1.4 GHz, Rinard *et al.*, **154**, 80
250 MHz to 9.1 GHz, Rinard *et al.*, **156**, 113
- Frequency modulation
1.1-GHz continuous-wave EPR spectroscopy method with, Hirata *et al.*, **155**, 140
gradient-induced, removal by modulus signal selection: localized ¹H MRS without water suppression, Serrai *et al.*, **154**, 53
- Frequency-selective algorithm
time domain quantification, Romano *et al.*, **155**, 226
- Frequency-switched Lee–Goldburg
effect of Hartmann–Hahn mismatching on polarization inversion spin exchange at magic angle, Fu *et al.*, **159**, 167
pulse sequence, NMR spin locking of proton magnetization under, Fu, Tian, and Cross, **154**, 130
- Fullerene bisadducts
radical anions of, multifrequency CW-EPR study, Zoleo *et al.*, **159**, 226
- Fullerenes
endohedral, N@C₆₀ and N@C₇₀, in liquid crystal, EPR study, Jakes *et al.*, **156**, 303
- Fusion peptides
membrane-bound, filtered MAS observation of backbone carbons, with REDOR subtraction, Yang *et al.*, **159**, 101
- G**
- Gamma-encoding NMR
2D chemical-shift, ¹³C NMR spectral assignments for nematic liquid crystals by, Nishiyama, Kubo, and Terao, **158**, 60
- Gamma-irradiation
polycrystalline solids, radicals with two methyl groups generated by, electron spin-lattice relaxation, Harbridge, Eaton, and Eaton, **159**, 195
- GAMMA simulations
of stray-field responses: slice thickness and pulse calibration, Kinchesch *et al.*, **154**, 252
- Gas diffusion NMR
xenon, of model porous media, narrow pulse approximation and long length scale determination in, Mair *et al.*, **156**, 202
- Gradient
induced frequency modulations, removal by modulus signal selection: localized ¹H MRS without water suppression, Serrai *et al.*, **154**, 53
- Gradient coils
short planar, for MR microscopy using concentric return paths, Dodd and Ho, **156**, 1
- Gradient echo imaging
flowing hyperpolarized nuclei, Johansson *et al.*, **159**, 69
- Gradient shimming
variable temperature, method for, Evans, Morris, and Davis, **154**, 325
- Graphical analysis
relative orientation of molecular alignment tensors for protein dissolved in different anisotropic media, Nomura and Kainosho, **154**, 146
- g-tensor
nuclear frequencies at canonical orientations of, determination of hyperfine tensor components from, Shubin and Dikanov, **155**, 100
- H**
- ¹H
¹H-detected ¹³C echo planar imaging, Hudson *et al.*, **155**, 64
high-resolution 2D ¹H–¹³C correlation of cholesterol in model membrane, Soubias *et al.*, **158**, 143
- ²H
2D exchange NMR analysis of large-angle jumps of small molecules in amorphous matrices, Medick, Vogel, and Rössler, **159**, 126
amide proton relaxation and chemical shift in perdeuterated fibronectin, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181
- Hartmann–Hahn
homonuclear, and COSY-type mixing sequences, efficiency in presence of scalar and residual dipolar couplings, Kramer and Glaser, **155**, 83
mismatch
cross-polarization from dipolar reservoir under, Venkatraman, Sinha, and Ramanathan, **157**, 137
effect on polarization inversion spin exchange at magic angle, Fu *et al.*, **159**, 167
- HCC–TOCSY
tailored experiment for resonance assignment in proximity of paramagnetic center, Piccioli and Poggi, **155**, 236
- HCCN dipolar correlation experiment
torsion angle Ψ determination in α -helical peptides and proteins, Ladizhansky, Veshtort, and Griffin, **154**, 317
- Head pitch
 B_0 homogeneity variation with, quantification by registered 3D field mapping, Tyszka and Mamelak, **159**, 213
- Heart
canine, MR tagging using tandem of two adiabatic DANTE inversion sequences, Tsekos, Garwood, and Ugurbil, **156**, 187
metabolite quantitation errors due to saturation factor changes in MRS, optimized pulse parameters for reducing, Galbán and Spencer, **156**, 161
- Helium
laser-polarized, in molecules, materials, and organisms, NMR, review, Goodson, **155**, 157

- Heterogeneous systems
multiple spin echoes: origin of observed dips, Alessandri, Capuani, and Maraviglia, **156**, 72
- Heteronuclear dipolar interactions
amplified, 2D MAS solid-state NMR method to recover, Wei *et al.*, **158**, 23
- Heteronuclear *IS* spin systems
time-optimal coherence-order-selective transfer of in-phase coherence in, Reiss, Khaneja, and Glaser, **154**, 192
- Heteronuclear Nuclear Overhauser effect
two-dimensional artifact from asynchronous decoupling, van Ingen, Vuister, and Tessari, **156**, 258
- Heteronuclear shift correlation
long-range, detection with LR-*J*-HSMQC, Torres, Bubb, and Kuche, **156**, 249
- Hexamethyl benzene
in amorphous matrices, large-angle jumps of, 2D exchange NMR analysis, Medick, Vogel, and Rössler, **159**, 126
- Hexamethylenetetramine
¹³C relaxation: dipole-dipole coupling constant for directly bonded CH pair, Kowalewski, Effemey, and Jokisaari, **157**, 171
- High field
MAS NMR 2D dipolar correlation spectroscopy, model for aggregated chlorophyll *a*/H₂O, van Rossum *et al.*, **155**, 1
- High-field EPR
130 GHz ESEEM induced by electron–electron interaction in biradical, Kulik, Paschenko, and Dzuba, **159**, 237
- High-frequency magic-angle spinning
double-quantum dipolar recoupling at, Kristiansen, Mitchell, and Evans, **157**, 253
- High-molecular-weight systems
coherence transfer in, evaluation and optimization, Permi and Annila, **155**, 123
- High-resolution DOSY NMR
with spins in different chemical surroundings, effect of particle exchange, Cabrita *et al.*, **157**, 124
- High-resolution magic-angle spinning NMR
and cell uptake of MRI contrast agents, Calabi *et al.*, **156**, 222
line broadening of organic molecules grafted onto polystyrene resin in, and resin dynamics, Lippens, Chessari, and Wieruszkeski, **156**, 242
optimizing delays in MBOB, broadband HMBC, and broadband XLOC pulse sequences, Schulte-Herbrüggen *et al.*, **156**, 282
proton dipolar recoupling in resin-bound peptides under, Raya *et al.*, **157**, 43
- High-resolution NMR
920-MHz, Hashi *et al.*, **156**, 318
solid-state, sample restriction using RF selective pulses in, Charmont, Sakellariou, and Emsley, **54**, 136
subband ARMA modeling approach, Tomczak and Djermoune, **158**, 86
two-dimensional, in matched *B*₀ and *B*₁ field gradients, Heise *et al.*, **156**, 146
- HMBC
MBOB, broadband HMBC, and broadband XLOC pulse sequences, optimizing delays in, Schulte-Herbrüggen *et al.*, **156**, 282
- HNCA
intraresidue, experiments for protein backbone assignment, Brutscher, **156**, 155
- HN(CO)CANH
evaluation and optimization of coherence transfer in high-molecular-weight systems, Permi and Annila, **155**, 123
- ²H NMR
¹H-detected, membrane peptide orientation by, Yamaguchi and Hong, **155**, 244
- Homonuclear dipolar recoupling
zero-quantum, in fast MAS NMR, Brinkmann, Schmedt auf der Günne, and Levitt, **156**, 79
- Homonuclear Hartmann–Hahn
and COSY-type mixing sequences, efficiency in presence of scalar and residual dipolar couplings, Kramer and Glaser, **155**, 83
- Homonuclear zero-quantum recoupling
dipolar, in fast magic-angle spinning NMR, Brinkmann, Schmedt auf der Günne, and Levitt, **156**, 79
- HSQC
3D HSQC-HSQMBC, increasing resolution of long-range proton–carbon correlation experiments, Uhrín, **159**, 145
cosine-modulated, rapid determination of ³*J*_{H_NH_α} scalar couplings in ¹⁵N-labeled proteins, Petit, Vincent, and Zwanen, **156**, 313
J-modulated 2D, small one-bond proton-carbon residual dipolar coupling constants in ¹³C natural abundance carbohydrates, Pham *et al.*, **157**, 200
selective detection of ¹H NMR spectra of molecules with rare spins in liquid crystal samples, Emsley *et al.*, **154**, 303
sensitivity-enhanced E.COSY-type, for protein one-bond ¹⁵N–¹H^N and ¹⁵N–¹³C' and two-bond ¹³C'–¹H^N residual dipolar couplings, Ding and Gronenborn, **158**, 173
- HSQMBC
3D HSQC-HSQMBC, increasing resolution of long-range proton–carbon correlation experiments, Uhrín, **159**, 145
- Hydrated iron oxide
particle suspensions, cesium adsorption in, NMR study, Gossuin *et al.*, **157**, 132
- Hydrogen bonds
in human ubiquitin, and temperature coefficients of amide protons, Cierpicki *et al.*, **157**, 178
nitroxides with, polarity contributions to hyperfine splittings of, Marsh, **157**, 114
- Hydrophobic mismatch
and study methods for transmembrane peptides in bicelles, **158**, 149
- Hyperfine interaction
and tailored HCCH–TOCSY experiment for resonance assignment in proximity of paramagnetic center, Piccioli and Poggi, **155**, 236
- Hyperfine splittings
hydrogen-bonded nitroxides, polarity contributions, Marsh, **157**, 114
- Hyperfine tensor
components, determination from nuclear frequencies at canonical orientations of *g*-tensor, Shubin and Dikanov, **155**, 100
- Hyperpolarized nuclei
flowing, gradient echo imaging, Johansson *et al.*, **159**, 69
- Hyperpolarized ¹²⁹Xe (*see* ¹²⁹Xe, hyperpolarized)
- HYSORE
spectra of *S* > 1/2 centers in orientationally disordered systems, Benetis, Dave, and Goldfarb, **158**, 126
- |
- Imaging (*see also* Magnetic resonance imaging; Magnetic resonance spectroscopic imaging)
1D, water content profiles with 1D centric SPRITE acquisition, Mastikhin *et al.*, **156**, 122
contrast solid-state, spin interaction filter in, Casieri and De Luca, **159**, 63
echo planar, ¹H-detected ¹³C, Hudson *et al.*, **155**, 64
endovascular, opposed-solenoid MRI probes for, reciprocity and sensitivity, Crottet *et al.*, **159**, 219
EPR, with modulated field gradient, kinetic measurements using, Herrling, Fuchs, and Groth, **154**, 6
free radicals *in vivo*, comparison of pulsed and continuous-wave RF EPR techniques for, Yamada *et al.*, **154**, 287
gradient echo, flowing hyperpolarized nuclei, Johansson *et al.*, **159**, 69
multidimensional, using combined stray-field and pulsed gradients, Godward *et al.*, **155**, 92
NMR, contrast parameter for visualization of crosslink density in rubber based on dipolar-correlation effect, Grinberg, Heidenreich, and Kuhn, **159**, 88

- pulsed EPR, combined k-space q-space, Feintuch *et al.*, **157**, 69
 quantitative MRI, Euclidian distance-weighted smoothing for, Bonny and Renou, **159**, 183
 rotating-frame, ring down elimination in rapid imaging pulse trains—RIDE'n RIPT, Woelk *et al.*, **159**, 207
 solid-state NMR, spin interaction filter in, Casieri and De Luca, **159**, 63
 void-space, materials and organisms: principles and MR applications of laser-polarized noble gases, Goodson, **155**, 157
 whole-body small animal, double end-cap birdcage coil for, Gulsen, Muftuler, and Nalcioğlu, **156**, 309
- Immobilization
 spin label, structural determination, Sale *et al.*, **156**, 104
- Immobilized peptides and proteins
 enhanced spectral resolution by combining chemical shift sum and difference spectroscopy, Luca and Baldus, **159**, 243
- Indene
 partially oriented, multiple-quantum NMR spectra, Field, Ramadan, and Pierens, **156**, 64
- Inductive shielding
 NMR phase noise, Sigmund *et al.*, **159**, 190
- Inhomogeneous fields
 quantitative measurement of 2D distribution functions of diffusion and relaxation in, Hürlimann and Venkataraman, **157**, 31
- Inorganic nitrates
¹⁴N chemical shifts and quadrupole coupling constants, Marburger, Fung, and Khitritin, **154**, 205
- Interaction tensors
 orientation, sensitivity of DQ solid state NMR spectra of ¹³C spin pairs coupled to ¹⁴N, Hughes *et al.*, **159**, 25
- Intermediate motional regime
 application of MAS recoupling methods in, Saalwächter and Fischbach, **157**, 17
- Intermolecular cross-saturation NMR
 and STD-NMR: complete relaxation and conformational exchange matrix, Jayalakshmi and Krishna, **155**, 106
- Intermolecular distance restraints
 long-range, in ultra-high magnetic field: 2D ¹³C–¹³C MAS NMR correlation spectroscopy, de Boer *et al.*, **157**, 286
- Intermolecular saturation transfer effects
 in reversibly forming ligand–receptor complexes, CORCEMA analysis, Jayalakshmi and Krishna, **155**, 106
- Internal field gradients
 enhanced transverse relaxation in porous media due to, Dunn, **156**, 171
- Internuclear distance
 $S = 1$, $I = 1/2$ spin pairs, determination by REAPDOR NMR, Hughes *et al.*, **156**, 230
- Interventional MRI
 use of Overhauser enhancement to assist with needle and catheter placement during, Vahala *et al.*, **157**, 298
- Intervoxel anisotropy index mapping
 with DTI, application of Euclidian distance-weighted smoothing for quantitative MRI, Bonny and Renou, **159**, 183
- Intramyocellular lipids
 and extramyocellular lipids, signal separation in proton MR spectra by magnetic field distribution, Steidle *et al.*, **154**, 228
- Inversion sequences
 MR tagging using tandem of two adiabatic DANTE inversion sequences, Tsekos, Garwood, and Ugurbil, **156**, 187
- In vivo* electron paramagnetic resonance spectroscopy
 free radical detection/imaging, pulsed and continuous-wave RF EPR techniques, comparison, Yamada *et al.*, **154**, 287
- In vivo* magnetic resonance microscopy
 brain structure in unanesthetized flies, Jasanoff and Sun, **158**, 79
- In vivo* NMR
¹H, magnitude spectra, automated artificial neural network quantification of human brain metabolites, Hiltunen *et al.*, **154**, 1
- Ion-sensitive electrodes
 in device for aeration and mixing of cell and organelle suspensions during NMR, Bourne and Kuchel, **159**, 158
- Iridium
 single-crystal surfaces, multilayer experiments of T_1 relaxation of ¹²⁹Xe on, and monolayer considerations, Stahl *et al.*, **159**, 1
- Iron
 Fe³⁺, in natrolite, EPR: low-symmetry spin Hamiltonian and crystal field analysis, Vinokurov *et al.*, **155**, 57
 Fe³⁺ magnetic center in kaolin, EPR spectrum, temperature-induced changes, Vasyukov *et al.*, **154**, 15
- Isotopic labeling
 correlation of backbone amide and side-chain ¹³C resonances in perdeuterated proteins, Löhner and Rüterjans, **156**, 10
- J
- J*-compensation
 LR-*J*-HSMQC experiments to detect long-range heteronuclear shift correlations: Torres, Bubb, and Kuchel, **156**, 249
- J*-HSMQC
 detection of long-range heteronuclear shift correlations: LR-*J*-HSMQC, Torres, Bubb, and Kuchel, **156**, 249
- J*-modulated HSQC
 small one-bond proton-carbon residual dipolar coupling constants in ¹³C natural abundance carbohydrates, Pham *et al.*, **157**, 200
- K
- Kaolin
 magnetic center, temperature-induced changes in EPR spectrum, Vasyukov *et al.*, **154**, 15
- k-Space
 combined k-space q-space pulsed ESR imaging, Feintuch *et al.*, **157**, 69
- k-Space sampling
 and 2D prolate spheroidal wave functions for MRI, Yang *et al.*, **158**, 43
- L
- Lanczos algorithms
 improved, for blackbox MRS data quantitation, Laudadio *et al.*, **157**, 292
- Laplace inversion
 fast two-dimensional, T_2 – T_2 correlation spectra obtained using, Song *et al.*, **154**, 261
- Laser polarization
 NMR of laser-polarized noble gases in molecules, materials, and organisms, review, Goodson, **155**, 157
- Laser-polarized MRI
¹²⁹Xe, at ultralow magnetic fields, Wong-Foy *et al.*, **157**, 235
- Laser-polarized NMR
¹²⁹Xe, at ultralow magnetic fields, Wong-Foy *et al.*, **157**, 235
- Lateral lipid domains
 in erythrocyte ghost membranes, properties, quantification using EPR-spectra decomposition, Arsov, Schara, and Štrancar, **157**, 52
- Ligand–receptor complexes
 reversibly forming, intermolecular saturation transfer effects in, CORCEMA analysis, Jayalakshmi and Krishna, **155**, 106
- Line-narrowing effect
 in triple-quantum resonance in two-level NMR system, Hatanaka and Tabuchi, **155**, 119
- Lineshape
 variations, principal component analysis procedure for determining, Stoyanova and Brown, **154**, 163
- Linewidth
 NMR line broadening of organic molecules grafted onto polystyrene resin, and resin dynamics, Lippens, Chessari, and Wieruszkeski, **156**, 242

- Lipid bilayers
 cholesterol in, high-resolution 2D ^1H – ^{13}C correlation, Soubias *et al.*, **158**, 143
 macroscopically oriented, anisotropic water diffusion in, pulsed magnetic field gradient NMR, Wästerby, Orädd, and Lindblom, **157**, 156
- Lipids
 intra- and extramyocellular, signal separation in proton MR spectra by magnetic field distribution, Steidle *et al.*, **154**, 228
- Liquid crystalline samples
 molecules with natural abundant rare spins in, selective detection of ^1H NMR spectra, Emsley *et al.*, **154**, 303
- Liquid crystalline solvents
 nematic, dipolar couplings in partially oriented molecules in, local field NMR with low-power decoupling, Marjanska *et al.*, **158**, 52
- Liquid crystals
 2D NMR correlations using switched angle spinning, Havlin, Park, and Pines, **157**, 163
 endohedral fullerenes $\text{N}@C_{60}$ and $\text{N}@C_{70}$ in, EPR study, Jakes *et al.*, **156**, 303
 nematic
para-chlorotoluene dissolved in, simultaneous selective detection of MQ spectra, Syvitski *et al.*, **155**, 251
 ^{13}C NMR spectral assignments by 2D chemical-shift γ -encoding NMR, Nishiyama, Kubo, and Terao, **158**, 60
 solvent, order estimation: MQ NMR spectra of partially oriented indene, Field, Ramadan, and Pierens, **156**, 64
- Liquid crystal solvents
 weakly oriented chiral, enantiomeric excess measurements using 2D ^1H selective refocusing, Farjon *et al.*, **158**, 169
- Liquid films
 falling, NMR velocimetry, Heine *et al.*, **154**, 311
- Liquids
 NMR, adjustable broadband selective excitation with uniform phase, Cano, Smith, and Shaka, **155**, 131
- Liquid xenon polarization
 rapid precise probe for measurement, Verhulst *et al.*, **155**, 145
- Local field spectroscopy
 with low-power decoupling, dipolar couplings in partially oriented molecules, Marjanska *et al.*, **158**, 52
- Localized spectroscopy
 proton, without water suppression: frequency demodulation by modulus signal selection, Serrai *et al.*, **154**, 53
- Localized tagging
 SPAMM 1–1, rotational approach to, Ikonomidou and Sergiadis, **157**, 218
- Longitudinal magnetization
 dipolar-encoded, NMR of multipolar spin states excited in strongly inhomogeneous magnetic fields, Wiesmath *et al.*, **154**, 60
- Long length scale
 determination in xenon gas diffusion NMR of porous media, Mair *et al.*, **156**, 202
- Long-range correlations
 heteronuclear shift, LR-*J*-HSMQC experiments for detection: Torres, Bubb, and Kuchel, **156**, 249
 proton–carbon, increasing resolution of, 3D HSQC-HSQMBC for, Uhrin, **159**, 145
- Long-range dipolar interactions
 origins of dips in multiple spin echoes in heterogeneous systems, Alessandri, Capuani, and Maraviglia, **156**, 72
- Low-field MRI
 laser-polarized noble gases in molecules, materials, and organisms, review, Goodson, **155**, 157
- Low-field NMR
 laser-polarized noble gases in molecules, materials, and organisms, review, Goodson, **155**, 157
 relaxation data, rapid unique curve resolution, Pedersen, Bro, and Engelsen, **157**, 141
- Low-field NMR relaxation
in vitro, changes in porcine muscle water characteristics during growth, Bertram *et al.*, **157**, 267
- Low-symmetry effect
 spin Hamiltonian and crystal field, Fe^{3+} in natrolite, Vinokurov *et al.*, **155**, 57
- Low temperatures
 multiple-quantum dynamics in linear chains and rings of nuclear spins in solids at, Fel'dman and Maximov, **157**, 106
- LR-*J*-HSMQC
 detection of long-range heteronuclear shift correlations: Torres, Bubb, and Kuchel, **156**, 249

M

- Magic angle
 NMR spin locking of proton magnetization under frequency-switched Lee–Goldburg pulse sequence, Fu, Tian, and Cross, **154**, 130
- Magic-angle spinning (*see also* High-resolution magic-angle spinning NMR; Multiple-quantum MAS NMR)
 2D ^{13}C – ^{13}C MAS NMR correlation spectroscopy with mixing by true ^1H spin diffusion, de Boer *et al.*, **157**, 286
 2D dipolar correlation high-field, 3D structural model of aggregated chlorophyll *a*/H₂O, van Rossum *et al.*, **155**, 1
 2D MAS NMR method to recover amplified heteronuclear dipolar and CSA interactions, Wei *et al.*, **158**, 23
 2D separation of undistorted chemical-shift anisotropy powder patterns, Liu, Mao, and Schmidt-Rohr, **155**, 15
 conditions, enhanced spectral resolution of immobilized peptides and proteins by combining chemical shift sum and difference spectroscopy, Luca and Baldus, **159**, 243
 cross-polarization
 4-mm probe for ^{13}C CP/MAS NMR of solids at 21.15 T, Jakobsen *et al.*, **156**, 152
 and hyperpolarized ^{129}Xe , for spin polarization transfer to surface nuclei, Smith *et al.*, **159**, 111
 fast, homonuclear zero-quantum recoupling in, Brinkmann, Schmedt auf der Günne, and Levitt, **156**, 79
 filtered, observation of backbone carbons of membrane-bound fusion peptides, with REDOR subtraction, Yang *et al.*, **159**, 101
 high rates, double-quantum dipolar recoupling at, Kristiansen, Mitchell, and Evans, **157**, 253
 improved narrowband dipolar coupling for homonuclear distance measurements in rotating solids, Goobes and Vega, **154**, 236
 recoupling methods, application in intermediate motional regime, Saalwächter and Fischbach, **157**, 17
 satellite-transition
 optimizing, Huguenard *et al.*, **156**, 131
 spin $I = 3/2, 5/2, 7/2,$ and $9/2$ nuclei, Ashbrook and Wimperis, **156**, 269
 spectra enhancement, influencing satellite transitions of half-integer quadrupolar nuclei for, Iuga and Kentgens, **158**, 65
 SPINOE under: review of NMR of laser-polarized noble gases, Goodson, **155**, 157
 techniques, evaluation at low spinning rates in aqueous samples and in presence of magnetic susceptibility gradients, Hu and Wind, **159**, 93
- Magic-angle turning
 phase-corrected (*see* PHORMAT)
- Magnesium chloride
 ^{13}C NMR relaxation agent for amino acids and carboxylic acids, Tian *et al.*, **159**, 137
- Magnet
 open, utilizing ferro-refraction current magnetization, Pulyer and Hrovat, **154**, 298
- Magnetic alignment
 phage Pf1, capture and manipulation with aqueous polymer gel, Riley, Giuliani, and Augustine, **159**, 83

- Magnetic center
 Fe³⁺, in kaolin, EPR spectrum, temperature-induced changes, Vasyukov *et al.*, **154**, 15
- Magnetic field distribution
 separation of intra- and extramyocellular lipid signals in ¹H MR spectra by, Steidle *et al.*, **154**, 228
- Magnetic field gradient
 and B₁ field gradient, matched, 2D high-resolution NMR spectra in, Heise *et al.*, **156**, 146
 multiple-quantum NMR spectra of partially oriented indene acquired using, Field, Ramadan, and Pierens, **156**, 64
- Magnetic field homogeneity
 B₀, variation with head pitch, quantification by registered 3D field mapping, Tyszka and Mamelak, **159**, 213
- Magnetic fields
 strongly inhomogeneous, NMR of multipolar spin states excited in, Wiesmath *et al.*, **154**, 60
 and velocity field, spatial correlations, and PFG NMR measurements of flow through porous media, Lebon and Leblond, **159**, 13
- Magnetic resonance force microscopy
 probe–sample coupling in, Suter *et al.*, **154**, 210
- Magnetic resonance imaging
 automatic tuned MRI RF coil for multinuclear imaging of small animals at 3T, Muftuler *et al.*, **155**, 39
 contrast agents for
 cell uptake, high-resolution MAS NMR, Calabi *et al.*, **156**, 222
 relaxation enhancement by, role of diffusion and exchange, Gossein *et al.*, **158**, 36
 diffusion tensor, discrete data, continuous tensor field approximation, Pajevic, Aldroubi, and Bassar, **154**, 85
 double end-cap birdcage coil for small animal whole-body imaging, Gulsen, Muftuler, and Nalcioğlu, **156**, 309
 high-capacity production of >65% spin-polarized ¹²⁹Xe for, Zook, Adhyaru, and Bowers, **159**, 175
 interventional, use of Overhauser enhancement to assist with needle and catheter placement during, Vahala *et al.*, **157**, 298
 laser-polarized ¹²⁹Xe, at ultralow magnetic fields, Wong-Foy *et al.*, **157**, 235
 living tissues, application of laser-polarized noble gases, Goodson, **155**, 157
 multiple spectra in, quantitation of single resonances via complex principal component analysis, Van Huffel *et al.*, **158**, 1
 opposed-solenoid endovascular probes for, reciprocity and sensitivity, Crottet *et al.*, **159**, 219
 quantitative, Euclidian distance-weighted smoothing for, Bonny and Renou, **159**, 183
 two-dimensional prolate spheroidal wave functions for, Yang *et al.*, **158**, 43
 water content profiles with 1D centric SPRITE acquisition, Mastikhin *et al.*, **156**, 122
- Magnetic resonance imaging microscopy
 3D, with resolution 3.7 μm by 3.3 μm by 3.3 μm, Ciobanu, Seeber, and Pennington, **158**, 178
- Magnetic resonance microscopy
 3D, with resolution 3.7 μm by 3.3 μm by 3.3 μm, Ciobanu, Seeber, and Pennington, **158**, 178
in vivo, brain structure in unanesthetized flies, Jasanoff and Sun, **158**, 79
 short planar gradients coils using concentric return paths for, Dodd and Ho, **156**, 1
- Magnetic resonance signal formation
 effects of restricted diffusion, Sukstanskii and Yablonskiy, **157**, 92
- Magnetic resonance spectrometers
 complex response function of, Annino *et al.*, **157**, 74
- Magnetic resonance spectroscopic imaging
 automated correction of unwanted phase jumps in reference signals corrupting MRSI spectra after eddy current correction, Simonetti *et al.*, **159**, 151
 multiple spectra in, quantitation of single resonances via complex principal component analysis, Van Huffel *et al.*, **158**, 1
- Magnetic resonance spectroscopy (*see* NMR)
- Magnetic resonance tagging
 localized SPAMM 1–1, rotational approach to, Ikonomidou and Sergiadis, **157**, 218
 using tandem of two adiabatic DANTE inversion sequences, Tsekos, Garwood, and Ugurbil, **156**, 187
- Magnetic susceptibility gradients
 evaluation of MAS techniques at low spinning rates in aqueous samples in presence of, Hu and Wind, **159**, 93
- Magnetization
 ferro-refraction current, open magnet utilizing, Pulyer and Hrovat, **154**, 298
 longitudinal, dipolar-encoded, NMR of multipolar spin states excited in strongly inhomogeneous magnetic fields, Wiesmath *et al.*, **154**, 60
 OH and NH protons, selective dephasing based on ¹H chemical-shift anisotropy recoupling, Schmidt-Rohr and Mao, **157**, 210
 proton, NMR spin locking under frequency-switched Lee–Goldburg pulse sequence, Fu, Tian, and Cross, **154**, 130
- Magnitude spectra
¹H NMR, automated artificial neural network quantification of *in vivo* human brain metabolites from, Hiltunen *et al.*, **154**, 1
- Master equations
 for linear and partial adiabatic pulses, Tesiram and Bendall, **156**, 26
- Materials
 NMR of laser-polarized noble gases in, review, Goodson, **155**, 157
- Matrices
 amorphous, large-angle jumps of small molecules in, 2D exchange NMR analysis, Medick, Vogel, and Rössler, **159**, 126
- MBOB
 pulse sequence, optimizing delays in, Schulte-Herbrüggen *et al.*, **156**, 282
- Mean squared dipolar fluctuation
 contrast parameter for visualization of crosslink density in rubber based on dipolar-correlation effect, Grinberg, Heidenreich, and Kuhn, **159**, 88
- Mechanical vibrations
 magnetic resonance detection using radiofrequency field gradient, Baril *et al.*, **154**, 22
- Membrane peptides
 fusion, filtered MAS observation of backbone carbons, with REDOR subtraction, Yang *et al.*, **159**, 101
 orientation, from ¹H-detected ²H NMR, Yamaguchi and Hong, **155**, 244
- Membranes
 model, cholesterol in, high-resolution 2D ¹H–¹³C correlation, Soubias *et al.*, **158**, 143
- Metabolite quantitation
 errors due to saturation factor changes in MRS, optimized pulse parameters for reducing, Galbán and Spencer, **156**, 161
- Metabolites
 brain, automated artificial neural network quantification from *in vivo* ¹H NMR magnitude spectra, Hiltunen *et al.*, **154**, 1
- Metal surfaces
 single-crystal, T₁ relaxation of ¹²⁹Xe on, Stahl *et al.*, **159**, 1
- Methyl groups
 Curie spin relaxation in, cross-correlation effects involving, Madhu, Mandal, and Müller, **155**, 29
- Methyl rotation effects
 on electron spin-lattice relaxation in radicals with two methyl groups in γ-irradiated polycrystalline solids, Harbridge, Eaton, and Eaton, **159**, 195
- Micron resolution
 3D MR microscopy with resolution 3.7 μm by 3.3 μm by 3.3 μm, Ciobanu, Seeber, and Pennington, **158**, 178

- Microprocessor control
for portable NMR relaxometer: thermostatic control system, Anisimov, Teplov, and Silkin, **154**, 176
- Mixing
and aeration, cell and organelle suspensions during NMR experiments, device for, Bourne and Kuchel, **159**, 158
- Mixtures
resolving diffusion coefficients: 2D-J-DOSY experiment, Lucas, Otto, and Larive, **156**, 138
- Modulated field gradient
EPR imaging with, kinetic measurements using, Herrling, Fuchs, and Groth, **154**, 6
PFGSE NMR study of restricted self-diffusion of water in highly concentrated W/O emulsion, Topgaard, Melander, and Söderman, **156**, 195
- Modulus signal selection
removal of gradient-induced frequency modulations: localized ^1H MRS without water suppression, Serrai *et al.*, **154**, 53
- Monitoring
design and implementation of protection system for NMR spectrometers, Laurent and Guéron, **156**, 295
- Monolayer
considerations, and T_1 relaxation of ^{129}Xe on single-crystal metal surfaces, Stahl *et al.*, **159**, 1
- Monte Carlo minimization approach
structural determination of spin label immobilization and orientation, Sale *et al.*, **156**, 104
- MP-HNCA
evaluation and optimization of coherence transfer in high-molecular-weight systems, Permi and Annala, **155**, 123
- Multi-compartment systems
with cylindrical geometry, modeling of self-diffusion and relaxation time NMR in, van der Weerd *et al.*, **156**, 213
- Multidimensional electrophoretic NMR
constant-time, Li and He, **156**, 181
- Multidimensional NMR
achievable accuracy of parameter estimation for, Ober *et al.*, **157**, 1
- Multifrequency EPR
CW-EPR study of radical anions of fullerene bisadducts, Zoleo *et al.*, **159**, 226
- Multilayer experiments
 T_1 relaxation of ^{129}Xe on single-crystal metal surfaces, Stahl *et al.*, **159**, 1
- Multiple-quantum MAS NMR
modified RIACT, for $I = 3/2$ nuclei, efficient triple-quantum excitation in, Lim, Charpentier, and Pines, **154**, 196
rotor-encoded heteronuclear, of half-integer quadrupolar and spin $I = 1/2$ nuclei, Lupulescu, Brown, and Spiess, **154**, 101
spin-1/2 quadrupolar nuclei, signal enhancement in 5QMAS spectra, Goldbourt and Vega, **154**, 280
triple-quantum, spins-3/2 in solids, signal enhancement, Madhu and Levitt, **155**, 150
- Multiple-quantum NMR
3D, simultaneous selective detection of MQ spectra, Syvitski *et al.*, **155**, 251
line-narrowing effect in triple-quantum resonance in two-level NMR system, Hatanaka and Tabuchi, **155**, 119
multiple-quantum dynamics in linear chains and rings of nuclear spins in solids at low temperatures, Fel'dman and Maximov, **157**, 106
sensitivity enhancement with CPMG detection in, Lim *et al.*, **157**, 160
spectra of partially oriented indene, Field, Ramadan, and Pierens, **156**, 64
- Multiple spin echoes
in heterogeneous systems: origin of observed dips, Alessandri, Capuani, and Maraviglia, **156**, 72
- Multipolar spin states
excited in strongly inhomogeneous magnetic fields, NMR, Wiesmath *et al.*, **154**, 60
- Multispin labeling
and MAS NMR 2D dipolar correlation, model for aggregated chlorophyll *a*/H₂O from, van Rossum *et al.*, **155**, 1
- ## N
- ^{14}N
dipolar coupling to, effect on ^{13}C FIREMAT experiments, Strohmeier, Alderman, and Grant, **155**, 263
isotropic chemical shifts and quadrupole coupling constants, for inorganic nitrates, Marburger, Fung, and Khitritin, **154**, 205
- ^{15}N
 ^{15}N , ^{13}C -labeled proteins
backbone assignment, intraresidue HNCA and COHNCA experiments for, Brutscher, **156**, 155
perdeuterated, correlation of backbone amide and side-chain ^{13}C resonances in, Löhr and Rüterjans, **156**, 10
 ^{15}N -labeled proteins, $^3J_{\text{HNH}\alpha}$ scalar couplings in, rapid determination with cosine-modulated HSQC, Petit, Vincent, and Zwahlen, **156**, 313
one-bond ^{15}N - ^1H and ^{15}N - ^{13}C and two-bond ^{13}C - ^1H residual dipolar couplings in proteins, sensitivity-enhanced E.COSY-type HSQC for, Ding and Gronenborn, **158**, 173
 $\text{U-}^{13}\text{C}$, ^{15}N -labeled peptides and proteins, *soft*-triple-resonance solid-state NMR for assignments of, Astrof and Griffin, **158**, 157
- Narrowband dipolar recoupling
improved, for homonuclear distance measurements in rotating solids, Goobes and Vega, **154**, 236
- Narrow pulse approximation
in xenon gas diffusion NMR of porous media, Mair *et al.*, **156**, 202
- Natrolite
 Fe^{3+} , EPR: low-symmetry spin Hamiltonian and crystal field analysis, Vinokurov *et al.*, **155**, 57
- Needle placement
during interventional MRI, use of Overhauser enhancement to assist with, Vahala *et al.*, **157**, 298
- Neural networks
use for ^{13}C NMR chemical shift prediction, Meiler *et al.*, **157**, 242
- NH
and OH, proton magnetization, selective dephasing based on ^1H chemical-shift anisotropy recoupling, Schmidt-Rohr and Mao, **157**, 210
- Nitrates
inorganic, ^{14}N chemical shifts and quadrupole coupling constants, Marburger, Fung, and Khitritin, **154**, 205
- Nitric oxide
adsorption and desorption behavior on H-ZSM-5, Na-ZSM-5, Na-A, EPR study, Rudolf, Böhlmann, and Pöpl, **155**, 45
- Nitroxide biradicals
electron–electron interaction in, 130 GHz ESEEM induced by, Kulik, Paschenko, and Dzuba, **159**, 237
field-step ELDOR, electron dipole–dipole ESEEM in, Kulik *et al.*, **157**, 61
- Nitroxides
hydrogen-bonded, polarity contributions to hyperfine splittings of, Marsh, **157**, 114
- NMR
aeration and mixing of cell and organelle suspensions during, device for, Bourne and Kuchel, **159**, 158
automated spectral normalization, subspace time-domain algorithm for, Lemmerling *et al.*, **157**, 190
blackbox MRS data quantitation, improved Lanczos algorithms for, Laudadio *et al.*, **157**, 292

- cesium adsorption in hydrated iron oxide particle suspensions, Gossuin *et al.*, **157**, 132
- detection of mechanical vibrations using radiofrequency field gradient, Baril *et al.*, **154**, 22
- diffusion measurements, analysis of protein–ligand interactions with, Derrick, McCord, and Larive, **155**, 217
- flow-compensating NMR-PFG techniques, sizing of emulsion droplets under flow using, Johns and Gladden, **154**, 142
- ¹H (see Proton NMR)
- ²H, ¹H-detected, membrane peptide orientation by, Yamaguchi and Hong, **155**, 244
- high-capacity production of >65% spin-polarized ¹²⁹Xe for, Zook, Adhyaru, and Bowers, **159**, 175
- high-resolution (see High-resolution NMR)
- laser-polarized noble gases in molecules, materials, and organisms, review, Goodson, **155**, 157
- laser-polarized ¹²⁹Xe, at ultralow magnetic fields, Wong-Foy *et al.*, **157**, 235
- liquids, adjustable broadband selective excitation with uniform phase, Cano, Smith, and Shaka, **155**, 131
- multidimensional, achievable accuracy of parameter estimation for, Ober *et al.*, **157**, 1
- multiple-quantum (see Multiple-quantum NMR)
- multipolar spin states excited in strongly inhomogeneous magnetic fields, Wiesmath *et al.*, **154**, 60
- phase noise, inductive shielding, Sigmund *et al.*, **159**, 190
- quantitation errors due to saturation factor changes, optimized pulse parameters for reducing, Galbán and Spencer, **156**, 161
- quantum information processing by, implementation of conditional phase-shift gate for, using transition-selective pulses, Das, Mahesh, and Kumar, **159**, 46
- single-crystal metal surfaces, *T*₁ relaxation of ¹²⁹Xe, Stahl *et al.*, **159**, 1
- spectra
- automatic phase correction, efficient algorithm based on entropy minimization, Chen *et al.*, **158**, 164
 - improved solvent suppression using wavelet transforms: WAVE-MAT, Günther, Ludwig, and Rüterjans, **156**, 19
 - quantitation by principal component analysis: procedure for determining lineshape variations, Stoyanova and Brown, **154**, 163
- time-domain frequency-selective quantification algorithm, Romano *et al.*, **155**, 226
- torsion angle Ψ determination in α -helical peptides and proteins, Ladizhansky, Veshtort, and Griffin, **154**, 317
- two-level system, line-narrowing effect in triple-quantum resonance in, Hatanaka and Tabuchi, **155**, 119
- NMR difference probe
- dual-coil, Macnaughtan *et al.*, **156**, 97
- NMR imaging
- contrast parameter for visualization of crosslink density in rubber based on dipolar-correlation effect, Grinberg, Heidenreich, and Kuhn, **159**, 88
- NMR-MOUSE
- analysis of excitation of multipolar spin states in strongly inhomogeneous magnetic fields, Wiesmath *et al.*, **154**, 60
- NMR Nomenclature
- Nuclear Spin Properties and Conventions for Chemical Shifts—IUPAC Recommendations, announcement, **156**, 323
- NMR relaxation
- paramagnetic enhancement, application of closed-form expressions for level-averaged ESR times outside Zeeman limit, Sharp, **154**, 269
- NMR relaxometer
- portable, microprocessor control for, Anisimov, Teplov, and Silkin, **154**, 176
- NMR scattering
- scaling laws via dipolar fields, Brown, Sen, and Cory, **154**, 154
- NMR spin locking
- proton magnetization under frequency-switched Lee–Goldburg pulse sequence, Fu, Tian, and Cross, **154**, 130
- NMR velocimetry
- falling liquid films, Heine *et al.*, **154**, 311
- Noble gases
- laser-polarized, in molecules, materials, and organisms, NMR: review, Goodson, **155**, 157
- Noise spectroscopy
- suppression of ring-down in, Yang and Zax, **158**, 73
- Nonuniform averaging
- in indirect dimension: achievable accuracy of parameter estimation for multidimensional NMR, Ober *et al.*, **157**, 1
- Normalization
- automated NMR spectral, subspace time-domain algorithm for, Lemmerling *et al.*, **157**, 190
- Nuclear frequencies
- at canonical orientations of *g*-tensor, determination of hyperfine tensor components from, Shubin and Dikanov, **155**, 100
- Nuclear spins
- in solids at low temperatures, multiple-quantum dynamics in linear chains and rings of, Fel'dman and Maximov, **157**, 106
- Numerical modeling
- self-diffusion and relaxation time NMR in multi-compartment systems with cylindrical geometry, van der Weerd *et al.*, **156**, 213
- Numerical simulations
- biological solid-state NMR, and specification/visualization of polypeptide anisotropic interaction tensors, Bak *et al.*, **154**, 28
 - one- and two-dimensional ESEEM experiments, Madi, Van Doorslaer, and Schweiger, **154**, 181
- O
- Offset effects
- and chemical-shift anisotropy effects, in cross-polarization solid-state NMR, Shekar, Lee, and Ramamoorthy, **157**, 223
- OH
- and NH, proton magnetization, selective dephasing based on ¹H chemical-shift anisotropy recoupling, Schmidt-Rohr and Mao, **157**, 210
- Oligosaccharides
- ¹³C natural abundance, partially oriented, measurement of small one-bond proton–carbon residual dipolar coupling constants, Pham *et al.*, **157**, 200
- One-dimensional imaging
- water content profiles with 1D centric SPRITE acquisition, Mastikhin *et al.*, **156**, 122
- One-dimensional organic conductors
- restricted diffusion mapping in (FA)₂PF₆: combined k-space q-space pulsed ESR imaging, Feintuch *et al.*, **157**, 69
- ¹⁷O NMR
- cristobalite, central transition NMR spectroscopy: anisotropic relaxation and motion of half-integer quadrupolar nuclei, Kristensen and Farnan, **158**, 99
- Open magnet
- utilizing ferro-refraction current magnetization, Pulyer and Hrovat, **154**, 298
- Optical pumping
- noble gases, Goodson, **155**, 157
- Optimal control theory
- time-optimal coherence-order-selective transfer of in-phase coherence in heteronuclear *IS* spin systems, Reiss, Khaneja, and Glaser, **154**, 192
- Order parameters
- in nematic phase, estimation: MQ NMR spectra of partially oriented indene, Field, Ramadan, and Pierens, **156**, 64
- Organelles
- suspensions, aeration and mixing during NMR experiments, device for, Bourne and Kuchel, **159**, 158
- Organic molecules
- grafted onto polystyrene resins, NMR line broadening, role of resin dynamics, Lippens, Chessari, and Wieruszkeski, **156**, 242

Organisms

NMR of laser-polarized noble gases in, review, Goodson, **155**, 157

Orientation

canonical, *g*-tensor, determination of hyperfine tensor components from nuclear frequencies at, Shubin and Dikanov, **155**, 100

enantiomeric excess measurements in weakly oriented chiral liquid crystal solvents using 2D ^1H selective refocusing, Farjon *et al.*, **158**, 169

interaction tensors, sensitivity of DQ solid state NMR spectra of ^{13}C spin pairs coupled to ^{14}N , Hughes *et al.*, **159**, 25

membrane peptide, from ^1H -detected ^2H NMR, Yamaguchi and Hong, **155**, 244

relative

alignment tensors for protein dissolved in two anisotropic media, graphical analysis, Nomura and Kainosho, **154**, 146

CSA–dipolar, from REDOR sidebands, O'Connor and Schaefer, **154**, 46

two domains from dipolar couplings, by conjoined rigid body/torsion angle simulated annealing, Clore and Bewley, **154**, 329

spin label, structural determination, Sale *et al.*, **156**, 104

Orientationally disordered systems

S > 1/2 centers in, ESEEM and HYSOCORE spectra, Benetis, Dave, and Goldfarb, **158**, 126

Overhauser enhancement

use to assist with needle and catheter placement during interventional MRI, Vahala *et al.*, **157**, 298

Oxygen

dissolved, effects on amide proton relaxation and chemical shift in perdeuterated protein, Ulmer, Campbell, and Boyd, **157**, 181

P

Paramagnetic center

resonance assignment in proximity of, tailored HCCH–TOCSY experiment for, Piccioli and Poggi, **155**, 236

Paramagnetic NMR relaxation

application of closed-form expressions for level-averaged ESR times outside Zeeman limit, Sharp, **154**, 269

Parameter estimation

for multidimensional NMR, achievable accuracy of, Ober *et al.*, **157**, 1

PASS

2D, evaluation at low spinning rates in aqueous samples and in presence of magnetic susceptibility gradients, Hu and Wind, **159**, 93

PELDOR

at S- and X-band frequencies, and separation of exchange coupling from dipolar coupling, Weber *et al.*, **157**, 277

Peltier thermopiles

thermostatic control system, microprocessor control for portable NMR relaxometer, Anisimov, Teplov, and Silkin, **154**, 176

Penetration enhancement

skin: kinetic measurements by EPR imaging with modulated field gradient, Herrling, Fuchs, and Groth, **154**, 6

Peptides

fusion, membrane-bound, filtered MAS observation of backbone carbons, with REDOR subtraction, Yang *et al.*, **159**, 101

immobilized, enhanced spectral resolution by combining chemical shift sum and difference spectroscopy, Luca and Baldus, **159**, 243

membrane, orientation from ^1H -detected ^2H NMR, Yamaguchi and Hong, **155**, 244

resin-bound, proton dipolar recoupling under high-resolution MAS in, Raya *et al.*, **157**, 43

transmembrane, in bicelles, study methods: consequences of hydrophobic mismatch and peptide sequence, Whiles *et al.*, **158**, 149

$\text{U-}^{13}\text{C}$, ^{15}N -labeled, *soft*-triple-resonance solid-state NMR for assignments of, Astrof and Griffin, **158**, 157

Perdeuterated proteins

amide proton relaxation and chemical shift in, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181

correlation of backbone amide and side-chain ^{13}C resonances in, Löhner and Rüterjans, **156**, 10

Pfl

magnetically aligned, capture and manipulation with aqueous polymer gel, Riley, Giuliani, and Augustine, **159**, 83

Phantom studies

gradient echo imaging of flowing hyperpolarized nuclei: ^{129}Xe dissolved in ethanol, Johansson *et al.*, **159**, 69

Phase correction

automatic, NMR spectra, efficient algorithm based on entropy minimization, Chen *et al.*, **158**, 164

Phase cycling

cogwheel, description of method, Levitt, Madhu, and Hughes, **155**, 300

Phase-incremented pulses

coherent excitation with, Zhang and Gorenstein, **154**, 73

Phase noise

NMR, inductive shielding, Sigmund *et al.*, **159**, 190

PHORMAT

evaluation at low spinning rates in aqueous samples and in presence of magnetic susceptibility gradients, Hu and Wind, **159**, 93

Picoliter ^1H NMR

theoretical basis review and demonstration, Minard and Wind, **154**, 336

PISEMA

effect of Hartmann–Hahn mismatching, Fu *et al.*, **159**, 167

NMR spin locking of proton magnetization, frequency-switched Lee–Goldburg pulse sequence, Fu, Tian, and Cross, **154**, 130

Pluronic F-127

gel, capture and manipulation of magnetically aligned phage Pfl with, Riley, Giuliani, and Augustine, **159**, 83

Polarity

role in hyperfine splittings of hydrogen-bonded nitroxides, Marsh, **157**, 114

Polarization

liquid ^{129}Xe , rapid precise probe for measurement, Verhulst *et al.*, **155**, 145

Polycrystalline solids

γ -irradiated, radicals with two methyl groups in, electron spin-lattice relaxation, Harbridge, Eaton, and Eaton, **159**, 195

Polymer gel

aqueous, capture and manipulation of magnetically aligned phage Pfl with, Riley, Giuliani, and Augustine, **159**, 83

Polymeric composites

temperature-induced changes in EPR spectrum of magnetic center in kaolin, Vasyukov *et al.*, **154**, 15

Polypeptides

2D MAS NMR method to recover amplified heteronuclear dipolar and CSA interactions, Wei *et al.*, **158**, 23

anisotropic interaction tensors in, specification and visualization, and numerical simulations in biological solid-state NMR, Bak *et al.*, **154**, 28

Polystyrene resin

NMR line broadening of organic molecules grafted onto, and resin dynamics, Lippens, Chessari, and Wieruszkeski, **156**, 242

Porous media

enhanced transverse relaxation due to internal field gradients, Dunn, **156**, 171

flow through, PFG NMR measurements: effects of spatial correlations of magnetic field and velocity field, Lebon and Leblond, **159**, 13

multiple spin echoes, physical origin of observed dips, Alessandri, Capuani, and Maraviglia, **156**, 72

water content profiles with 1D centric SPRITE acquisition, Mastikhin *et al.*, **156**, 122

xenon gas diffusion NMR, narrow pulse approximation and long length scale determination in, Mair *et al.*, **156**, 202

Powder patterns

undistorted chemical-shift anisotropy, 2D separation in MAS NMR, Liu, Mao, and Schmidt-Rohr, **155**, 15

- Principal component analysis
 complex, automatic frequency alignment and quantitation of single resonances in multiple MR spectra via, Van Huffel *et al.*, **158**, 1
 NMR spectral quantitation by, procedure for determining lineshape variations, Stoyanova and Brown, **154**, 163
- Probes
 opposed-solenoid endovascular MRI, reciprocity and sensitivity, Crottet *et al.*, **159**, 219
 rapid precise, for liquid ^{129}Xe polarization measurement, Verhulst *et al.*, **155**, 145
- Probe-sample interactions
 in magnetic resonance force microscopy, Suter *et al.*, **154**, 210
- Prolate spheroidal wave functions
 two-dimensional, for MRI, Yang *et al.*, **158**, 43
- Propagator
 determination, and spectrally resolved velocity exchange spectroscopy of two-phase flow, Khrapitchev *et al.*, **159**, 36
- Protection system
 for NMR spectrometers, design and implementation, Laurent and Guéron, **156**, 295
- Protein background
 eliminating, and analysis of protein–ligand interactions with NMR diffusion measurements, Derrick, McCord, and Larive, **155**, 217
- Protein content
 and changes in porcine muscle water characteristics during growth, *in vitro* low-field ^1H NMR, Bertram *et al.*, **157**, 267
- Protein domains
 covalently linked, from dipolar couplings, relative orientation by conjoined rigid body/torsion angle simulated annealing, Clore and Bewley, **154**, 329
- Protein–ligand interactions
 analysis with NMR diffusion measurements, Derrick, McCord, and Larive, **155**, 217
- Proteins
 backbone assignment, intraresidue HNCA and COHNCA experiments for, Brutscher, **156**, 155
 dissolved in different anisotropic media, graphical analysis of relative orientation of molecular alignment tensors for, Nomura and Kainosho, **154**, 146
 immobilized, enhanced spectral resolution by combining chemical shift sum and difference spectroscopy, Luca and Baldus, **159**, 243
 with known structure, assignment strategy for, Hus, Prompers, and Brüschweiler, **157**, 119
 ^{15}N -labeled, $^3J_{\text{HNH}\alpha}$ scalar couplings in, rapid determination with cosine-modulated HSQC, Petit, Vincent, and Zwanen, **156**, 313
 one-bond ^{15}N - ^1H and ^{15}N - ^{13}C and two-bond ^{13}C - ^1H residual dipolar couplings in, sensitivity-enhanced E.COSY-type HSQC for, Ding and Gronenborn, **158**, 173
 sequential resonance assignment, 2D triple-resonance NMR experiments for, Ding and Gronenborn, **156**, 262
 U- ^{13}C , ^{15}N -labeled, *soft*-triple-resonance solid-state NMR for assignments of, Astrof and Griffin, **158**, 157
- Proton–carbon long-range correlation
 increasing resolution of, 3D HSQC-HSQMBC for, Uhrin, **159**, 145
- Proton-detected local field spectroscopy
 with low-power decoupling, measurement of dipolar couplings in partially oriented molecules, Marjanska *et al.*, **158**, 52
- Proton dipolar recoupling
 in resin-bound peptides under high-resolution MAS, Raya *et al.*, **157**, 43
- Proton magnetization
 NMR spin locking under frequency-switched Lee–Goldburg pulse sequence, Fu, Tian, and Cross, **154**, 130
 OH and NH, selective dephasing based on ^1H chemical-shift anisotropy recoupling, Schmidt-Rohr and Mao, **157**, 210
- Proton NMR
 enantiomeric excess measurements in weakly oriented chiral liquid crystal solvents using 2D ^1H selective refocusing, Farjon *et al.*, **158**, 169
 ^1H magnitude spectra, automated artificial neural network quantification of *in vivo* human brain metabolites, Hiltunen *et al.*, **154**, 1
 localized, without water suppression: frequency demodulation by modulus signal selection, Serrai *et al.*, **154**, 53
 picoliter, Minard and Wind, **154**, 336
 separation of intra- and extramyocellular lipid signals by magnetic field distribution, Steidle *et al.*, **154**, 228
 spectra of molecules with rare spins at natural abundance in nematic solvent, selective detection, Emsley *et al.*, **154**, 303
- Protons
 amide, temperature coefficients, location of hydrogen bonds in human ubiquitin using, Cierpicki *et al.*, **157**, 178
- Pulse calibration
 and GAMMA simulations of stray-field responses, Kinchesch *et al.*, **154**, 252
- Pulsed ESR imaging
 combined k-space q-space, Feintuch *et al.*, **157**, 69
- Pulsed-field gradient
 flow-compensating NMR-PFG techniques, sizing of emulsion droplets under flow using, Johns and Gladden, **154**, 142
- Pulsed-field gradient NMR
 analysis of protein–ligand interactions with NMR diffusion measurements, Derrick, McCord, and Larive, **155**, 217
 anisotropic water diffusion in macroscopically oriented lipid bilayers, Wästerby, Orädd, and Lindblom, **157**, 156
 measurements of flow through porous media, effects of spatial correlations of magnetic field and velocity field, Lebon and Leblond, **159**, 13
- Pulsed-field gradient spin-echo NMR
 modulated, study of restricted self-diffusion of water in highly concentrated W/O emulsion, Topgaard, Melander, and Söderman, **156**, 195
- Pulsed gradients
 and stray-field gradients, combined, multidimensional imaging using, Godward *et al.*, **155**, 92
- Pulsed gradient spin-echo NMR
 xenon gas diffusion NMR of porous media, Mair *et al.*, **156**, 202
- Pulsed spin locking
 sensitivity enhancement in structural measurements by solid-state NMR, Petkova and Tycko, **155**, 293
- Pulse parameters
 optimized, for reducing quantitation errors due to saturation factor changes in MRS, Galbán and Spencer, **156**, 161
- Pulse sequence
 frequency-switched Lee–Goldburg, NMR spin locking of proton magnetization under, Fu, Tian, and Cross, **154**, 130
 method for variable temperature gradient shimming, Evans, Morris, and Davis, **154**, 325
 RIDE'n RIPT—ring down elimination in rapid imaging pulse trains, Woelk *et al.*, **159**, 207

Q

- q-space
 combined k-space q-space pulsed ESR imaging, Feintuch *et al.*, **157**, 69
- Quadrupolar nuclei
 half-integer
 anisotropic relaxation and motion, central transition NMR spectroscopy, Kristensen and Farnan, **158**, 99
 rotor-encoded heteronuclear MQ MAS NMR, Lupulescu, Brown, and Spiess, **154**, 101
 satellite transitions of, influencing for enhancement of MAS NMR spectra, Iuga and Kentgens, **158**, 65
 $I = 3/2$, modified RIACT MQMAS NMR for, efficient triple-quantum excitation in, Lim, Charpentier, and Pines, **154**, 196

- optimizing STMAS, Huguenard *et al.*, **156**, 131
 spin-5/2, signal enhancement in 5QMAS spectra, Goldbourt and Vega, **154**, 280
- Quadrupole coupling constants
 for ^{14}N , of inorganic nitrates, Marburger, Fung, and Khitritin, **154**, 205
- Quadrupole MRI probes
 opposed-solenoid endovascular, reciprocity and sensitivity, Crottet *et al.*, **159**, 219
- Quantification algorithm
 time-domain frequency-selective, Romano *et al.*, **155**, 226
- Quantitation
 errors due to saturation factor changes in MRS, optimized pulse parameters for reducing, Galbán and Spencer, **156**, 161
 single resonances in multiple MR spectra, via complex principal component analysis, Van Huffel *et al.*, **158**, 1
- Quantitative data analysis
 improved Lanczos algorithms for blackbox MRS data quantitation, Laudadio *et al.*, **157**, 292
- Quantitative imaging
 MRI, Euclidian distance-weighted smoothing for, Bonny and Renou, **159**, 183
- Quantum information processing
 by NMR, implementation of conditional phase-shift gate for, using transition-selective pulses, Das, Mahesh, and Kumar, **159**, 46
- R**
- Radical anions
 fullerene bisadducts, multifrequency CW-EPR study, Zoleo *et al.*, **159**, 226
- Radical pair
 130 GHz ESEEM induced by electron–electron interaction in biradical, Kulik, Paschenko, and Dzuba, **159**, 237
- Radicals
 with two methyl groups, in γ -irradiated polycrystalline solids, electron spin–lattice relaxation, Harbridge, Eaton, and Eaton, **159**, 195
- Radiofrequency coil
 automatic tuned MRI, for multinuclear imaging of small animals at 3T, Muftuler *et al.*, **155**, 39
 double end-cap birdcage coil for small animal whole-body imaging, Gulsen, Muftuler, and Nalcioglu, **156**, 309
- Radiofrequency-driven dipolar recoupling
 use in recovering proton dipolar interaction in resin-bound peptides under high-resolution MAS, Raya *et al.*, **157**, 43
- Radiofrequency field gradient
 and B_0 field gradient, matched, 2D high-resolution NMR spectra in, Heise *et al.*, **156**, 146
 magnetic resonance detection of mechanical vibrations using, Baril *et al.*, **154**, 22
- Radiofrequency field pulses
 selective, sample restriction in high-resolution solid-state NMR using, Charmont, Sakellariou, and Emsley, **54**, 136
- Radiofrequency offset
 and chemical-shift anisotropy, effects in cross-polarization solid-state NMR, Shekar, Lee, and Ramamoorthy, **157**, 223
- Radiofrequency power
 feedback: active resonator system for CW-ESR operating at 700 MHz, Sato *et al.*, **159**, 161
- Rapid imaging pulse trains
 ring down elimination in—RIDE'n RIPT, Woelk *et al.*, **159**, 207
- Rapid-passage EPR
 out-of-phase, for samples with long electron spin relaxation times, enhanced signal intensities, Harbridge *et al.*, **156**, 41
- REAPDOR
 internuclear distance determination of $S = 1$, $I = 1/2$ spin pairs, Hughes *et al.*, **156**, 230
- Reciprocity
 opposed-solenoid endovascular MRI probes, Crottet *et al.*, **159**, 219
- Recoupling
 double-quantum dipolar, at high magic-angle spinning rates, Kristiansen, Mitchell, and Evans, **157**, 253
 ^1H chemical-shift anisotropy, selective dephasing of OH and NH proton magnetization based on, Schmidt-Rohr and Mao, **157**, 210
 MAS, methods, application in intermediate motional regime, Saalwächter and Fischbach, **157**, 17
- REDOR
 implications of application of MAS recoupling methods in intermediate motional regime, Saalwächter and Fischbach, **157**, 17
 sidebands, relative CSA–dipolar orientation from, O'Connor and Schaefer, **154**, 46
 structural measurements, sensitivity enhancement by pulsed spin locking, Petkova and Tycko, **155**, 293
- REDOR subtraction
 filtered MAS observation of backbone carbons of membrane-bound fusion peptides using, Yang *et al.*, **159**, 101
- Reference deconvolution
 automated correction of unwanted phase jumps in reference signals corrupting MRSI spectra after eddy current correction, Simonetti *et al.*, **159**, 151
- Relaxation
 amide proton, in perdeuterated protein, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181
 ^{13}C , hexamethylenetetramine: dipole-dipole coupling constant for directly bonded CH pair, Kowalewski, Effemey, and Jokisaari, **157**, 171
 and diffusion, 2D distribution functions, quantitative measurement in grossly inhomogeneous fields, Hürlimann and Venkataraman, **157**, 31
 enhancement by MRI contrast agents, role of diffusion and exchange, Gossuin *et al.*, **158**, 36
 $R_{1\rho}$, outside of fast-exchange limit, Trott and Palmer, **154**, 157
- Relaxation agent
 magnesium chloride as, for ^{13}C NMR of amino acids and carboxylic acids, Tian *et al.*, **159**, 137
- Relaxation time NMR
 modeling in multi-compartment systems with cylindrical geometry, van der Weerd *et al.*, **156**, 213
- Residual dipolar coupling constants
 small one-bond proton–carbon, measurement in partially oriented ^{13}C natural abundance oligosaccharides, Pham *et al.*, **157**, 200
- Residual dipolar couplings
 assignment strategy for proteins with known structure using, Hus, Prompers, and Brüschweiler, **157**, 119
 efficiency of homonuclear Hartmann-Hahn and COSY-type mixing sequences in presence of, Kramer and Glaser, **155**, 83
 graphical analysis of relative orientation of protein alignment tensors in two anisotropic media, Nomura and Kainosho, **154**, 146
 one-bond ^{15}N - ^1H and ^{15}N - ^{13}C and two-bond ^{13}C - ^1H in proteins, sensitivity-enhanced E.COSY-type HSQC, Ding and Gronenborn, **158**, 173
- Resin-bound peptides
 proton dipolar recoupling in, under high-resolution MAS, Raya *et al.*, **157**, 43
- Resin dynamics
 and NMR line broadening of organic molecules grafted onto polystyrene resin, Lippens, Chessari, and Wieruszski, **156**, 242
- Resins
 polystyrene, NMR line broadening of organic molecules grafted onto, and resin dynamics, Lippens, Chessari, and Wieruszski, **156**, 242
- Resonance assignment
 in proximity of paramagnetic center, tailored HCCH–TOCSY experiment for, Piccioli and Poggi, **155**, 236

- sequential, of proteins, 2D triple-resonance NMR experiments for, Ding and Gronenborn, **156**, 262
- strategy for proteins with known structure, Hus, Prompers, and Brüschweiler, **157**, 119
- U-¹³C, ¹⁵N-labeled peptides and proteins, *soft*-triple-resonance solid-state NMR, Astrof and Griffin, **158**, 157
- Resonator
active resonator system for CW-ESR measurement operating at 700 MHz, Sato *et al.*, **159**, 161
- Response function
complex, of magnetic resonance spectrometers, Annino *et al.*, **157**, 74
- Restricted diffusion
and categories of coherence pathways for CPMG sequence, Song, **157**, 82
- effects on MR signal formation, Sukstanskii and Yablonskiy, **157**, 92
- mapping in (FA)₂PF₆: combined k-space q-space pulsed ESR imaging, Feintuch *et al.*, **157**, 69
- Restricted self-diffusion
of water in highly concentrated W/O emulsion, modulated gradient spin-echo NMR, Topgaard, Melander, and Söderman, **156**, 195
- RIACT
FAM–RIACT–FAM, signal enhancement in triple-quantum MAS NMR of spins-3/2 in solids, Madhu and Levitt, **155**, 150
- modified RIACT MQMAS NMR for $I = 3/2$ nuclei, efficient triple-quantum excitation in, Lim, Charpentier, and Pines, **154**, 196
- RIDE'n RIPT
ring down elimination in rapid imaging pulse trains, Woelk *et al.*, **159**, 207
- Ring-down elimination
in noise spectroscopy, Yang and Zax, **158**, 73
- in rapid imaging pulse trains—RIDE'n RIPT, Woelk *et al.*, **159**, 207
- Rotating-frame imaging
ring down elimination in rapid imaging pulse trains—RIDE'n RIPT, Woelk *et al.*, **159**, 207
- Rotating-frame relaxation
 $R_{1\rho}$ relaxation outside of fast-exchange limit, Trott and Palmer, **154**, 157
- Rotation
chemical shielding tensor orientation from two or one rotations of single crystal, Shekar, Ramamoorthy, and Wittebort, **155**, 257
- Rotational resonance
improved narrowband dipolar coupling for homonuclear distance measurements in rotating solids, Goobes and Vega, **154**, 236
- use in recovering proton dipolar interaction in resin-bound peptides under high-resolution MAS, Raya *et al.*, **157**, 43
- Rotor-encoded spinning-sideband patterns
heteronuclear MQ MAS NMR, half-integer quadrupolar and spin $I = 1/2$ nuclei, Lupulescu, Brown, and Spiess, **154**, 101
- Rubber
crosslink density in, visualization based on dipolar-correlation effect, contrast parameter for, Grinberg, Heidenreich, and Kuhn, **159**, 88
- natural, multipolar spin states, effects of crosslinking and strain, Wiesmath *et al.*, **154**, 60
- S**
- Sample restriction
using RF selective pulses, in high-resolution solid-state NMR, Charmont, Sakellariou, and Emsley, **54**, 136
- Satellite-transition MAS NMR
optimizing, Huguenard *et al.*, **156**, 131
- spin $I = 3/2, 5/2, 7/2,$ and $9/2$ nuclei, Ashbrook and Wimperis, **156**, 269
- Satellite transitions
half-integer quadrupolar nuclei, influencing for enhancement of MAS NMR spectra, Iuga and Kentgens, **158**, 65
- Saturation factor
changes, quantitation errors in MRS due to, optimized pulse parameters for reducing, Galbán and Spencer, **156**, 161
- Saturation transfer difference NMR
with ICS-NMR: complete relaxation and conformational exchange matrix, Jayalakshmi and Krishna, **155**, 106
- Saturation transfer effects
intermolecular, in reversibly forming ligand–receptor complexes, CORCEMA analysis, Jayalakshmi and Krishna, **155**, 106
- S-band EPR
PELDOR at S- and X-band frequencies and separation of exchange coupling from dipolar coupling, Weber *et al.*, **157**, 277
- Scalar coupling
³J_{HNH₂}, in ¹⁵N-labeled proteins, rapid determination with cosine-modulated HSQC, Petit, Vincent, and Zwahlen, **156**, 313
- dipolar, efficiency of homonuclear Hartmann-Hahn and COSY-type mixing sequences in presence of, Kramer and Glaser, **155**, 83
- Scaling laws
in NMR scattering, via dipolar fields, Brown, Sen, and Cory, **154**, 154
- Selective excitation
pulses, effect on localized SPAMM 1–1 sequence, Ikonomidou and Sergiadis, **157**, 218
- Selective pulse
adjustable broadband selective excitation with uniform phase, Cano, Smith, and Shaka, **155**, 131
- Selective refocusing
2D ¹H, enantiomeric excess measurements in weakly oriented chiral liquid crystal solvents using, Farjon *et al.*, **158**, 169
- Self-diffusion
modeling in multi-compartment systems with cylindrical geometry, van der Weerd *et al.*, **156**, 213
- restricted, of water in highly concentrated W/O emulsion, modulated gradient spin-echo NMR, Topgaard, Melander, and Söderman, **156**, 195
- Sensitivity
opposed-solenoid endovascular MRI probes, Crottet *et al.*, **159**, 219
- satellite-transition MAS NMR of spin $I = 3/2, 5/2, 7/2,$ and $9/2$ nuclei, Ashbrook and Wimperis, **156**, 269
- Sensitivity enhancement
MAS spectra, influencing satellite transitions of half-integer quadrupolar nuclei for, Iuga and Kentgens, **158**, 65
- membrane peptide orientation from ¹H-detected ²H NMR, Yamaguchi and Hong, **155**, 244
- in multiple-quantum NMR experiments with CPMG detection, Lim *et al.*, **157**, 160
- structural measurements by solid-state NMR, via pulsed spin locking, Petkova and Tycko, **155**, 293
- time-optimal coherence-order-selective transfer of in-phase coherence in heteronuclear IS spin systems, Reiss, Khaneja, and Glaser, **154**, 192
- via polarization transfer, in NMR of laser-polarized noble gases, Goodson, **155**, 157
- Separation of undistorted powder patterns by effortless recoupling (SUPER)
2D separation of undistorted CSA powder patterns in MAS NMR, Liu, Mao, and Schmidt-Rohr, **155**, 15
- Sequential assignment
proteins, 2D triple-resonance NMR experiments for, Ding and Gronenborn, **156**, 262
- Shielding
inductive, of NMR phase noise, Sigmund *et al.*, **159**, 190
- Shielding tensor
orientation, from two or one rotations of single crystal, Shekar, Ramamoorthy, and Wittebort, **155**, 257
- Shimming
gradient, variable temperature, method for, Evans, Morris, and Davis, **154**, 325

- Sidebands
REDOR, relative CSA–dipolar orientation from, O'Connor and Schaefer, **154**, 46
- Side chains
¹³C resonances, correlation with backbone amide resonances in perdeuterated proteins, Löhner and Rüterjans, **156**, 10
- Signal enhancement
in 3QMAS NMR of spins-3/2 in solids, Madhu and Levitt, **155**, 150
in 5QMAS spectra of spin-5/2 quadrupolar nuclei, Goldbourt and Vega, **154**, 280
out-of-phase rapid-passage EPR for samples with long electron spin relaxation times, Harbridge *et al.*, **156**, 41
- Signal intensity
enhanced, by out-of-phase rapid-passage EPR for samples with long electron spin relaxation times, Harbridge *et al.*, **156**, 41
EPR, frequency dependence
248 MHz to 1.4 GHz, Rinard *et al.*, **154**, 80
250 MHz to 9.1 GHz, Rinard *et al.*, **156**, 113
- Signal-to-noise ratio
prediction, picoliter ¹H NMR, Minard and Wind, **154**, 336
- Signal processing
biomedical, improved Lanczos algorithms for blackbox MRS data quantitation, Laudadio *et al.*, **157**, 292
two-dimensional prolate spheroidal wave functions for MRI, Yang *et al.*, **158**, 43
- SIMMOL
specification and visualization of anisotropic interaction tensors in polypeptides and numerical simulations in biological solid-state NMR, Bak *et al.*, **154**, 28
- Simulated annealing
conjoined rigid body/torsion angle, determination of relative orientation of two domains from dipolar couplings, Clore and Bewley, **154**, 329
- Single crystal
chemical shielding tensor orientation from two or one rotations of, Shekar, Ramamoorthy, and Wittebort, **155**, 257
metal surfaces, *T*₁ relaxation of ¹²⁹Xe on, Stahl *et al.*, **159**, 1
- Size distribution
emulsion droplets under flow, determination using flow-compensating NMR-PFG techniques, Johns and Gladden, **154**, 142
- Skeletal muscle
intra- and extramyocellular lipids, signal separation in proton MR spectra by magnetic field distribution, Steidle *et al.*, **154**, 228
metabolite quantitation errors due to saturation factor changes in MRS, optimized pulse parameters for reducing, Galbán and Spencer, **156**, 161
porcine, water characteristics, changes during growth, *in vitro* low-field NMR relaxation study, Bertram *et al.*, **157**, 267
- Skin
kinetic measurements using EPR imaging with modulated field gradient, Herrling, Fuchs, and Groth, **154**, 6
- Slice thickness
and GAMMA simulations of stray-field responses, Kinchesch *et al.*, **154**, 252
- SLICING
trilinear, for 2D noniterative *T*₂ decomposition of low-field NMR relaxation data, Pedersen, Bro, and Engelsen, **157**, 141
- Small animals
multinuclear imaging at 3T, automatic tuned MRI RF coil for, Muftuler *et al.*, **155**, 39
whole-body imaging, double end-cap birdcage coil for, Gulsen, Muftuler, and Nalcioğlu, **156**, 309
- Small molecules
in amorphous matrices, large-angle jumps of, 2D exchange NMR analysis, Medick, Vogel, and Rössler, **159**, 126
- Smoothing
Euclidian distance-weighted, for quantitative MRI, Bonny and Renou, **159**, 183
- Solid-phase resins
resin dynamics and NMR line broadening of organic molecules grafted onto polystyrene resin, Lippens, Chessari, and Wieruszkeski, **156**, 242
- Solids
at low temperatures, multiple-quantum dynamics in linear chains and rings of, Fel'dman and Maximov, **157**, 106
polycrystalline, γ -irradiated, radicals with two methyl groups in, electron spin-lattice relaxation, Harbridge, Eaton, and Eaton, **159**, 195
rotating, improved narrowband dipolar coupling for homonuclear distance measurements in, Goobes and Vega, **154**, 236
spins-3/2 in, triple-quantum MAS NMR, signal enhancement, Madhu and Levitt, **155**, 150
- Solid-state NMR
biological, numerical simulations, and specification and visualization of anisotropic interaction tensors in polypeptides, Bak *et al.*, **154**, 28
cross-polarization, chemical-shift anisotropy and offset effects in, Shekar, Lee, and Ramamoorthy, **157**, 223
double-quantum, ¹³C spin pairs coupled to ¹⁴N, Hughes *et al.*, **159**, 25
²H, transmembrane peptides in bicelles, and hydrophobic mismatch and peptide sequence, Whiles *et al.*, **158**, 149
high-resolution, sample restriction using RF selective pulses in, Charmont, Sakellariou, and Emsley, **54**, 136
soft-triple-resonance, for assignments of U-¹³C, ¹⁵N-labeled peptides and proteins, Astrof and Griffin, **158**, 157
structural measurements, sensitivity enhancement by pulsed spin locking, Petkova and Tycko, **155**, 293
- Solid-state NMR imaging
spin interaction filter in, Casieri and De Luca, **159**, 63
- Solution-state NMR
spectrometers, temperature accuracy and temperature gradients in, Loening and Keeler, **159**, 55
- Solvent suppression
improvement in NMR spectra using wavelet transforms: WAVEMAT, Günther, Ludwig, and Rüterjans, **156**, 19
NMR difference probe applicable to, Macnaughtan *et al.*, **156**, 97
- Solvent-suppression pulses
simple scheme for design of, Prost *et al.*, **159**, 77
- SPAMM sequence
rotational approach to localized SPAMM 1–1 tagging, Ikonomidou and Sergiadis, **157**, 218
- Spatial correlations
magnetic field and velocity field, and PFG NMR measurements of flow through porous media, Lebon and Leblond, **159**, 13
- Spectral assignments
¹³C NMR, for nematic liquid crystals, by 2D chemical-shift γ -encoding NMR, Nishiyama, Kubo, and Terao, **158**, 60
- Spectral decomposition
EPR spectra, quantification of lateral lipid domain properties of erythrocyte ghosts using, Arsov, Schara, and Štrancar, **157**, 52
- Spectral resolution
enhanced, in immobilized peptides and proteins, by combining chemical shift sum and difference spectroscopy, Luca and Baldus, **159**, 243
- Spectrometers
design and implementation of protection system for, Laurent and Guéron, **156**, 295
solution-state NMR, temperature accuracy and temperature gradients in, Loening and Keeler, **159**, 55
- Spin diffusion
true ¹H, 2D ¹³C–¹³C MAS NMR correlation spectroscopy with, de Boer *et al.*, **157**, 286

- Spin echoes
 multiple, in heterogeneous systems: origin of observed dips, Alessandri, Capuani, and Maraviglia, **156**, 72
 non-CPMG fast spin echo with full signal, Le Roux, **155**, 278
 signal behavior in presence of magnetic field gradients, theoretical framework for analysis, Sukstanskii and Yablonskiy, **157**, 92
- Spin Hamiltonian tensors
 low-symmetry, analysis: Fe³⁺ in natrolite, Vinokurov *et al.*, **155**, 57
- Spin interaction filter
 in solid-state NMR imaging, Casieri and De Luca, **159**, 63
- Spin labels
 immobilization and orientation, structural determination, Sale *et al.*, **156**, 104
 microenvironment: polarity contributions to hyperfine splittings of nitroxides, Marsh, **157**, 114
 multispin labeling and MAS NMR 2D dipolar correlation, model for aggregated chlorophyll *a*/H₂O from, van Rossum *et al.*, **155**, 1
- Spin–lattice relaxation
 amide proton, in perdeuterated protein, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181
 effects on metabolite quantitation: optimized pulse parameters for error reduction due to saturation factor changes in MRS, Galbán and Spencer, **156**, 161
 with *T*₁ measurements, deconvolution of compartmental water diffusion coefficients in yeast cell suspensions, Silva *et al.*, **156**, 52
*T*₂–*T*₂ correlation spectra using fast 2D Laplace inversion, Song *et al.*, **154**, 261
¹²⁹Xe on metal single-crystal surfaces, Stahl *et al.*, **159**, 1
- Spin locking
 NMR, proton magnetization under frequency-switched Lee–Goldburg pulse sequence, Fu, Tian, and Cross, **154**, 130
- Spinning frequency
 low, aqueous samples in presence of magnetic susceptibility gradients at, evaluation of MAS techniques in, Hu and Wind, **159**, 93
- Spinning sidebands
 rotor-encoded heteronuclear MQ MAS NMR of half-integer quadrupolar and spin *I* = 1/2 nuclei, Lupulescu, Brown, and Spiess, **154**, 101
- Spin pairs
¹³C, coupled to ¹⁴N, double-quantum solid-state NMR, Hughes *et al.*, **159**, 25
S = 1, *I* = 1/2, internuclear distance determination by REAPDOR NMR, Hughes *et al.*, **156**, 230
- Spin polarization
 high-capacity production of >65% spin-polarized ¹²⁹Xe for NMR spectroscopy and MRI, Zook, Adhyaru, and Bowers, **159**, 175
- Spin polarization-induced Nuclear Overhauser effect
 NMR of laser-polarized noble gases, review, Goodson, **155**, 157
- Spin polarization transfer
 to surface nuclei, hyperpolarized ¹²⁹Xe and CPMAS for, Smith *et al.*, **159**, 111
- Spin relaxation
 Curie, in methyl groups, cross-correlation effects involving, Madhu, Mandal, and Müller, **155**, 29
- Spins
 rare, at natural abundance, molecules in nematic solvent with, selective detection of ¹H NMR spectra, Emsley *et al.*, **154**, 303
S > 1/2 centers in orientationally disordered systems, ESEEM and HYSCORE spectra, Benetis, Dave, and Goldfarb, **158**, 126
 spin *I* = 1/2 nuclei, rotor-encoded heteronuclear MQ MAS NMR, Lupulescu, Brown, and Spiess, **154**, 101
 spin *I* = 3/2, 5/2, 7/2, and 9/2 nuclei, satellite-transition MAS NMR, Ashbrook and Wimperis, **156**, 269
 spins-3/2 in solids, triple-quantum MAS NMR, signal enhancement, Madhu and Levitt, **155**, 150
- Spin–spin relaxation
 amide proton, in perdeuterated protein, effects of dissolved oxygen, Ulmer, Campbell, and Boyd, **157**, 181
 enhanced, in porous media due to internal field gradients, Dunn, **156**, 171
 low-field ¹H NMR *in vitro*, porcine muscle, study of water characteristics during growth, Bertram *et al.*, **157**, 267
 rapid unique curve resolution of low-field NMR relaxation data, Pedersen, Bro, and Engelsen, **157**, 141
*T*₂–*T*₂ correlation spectra obtained using fast 2D Laplace inversion, Song *et al.*, **154**, 261
- Spin states
 multipolar, excited in strongly inhomogeneous magnetic fields, NMR, Wiesmath *et al.*, **154**, 60
- Spin systems
 evolution, calculating: coherent excitation with phase-incremented pulses, Zhang and Gorenstein, **154**, 73
 heteronuclear *IS*, time-optimal coherence-order-selective transfer of in-phase coherence in, Reiss, Khaneja, and Glaser, **154**, 192
- SPRITE
 water content profiles with 1D centric SPRITE acquisition, Mastikhin *et al.*, **156**, 122
- Strain
 crosslinked natural rubber, effects on multipolar spin states, Wiesmath *et al.*, **154**, 60
- Stray-field gradients
 and pulsed gradients, combined, multidimensional imaging using, Godward *et al.*, **155**, 92
- Stray-field responses
 GAMMA simulations: slice thickness and pulse calibration, Kinchesch *et al.*, **154**, 252
- Structural model
 3D, solid self-assembled chlorophyll *a*/H₂O, van Rossum *et al.*, **155**, 1
- Structure
 from ¹³C–¹⁴N systems, using ¹³C FIREMAT experiments, Strohmeier, Alderman, and Grant, **155**, 263
 long length scale determination in xenon gas diffusion NMR of porous media, Mair *et al.*, **156**, 202
 measurement by solid-state NMR, sensitivity enhancement via pulsed spin locking, Petkova and Tycko, **155**, 293
 of spin label immobilization and orientation, determination, Sale *et al.*, **156**, 104
- Subband decomposition
 subband ARMA modeling approach to high-resolution NMR, Tomczak and Djermoune, **158**, 86
- Subspace time-domain algorithm
 for automated NMR spectral normalization, Lemmerling *et al.*, **157**, 190
- Sudden passage
 condition for adiabatic passage in Earth's-field NMR technique, Melton and Pollak, **158**, 15
- Superconducting magnet
 achievement of 920-MHz high-resolution NMR, Hashi *et al.*, **156**, 318
- Surface nuclei
 spin polarization transfer to, hyperpolarized ¹²⁹Xe and CPMAS for, Smith *et al.*, **159**, 111
- Suspensions
 cell and organelle, aeration and mixing during NMR experiments, device for, Bourne and Kuchel, **159**, 158
 yeast-cell, deconvolution of compartmental water diffusion coefficients, combined *T*₁ and *D* measurements, Silva *et al.*, **156**, 52
- Switched angle spinning
 two-dimensional NMR correlations of liquid crystals using, Havlin, Park, and Pines, **157**, 163
- T
- T*₁ (see Spin–lattice relaxation)
*T*₂ (see Spin–spin relaxation)
- Tagging
 of magnetization, using tandem of two adiabatic DANTE inversion sequences, Tsekos, Garwood, and Ugurbil, **156**, 187

- Temperature
 accuracy and gradients, in solution-state NMR spectrometers, Loening and Keeler, **159**, 55
 induced changes in EPR spectrum of magnetic center in kaolin, Vasyukov *et al.*, **154**, 15
- Temperature coefficients
 amide proton, location of hydrogen bonds in human ubiquitin using, Cierpicki *et al.*, **157**, 178
- Temperature gradients
 and temperature accuracy, in solution-state NMR spectrometers, Loening and Keeler, **159**, 55
- Tensor field approximation
 continuous, of discrete DT-MRI data, Pajevic, Aldroubi, and Bassar, **154**, 85
- Thermopiles
 Peltier, thermostatic control system, microprocessor control for portable NMR relaxometer, Anisimov, Teplov, and Silkin, **154**, 176
- Thermostatic control system
 Peltier thermopiles, microprocessor control for portable NMR relaxometer, Anisimov, Teplov, and Silkin, **154**, 176
- Three-dimensional field mapping
 registered, quantification of B_0 homogeneity variation with head pitch by, Tyszka and Mamelak, **159**, 213
- Three-dimensional NMR
 HSQC-HSQMBC, for increasing resolution of long-range proton-carbon correlation experiments, Uhrin, **159**, 145
- Time-domain algorithm
 frequency-selective quantification, Romano *et al.*, **155**, 226
 subspace-based, for automated NMR spectral normalization, Lemmerling *et al.*, **157**, 190
- Time evolution data
 EPR dipolar, direct conversion to distance distributions, Jeschke *et al.*, **155**, 72
- Tissue
 microstructure/architecture, continuous tensor field approximation of DT-MRI data, Pajevic, Aldroubi, and Bassar, **154**, 85
- TOCSY
 tailored HCCH-TOCSY experiment for resonance assignment in proximity of paramagnetic center, Piccioli and Poggi, **155**, 236
- Torsion angle
 conjoined rigid body/torsion angle simulated annealing for orientation of covalently linked domains from dipolar couplings, Clore and Bewley, **154**, 329
- Ψ , NMR determination in α -helical peptides and proteins, Ladizhansky, Veshtort, and Griffin, **154**, 317
- sensitivity of DQ solid state NMR spectra of ^{13}C spin pairs coupled to ^{14}N , Hughes *et al.*, **159**, 25
- TOSS
 evaluation at low spinning rates in aqueous samples and in presence of magnetic susceptibility gradients, Hu and Wind, **159**, 93
- Transition-selective pulses
 implementation of conditional phase-shift gate for quantum information processing by NMR using, Das, Mahesh, and Kumar, **159**, 46
- Transmembrane peptides
 in bicelles, study methods: consequences of hydrophobic mismatch and peptide sequence, Whiles *et al.*, **158**, 149
- Triple-quantum excitation
 efficient, in modified RIACT MQMAS NMR for $I = 3/2$ nuclei, Lim, Charpentier, and Pines, **154**, 196
- Triple-quantum resonance
 in two-level NMR system, line-narrowing effect in, Hatanaka and Tabuchi, **155**, 119
- Triple-resonance NMR
 2D experiments for sequential resonance assignments of proteins, Ding and Gronenborn, **156**, 262
soft-, solid-state, for assignments of U- ^{13}C , ^{15}N -labeled peptides and proteins, Astrof and Griffin, **158**, 157
- TROSY
 evaluation and optimization of coherence transfer in high-molecular-weight systems, Permi and Annala, **155**, 123
- Two-dimensional distribution functions
 diffusion and relaxation, quantitative measurement in grossly inhomogeneous fields, Hürlimann and Venkataraman, **157**, 31
- Two-dimensional D - J -DOSY
 resolving diffusion coefficients in mixtures, Lucas, Otto, and Larive, **156**, 138
- Two-dimensional electrophoretic NMR
 constant-time, Li and He, **156**, 181
- Two-dimensional exchange NMR
 amorphous matrices analyzed by, large-angle jumps of small molecules in, Medick, Vogel, and Rössler, **159**, 126
- Two-dimensional NMR
 ^{13}C - ^{13}C MAS NMR correlation spectroscopy with mixing by true ^1H spin diffusion, de Boer *et al.*, **157**, 286
 correlations of liquid crystals using switched angle spinning in, Havlin, Park, and Pines, **157**, 163
 high-resolution spectra in matched B_0 and B_1 field gradients, Heise *et al.*, **156**, 146
- Two-dimensional triple-resonance NMR
 for sequential resonance assignments of proteins, Ding and Gronenborn, **156**, 262
- Two-phase flow
 spectrally resolved velocity exchange spectroscopy, Khrapitchev *et al.*, **159**, 36

U

- Ubiquitin
 hydrogen bonds in, and temperature coefficients of amide protons, Cierpicki *et al.*, **157**, 178
- Ultra-high magnetic field
 long-range intermolecular distance restraints: 2D ^{13}C - ^{13}C MAS NMR correlation spectroscopy, de Boer *et al.*, **157**, 286
- Ultralow fields
 laser-polarized ^{129}Xe NMR and MRI at, Wong-Foy *et al.*, **157**, 235
- Uniform phase excitation
 adjustable broadband selective excitation with uniform phase, Cano, Smith, and Shaka, **155**, 131

V

- Variable temperature gradient shimming
 method for, Evans, Morris, and Davis, **154**, 325
- Vectorial analysis
 magnetic resonance spectrometers: complex response function, Annino *et al.*, **157**, 74
- Velocimetry
 NMR, of falling liquid films, Heine *et al.*, **154**, 311
- Velocity exchange spectroscopy
 spectrally resolved, of two-phase flow, Khrapitchev *et al.*, **159**, 36
- Velocity field
 and magnetic field, spatial correlations, and PFG NMR measurements of flow through porous media, Lebon and Leblond, **159**, 13
- Vibrations
 mechanical, magnetic resonance detection using radiofrequency field gradient, Baril *et al.*, **154**, 22
- Visualization
 anisotropic interaction tensors in polypeptides, and numerical simulations in biological solid-state NMR, Bak *et al.*, **154**, 28
 anisotropic interaction tensors in polypeptides: SIMMOL, Bak *et al.*, **154**, 28

Void-space imaging

materials and organisms: principles and MR applications of laser-polarized noble gases, Goodson, **155**, 157

W

Water

anisotropic diffusion in macroscopically oriented lipid bilayers, pulsed magnetic field gradient NMR, Wästerby, Orådd, and Lindblom, **157**, 156

mobility and distribution in porcine muscle, changes during growth, *in vitro* low-field NMR relaxation study, Bertram *et al.*, **157**, 267

restricted self-diffusion in highly concentrated W/O emulsion, modulated gradient spin-echo NMR, Topgaard, Melander, and Söderman, **156**, 195

Water content

profiles, with 1D centric SPRITE acquisition, Mastikhin *et al.*, **156**, 122

Water diffusion coefficients

compartmentalized, deconvolution in yeast cell suspensions, combined T_1 and D measurements, Silva *et al.*, **156**, 52

Water-in-oil emulsion

highly concentrated, restricted self-diffusion of water in, modulated gradient spin-echo NMR, Topgaard, Melander, and Söderman, **156**, 195

Water suppression

improvement in NMR spectra using wavelet transforms: WAVEMAT, Günther, Ludwig, and Rüterjans, **156**, 19

localized ^1H MRS water suppression without: frequency demodulation by modulus signal selection, Serrai *et al.*, **154**, 53

Wavelet transforms

improved solvent suppression in NMR spectra using: WAVEMAT, Günther, Ludwig, and Rüterjans, **156**, 19

WAVEWAT

improved solvent suppression in NMR spectra using wavelet transforms, Günther, Ludwig, and Rüterjans, **156**, 19

Whole-body imaging

small animal, double end-cap birdcage coil for, Gulsen, Muftuler, and Nalcioglu, **156**, 309

X

X-band EPR

PELDOR at S- and X-band frequencies and separation of exchange coupling from dipolar coupling, Weber *et al.*, **157**, 277

 ^{129}Xe

dissolved in ethanol, flowing, gradient echo imaging: theory and phantom studies, Johansson *et al.*, **159**, 69

gas diffusion NMR, of model porous media, narrow pulse approximation and long length scale determination in, Mair *et al.*, **156**, 202

>65% spin-polarized, high-capacity production for NMR spectroscopy and imaging, Zook, Adhyaru, and Bowers, **159**, 175

hyperpolarized, and CPMAS, for spin polarization transfer to surface nuclei, Smith *et al.*, **159**, 111

laser-polarized ^{129}Xe NMR and MRI at ultralow magnetic fields, Wong-Foy *et al.*, **157**, 235

liquid, polarization, rapid precise probe for measurement, Verhulst *et al.*, **155**, 145

T_1 relaxation on metal single-crystal surfaces, Stahl *et al.*, **159**, 1

Xenon gas

laser-polarized, in molecules, materials, and organisms, NMR, review, Goodson, **155**, 157

Y

Yeast-cell suspension

deconvolution of compartmental water diffusion coefficients, combined T_1 and D measurements, Silva *et al.*, **156**, 52

Z

Zeeman limit

closed-form expressions for level-averaged ESR times outside of, application to paramagnetic NMR relaxation, Sharp, **154**, 269

Zeolites

H-ZSM-5, Na-ZSM-5, and Na-A, NO adsorption/desorption behavior, EPR, Rudolf, Böhlmann, and Pöppel, **155**, 45