

Corrigendum

Corrigendum to “Residue-specific NH exchange rates studied by NMR diffusion experiments” [J. Magn. Reson. 187 (2007) 97–104]

Torsten Brand ^a, Eurico J. Cabrita ^b, Gareth A. Morris ^c, Robert Günther ^d,
Hans-Jörg Hofmann ^d, Stefan Berger ^{a,*}

^a *Institute of Analytical Chemistry, University of Leipzig, Linnéstraße 3, Leipzig 04103, Germany*

^b *REQUIMTE/CQFB, Dep. de Química, Faculdade de Ciências e Tecnologia, Univ. Nova de Lisboa, 2829-516 Caparica, Portugal*

^c *School of Chemistry, University of Manchester, Oxford Road, Manchester M13 9PL, UK*

^d *Institute of Biochemistry, University of Leipzig, Brüderstraße 34, Leipzig 04103, Germany*

The authors apologize for having unfortunately omitted the reference to:

- [31a] Hadj B. Seba, Pierre Thureau, Bernard Ancian, André Thévand, Combined use of Overhauser spectroscopy and NMR diffusion experiments for mapping the hydration structure of nucleosides: structure and dynamics of uridine in water, *Magn. Reson. Chem.* 44 (2006) 1109–1117.

DOI of original article: [10.1016/j.jmr.2007.03.021](https://doi.org/10.1016/j.jmr.2007.03.021)

* Corresponding author. Fax: +49 341 97 36115.

E-mail address: stberger@rz.uni-leipzig.de (S. Berger).