



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Journal of Magnetic Resonance

journal homepage: www.elsevier.com/locate/jmr



Corrigendum

Corrigendum to “Theoretical prediction and experimental measurement of the field dependence of the apparent transverse relaxation of hyperpolarized noble gases in lungs” [J. Magn. Res. 192 (2008) 85–91]

Juan Parra-Robles^a, William Dominguez Viqueira^a, Xiaojun Xu^a, Alexei Ouriadov^a, Giles E. Santyr^{a,b,c,*}

^a *Imaging Research Laboratories, Robarts Research Institute, P.O. Box 5015, 100 Perth Drive, London, Ont., Canada N6A 5K8*

^b *Department of Medical Biophysics, The University of Western Ontario, London, Ont., Canada N6A 5C1*

^c *Department of Diagnostic Radiology and Nuclear Medicine, The University of Western Ontario, London, Ont., Canada N6A 5A5*

On page 86, in the second full paragraph, the third sentence contains the erroneous statement that “the model fails to demonstrate the substantial differences in ADC measured between normal and emphysematous lungs at the acinar level.”

The correct entire third sentence should read: “This model accounts for the anisotropy of diffusion in the lungs and allows

the estimation of the average radii of the alveolar ducts and two orientation-dependent diffusion coefficients (longitudinal and transverse ADC). Their results demonstrated substantial differences between normal and emphysematous lungs at the acinar level.”

DOI of original article: [10.1016/j.jmr.2008.02.009](https://doi.org/10.1016/j.jmr.2008.02.009).

* Corresponding author. Fax: +1 519 663 3900.

E-mail address: gsantyr@imaging.robarts.ca (G.E. Santyr).