



DNP enhanced frequency-selective TEDOR experiments in bacteriorhodopsin

Vikram S. Bajaj^{a,1}, Melody L. Mak-Jurkauskas^{a,b}, Marina Belenky^b, Judith Herzfeld^b, Robert G. Griffin^{a,*}

^a Francis Bitter Magnet Laboratory, Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, United States

^b Department of Chemistry, Brandeis University, Waltham, MA 02454, United States

ARTICLE INFO

Article history:

Received 29 June 2009

Revised 27 August 2009

Available online 9 September 2009

Keywords:

Heteronuclear distance measurement

Chemical shift assignments

TEDOR

REDOR

DNP

Dynamic nuclear polarization

Bacteriorhodopsin

Magic angle spinning

ABSTRACT

We describe a new approach to multiple ^{13}C – ^{15}N distance measurements in uniformly labeled solids, frequency-selective (FS) TEDOR. The method shares features with FS-REDOR and ZF- and BASE-TEDOR, which also provide quantitative ^{15}N – ^{13}C spectral assignments and distance measurements in $[\text{U-}^{13}\text{C},^{15}\text{N}]$ samples. To demonstrate the validity of the FS-TEDOR sequence, we measured distances in $[\text{U-}^{13}\text{C},^{15}\text{N}]$ -asparagine which are in good agreement with other methods. In addition, we integrate high frequency dynamic nuclear polarization (DNP) into the experimental protocol and use FS-TEDOR to record a resolved correlation spectrum of the Arg- $^{13}\text{C}_\gamma$ – $^{15}\text{N}_\epsilon$ region in $[\text{U-}^{13}\text{C},^{15}\text{N}]$ -bacteriorhodopsin. We resolve six of the seven cross-peaks expected based on the primary sequence of this membrane protein.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

Most heteronuclear dipolar recoupling sequences produce Hamiltonians that are simple in comparison with their homonuclear counterparts in that they do not contain flip-flop terms; therefore, these recoupling experiments are not subject to the phenomenon of dipolar truncation [2]. Accordingly, successful approaches to heteronuclear distance measurements exploit this simplicity and are usually based on REDOR [3] and its variants such as TEDOR [4–6]. However, even though dipolar truncation is absent, there remain complications when distance measurements in uniformly ^{13}C – ^{15}N labeled systems are involved. In particular, evolution under the homonuclear ^{13}C – ^{13}C J-couplings imposes an overall modulation on the buildup or dephasing during the recoupling period. J-couplings can also generate anti-phase coherences, which in turn give rise to phase-twisted line shapes. In order to circumvent these limitations, Jaroniec et al. introduced J-decoupled [7] and frequency-selective REDOR [8] experiments in which the detrimental effects of J-coupling are removed through the application of frequency-selective pulses. These effects were also considered by Mehta and Schaefer [9] in the development of the RDX version of REDOR. Another logical extension of these techniques has been

the 3D ZF- and BASE-TEDOR experiments [1,10], in which multiple ^{13}C – ^{15}N distances are measured simultaneously in a broadband or a band selective manner in the context of a ^{13}C – ^{15}N correlation experiment. In these cases, the effects of J-coupling are removed through the application of a coherence filter or frequency-selective ^{13}C pulse, respectively. This family of sequences has been successfully employed in the *de novo* structural determination of a peptide [11], an amyloid fibril [12], and quantitative distance measurements in a membrane protein via MAS NMR [13], and is therefore an important part of the repertoire of experiments for determining structures of proteins with MAS NMR.

However, even where the recoupled dipolar Hamiltonian for individual spin pair interactions commutes, strong dipolar couplings can compromise the sensitivity of the experiment. For example, in the TEDOR experiment, the intensity corresponding to a single ^{15}N site is distributed over all ^{13}C – ^{15}N cross-peaks and is a function of the dipolar couplings, their mutual orientations, homonuclear J-couplings, and the mixing time. Thus, in the practically important case of distance measurements between backbone ^{15}N 's and ^{13}C 's of an amino acid side chain, the presence of strong $^{13}\text{C}_\beta$ – $^{15}\text{N}_\alpha$ couplings (2.5 Å) in the TEDOR dynamics reduces the maximum intensity of polarization transfer due to weaker couplings (~3.3 Å) to 1–3% of the initial intensity [12,14]. This sometimes compromises the application of broadband techniques to membrane proteins and other systems in which sensitivity is limited. This effect is also particularly acute in amino acids whose side chains contain nitrogen (e.g. asparagine, glutamine lysine, arginine), as a single ^{13}C site may be subject to several strong couplings

* Corresponding author. Fax: +1 617 253 5405.

E-mail address: rgg@mit.edu (R.G. Griffin).

¹ Present addresses: Department of Chemistry, University of California, Berkeley, United States; Materials Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, United States.

that dominate the spin dynamics [15]. However, when the backbone or side chain ^{15}N 's are resolved, then a ^{15}N -frequency-selective analog of the 3D TEDOR experiment can simplify the spin dynamics and allow high sensitivity measurement of weak couplings.

2. Results and discussion

These deficiencies and opportunities have stimulated us to develop the frequency-selective (FS) TEDOR experiment illustrated in Fig. 1 that shares many features with the FS-REDOR and BASE and ZF-TEDOR methods discussed above. Following ramped, ^1H - ^{13}C cross polarization, we apply a REDOR train to generate ^{13}C - ^{15}N anti-phase coherence which evolves under the heteronuclear dipolar couplings. During the excitation period, a frequency-selective pulse is applied on the ^{15}N channel, such that only those ^{15}N nuclei within the bandwidth contribute appreciably to the spin dynamics during the REDOR mixing period. Following the excitation period, optional frequency labeling with the ^{15}N chemical shift occurs during t_1 . A second, frequency-selective REDOR period converts the resulting coherence into observable magnetization on the ^{13}C channel, which is detected during t_2 . A delay, τ , is necessary to ensure that the time between REDOR periods is an integer number of rotor cycles during which the signal is transferred to the ^{13}C 's. Finally, ^{13}C - ^{13}C J-couplings give rise to undesired zero-quantum and double-quantum coherences that, after the reconversion period, result in severe anti-phase distortions to the line shape and spurious cross-peaks in 2D spectra. Two variable-length coherence filters (z-filters) are applied to dephase these undesired coherences, resulting in undistorted and purely absorptive spectra. TPPM and CW decoupling is used where appropriate [16]. In addition, for the experiments on the membrane protein bacteriorhodopsin (bR), we enhance the overall signal sensitivity of the experiments by integrating dynamic nuclear polarization (DNP) into the protocol, as illustrated in Fig. 1. In particular, we suspend the bR sample in 60% glycerol/ H_2O that forms a stable glass and serves to disperse the biradical polarizing agent TOTAPOL [17]. Irradiation with 250 GHz microwaves [18] leads to a polarization enhancement of ~ 43 in the signal intensities.

As an initial test of FS-TEDOR, all heteronuclear distances in $[\text{U-}^{13}\text{C}, ^{15}\text{N}]$ -asparagine were measured in two experiments, with a selective pulse applied at the frequency of the backbone ($^{-15}\text{NH}_3^+$) or side chain ^{15}N ($^{-15}\text{NH}_2$). The results are summarized in Fig. 2 and Table 1. All data were recorded on a 500 MHz (^1H) home-built spectrometer (Cambridge Instruments) using a Varian

triple-resonance MAS probe and in a sample of asparagine diluted to 10% in its natural abundance analog to avoid the confounding effects of intermolecular couplings. The data were fit using a multi-spin simulation performed with the program SPINEVOLUTION [19]. Our FS-TEDOR distance is in good agreement with the distance measured by frequency-selective REDOR [8] which is a true two-spin experiment and with the neutron diffraction structure. We note that the FS-REDOR and FS-TEDOR will likely be more accurate than SPECIFIC-CP [20] experiments for distance measurements. A careful comparison with results from PAIN-CP [21] is in progress.

To demonstrate the utility of FS-TEDOR as an approach for spectral editing in a more challenging and interesting case, we used the FS-TEDOR sequence, together with DNP, to record 2D ^{13}C - ^{15}N spectra of the seven arginine side chains in the membrane protein bacteriorhodopsin (bR), a light-driven ion pump that has been studied extensively by magnetic resonance [22–29]. Owing to their hydrophilicity, many of the lysine and arginine side chains in proteins are located at or near the surface. In contrast, those buried in the hydrophobic interior are frequently functionally important. bR contains a number of such examples, including Lys-216, to which the retinal chromophore is attached via a Schiff base linkage, and Arg-82, which is part of the complex counterion and is perturbed following deprotonation of the Schiff Base, a key step in the $\text{L} \rightarrow \text{M}_0$ photostate transition of bR [30]. In this and other systems, approaches to spectral editing based on selective cross polarization [20,31,32], magnetization preparation or selective dephasing [13], or both in combination [33], have been used to achieve similar ends. The optimal sequence to use in each case depends on the spin topology (e.g. ^{13}C coupled to multiple ^{15}N s or the reverse) and desired selectivity.

In this case, addition of DNP, which consists of CW irradiation of the EPR spectrum of TOTAPOL with 250 GHz microwaves to the FS-TEDOR experiment, enhances the signal sensitivity by a factor of ~ 43 . Together with the lower temperatures, we obtain a factor of ~ 90 increase in signal intensity over our previous experiments on bR at 200 K. Fig. 3 shows the 1D ^{15}N spectrum of U- $^{13}\text{C}, ^{15}\text{N}$ -bR.

We recorded correlation spectra, focusing on the $^{13}\text{C}_\gamma$ - $^{15}\text{N}_\epsilon$ region of arginine. Details of instrumentation [18,34–36], MAS probes [37], polarizing agents [38,39], and sample preparation for applications of DNP to bacteriorhodopsin [33,40,41] are published elsewhere and not reviewed here. We empirically and separately optimized the TPPM decoupling [16] during the inter-pulse periods of the REDOR sequence, the selective pulse, and detection, to minimize the transverse relaxation (T_2), avoid interference effects between the decoupling and multiple pulse sequence, and to maximize the sensitivity. We chose a Gaussian refocusing pulse




Fig. 1. Pulse sequence for ^{15}N -frequency-selective TEDOR transfer. Only ^{15}N spins within the bandwidth of the selective pulses contribute to the spin dynamics under REDOR. The sequence is described in the text. The REDOR pulses are phase cycled according to the XY-8 scheme and the overall phase cycle is documented elsewhere [1]. CW microwave irradiation was employed in the version of the experiment used to acquire the DNP enhanced spectrum.




Fig. 2. FS-TEDOR experiment in 10% U-[^{13}C , ^{15}N]-Asparagine. We employed 100 kHz TPPM decoupling during REDOR and other periods, a 1 ms Gaussian refocusing pulse (switched between the NH_3 and NH_2 sites), 50 kHz REDOR pulses. $\omega_r 2\pi = 10$ kHz.

Table 1

Heteronuclear distances measured with FS-TEDOR experiment as applied to [U- ^{13}C , ^{15}N]-Asparagine.

	FS-TEDOR (Å)	FS-REDOR (Å)	Neutron diffraction (Å)
N-C $^\alpha$	1.50	1.50	1.49
N $^{\delta 2}$ -C $^\alpha$	3.56	3.58	3.75
N-C $^\beta$	2.46	2.49	2.48
N $^{\delta 2}$ -C $^\beta$	2.41	2.44	2.42




Fig. 3. 1D DNP enhanced ^{15}N spectra of bR illustrating the excellent S/N available with DNP that permits observation of the ^{15}N signal of the single Schiff Base (K216). The Arg resonances irradiated with the selective pulse are labeled and lie at ~ 80 ppm.

whose length was optimized in the range $1.98 \text{ ms} \pm 0.265 \text{ ms}$ [42] (see Fig. 4) and whose frequency was centered on the arginine side chain region (~ 90 ppm for $^{15}\text{N}_\epsilon$) (Fig. 4). A pulse of this length sufficiently suppresses the amide backbone resonances, which otherwise would compromise the dynamic range of the experiment and obscure the arginine cross-peaks, which are an order of magnitude less intense. Further, we selected a TEDOR mixing time of 4 ms, in which polarization transfer over two-bond distances (e.g. ~ 2.4 – 2.5 Å between Arg- $^{13}\text{C}_\gamma$ - $^{15}\text{N}_\epsilon$) in U- ^{13}C , ^{15}N compounds is maximized.




Fig. 4. 2D ^{15}N - ^{13}C selective correlation experiment focused on the Arg- $^{13}\text{C}_\gamma$ - $^{15}\text{N}_\epsilon$ of bR. The sequence described in Fig. 1 was employed with a 1.98 ms Gaussian refocusing pulse, and TPPM decoupling optimized to reduce transverse dephasing. The REDOR mixing time of 4 ms was chosen to favor short distances (~ 2.4 – 2.5 Å). The spinning frequency was 7.576 kHz, and all DNP experiments were conducted in dark-adapted bacteriorhodopsin (90 K), as described elsewhere.

The spectrum in Fig. 4 reveals ^{15}N - ^{13}C correlations which we assign to six of the seven expected Arg- $^{13}\text{C}_\gamma$ - $^{15}\text{N}_\epsilon$ cross-peaks. Our assignment is based on our choice of parameters (a selective pulse centered at 90 ppm and a 4 ms TEDOR recoupling time) which favor Arg- $^{13}\text{C}_\gamma$ - $^{15}\text{N}_\epsilon$ magnetization transfer over Arg- $^{13}\text{C}_\delta$ - $^{15}\text{N}_{\epsilon, \eta 1, \eta 2}$, due to the mixing time and the presence of multiple $^{13}\text{C}_\delta$ - ^{15}N couplings. In addition, the chemical shifts observed (see Table 2) agree with typical values for Arg- $^{15}\text{N}_\epsilon$ and Arg- $^{13}\text{C}_\gamma$, indicated by amino acid chemical shift statistics for proteins [43,44] which show Arg- $^{15}\text{N}_\epsilon$ chemical shifts as 90 ± 19 ppm vs. 75 ± 14 ppm and 75 ± 16 ppm for Arg- $^{15}\text{N}_{\eta 1}$ and Arg- $^{15}\text{N}_{\eta 2}$, respectively, and 27.2 ± 2.0 ppm for Arg- $^{13}\text{C}_\gamma$ vs. 43.2 ± 1.8 ppm for Arg- $^{13}\text{C}_\delta$. However, while both the ^{15}N and ^{13}C chemical shifts

Table 2

Chemical shifts and intensities of crosspeaks in the Arg-¹³C_γ-¹⁵N_ε region of 2D fs-TEDOR spectra bR (Fig. 4). The chemical shifts and intensities were fit using SPARKY [45] following processing in NMRPipe [46].

Peak	¹⁵ N chemical shift (ppm)	¹³ C chemical shift (ppm)	Peak volume (arbitrary units)
A	82.9	30.9	2.9
B	87.3	31.7	2.7
C ^a	87.8	31.0	2.7
D	89.9	30.7	3.1
E	89.4	29.3	3.0
F	87.6	27.9	3.5
G	82.1	26.0	3.1

^a Cross-peak C is not resolved from D.

are only consistent with the Arg-¹³C_γ-¹⁵N_ε assignments, their eventual validation will require site-specific assignment of all the Arg side chains, which we did not attempt here.

3. Conclusions

In summary, we have demonstrated that the FS-TEDOR experiment, a close analog of the frequency-selective REDOR experiment, can be used quantitatively and qualitatively for ¹⁵N-¹³C correlation spectroscopy in crystalline solids and membrane proteins. Using DNP, we have recorded resolved Arg-¹³C_γ-¹⁵N_ε correlation spectra, tentatively identifying six of the seven expected resonances. Similar methods might be contemplated in lieu of specific isotopic labeling or suppression [47] to simplify the spin dynamics, or in contexts where quantitative distance information is required from a resolved ¹⁵N site, such as in distance measurements from the Schiff base of bR and other rhodopsins, or in peptides containing ¹⁵N-rich side chains [15,48]. The decision to use FS-TEDOR over other methods depends on the topology of the spin system under consideration, the degree of selectivity desired, and the dephasing of transverse magnetization during the evolution times of the experiment.

Acknowledgments

The authors acknowledge invaluable technical assistance from Dr. D. Ruben, A. Thakkar, J. Bryant, Dr. J. Sirigiri, Dr. R.J. Temkin, and discussions with Professor C.P. Jaroniec. V.S.B. acknowledges the receipt of a PGS Fellowship from the Natural Sciences and Engineering Research Council of Canada. This research was supported by the National Institutes of Biomedical Imaging and Bioengineering through Grants EB-001960, EB-002804, EB-001035, and EB-002026.

References

- [1] C.P. Jaroniec, C. Filip, R.G. Griffin, 3D TEDOR NMR experiments for the simultaneous measurement of multiple carbon–nitrogen distances in uniformly C-13, N-15-labeled solids, *J. Am. Chem. Soc.* 124 (2002) 10728–10742.
- [2] M.J. Bayro, M. Huber, R. Ramachandran, T.C. Davenport, B.H. Meier, M. Ernst, R.G. Griffin, Dipolar truncation in magic-angle spinning NMR recoupling experiments, *J. Chem. Phys.* 130 (2009) 114506.
- [3] T. Gullion, J. Schaefer, Rotational-echo double-resonance NMR, *J. Magn. Reson.* 81 (1989) 196–200.
- [4] A.W. Hing, S. Vega, J. Schaefer, Transfer echo double resonance NMR, *J. Magn. Reson.* 96 (1992) 205–209.
- [5] Y. Li, R.J. Appleyard, W.A. Shuttlesworth, J.N.S. Evans, Time-resolved solid-state REDOR NMR measurements of 5-enolpyruvylshikimate 3-phosphate synthase, *J. Am. Chem. Soc.* 116 (1994) 10799–10800.
- [6] C.A. Michal, L.W. Jelinski, REDOR 3D: heteronuclear distance measurements in uniformly labeled and natural abundance solids, *J. Am. Chem. Soc.* 119 (1997) 9059–9060.

- [7] C.P. Jaroniec, B.A. Tounge, C.M. Rienstra, J. Herzfeld, R.G. Griffin, Measurement of C-13–N-15 distances in uniformly C-13 labeled biomolecules: J-decoupled REDOR, *J. Am. Chem. Soc.* 121 (1999) 10237–10238.
- [8] C.P. Jaroniec, B.A. Tounge, J. Herzfeld, R.G. Griffin, Frequency selective heteronuclear dipolar recoupling in rotating solids: Accurate C-13–N-15 distance measurements in uniformly C-13, N-15-labeled peptides, *J. Am. Chem. Soc.* 123 (2001) 3507–3519.
- [9] A.K. Mehta, J. Schaefer, Rotational-echo double resonance of uniformly labeled C-13 clusters, *J. Magn. Reson.* 163 (2003) 188–191.
- [10] J.J. Helmus, P.S. Nadaud, N. Hofer, C.P. Jaroniec, Determination of methyl C-13–N-15 dipolar couplings in peptides and proteins by three-dimensional and four-dimensional magic-angle spinning solid-state NMR spectroscopy, *J. Chem. Phys.* 128 (2008) 052314.
- [11] C.M. Rienstra, L. Tucker-Kellogg, C.P. Jaroniec, M. Hohwy, B. Reif, M.T. McMahon, B. Tidor, T. Lozano-Perez, R.G. Griffin, De novo determination of peptide structure with solid-state magic-angle spinning NMR spectroscopy, *Proc. Natl. Acad. Sci. USA* 99 (2002) 10260–10265.
- [12] C.P. Jaroniec, C.E. MacPhee, V.S. Bajaj, M.T. McMahon, C.M. Dobson, R.G. Griffin, High resolution molecular structure of a peptide in an amyloid fibril determined by MAS NMR spectroscopy, *Proc. Natl. Acad. Sci. USA* 101 (2004) 711–716.
- [13] C.P. Jaroniec, J.C. Lansing, B.A. Tounge, M. Belenky, J. Herzfeld, R.G. Griffin, Measurement of dipolar couplings in a uniformly C-13, N-15- labeled membrane protein: distances between the Schiff base and aspartic acids in the active site of bacteriorhodopsin, *J. Am. Chem. Soc.* 123 (2001) 12929–12930.
- [14] C.P. Jaroniec, C.E. MacPhee, N.S. Astrof, C.M. Dobson, R.G. Griffin, Molecular conformation of a peptide fragment of transthyretin in an amyloid fibril, *Proc. Natl. Acad. Sci. USA* 99 (2002) 16748–16753.
- [15] P.C.A. van der Wel, J.R. Lewandowski, R.G. Griffin, Solid-state NMR study of amyloid nanocrystals and fibrils formed by the peptide GNNQQNY from yeast prion protein Sup35p, *J. Am. Chem. Soc.* 129 (2007) 5117–5130.
- [16] A.E. Bennett, C.M. Rienstra, M. Auger, K.V. Lakshmi, R.G. Griffin, Heteronuclear decoupling in rotating solids, *J. Chem. Phys.* 103 (1995) 6951.
- [17] C. Song, K.-N. Hu, C.-G. Joo, T.M. Swager, R.G. Griffin, TOTAPOL – A biradical polarizing agent for dynamic nuclear polarization experiments in aqueous media, *J. Am. Chem. Soc.* 128 (2006) 11385–11390.
- [18] V.S. Bajaj, M.K. Hornstein, K.E. Kreisler, J.R. Sirigiri, P.P. Woskov, M.L. Mak-Jurkaskas, J. Herzfeld, R.J. Temkin, R.G. Griffin, 250 GHz CW gyrotron oscillator for dynamic nuclear polarization in biological solid state NMR, *J. Magn. Reson.* 189 (2007) 251–279.
- [19] M. Veshkort, R.G. Griffin, SPINEVOLUTION: a powerful tool for simulations of solid and liquid state NMR experiments, *J. Magn. Reson.* 178 (2006) 248–282.
- [20] M.A. Baldus, A.T. Petkova, J. Herzfeld, R.G. Griffin, Cross polarization in the tilted frame: assignment and spectral simplification in heteronuclear spin systems, *Mol. Phys.* 95 (1998) 1197–1207.
- [21] J.R. Lewandowski, G.d. Paepe, R.G. Griffin, Proton assisted insensitive nuclei cross polarization, *J. Am. Chem. Soc.* 129 (2007) 728–729.
- [22] J.K. Lanyi, B. Schobert, Mechanism of proton transport in bacteriorhodopsin from crystallographic structures of the K, L, M1, M2, and M2' intermediates of the photocycle, *J. Mol. Biol.* 328 (2003) 439–450.
- [23] H. Luecke, B. Schobert, H.T. Richter, J.P. Cartailler, J.K. Lanyi, Structure of bacteriorhodopsin at 1.55 angstrom resolution, *J. Mol. Biol.* 291 (1999) 899–911.
- [24] J. Lugtenburg, M. Muradinszweykowska, C. Heeremans, J.A. Pardo, G.S. Harbison, J. Herzfeld, R.G. Griffin, S.O. Smith, R.A. Mathies, Mechanism for the opsin shift of retinals absorption in bacteriorhodopsin, *J. Am. Chem. Soc.* 108 (1986) 3104–3105.
- [25] J. Herzfeld, J.G. Hu, M.E. Hatcher, M. Belenky, P. Verdegem, J. Lugtenburg, R.G. Griffin, NMR evidence for an electrostatically controlled, torsion-based switch in light-driven proton transport by bacteriorhodopsin, *Biophys. J.* 80 (2001) 85.
- [26] J. Herzfeld, J.G. Hu, B.Q. Sun, A. Petkova, M.E. Hatcher, M. Bizounok, R.G. Griffin, Solid-state NMR studies of energy transduction in bacteriorhodopsin, *Biophys. J.* 74 (1998) A137.
- [27] J. Herzfeld, J.C. Lansing, Magnetic resonance studies of the bacteriorhodopsin pump cycle, *Annu. Rev. Biophys. Biomol. Struct.* 31 (2002) 73–95.
- [28] J. Herzfeld, B. Tounge, NMR probes of vectoriality in the proton-motive photocycle of bacteriorhodopsin: evidence for an 'electrostatic steering' mechanism, *Biochim. Biophys. Acta Bioenerg.* 1460 (2000) 95–105.
- [29] W. Stoeckenius, R.H. Lozier, W. Niederberger, Photoreactions of bacteriorhodopsin, *Biophys. Struct. Mech.* 3 (1977) 65–68.
- [30] A.T. Petkova, J.G.G. Hu, M. Bizounok, M. Simpson, R.G. Griffin, J. Herzfeld, Arginine activity in the proton-motive photocycle of bacteriorhodopsin: solid-state NMR studies of the wild-type and D85N proteins, *Biochemistry* 38 (1999) 1562–1572.
- [31] S. Jehle, K. Rehbein, A. Diehl, B.J. van Rossum, Amino-acid selective experiments on uniformly C-13 and N-15 labeled proteins by MAS NMR: filtering of lysines and arginines, *J. Magn. Reson.* 183 (2006) 324–328.
- [32] A.T. Petkova, M. Baldus, M. Belenky, M. Hong, R.G. Griffin, J. Herzfeld, Backbone and sidechain assignment strategies for multiply labeled membrane peptides and proteins, *J. Magn. Reson.* 160 (2003) 1–12.
- [33] V.S. Bajaj, M.L. Mak-Jurkaskas, M. Belenky, J. Herzfeld, R.G. Griffin, Functional and shunt states of bacteriorhodopsin resolved by 250 GHz dynamic nuclear polarization-enhanced solid state NMR, *Proc. Natl. Acad. Sci. USA* 106 (2009) 9244–9249.

- [34] V.S. Bajaj, C.T. Farrar, M.K. Hornstein, I. Mastovsky, J. Vieregg, J. Bryant, B. Elena, K.E. Kreisler, R.J. Temkin, R.G. Griffin, Dynamic nuclear polarization at 9T using a novel 250 GHz gyrotron microwave source, *J. Magn. Reson.* 160 (2003) 85–90.
- [35] M.K. Hornstein, V.S. Bajaj, R.G. Griffin, R.J. Temkin, Continuous-wave operation of a 460-GHz second harmonic gyrotron oscillator, *IEEE Trans. Plasma Sci.* 34 (2006) 524–533.
- [36] P.W. Woskov, V.S. Bajaj, M.K. Hornstein, R.J. Temkin, R.G. Griffin, Corrugated waveguide and directional coupler for CW 250 GHz gyrotron DNP experiments, *IEEE Trans. Microw. Theory Tech.* 53 (2005) 1863–1869.
- [37] A.B. Barnes, M.L. Mak-Jurkauskas, Y. Matsuki, V.S. Bajaj, P.C.A.v.d. Wel, R. DeRocher, J. Bryant, J.R. Sirigiri, R.J. Temkin, J. Lugtenburg, J. Herzfeld, R.G. Griffin, Cryogenic sample exchange NMR probe for magic angle spinning dynamic nuclear polarization, *J. Magn. Reson.* 198 (2009) 261–270.
- [38] K.-N. Hu, V.S. Bajaj, M.M. Rosay, R.G. Griffin, High frequency dynamic nuclear polarization using mixtures of TEMPO and trityl radicals, *J. Chem. Phys.* 126 (2007) 044512.
- [39] K.N. Hu, H.H. Yu, T.M. Swager, R.G. Griffin, Dynamic nuclear polarization with biradicals, *J. Am. Chem. Soc.* 126 (2004) 10844–10845.
- [40] M.L. Mak-Jurkauskas, V.S. Bajaj, M.K. Hornstein, M. Belenky, R.G. Griffin, J. Herzfeld, Gradual winding of the bacteriorhodopsin chromophore in the first half of its ion-motive photocycle: a dynamic nuclear polarization-enhanced solid state NMR study, *Proc. Natl. Acad. Sci. USA* 105 (2008) 883–888.
- [41] M. Rosay, A.C. Zeri, N.S. Astrof, S.J. Opella, J. Herzfeld, R.G. Griffin, Sensitivity-enhanced NMR of biological solids: dynamic nuclear polarization of Y21M fd bacteriophage and purple membrane, *J. Am. Chem. Soc.* 123 (2001) 1010–1011.
- [42] Y. Li, B.J. Wylie, C.M. Rienstra, Selective refocusing pulses in magic-angle spinning NMR: characterization and applications to multi-dimensional protein spectroscopy, *J. Magn. Reson.* 179 (2006) 206–216.
- [43] J.L. Markley, E.L. Ulrich, H.M. Berman, K. Henrick, H. Nakamura, H. Akutsu, BioMagResBank (BMRB) as a partner in the Worldwide Protein Data Bank (wwPDB): new policies affecting biomolecular NMR depositions, *J. Biomol. NMR* 40 (2008) 153–155.
- [44] E.L. Ulrich, H. Akutsu, J.F. Doreleijers, Y. Harano, Y.E. Ioannidis, J. Lin, M. Livny, S. Mading, D. Maziuk, Z. Miller, E. Nakatani, C.F. Schulte, D.E. Tolmie, R.K. Wenger, H.Y. Yao, J.L. Markley, BioMagResBank, *Nucleic Acids Res.* 36 (2008) D402–D408.
- [45] T.D. Goddard, D.G. Kneller, SPARKY Version 3.115, University of California, San Francisco, 2008.
- [46] F. Delaglio, S. Grzesiek, G.W. Vuister, G. Zhu, J. Pfeifer, A. Bax, NMRPipe: a multidimensional spectral processing system based on UNIX pipes, *J. Biomol. NMR* 6 (1995) 277–293.
- [47] G.W. Vuister, S.J. Kim, C. Wu, A. Bax, 2D and 3D NMR study of phenylalanine residues in proteins by reserve isotopic labeling, *J. Am. Chem. Soc.* 116 (1994) 9206–9210.
- [48] P.C.A. van der Wel, K.N. Hu, J. Lewandowski, R.G. Griffin, Dynamic nuclear polarization of amyloidogenic peptide nanocrystals: GNNQQNY, a core segment of the yeast prion protein Sup35p, *J. Am. Chem. Soc.* 128 (2006) 10840–10846.