



## Corrigendum

## Corrigendum to “A conformational study of N-acetyl glucosamine derivatives utilizing residual dipolar couplings” [J. Magn. Reson. 212 (2011) 174–185]

Markus Kramer, Erich Kleinpeter \*

University of Potsdam, Department of Chemistry, Karl-Liebknecht-Str. 24-25, 14476 Potsdam/Golm, Germany

In Fig. 4 the denomination of the regions for the substances 1, 2, 3 and 5 has to be changed to that shown in the corrected figure. Due to the symmetric aglycones of that substances the conformers

represented by the regions “A” and “B” as well as “C” and “D” are equal.

Consequently, in Table 5 in the column “Region” for substance 5 “A/B” changes to “A”.

**Table 5**  
Dihedral angles of the conformers obtained from the MD simulations.


Substance	Region <sup>a</sup>	$\Phi$ (°)	$\Psi$ (°)
1	A	−87.3	172.9
	B	63.8	172.1
2	A	−45.2	−164.3
	B	74.4	−34.4
3	A	−56.1	173.7
	B	43.8	−164.6
4	A	−51.1	−0.6
	B	−68.4	164.1
	C	66.1	−37.9
	D	58.0	171.0
5	A	−79.2	173.4
6	A	−62.1	81.4
	B	−65.5	−105.1

<sup>a</sup> The letters in this column correspond to the letters shown in the  $\Phi/\Psi$ -plots.

DOI of original article: [10.1016/j.jmr.2011.06.029](https://doi.org/10.1016/j.jmr.2011.06.029).

\* Corresponding author. Fax: +49 331 977 5064.

E-mail address: [ekleinp@uni-potsdam.de](mailto:ekleinp@uni-potsdam.de) (E. Kleinpeter).

Fig. 4.  $\Phi/\Psi$ -plots from the MD simulations.