

# Supersymmetric solutions of PT-/non-PT-symmetric and non-Hermitian central potentials via Hamiltonian hierarchy method

Gholamreza Faridfathi, Ramazan Sever\*, and Metin Aktaş

Department of Physics, Middle East Technical University, 06531 Ankara, Turkey  
E-mail: sever@metu.edu.tr

Received 6 December 2004; revised 16 February 2005

The supersymmetric solutions of PT-/non-PT-symmetric and non-Hermitian deformed Morse and Pöschl-Teller potentials are obtained by solving the Schrödinger equation. The Hamiltonian hierarchy method is used to get the real energy eigenvalues and corresponding eigenfunctions.

**KEY WORDS:** supersymmetric quantum mechanics, Hamiltonian hierarchy method, Morse potential, Pöschl-Teller potential

**PACS:** 05.20.-y, 05.30.-d, 05.70. Ce; 03.65.-w

## 1. Introduction

PT-symmetric quantum systems have generated much interest in recent years [1]. About 10 years ago, Basis suggested that the eigenvalue spectrum of complex-valued Hamiltonians is real and positive. Bender and Boettcher claimed that this result is due to PT-symmetry where P and T are the parity and time reversal operators, respectively. It is neither a necessary nor a sufficient condition for a Hamiltonian to have real spectrum. In particular, the spectrum of the Hamiltonian is real if PT-symmetry is not spontaneously broken. Thus, the property of exactness guarantees the real eigenvalues. Recently, Mostafazadeh introduced another concept for a class of PT-invariant Hamiltonians called  $\eta$ -pseudo-Hermiticity [2]. In fact, Hamiltonians of this type satisfy the transformation  $\eta\hat{H}\eta^{-1} = \hat{H}^\dagger$  [3]. Moreover, completeness and orthonormality conditions for the eigenstates of such potentials are proposed [4]. Various techniques have been applied in the study of PT-invariant potentials such as variational methods [5], numerical approaches [6], Fourier analysis [7], semi-classical estimates [8], quantum field theory [9] and Lie group theoretical approaches [10–13]. In the applications, a generalization of

\*Corresponding author.

the symmetry concept is encountered in the supersymmetric quantum mechanics (SUSYQM) [14]. A variety of PT-symmetric examples can be found using the SUSYQM techniques [15–20]. Furthermore, one can get more examples of the PT-symmetric and non-PT-symmetric and also non-Hermitian potential cases such as oscillator type potentials [21], flat, step and double square-well like potentials within the framework of SUSYQM [22,23], exponential type screened potentials [24], quasi/conditionally exactly solvable ones [25], complex Hénon–Heiles potentials [26], periodic isospectral potentials [27] and some others [28,29].

The aim of the present work is to calculate the energy eigenvalues and the corresponding eigenfunctions of the deformed Morse and Pöschl-Teller potentials using the Hamiltonian hierarchy method [30] within the framework of the PT-SUSYQM. This method is also known as the factorization method introduced by Schrödinger [31] and later developed by Infeld and Hull [32]. It is useful to obtain the energy spectra for different potentials in non-relativistic quantum mechanics [33–35].

The organization of this paper is as follows: In Section 2 we introduce a brief review of the Hamiltonian hierarchy method. In Section 3 and 4 we present the supersymmetric solutions of PT-symmetric and Hermitian/non-Hermitian forms of the well-known potentials by using this method. We discuss the results in Section 5.

## 2. Hamiltonian hierarchy method

The radial Schrödinger equation for some specific potential energies can be solved analytically only for the states with zero angular momentum [36,37]. However, in supersymmetric quantum mechanics, one can get exact results with the hierarchy problem by using effective potentials for non-zero angular momentum states. In the framework of SUSYQM, this method provides an eigenvalue spectra for adjacent members of supersymmetric partner Hamiltonians. These Hamiltonians share the same eigenvalue spectra except for the missing ground state.

In the application of this method, we first look for an effective potential similar to the original specific potential and inspired by the SUSYQM to propose a superpotential, namely  $W_{(l+1)}(x)$ , as an ansatz, where  $(l + 1)$  denotes the partner number with  $l = 0, 1, 2, \dots$ . Substituting the proposed superpotential into the Riccati equation,

$$V_{(l+1)}(x) - E_{(l+1)}^0 = W_{(l+1)}^2(x) - \frac{dW_{(l+1)}(x)}{dx}, \quad (1)$$

the  $(l + 1)$ th member of the Hamiltonian hierarchy can be obtained. Taking into account the shape invariance requirement [14], the bound-state energies can be obtained through equation (1), and the corresponding eigenfunctions by means of

$$\Psi_{(l+1)}(x) = N \exp\left(-\int^x W_{(l+1)}(x')dx'\right). \tag{2}$$

### 3. Generalized Morse potential

The generalized Morse potential is given by [24]

$$V(x) = V_1e^{-2\alpha x} - V_2e^{-\alpha x}. \tag{3}$$

To apply the Hamiltonian hierarchy method, we shall take the coefficients as  $V_1 = V$ , and  $V_2/V_1 = q$ . Inspired by the SUSYQM, we propose an ansatz for the superpotential

$$W_{(l+1)}(x) = -\lambda e^{-\alpha x} + \left(\lambda q - \frac{2l + 1}{2}\right), \tag{4}$$

where,  $\lambda^2 = 2mV/a^2\hbar^2$ , and  $(2l + 1)$  denotes the partner number with  $l = 0, 1, 2, \dots$ , and the parameter  $m$  is the reduced mass of a diatomic molecule. The superpotential chosen in equation (4) leads to the  $(l + 1)$ th member of the Hamiltonian hierarchy through the Riccati equation as,

$$V_{(l+1)}(x) - E_{(l+1)}^0 = W_{(l+1)}^2(x) - \frac{1}{\alpha} \frac{dW_{(l+1)}(x)}{dx}, \tag{5}$$

which yields,

$$V_{(l+1)}(x) = \lambda^2(e^{-2\alpha x} - qe^{-\alpha x}) + 2l\lambda e^{-\alpha x}. \tag{6}$$

Now, using the shape invariance requirement, the energy eigenvalues for any  $n$ th state become,

$$E_{(l+1)}^n = -\left(\lambda q - \frac{2l + n + 1}{2}\right)^2, \quad n = 0, 1, 2, \dots \tag{7}$$

The corresponding eigenfunctions are obtained through equation (2) as

$$\Psi_{(l+1)}^{n=0}(x) = N \exp\left[-\frac{\lambda}{\alpha}e^{-\alpha x} - \left(\lambda q - \frac{2l + 1}{2}\right)x\right], \tag{8}$$

where,  $N$  is the normalization constant.

3.1. *Non-PT symmetric and non-Hermitian Morse case*

We define the potential parameters in equation (3) as  $V_1 = (A + iB)^2$ ,  $V_2 = (2C + 1)(A + iB)$ , and  $\alpha = 1$ .  $A$ ,  $B$  and  $C$  are real, and  $i = \sqrt{-1}$ . For simplicity we define  $A + iB = i\omega$ ,  $(A + iB)^2 = -\omega^2$ ,  $2C + 1 = K$ . Thus, the potential becomes

$$V(x) = -\frac{\omega^2}{K} \left[ K e^{-2x} - \frac{K^2}{i\omega} e^{-x} \right]. \tag{9}$$

To get the final compact form, we also define  $\frac{\omega^2}{K} = G$ , and  $\frac{K^2}{\omega} = t$ , and  $GK = D$ , and also  $\frac{t}{K} = P$ . As a result, we get,

$$V(x) = -D \left[ e^{-2x} + iPe^{-x} \right]. \tag{10}$$

We propose an ansatz for the superpotential as,

$$W_{(l+1)}(x) = -i\lambda e^{-x} + \left( \lambda - \frac{2l + 1}{2} \right), \tag{11}$$

where  $\lambda^2 = \frac{2mD}{\hbar^2}$ . Consequently, according to the Hamiltonian hierarchy method, we get,

$$V_{(l+1)}(x) = -\lambda^2(e^{-2x} + 2ie^{-x}) + 2i\lambda e^{-x}, \tag{12}$$

By substituting this into equation (1), the corresponding eigenvalues and eigenfunctions are obtained as

$$E_{(l+1)}^n = -\left( \lambda - \frac{n + 2l + 1}{2} \right)^2, \quad n = 0, 1, 2, \dots, \tag{13}$$

and

$$\Psi_{(l+1)}^{n-0}(x) = N \exp \left[ -i\lambda e^{-x} - \left( \lambda - \frac{2l + 1}{2} \right) x \right], \tag{14}$$

where,  $N$  is a normalization constant.

3.2. *The first type of PT-symmetric and non-Hermitian Morse case*

We take the coefficients of the generalized Morse potential as  $V_1 = (A + iB)^2$ ,  $V_2 = (2C + 1)(A + iB)$ , and  $\alpha = i$ . Following the same procedure, we propose an ansatz for the superpotential

$$W_{(l+1)}(x) = -\lambda e^{-ix} + \left( \lambda - \frac{2l + 1}{2} \right). \tag{15}$$

The Hamiltonian hierarchy method yields,

$$V_{(l+1)}(x) = \lambda^2(e^{-2ix} - e^{-ix}) + 2l\lambda e^{-ix}. \tag{16}$$

This form of potential gives the same eigenvalues as in equation (13).

### 3.3. The second type of PT-symmetric and non-Hermitian Morse case

Now, we take the parameters as  $V_1 = -\omega^2$ , and  $V_2 = D$ , and  $\alpha = i\alpha$  in equation (3), where  $V_1$  and  $V_2$  are real. For  $V_1 \implies 0$ , we get no real spectra for this kind of PT-symmetric Morse potentials. The superpotential can be proposed as,

$$W_{(l+1)}(x) = -e^{-i\alpha x} + \left(2l + 1 + \frac{D}{2\omega}\right). \tag{17}$$

By applying the Hamiltonian hierarchy method, we get the potential

$$V_{(l+1)}(x) = e^{-2i\alpha x} - 2 \left[ (2l + 1) + \frac{D}{2\omega} + \frac{i\alpha}{2} \right] e^{-i\alpha x}, \tag{18}$$

and the corresponding eigenvalues for any  $n$ th state are,

$$E_{(l+1)}^n = - \left( 2l + n + 1 + \frac{D}{2\omega} \right)^2. \tag{19}$$

## 4. Pöschl-Teller potential

The Pöschl-Teller potential is given as,

$$V(x) = -4V_0 \frac{e^{-2\alpha x}}{(1 + qe^{-2\alpha x})^2}. \tag{20}$$

In the framework of the SUSYQM, the corresponding superpotential can be proposed as,

$$W_{(l+1)}(x) = -\frac{\hbar}{\sqrt{2m}} \frac{(l + 1)e^{-2\alpha x}}{(1 + qe^{-2\alpha x})^2} + \sqrt{\frac{m}{2}} \frac{e^2}{\hbar} \left[ \frac{1}{(l + 1)} - \frac{(l + 1)}{2} \beta \right], \tag{21}$$

where,  $\beta = \hbar^2/m e^2$ , and  $l = 0, 1, 2, \dots$ . By applying the Hamiltonian hierarchy method we get,

$$V_{(l+1)}(x) = \frac{\hbar^2}{2m} \frac{e^{-4\alpha x}}{(1 + qe^{-2\alpha x})^4} l(l + 1) - e^2 \frac{e^{-2\alpha x}}{(1 + qe^{-2\alpha x})^2} \left[ 1 - l(l + 1) \frac{\beta}{2} \right]. \tag{22}$$

As a result, the corresponding eigenvalues of this potential for any  $n$ th state are,

$$E_{(l+1)}^n = \frac{q^2 m e^4}{2 \hbar^2} \left[ \frac{1}{(n+l+1)} - \frac{(n+l+1)}{2} \beta \right]^2. \tag{23}$$

The corresponding eigenfunctions are,

$$\Psi_{(l+1)}^{n=0}(x) = N (1 + q e^{-2\alpha x})^{l+1} \exp \left\{ -\frac{m e^2}{\hbar^2} \left[ \frac{1}{l+1} - \frac{(l+1)}{2} \beta \right] x \right\}, \tag{24}$$

where,  $N$  is a normalization constant.

#### 4.1. Non-PT symmetric and non-Hermitian Pöschl-Teller cases

Here,  $V_0$  and  $q$  are complex parameters:  $V_0 = V_{0R} + i V_{0I}$  and  $q = q_R + i q_I$ , but  $\alpha$  is a real parameter. Although the potential is complex and the corresponding Hamiltonian is non-Hermitian and also non-PT-symmetric, there may be real spectra if and only if  $V_{0I} q_R = V_{0R} q_I$ . When both parameters  $V_0$ , and  $q$  are taken pure imaginary, the potential turns out to be,

$$V(x) = -4V_0 \frac{2q e^{-4\alpha x} + i(1 - q^2 e^{-4\alpha x})}{(1 + q^2 e^{-4\alpha x})^2}. \tag{25}$$

For simplicity, we use the notation  $V_0$  and  $q$  instead of  $V_{0I}$  and  $q_I$ . To obtain the energy eigenvalues, we propose the superpotential

$$W_{(l+1)}(x) = -\frac{\hbar}{\sqrt{2m}} \frac{(l+1)q e^{-4\alpha x}}{(1 + q^2 e^{-4\alpha x})^2} + \sqrt{\frac{m}{2}} \frac{e^2}{\hbar} \left[ \frac{1}{(l+1)} - \frac{(l+1)}{2} \beta \right]. \tag{26}$$

Therefore, substituting this equation into the Riccati equation, we get the same potential as in equation (22), and also the same energy eigenvalues as in equation (23).

#### 4.2. PT-symmetric and non-Hermitian Pöschl-Teller cases

We choose the parameters  $V_0$  and  $q$  as real, and also  $\alpha \implies i\alpha$  in equation (25). Here, we propose the superpotential similar to this potential as,

$$W_{(l+1)}(x) = -\frac{\hbar}{\sqrt{2m}} \frac{(l+1)q e^{-4i\alpha x}}{(1 + q^2 e^{-4i\alpha x})^2} + \sqrt{\frac{m}{2}} \frac{e^2}{\hbar} \left[ \frac{1}{(l+1)} - \frac{(l+1)}{2} \beta \right]. \tag{27}$$

By applying the same procedure, we get the energy eigenvalues as in equation (23).

## 5. Conclusions

We have applied the *PT*-symmetric formulation to solve the Schrödinger equation for more general Morse and Pöschl-Teller potentials. The Hamiltonian hierarchy method within the framework of the SUSYQM is used. We have obtained the energy eigenvalues and the corresponding eigenfunctions for different forms of these potentials. The energy spectrum of the *PT*-invariant complex-valued non-Hermitian potentials may be real or complex depending on the parameter values. We have imposed some restrictions on the potential parameters to get the real spectra in *PT*-symmetric, or more generally, in non-Hermitian cases. It is also pointed out that the superpotentials, and partners must satisfy the *PT*-symmetry condition. Finally, we have pointed out that our exact results of complexified general Morse and Pöschl-Teller potentials may increase the number of applications in the study of different quantum systems.

## References

- [1] C.M. Bender and S. Boettcher, *Phys. Rev. Lett.* 80 (1998) 5243; C.M. Bender, *J. Math. Phys.* 40 (1999) 2201; C.M. Bender, D.C. Brody and H.F. Jones, *Phys. Rev. Lett.* 89 (2002) 270401; C.M. Bender, P.N. Meisinger and Q. Wang, *J. Phys. A* 36 (2003) 6791; C.M. Bender, D.C. Brody and H.F. Jones, *Am. J. Phys.* 71 (2003) 1095; C.M. Bender, P.N. Meisinger and Q. Wang, *J. Phys. A* 36 (2003) 1029; C.M. Bender, M.V. Berry and A. Mandilara, *J. Phys. A* 35 (2002) L467.
- [2] A. Mostafazadeh, *J. Math. Phys.* 43 (2002) 205, *ibid* 43 (2002) 2814, *ibid* 43 (2002) 3944; *J. Phys. A* 36 (2003) 7081.
- [3] Z. Ahmed, *Phys. Lett. A* 290 (2001) 19, *ibid* 310 (2003) 139, *J. Phys. A* 36 (2003) 10325.
- [4] S. Weigert, *Phys. Rev. A* 68 (2003) 062111.
- [5] C.M. Bender, F. Cooper, P.N. Meisinger and M.V. Savage, *Phys. Lett. A* 259 (1999) 224.
- [6] C.M. Bender, G.V. Dunne and P.N. Meisinger, *Phys. Lett. A* 252 (1999) 272.
- [7] V. Buslaev and V. Grecchi, *J. Phys. A* 26 (1993) 5541.
- [8] E. Delabaere and F. Pham, *Phys. Lett. A* 250 (1998) 25, *ibid*, 250 (1998) 29.
- [9] C.M. Bender, K.A. Milton and V.M. Savage, *Phys. Rev. D* 62 (2000) 085001; C.M. Bender, S. Boettcher, H.F. Jones and P.N. Meisinger, *J. Math. Phys.* 42 (2001) 1960; C.M. Bender, S. Boettcher, H.F. Jones, P.N. Meisinger and M. Şimşek, *Phys. Lett. A* 291 (2001) 197; C.M. Bender, *Czech. J. Phys.* 54 (2004) 13.
- [10] B. Bagchi and C. Quesne, *Phys. Lett. A* 273 (2000) 285.
- [11] B. Bagchi and C. Quesne, *Phys. Lett. A* 300 (2002) 18.
- [12] G. Lévai, F. Cannata and A. Ventura, *J. Phys. A* 34 (2001) 839.
- [13] G. Lévai, F. Cannata and A. Ventura, *J. Phys. A* 35 (2002) 5041.
- [14] F. Cooper, A. Khare and U. Sukhatme, *Phys. Rep.* 251 (1995) 267, hep-th/0209062.
- [15] F. Cannata, G. Junker and J. Trost, *Phys. Lett. A* 246 (1998) 219.
- [16] B. Bagchi, S. Mallik and C. Quesne, *Mod. Phys. Lett. A* 17 (2002) 1651.
- [17] B. Bagchi and R. Roychoudhury, *J. Phys. A* 33, (2000) L1.
- [18] A. A. Adrianov, et al., *Int. J. Mod. Phys. A* 14 (1999) 2675.
- [19] S. Vincenzo and V. Alonso, *Phys. Lett. A* 298 (2002) 98.
- [20] J.S. Petrović, V. Milanović and Z. Ikonić, *Phys. Lett. A* 300 (2002) 595.

- [21] F.M. Fernandez, R. Guardiola, J. Ros and M. Znojil, *J. Phys. A* 31 (1998) 10105; G.A. Merzinescu, *J. Phys. A* 33 (2000) 4911; O.Mustafa and M. Znojil, *J. Phys. A* 35 (2002) 8929; M. Znojil, F. Gemperle and O. Mustafa, *J. Phys. A* 35 (2002) 5781; M. Znojil, *J. Phys. A* 35 (2002) 2341; M. Znojil, *Phys. Lett. A* 271 (2000) 327.
- [22] V. Milanović, Z. Ikonić, *Phys. Lett. A* 293 (2002) 29.
- [23] C.S. Jia, X.L. Zeug and L.T. Sun, *Phys. Lett. A* 294 (2002) 185; M. Znojil, *J. Phys. A* 36 (2003) 7639; M. Znojil, *J. Phys. A* 36 (2003) 7825; M. Znojil and G. Lévai, *Mod. Phys. Lett. A* 16 (2001) 2273; M. Znojil, *Phys. Lett. A* 285 (2001) 7, hep-th/0404213.
- [24] H. Taşeli, *J. Phys. A* 31 (1998) 779; M. Znojil, *Phys. Lett. A* 264 (1999) 108; Ö. Yeşiltaş, M. Şimşek, R. Sever and C. Tezcan, *Phys. Scripta* T67 (2003) 472; D.T. Barclay, R. Dutt, A. Gangopadhyaya, A. Khare, A. Pagnamenta, C.S. Jia, Y. Sun and Y. Li, *Phys. Lett. A* 305 (2002) 231; G. Lévai and M. Znojil, *J. Phys. A* 35 (2002) 8793.
- [25] C.M. Bender and S. Boettcher, *J. Phys. A* 31 (1998) L273; C.M. Bender, S. Boettcher, H.F. Jones and V.M. Savage, *J. Phys. A* 32 (1999) 6771, M. Znojil, *J. Phys. A* 33 (2000) 4203; A. Sinha, G. Lévai and P. Roy, *Phys. Lett. A* 322 (2004) 78.
- [26] C.M. Bender, G.V. Dunne, P.N. Meisinger and M. Şimşek, *Phys. Lett. A* 281 (2001) 311.
- [27] A. Khare and U. Sukhatme, *Phys. Lett. A* 324 (2004) 406.
- [28] G. Lévai and M. Znojil, *J. Phys. A* 33 (2000) 7165.
- [29] D. Parthasarathi, D. Parashar and R.S. Kaushal, *J. Phys. A* 37 (2004) 781.
- [30] B. Gönül, O. Özer, Y. Cañçelik and M. Koçak, *Phys. Lett. A* 275 (2000) 238.
- [31] E. Schrödinger, *Proc. R. Irish Acad. A* 46 (1940) 9; *ibid*, 46 (1940) 183; *ibid*, 47 (1941) 53.
- [32] L. Infeld and T.E. Hull, *Mod. Phys.* 23 (1951) 21.
- [33] M. Ioffe, A.I. Neelov, *J. Phys. A* 36 (2003) 2493.
- [34] V.M. Tkachuk and T.V. Fityo, *J. Phys. A* 34 (2001) 8673.
- [35] E.D. Pilho and R.M. Ricotta, *Phys. Lett. A* 269 (2000) 269.
- [36] C.S. Lam and Y.P. Varshni, *Phys. Rev. A* 4 (1971) 1875.
- [37] S. Flügge, *Practical Quantum Mechanics* (Springer, Berlin 1974).