

# Stability in generic mitochondrial models

Pete Donnell · Murad Banaji · Stephen Baigent

Received: 7 March 2008 / Accepted: 2 September 2008 / Published online: 23 September 2008  
© Springer Science+Business Media, LLC 2008

**Abstract** In this paper, we use a variety of mathematical techniques to explore existence, local stability, and global stability of equilibria in abstract models of mitochondrial metabolism. The class of models constructed is defined by the biochemical description of the system—an electron transport chain coupled to a process of charge translocation across a membrane. The conclusions are based on the reaction network structure, and we make minimal assumptions on the kinetics of the reactions involved. In the absence of charge translocation these models have previously been shown to behave in a very simple manner with a single, globally stable equilibrium. We show that with charge translocation the conclusion about a unique equilibrium remains true, but local and global stability do not necessarily follow. The length of the chains proves to be important: For short electron transport chains it is possible to make claims about local and global stability of the equilibrium which are no longer valid for longer chains. Some particular conditions which ensure stability of the equilibrium for chains of arbitrary length are presented.

**Keywords** Mitochondria · Electron transport · Stability · Logarithmic norms

## 1 Introduction

The processes of electron transport and oxidative phosphorylation in mitochondria are of vital biological importance, being central to cellular respiration and hence energy

---

Funded by an EPSRC/MRC grant to the MIAS IRC (Grant Ref: GR/N14248/01).

P. Donnell  
Department of Medical Physics and Bioengineering, University College London, Gower Street,  
London WC1E 6BT, UK

M. Banaji · S. Baigent (✉)  
Department of Mathematics, University College London, Gower Street, London WC1E 6BT, UK  
e-mail: s.baigent@ucl.ac.uk

production in most eukaryotic cells. More than 95% of oxygen consumption in the human body occurs in mitochondria [1]. The basic features of mitochondrial electron transport and oxidative phosphorylation are now well understood: Summaries can be found in textbooks such as [2] or [3], with more detail in [4]. Elucidation of the detailed mechanisms however, particularly those involving proton pumping, is still in progress [5].

Mitochondrial electron transport occurs via a series of coupled redox reactions in the mitochondrial inner membrane. After the initial reduction of a first electron donor (e.g. NADH or FADH<sub>2</sub> produced by glycolysis and the TCA cycle) electrons are transferred from substrate to substrate, finally being accepted by oxygen. During some of these electron transfers a second process takes place—protons are pumped across the mitochondrial inner membrane producing a proton gradient across this membrane. These protons then return down their gradient, either through leak channels or through a particular enzyme, ATP synthase, leading to the phosphorylation of ADP.

Generic models of electron transport chains were explored in [6], where the main emphasis was on the input–output response of such models. In the simplest case, where the proton gradient across the membrane was ignored, these models were found to have very simple behaviour—at all physically meaningful parameter values there was a single, globally stable, equilibrium. In [7], this result was shown using different mathematical techniques to generalise to the case of electron transfer networks with more general topology than a chain.


From energetic considerations we expect the build up of a proton gradient across the membrane to have an inhibitory effect on electron transport. Models which take this into account thus have extra negative feedback loops. Intuitively we might expect that such feedback should not be destabilising. While this proves true for short chains, we find, surprisingly, that for longer chains this is no longer the case.

Before discussing generic models, it is worth mentioning that there are several detailed models of electron transport and oxidative phosphorylation such as [8–11]. These ordinary differential equation models have been designed with numerical data in mind, and reflecting the complexity of the processes involved, the functional forms are quite involved. There is an interesting discussion of how to ascribe kinetics to electron transport systems in [12]. Our interest in mitochondria was originally inspired by analysis and simulation of some of these numerical models, but the approach here is quite different, and more akin to work in [6, 7, 13]—because only very weak kinetic assumptions are made, the generic model we construct could be instantiated in a great variety of numerical models and the claims we make are valid for all possible instances of the generic model.


## 2 The model

### 2.1 The basic reaction scheme


The basic reaction scheme will be summarised here. Assume that there are  $n$  substrates, each of which can exist in an oxidised state  $A_i$  and a reduced state  $B_i$  so that




Further, assume that protons can exist in two compartments—the mitochondrial matrix (where they are termed  $H_m^+$ ), and the intermembrane space (where they are termed  $H_e^+$ )—with the possibility of transfers of the form




We are interested in reactions which are in general the combination of three processes, a reduction, an oxidation, and the transport of some protons across the membrane. So for example, if substrate  $A_i$  is reduced to  $B_i$ ,  $B_j$  is oxidised to  $A_j$ , and  $p$  protons are pumped across the mitochondrial membrane we get the half reactions



which combine to give



We also allow the possibility that a reducing/oxidising agent may be external to the model giving reactions such as




A set of redox reactions of the forms 1 and 2 can be termed an electron transfer network. A particular case is an electron transfer chain where in general the  $i$ th substrate is able only to exchange electrons with the  $(i + 1)$ th and  $(i - 1)$ th substrates, if these exist. A first and a last substrate can exchange electrons with the outside, reflecting the initial reduction of NADH or FADH<sub>2</sub> and the final oxidation of cytochrome  $a_3$  by O<sub>2</sub>.

We assume there are  $n$  substrates, and hence  $n + 1$  redox reactions. The  $i$ th reaction has forward rate  $f_i$ , which may be positive or negative, potentially allowing reactions to be reversible. For  $i \leq n$ , the  $i$ th reaction involves substrate  $i$ , and for  $i \geq 2$ , the  $i$ th reaction involves substrate  $i - 1$ . We define  $p_i$  as the number of protons pumped across the mitochondrial membrane by the  $i$ th reaction. Assuming that the quantities  $p_i$  are constant discounts the possibility of “redox slip” [14], which does not appear to be very important in normal circumstances [15]. A quantity  $\psi$  can be defined so that transfer of a single proton across the membrane creates one unit of  $\psi$ .  $\psi$  can take any real value and is a strictly increasing function of the proton motive force, i.e. the electrical/chemical gradient against which protons are pumped across the membrane.

Finally, reflecting the combined effect of proton leak and ADP phosphorylation, there is a process with rate  $L$  representing the “decay” of  $\psi$ . When there is no gradient, no protons leak through the membrane, so that  $L(0) = 0$ . Further  $L$  is assumed to be strictly increasing in  $\psi$ . The structure of the model is illustrated in Fig. 1.

Because the total quantity—oxidised plus reduced—of any substrate in the chain is conserved, reduced forms of the substrates are not explicitly introduced. Instead,



**Fig. 1** A schematic representation of the reaction network. The quantities  $A_i$  and  $B_i$  refer to oxidised and reduced states of the substrates. The functions  $f_i$  define the forward rates of reaction of the  $n + 1$  coupled redox reactions. The quantity  $\psi$  represents the electrical and chemical gradient across the mitochondrial membrane, which has an inhibitory effect on any redox reactions which involve proton pumping

the concentration of  $A_i$  is referred to as  $x_i$ , and the total concentration of  $A_i + B_i$  is assumed constant at  $m_i$ . We arrive at a model of the form:

$$\left. \begin{aligned} \dot{x}_1 &= -f_1(x_1, \psi) + f_2(x_1, x_2, \psi) \\ \dot{x}_i &= -f_i(x_{i-1}, x_i, \psi) + f_{i+1}(x_i, x_{i+1}, \psi) \quad i = 2, \dots, n - 1 \\ \dot{x}_n &= -f_n(x_{n-1}, x_n, \psi) + f_{n+1}(x_n, \psi) \\ \dot{\psi} &= \sum_{i=1}^{n+1} p_i f_i - L(\psi) \end{aligned} \right\} \quad (3)$$

The phase space of this system is defined by the equations:

$$0 \leq x_i \leq m_i \quad i = 1, \dots, n \quad -\infty < \psi < \infty$$

and is hence  $n + 1$  dimensional, being the product of a closed  $n$ -dimensional box and the real line.

### 2.2 Assumptions

All the functions  $f_i$ , along with  $L$ , are assumed to be  $C^1$  (once differentiable in all their arguments with continuous derivatives). The following notation is used for the derivatives of the functions  $f_i$ :

$$f_{ij} \equiv \frac{\partial f_i}{\partial x_j}, \quad F_{ij} \equiv -f_{ij}, \quad f_{i\psi} \equiv \frac{\partial f_i}{\partial \psi}, \quad F_{i\psi} \equiv -f_{i\psi} \quad (4)$$

At finite substrate concentrations, all reaction rates are finite, so that at any fixed  $\psi$  each  $f_i$  is bounded on its domain of definition.

Since  $\psi$  represents a potential against which some of the reactions must do work, the following relations are obtained:

$$f_{i\psi} < 0 \text{ if } p_i \neq 0 \quad \text{and} \quad f_{i\psi} = 0 \text{ if } p_i = 0 \quad (5)$$

If  $p_i \neq 0$ , then  $\psi$  inhibits the forward reaction and we assume that sufficiently large values of  $\psi$  make the reaction rate arbitrarily small or negative, i.e.

$$\begin{aligned} \lim_{\psi \rightarrow \infty} f_i(\cdot, \psi) &\leq 0 \quad i = 1, n + 1 \\ \lim_{\psi \rightarrow \infty} f_i(\cdot, \cdot, \psi) &\leq 0 \quad i = 2, \dots, n \end{aligned}$$

This reflects the fact that the energy required to pump a proton against a chemical and electrical gradient becomes large as the gradient increases. Similarly  $-\psi$  inhibits the backward reaction so that:

$$\begin{aligned} \lim_{\psi \rightarrow -\infty} f_i(\cdot, \psi) &\geq 0 \quad i = 1, n + 1 \\ \lim_{\psi \rightarrow -\infty} f_i(\cdot, \cdot, \psi) &\geq 0 \quad i = 2, \dots, n \end{aligned}$$

The following equations imply that no reaction can proceed in the absence of any of its substrates:

$$\left. \begin{aligned} f_1(0, \cdot) &= 0 \\ f_i(\cdot, 0, \cdot) &= 0 \quad i = 2, \dots, n \\ f_i(m_{i-1}, \cdot, \cdot) &= 0 \quad i = 2, \dots, n \\ f_{n+1}(m_n, \cdot) &= 0 \end{aligned} \right\} \quad (6)$$

The final set of conditions imply that increased substrate concentration increases the rate of reaction unless one of the substrates is entirely absent:

$$\left. \begin{aligned} f_{11} &> 0 \\ f_{ii} &\geq 0 \quad \text{and} \quad f_{ii} > 0 \text{ if } x_{i-1} < m_{i-1} \quad i = 2, \dots, n \\ f_{i+1,i} &\leq 0 \quad \text{and} \quad f_{i+1,i} > 0 \text{ if } x_{i+1} > 0 \quad i = 1, \dots, n - 1 \\ f_{n+1,n} &< 0 \end{aligned} \right\} \quad (7)$$

The fact that the first and final inequalities are always strict implies that there is always some electron donor to reduce the initial substrate, and some electron acceptor to oxidise the final substrate, and ensures nondegenerate behaviour. The assumptions from (7) mean that  $f_{ii}$ ,  $F_{ij}$  and  $F_i\psi$  as defined in (4) are all nonnegative. The definition of these nonnegative quantities is solely to simplify later arguments.

### 3 General behaviour of the system

In this section we outline some properties of the model that hold regardless of the number  $n$  of redox pairs.

### 3.1 Boundedness of solutions

It is convenient to define an  $n \times (n + 1)$  matrix which can be regarded as a stoichiometric matrix for the redox reactions:

$$S \equiv \begin{bmatrix} -1 & 1 & \dots & 0 & 0 \\ 0 & -1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & -1 & 1 \end{bmatrix}$$

Defining the vector of reactant concentrations  $\mathbf{x} = [x_1, x_2, \dots, x_n]^T$ , the vector of reaction rates  $\mathbf{v}(\mathbf{x}, \psi) = [f_1, f_2, \dots, f_{n+1}]^T$ , and the nonnegative vector  $P \equiv [p_1, \dots, p_{n+1}]^T$ , we can rewrite the system of Eq. 3 more briefly as

$$\begin{aligned} \dot{\mathbf{x}} &= S\mathbf{v}(\mathbf{x}, \psi) \\ \dot{\psi} &= P^T \mathbf{v}(\mathbf{x}, \psi) - L(\psi) \end{aligned}$$

We now show that all forward trajectories of the system are bounded. Since the phase space is bounded in  $\mathbf{x}$ , what needs to be shown is that all trajectories enter a bounded region in the  $\psi$  direction. This amounts to showing that  $\dot{\psi} > 0$  for  $\psi$  sufficiently large and negative, and that  $\dot{\psi} < 0$  for  $\psi$  sufficiently large and positive. By assumption, for any given  $i$ , either  $p_i = 0$  or  $f_{i\psi}$  is negative and  $\lim_{\psi \rightarrow \infty} f_i(\cdot, \cdot, \psi) \leq 0$ ,  $\lim_{\psi \rightarrow -\infty} f_i(\cdot, \cdot, \psi) \geq 0$ . This in turn implies that  $\lim_{\psi \rightarrow \infty} P^T \mathbf{v}(\mathbf{x}, \psi) \leq 0$  and  $\lim_{\psi \rightarrow -\infty} P^T \mathbf{v}(\mathbf{x}, \psi) \geq 0$ . In addition  $L$  is strictly increasing with  $L(0) = 0$ . Thus for any fixed value of  $\mathbf{x}$ ,  $\lim_{\psi \rightarrow \infty} P^T \mathbf{v}(\mathbf{x}, \psi) - L(\psi) < 0$  and  $\lim_{\psi \rightarrow -\infty} P^T \mathbf{v}(\mathbf{x}, \psi) - L(\psi) > 0$ . Define  $\psi_0(\mathbf{x})$  as the value of  $\psi$  at which  $P^T \mathbf{v}(\mathbf{x}, \psi) - L(\psi) = 0$ .  $\psi_0(\mathbf{x})$  is uniquely defined since  $P^T \mathbf{v}(\mathbf{x}, \psi) - L(\psi)$  is strictly decreasing. Since  $P^T \mathbf{v}(\mathbf{x}, \psi) - L(\psi)$  is a differentiable function of  $\psi$  with non-zero derivative, by the implicit function theorem,  $\psi_0(\mathbf{x})$  is differentiable. Since it has a compact domain,  $\psi_0(\mathbf{x})$  achieves a maximum value which we call  $\psi_{\max}$ , and a minimum value which we call  $\psi_{\min}$ . By these definitions,  $\dot{\psi}(\psi, \mathbf{x}) < 0$  for all  $\psi > \psi_{\max}$ , and  $\dot{\psi}(\psi, \mathbf{x}) > 0$  for all  $\psi < \psi_{\min}$ .

Thus all trajectories enter a closed box,  $\mathcal{B}$ , bounded by the hyperplanes  $x_i = 0$ ,  $x_i = m_i$ ,  $\psi = \psi_{\min}$  and  $\psi = \psi_{\max}$ , and this box forms a trapping region for the system in all dimensions.

### 3.2 The Jacobian

Direct calculation gives that the Jacobian,  $J$ , of the system is:

$$J = \begin{bmatrix} -f_{11} - F_{21} & f_{22} & \dots & 0 & F_{1\psi} - F_{2\psi} \\ F_{21} & -f_{22} - F_{32} & \dots & 0 & F_{2\psi} - F_{3\psi} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & -f_{nn} - F_{n+1,n} & F_{n\psi} - F_{n+1,\psi} \\ p_1 f_{11} - p_2 F_{21} & p_2 f_{22} - p_3 F_{32} & \dots & p_n f_{nn} - p_{n+1} F_{n+1,n} & -L_\psi - \sum_{i=1}^{n+1} p_i F_{i\psi} \end{bmatrix}$$

Here  $L_\psi \equiv \frac{dL}{d\psi}$ . The structure of this Jacobian can be made clearer by defining two further quantities: A nonnegative vector in  $\mathbb{R}^n$ ,  $F \equiv [F_{1\psi}, \dots, F_{n\psi}]^T$ ; and an  $(n+1) \times n$  matrix

$$V \equiv \frac{\partial \mathbf{v}}{\partial \mathbf{x}} = \begin{bmatrix} f_{11} & 0 & 0 & \cdots & 0 \\ -F_{21} & f_{22} & 0 & \cdots & 0 \\ 0 & -F_{32} & f_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & f_{nn} \\ 0 & 0 & 0 & \cdots & -F_{n+1,n} \end{bmatrix}$$

Then the Jacobian can be written in the block form:

$$J = \begin{bmatrix} SV & SF \\ P^T V & -P^T F - L_\psi \end{bmatrix} \quad (8)$$

$SV$  is the Jacobian of the system without feedback, which is tridiagonal, and can easily be shown to have real negative eigenvalues [6]. It was shown in [16] that the structures of  $S$  and  $V$  along with the nonnegativity of  $P$  and  $F$  imply that  $J$  is a so called  $P^{(-)}$  matrix (see Appendix A for the definition).<sup>1</sup> This result is independent of  $n$ , the length of the chain. It has the consequence that the system is injective; this is discussed further in the next section.

The fact that  $J$  is a  $P^{(-)}$  matrix has another consequence of importance to us: It means that its eigenvalues are excluded from a certain wedge around the positive real axis: If  $\lambda = re^{i\theta}$  is an eigenvalue of an  $m \times m$   $P$  matrix, then it is proved in [17] that:

$$|\theta - \pi| > \pi/m$$

and equivalently for a  $P^{(-)}$  matrix,

$$|\theta| > \pi/m$$

Clearly when  $m = 2$ , this means that both eigenvalues lie in the left half plane, so that  $2 \times 2$   $P^{(-)}$  matrices are Hurwitz stable (see Appendix A for a definition of “Hurwitz stable” which we will abbreviate to “Hurwitz”). However for  $m > 2$ ,  $P^{(-)}$  matrices may be unstable.

### 3.3 A unique equilibrium

The existence of a unique equilibrium for this system was shown in [6] by a direct method. It also follows from the arguments presented above. The existence of a compact, convex, trapping region  $\mathcal{B}$  constructed above implies, by the Brouwer fixed point

<sup>1</sup> The nondegeneracy conditions presented in [16] are met because the  $n$ th substrate is terminal, and all substrates are able to transfer electrons along the chain to the  $n$ th substrate.

theorem, that an equilibrium exists. That this equilibrium must be unique follows from the fact that the Jacobian is a  $P^{(-)}$  matrix, and hence the system is injective [18]. Thus as our first result we can state that

Electron transport chains coupled to charge translocation across a membrane have exactly one equilibrium.

It is interesting that the possibility of multistability is immediately ruled out. However this in itself does not tell us whether all trajectories must necessarily converge to the unique equilibrium, or whether periodic or chaotic behaviour is still possible.

## 4 Stability of the equilibrium

In this section, we explore local and global stability of the unique equilibrium, starting with low dimensions (i.e. short chains). For two dimensions we prove that the equilibrium is globally asymptotically stable. In three dimensions we show that the addition of an extra, reasonable, constraint implies that the equilibrium is locally stable, and further constraints ensure that it is globally stable. We then demonstrate that these constraints do not suffice to guarantee stability in four dimensions and higher. Finally, we outline some additional special conditions that guarantee the Jacobian is Hurwitz in all dimensions.

### 4.1 The system in two dimensions

The system in 2D consists of a single redox pair subject to a reduction process and an oxidation process, both possibly coupled to proton translocation across the membrane. It takes the form

$$\begin{aligned}\dot{x}_1 &= -f_1(x_1, \psi) + f_2(x_1, \psi) \\ \dot{\psi} &= p_1 f_1 + p_2 f_2 - L(\psi)\end{aligned}$$

The Jacobian of the system in this case is:

$$J_2 = \begin{bmatrix} -f_{11} - F_{21} & F_{1\psi} - F_{2\psi} \\ p_1 f_{11} - p_2 F_{21} & -L_\psi - p_1 F_{1\psi} - p_2 F_{2\psi} \end{bmatrix} \quad (9)$$

We have already mentioned that 2D  $P^{(-)}$  matrices are Hurwitz stable, and it follows that the matrices  $J_2$  are Hurwitz stable. (This can also be shown with a direct calculation.)

Since  $J_2$  is Hurwitz stable *everywhere*, not just at the unique equilibrium, the Markus-Yamabe Theorem (e.g. [19–21]) ensures that the equilibrium is globally stable. We also offer an alternative, elementary, proof of global stability. By the Poincaré-Bendixson Theorem (see, for example, [22]),  $\omega$ -limit sets of a flow on compact subsets of  $\mathbb{R}^2$  must either contain equilibria or consist of a periodic orbit. In this



case we can rule out the possibility of periodic orbits: The divergence of the vector field is equal to

$$\text{Tr}(J) = -f_{11} - F_{21} - p_1 F_{1\psi} - p_2 F_{2\psi} - L_\psi$$

which is negative. Thus the vector field satisfies the Dulac criterion (e.g. [23]) and there are no periodic orbits. We know that there is only one equilibrium, which is locally stable, and therefore there are no heteroclinic or homoclinic orbits either. Since every forward trajectory enters the box  $B$ , the unique equilibrium must be the  $\omega$ -limit of every trajectory, and is hence globally stable.

#### 4.2 The system in three dimensions

Slightly more complex than the two dimensional system is the system in three dimensions which takes the form

$$\begin{aligned} \dot{x}_1 &= -f_1(x_1, \psi) + f_2(x_1, x_2, \psi) \\ \dot{x}_2 &= -f_2(x_1, x_2, \psi) + f_3(x_2, \psi) \\ \dot{\psi} &= p_1 f_1 + p_2 f_2 + p_3 f_3 - L(\psi) \end{aligned}$$

with Jacobian

$$J_3 = \begin{bmatrix} -f_{11} - F_{21} & f_{22} & F_{1\psi} - F_{2\psi} \\ F_{21} & -f_{22} - F_{32} & F_{2\psi} - F_{3\psi} \\ p_1 f_{11} - p_2 F_{21} & p_2 f_{22} - p_3 F_{32} & -L_\psi - p_1 F_{1\psi} - p_2 F_{2\psi} - p_3 F_{3\psi} \end{bmatrix} \quad (10)$$

As it stands,  $J_3$  is not always Hurwitz. For example, the Jacobian constructed using the following values:  $p_1 = 3$ ,  $p_2 = 0$ ,  $p_3 = 88$ ,  $F_{1\psi} = 33$ ,  $F_{2\psi} = 4$ ,  $F_{3\psi} = 0.6$ ,  $f_{11} = 23$ ,  $f_{22} = 3$ ,  $F_{21} = 94$ ,  $F_{32} = 76$ ,  $L_\psi = 6$  has two eigenvalues with positive real part.

$J_3$  can be shown to be Hurwitz everywhere provided one extra condition is met:  $p_1$  and  $p_3$  must have the same ordering as  $F_{1\psi}$  and  $F_{3\psi}$ . For a real number  $z$ , define the sign function in the usual way:

$$\text{sign}(z) \equiv \begin{cases} 1 & (z > 0) \\ 0 & (z = 0) \\ -1 & (z < 0) \end{cases} \quad (11)$$

Then the ordering assumption translates to the following statement:

$$\text{sign}(F_{3\psi} - F_{1\psi}) = \text{sign}(p_3 - p_1) \quad (12)$$

With this assumption, the Jacobian is everywhere Hurwitz, and hence the equilibrium is locally asymptotically stable. The proof is simple but requires some lengthy evaluations, and the details are presented in Appendix B.

Unlike in the 2D case it does not follow that the equilibrium is globally stable, since the Markus-Yamabe conjecture does not hold in dimensions greater than 2 [24]. However we can prove global stability in this case too subject to a strengthened version of the ordering assumption on the quantities  $p_i$  and  $F_{i\psi}$ . We now require

$$\text{sign}(F_{i\psi} - F_{j\psi}) = \text{sign}(p_i - p_j) \tag{13}$$

for  $i, j \in \{1, 2, 3\}$ .

With this assumption we are able to use a version of Li and Muldowney’s autonomous convergence theorem (Theorem 4.1 in [25]) to show that the unique equilibrium is globally stable. In order to use this theorem two concepts are needed:

1. The *second additive compound* of a matrix
2. *Logarithmic norms* of a matrix.

Both quantities are defined for square matrices. The second additive compound matrix of any  $n \times n$  matrix  $J$  is a square matrix of dimension  ${}^n C_2$  which we will term  $J^{[2]}$ . Logarithmic norms are scalar quantities, and corresponding to any given matrix norm, there is a logarithmic norm. Unlike matrix norms, however, logarithmic norms may take negative values. The definitions are given in Appendix A.

Consider a dynamical system with Jacobian  $J(x)$  at some point of phase space  $x$ . Define  $\mathbf{J}$  to be the set of all these Jacobians. For our purposes, the autonomous convergence theorem states the following: If a logarithmic norm  $\mu$  can be found such that

$$\mu(J^{[2]}) < 0 \text{ for all } J \in \mathbf{J} \tag{14}$$

then the limit set of each bounded semi-trajectory of the dynamical system is an equilibrium.

Since all trajectories enter the trapping region  $\mathcal{B}$  in our system, and since  $\mathcal{B}$  contains a unique equilibrium, finding a suitable logarithmic norm satisfying (14) will suffice to prove global stability of the equilibrium.

The second additive compound in this case is:

$$J_3^{[2]} = \begin{bmatrix} -f_{11} - F_{21} - f_{22} - F_{32} & F_{2\psi} - F_{3\psi} & -(F_{1\psi} - F_{2\psi}) \\ p_2 f_{22} - p_3 F_{32} & -f_{11} - F_{21} - L_\psi - \sum_{i=1}^3 p_i F_{i\psi} & f_{22} \\ -(p_1 f_{11} - p_2 F_{21}) & F_{21} & -f_{22} - F_{32} - L_\psi - \sum_{i=1}^3 p_i F_{i\psi} \end{bmatrix}$$

We will construct a logarithmic norm  $\mu_T$  such that  $\mu_T(J_3^{[2]}) < 0$ . For a real  $n \times n$  matrix, the logarithmic norm corresponding the usual  $\|\cdot\|_1$  norm takes the form:

$$\mu_1 = \max_{i \in \{1, \dots, n\}} \left( x_{ii} + \sum_{k \neq i} |x_{ki}| \right)$$

From the definition it is clear that a matrix has negative logarithmic norm  $\mu_1$  if and only if every diagonal entry is negative and it is strictly diagonally dominant in every column. Next we define a constant diagonal coordinate transformation

$$T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{p_{\max}} & 0 \\ 0 & 0 & \frac{1}{p_{\max}} \end{pmatrix}$$

where  $p_{\max} = \max_{i \in \{1,2,3\}} (p_i)$ .

According to Lemma 2.2 of [26], given any invertible transformation  $T$ ,  $\mu_T(M) \equiv \mu_1(TMT^{-1})$  defines a new logarithmic norm. In this case, since  $T$  is a diagonal matrix, the diagonal entries of  $M$  are the same as those of  $TMT^{-1}$ . Thus in order to prove that  $\mu_T(J_3^{[2]}) < 0$ , we need to show that  $J' \equiv TJ_3^{[2]}T^{-1}$  is strictly diagonally dominant in every column.

For the first column, we have

$$J'_{11} + |J'_{21}| + |J'_{31}| = -f_{22} - F_{32} - f_{11} - F_{21} + \left| \frac{p_2}{p_{\max}} f_{22} - \frac{p_3}{p_{\max}} F_{32} \right| + \left| \frac{p_2}{p_{\max}} F_{21} - \frac{p_1}{p_{\max}} f_{11} \right|$$

It can easily be seen that the term on the right hand side is negative since for any two nonnegative scalars  $|a - b| \leq \max\{|a|, |b|\}$ .

For the second column, we have

$$J'_{22} + |J'_{12}| + |J'_{32}| = - \sum_{i=1}^3 p_i F_{i\psi} - L_\psi - f_{11} + p_{\max} |F_{2\psi} - F_{3\psi}|$$

For the final column, we have

$$J'_{33} + |J'_{13}| + |J'_{23}| = - \sum_{i=1}^3 p_i F_{i\psi} - L_\psi - F_{32} + p_{\max} |F_{2\psi} - F_{1\psi}|$$

In order to show that the right hand sides of the last two expressions are negative we need to show in each case that our ordering assumption (13) implies that the final term (which may be positive) is dominated in magnitude by the other terms.

Note that  $|F_{i\psi} - F_{j\psi}| \leq \max\{F_{i\psi}, F_{j\psi}\} \leq \max_{k \in \{1,2,3\}} (F_{k\psi})$ . Then there are only three cases:

1. if  $p_{\max} = p_1$ , then  $p_{\max} |F_{2\psi} - F_{3\psi}| \leq p_1 F_{1\psi}$ , and  $p_{\max} |F_{2\psi} - F_{1\psi}| \leq p_1 F_{1\psi}$ .
2. if  $p_{\max} = p_2$ , then  $p_{\max} |F_{2\psi} - F_{3\psi}| \leq p_2 F_{2\psi}$ , and  $p_{\max} |F_{2\psi} - F_{1\psi}| \leq p_2 F_{2\psi}$ .
3. if  $p_{\max} = p_3$ , then  $p_{\max} |F_{2\psi} - F_{3\psi}| \leq p_3 F_{3\psi}$ , and  $p_{\max} |F_{2\psi} - F_{1\psi}| \leq p_3 F_{3\psi}$ .

Each of these possibilities leads to the same conclusion—that  $J'_{ii} + \sum_{k \neq i} |J'_{ki}| < 0$  for

each  $i$ . Hence we have  $\mu_T(J_3^{[2]}) < 0$ .

This result means that if the ordering assumption (13) holds, then the unique equilibrium is globally stable. The ordering assumption itself has the following reasonable physical meaning which we would expect to be fulfilled in practice: If redox reaction  $i$  is involved in pumping more protons across the membrane than redox reaction  $j$ , then reaction  $i$  is correspondingly more inhibited by  $\psi$  than reaction  $j$ . It is interesting to note however that this assumption is not necessary to prove global stability in the 2D case. It is also unknown to us whether the weaker assumption (12), which guarantees that the Jacobian is everywhere Hurwitz, actually guarantees global stability in 3D.

### 4.3 Unstable examples in higher dimensions

The ordering assumption (13) does not guarantee global or even local stability of the equilibrium in dimensions greater than 3. It is easy to construct counterexamples. For example, in four dimensions, the Jacobian constructed by choosing  $p_1 = 2$ ,  $p_2 = p_3 = 0$ ,  $p_4 = 73$ ,  $F_{1\psi} = 167$ ,  $F_{2\psi} = F_{3\psi} = 0$ ,  $F_{4\psi} = 176$ ,  $f_{11} = 4$ ,  $f_{22} = 7$ ,  $f_{33} = 1$ ,  $F_{21} = 32$ ,  $F_{32} = 64$ ,  $F_{43} = 174$ ,  $L_\psi = 33$ , satisfies all the constraints, including the ordering assumption on the values of  $p_i$  and  $F_{i\psi}$ . However it has two eigenvalues with positive real part.

We make the following remarks:


1. By continuity, the fact that a non-Hurwitz Jacobian can be constructed in four dimensions guarantees that such examples also exist in all higher dimensions.
2. Systems with non-Hurwitz Jacobian satisfying the ordering assumption (13) seem to be rare. A script implemented in the open source numerical computation program Scilab [27], was able to find counterexamples in dimension 4 by randomly choosing values for the different terms in the Jacobian, such that all the assumptions were satisfied. Out of hundreds of millions of sets of values, less than 10 were non-Hurwitz.
3. The counterexamples found appear always to be close to breaking the ordering assumption. For instance, in the example shown,  $p_4$  is much greater than  $p_1$ , whereas  $F_{4\psi}$  is close in magnitude to  $F_{1\psi}$ .

### 4.4 A special case: reaction rates dependent on potentials


In this section we consider an interesting assumption which ensures that the Jacobian is Hurwitz everywhere (and hence the unique equilibrium is locally stable). The assumption is as follows:

1. Associated with each half reaction is some “potential”: In the case of a redox reaction of the form  $A_i + e^- \rightleftharpoons B_i$ , a potential means any strictly increasing scalar function of  $[A_i]$ , such as a redox potential; In the case of a charge transfer across a membrane a potential means any strictly increasing scalar function of  $\psi$ .
2. The rate of any full reaction depends only on the *sum* of the potentials for the half reactions involved, and is a strictly decreasing function of this sum.

This assumption can be interpreted, loosely, as saying that the energetics of the system determines the kinetics. For example, consider the electron transfer coupled to some proton pumping



derived from the half reactions



In this case, the assumption would imply that the forward rate of the combined reaction can be written  $f(-g_j(x_j) + g_i(x_i) - pg_\psi(\psi))$  where the only stipulation is that  $f$ ,  $g_i$ ,  $g_j$  and  $g_\psi$  are strictly increasing in their arguments. When this assumption is made about all reaction rates in the system, the full system becomes:

$$\begin{aligned} \dot{x}_1 &= -f_1(g_1(x_1) - p_1g_\psi(\psi)) + f_2(-g_1(x_1) + g_2(x_2) - p_2g_\psi(\psi)) \\ \dot{x}_i &= -f_i(-g_{i-1}(x_{i-1}) + g_i(x_i) - p_i g_\psi(\psi)) + f_{i+1}(-g_i(x_i) + g_{i+1}(x_{i+1}) \\ &\quad - p_{i+1}g_\psi(\psi)) \quad i = 2, \dots, n \\ \dot{x}_n &= -f_n(-g_{n-1}(x_{n-1}) + g_n(x_n) - p_n g_\psi(\psi)) + f_{n+1}(-g_n(x_n) \\ &\quad - p_{n+1}g_\psi(\psi)) \\ \dot{\psi} &= \sum_{i=1}^{n+1} p_i f_i - L(\psi) \end{aligned}$$

The term  $f_i(-g_{i-1}(x_{i-1}) + g_i(x_i) - p_i g_\psi(\psi))$  represents the rate at which the  $i$ th substrate receives electrons from the  $(i - 1)$ th substrate. Denoting by  $f'_i$ ,  $g'_i$  and  $g'_\psi$  the derivatives of the functions  $f_i$ ,  $g_i$  and  $g_\psi$ , the Jacobian of this system can be written  $J = J_0 D$  where  $J_0$  is the symmetric matrix

$$J_0 = \begin{bmatrix} -(f'_1 + f'_2) & f'_2 & \cdots & 0 & p_1 f'_1 - p_2 f'_2 \\ f'_2 & -(f'_2 + f'_3) & \cdots & 0 & p_2 f'_2 - p_3 f'_3 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & -(f'_n + f'_{n+1}) & p_n f'_n - p_{n+1} f'_{n+1} \\ p_1 f'_1 - p_2 f'_2 & p_2 f'_2 - p_3 f'_3 & \cdots & p_n f'_n - p_{n+1} f'_{n+1} & -\sum_{i=1}^{n+1} p_i^2 f'_i - \frac{L'_\psi}{g'_\psi} \end{bmatrix} \quad (15)$$

and  $D$  is the positive diagonal matrix

$$D = \begin{bmatrix} g'_1 & 0 & \cdots & 0 & 0 \\ 0 & g'_2 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & g'_n & 0 \\ 0 & 0 & \cdots & 0 & g'_\psi \end{bmatrix} \quad (16)$$

From the discussions earlier,  $J_0$  is a  $P^{(-)}$  matrix. Further it is symmetric, and hence sign symmetric (see Appendix A for a definition of sign symmetry). This implies [28] that  $J_0$  is  $D$ -stable, i.e. the product of  $J_0$  with any positive diagonal matrix is Hurwitz. Hence  $J$  is Hurwitz. Thus the assumption that reaction rates depend on the sum of potentials of the half reactions ensures that the Jacobian of the system is everywhere Hurwitz.

## 5 Discussion and conclusions

We have analysed in some detail, and using a variety of mathematical techniques, the behaviour of electron transport chains coupled to a charge translocation process. In all cases trajectories are bounded, and there is a unique equilibrium, but questions about the stability of this equilibrium have proved harder. Where the chain consists of a single redox pair, the unique equilibrium is globally stable. When there are two redox pairs the same conclusions can be reached subject to some extra conditions on the feedback process. In higher dimensions no such general conditions can easily be found.

It is surprising that the coupling of electron transfer to a membrane potential—a negative feedback loop—can serve to destabilise the unique equilibrium for sufficiently long chains. Interestingly, when the reaction rates are monotonic functions of a sum of potentials, then the system in any dimension could be proved to be everywhere Hurwitz. Reaction rates cannot in general be seen in this way, but in the case of reactions which are primarily about charge transfer, the assumption could be reasonable. Certainly some of the choices of reaction rates in numerical models such as [8] satisfy this assumption.

It is interesting to note that the system without feedback is cooperative in the sense of [29] for any fixed values of  $\psi$ . Moreover, when  $\psi$  is seen as an input to the model, it follows from results in [6] that it has a well defined input to state static characteristic [30]—i.e. a single globally asymptotically stable, non-degenerate equilibrium for each value of  $\psi$ . These facts suggest that it should be possible to apply the so-called “small gain theorems” developed by Sontag, Angeli and others [31,30] in order to make claims about the behaviour of the system with feedback. The difficulty which arises is that the system as written is not monotone in its inputs (for any order) because the quantities  $F_{i,\psi} - F_{i+1,\psi}$  can take any real value. On the other hand the quantities  $F_{i,\psi}$  are of defined sign, suggesting that a clever recoordination might make the system monotone in its inputs, allowing application of small gain theorems. We consider this a very promising avenue for future work.

The work also raises other interesting open questions, both biological and mathematical. From a biological point of view, it would be interesting to see how additional processes such as transport processes in the full numerical models ([8, 11] for example) affect the conclusions presented here.

We note that since the Markus-Yamabe conjecture does not hold in dimensions greater than 2 [24], a Jacobian which is everywhere Hurwitz does not necessarily imply global stability of an equilibrium. An open mathematical question is whether there are equivalent conditions to the ordering condition in 3D which ensure that the Jacobian of the system is Hurwitz in arbitrary dimension, or better still that the second additive compound has negative logarithmic norm, and hence the unique equilibrium is globally stable. If such conditions exist can they be given general biological meanings?

It would also be interesting to explore when the results presented here survive weakening of the assumption that electrons are transferred along a chain. Although electron transfers taking place in the mitochondrial membrane are often described via a chain it is likely that this description is to some extent a simplification. General electron transfer networks in the absence of a potential were analysed in [7] and found to have simple behaviour. Application of the theory presented in [16] should allow determination of when these networks give rise to  $P^{(-)}$  Jacobians when interacting with a membrane potential.

Finally, given our comment on the Markus-Yamabe conjecture above, although conditions ensuring sign-symmetry of the system imply that the Jacobian is everywhere Hurwitz, it is an open question as to whether a sign-symmetric  $P^{(-)}$  Jacobian implies global stability of the unique equilibrium. We conjecture that this is the case, but this remains to be proved.

## Appendix

### A Definitions

*Hurwitz stability of matrices* A square matrix is defined to be *Hurwitz stable* if all its eigenvalues lie in the open left half of the complex plane—i.e. the real parts of all its eigenvalues are negative.

*P matrices and related classes* For some  $n \times m$  matrix  $A$ ,  $A(\alpha|\gamma)$  will refer to the submatrix of  $A$  with rows indexed by the set  $\alpha \subset \{1, \dots, n\}$  and columns indexed by the set  $\gamma \subset \{1, \dots, m\}$ . A *principal submatrix* of  $A$  is a submatrix containing columns and rows from the same index set, i.e. of the form  $A(\alpha|\alpha)$ . A *minor* is the determinant of any square submatrix of  $A$ . If  $A(\alpha|\gamma)$  is a square submatrix of  $A$  (i.e.  $|\alpha| = |\gamma|$ ), then  $A[\alpha|\gamma]$  will refer to the corresponding minor, i.e.  $A[\alpha|\gamma] = \det(A(\alpha|\gamma))$ . A *principal minor* of  $A$  is the determinant of a principal submatrix of  $A$ .

$P$  matrices are square matrices all of whose principal minors are positive. They are by definition nonsingular. If  $-A$  is a  $P$  matrix, then we will say that  $A$  is a  $P^{(-)}$  matrix. If  $A$  is a  $P^{(-)}$  matrix, this means that each  $k \times k$  principal minor of  $A$  has sign  $(-1)^k$ .

*Sign symmetry* An  $n \times n$  matrix is *sign-symmetric* if symmetrically placed minors have the same sign, i.e.  $A[\alpha|\gamma]A[\gamma|\alpha] \geq 0$  for every  $\alpha, \gamma \subset \{1, \dots, n\}$  with  $|\alpha| = |\gamma|$ .

*Second additive compound matrices* A brief definition of the second additive compound of any square matrix can be found in [32]. For a more detailed discussion see [33]. For a 3D matrix

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \tag{17}$$

the second additive compound takes the form<sup>2</sup>

$$A^{[2]} = \begin{pmatrix} a_{11} + a_{22} & a_{23} & -a_{13} \\ a_{32} & a_{11} + a_{33} & a_{12} \\ -a_{31} & a_{21} & a_{22} + a_{33} \end{pmatrix}$$

This second additive compound was constructed using the standard lexicographic ordering of basis vectors. Choices of ordering make no difference to the logarithmic norms of the matrix.

*Logarithmic norms* If  $\|\cdot\|$  denotes a vector norm on  $\mathbb{R}^n$ , and also the induced matrix norm on  $n \times n$  matrices, then the logarithmic norm [34], also known as a Lozinskiĭ measure, of an  $n \times n$  matrix  $A$  is defined by

$$\mu(A) = \lim_{h \rightarrow 0^+} \frac{\|I + hA\| - 1}{h} \tag{18}$$

### B Local stability in 3D

In this appendix we prove local stability of the equilibrium in three dimensions, subject to the assumption in (12), using the Routh-Hurwitz theorem. Consider the characteristic polynomial of a matrix  $A$ :

$$|\lambda I - A| = \lambda^n + b_1\lambda^{n-1} + \dots + b_{n-1}\lambda + b_n \tag{19}$$

In this equation,  $I$  is the  $n \times n$  identity matrix, and the coefficients  $b_i$  are the sums of all principal minors of  $-A$  of dimension  $i$ . For a  $P^{(-)}$  matrix,  $b_i > 0$  for all  $i$ . Now define  $b_k \equiv 0$  for all  $k > n$ , and construct a set of numbers  $\Delta_i$  as follows:

<sup>2</sup> In general, the second additive compound of a matrix  $A$  has dimension  ${}^d C_2$  where  $d = \dim(A)$ . When  $\dim(A) = 3$ , we get  $\dim(A^{[2]}) = 3$  also, but this is not generally the case.



$$\Delta_i = \begin{vmatrix} b_1 & 1 & 0 & 0 & 0 & 0 & \cdots & 0 \\ b_3 & b_2 & b_1 & 1 & 0 & 0 & \cdots & 0 \\ b_5 & b_4 & b_3 & b_2 & b_1 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & 0 \\ b_{2i-1} & b_{2i-2} & b_{2i-3} & b_{2i-4} & b_{2i-5} & b_{2i-6} & \cdots & b_i \end{vmatrix} \quad (20)$$

The Routh-Hurwitz theorem states that  $A$  is Hurwitz if and only if  $\Delta_i > 0$  for all  $i \leq n$ . In three dimensions, we need to check that the three quantities

$$\Delta_1 = b_1 \quad (21)$$

$$\Delta_2 = b_1 b_2 - b_3 \quad (22)$$

$$\Delta_3 = b_3(b_1 b_2 - b_3) = b_3 \Delta_2 \quad (23)$$

are all positive. Since all the  $b_i$  are positive, all three quantities are positive if and only if  $\Delta_2 > 0$ . This in turn follows (condition 12 in [28]) if

$$0 < a_{12} a_{23} a_{31} + a_{21} a_{32} a_{13} - 2a_{11} a_{22} a_{33}$$

where  $a_{ij}$  are elements of  $A$ . Substituting  $a_{ij}$  for the elements of the Jacobian and expanding using the open source symbolic algebra program Maxima [35] gives:

$$\begin{aligned} a_{12} a_{23} a_{31} + a_{21} a_{32} a_{13} - 2a_{11} a_{22} a_{33} = & F_{21} F_{32} (2p_3 F_{3\psi} + 2p_1 F_{1\psi} - p_3 F_{1\psi}) \\ & + f_{11} f_{22} (2p_3 F_{3\psi} + 2p_1 F_{1\psi} - p_1 F_{3\psi}) \\ & + \text{positive terms} \end{aligned}$$

With the ordering assumption (12), we get:

$$2p_3 F_{3\psi} + 2p_1 F_{1\psi} - p_3 F_{1\psi} \geq 0 \quad (24)$$

$$2p_3 F_{3\psi} + 2p_1 F_{1\psi} - p_1 F_{3\psi} \geq 0 \quad (25)$$

Thus the Jacobian is everywhere Hurwitz and hence the unique equilibrium of the system must be locally asymptotically stable. Note that the restriction (12) is stronger than necessary to ensure that  $J$  is Hurwitz, but no other set of conditions with a clear physical meaning that make the Jacobian Hurwitz have been discovered. Finding a set of necessary and sufficient conditions for  $J$  to be Hurwitz is a difficult problem.

## References

1. G.T. Babcock, M. Wikström, Oxygen activation and the conservation of energy in cell respiration. *Nature* **356**, 301 (1992)
2. R.H. Garrett, C.M. Grisham (eds.), *Biochemistry* (Saunders College Publishing, 1995)
3. N. Bhagavan, *Medical Biochemistry* (Harcourt/Academic Press, 2002)
4. D.G. Nicholls, S.J. Ferguson, *Bioenergetics 3* (Academic Press, 2002)
5. I. Belevich, M. Verkhovsky, M. Wikström, Proton-coupled electron transfer drives the proton pump of cytochrome *c* oxidase. *Nature* **440**(6), 829 (2006)

6. M. Banaji, A generic model of electron transport in mitochondria. *J. Theor. Biol.* **243**(4), 501 (2006)
7. M. Banaji, S. Baigent, Electron transfer networks. *J. Math. Chem.* **43**(4), 1355 (2008)
8. B. Korzeniewski, Simulation of oxidative phosphorylation in hepatocytes. *Biophys. Chem.* **58**, 215 (1996)
9. B. Korzeniewski, J.A. Zoladz, A model of oxidative phosphorylation in mammalian skeletal muscle. *Biophys. Chem.* **92**, 17 (2001)
10. A.D. Farmery, J.P. Whiteley, A mathematical model of electron transfer within the mitochondrial respiratory cytochromes. *J. Theor. Biol.* **213**, 197 (2001)
11. D.A. Beard, A biophysical model of the mitochondrial respiratory system and oxidative phosphorylation. *PLoS Comput. Biol.* **1**(4), e36 (2005)
12. Q. Jin, C.M. Bethke, Kinetics of electron transfer through the respiratory chain. *Biophys. J.* **83**(4), 1797 (2002)
13. P. De Leenheer, D. Angeli, E. Sontag, Monotone chemical reaction networks. *J. Math. Chem.* **41**(3), 295 (2007)
14. M.D. Brand, L. Chien, P. Diolecz, Experimental discrimination between proton leak and redox slip during mitochondrial electron transport. *Biochem. J.* **297**(1), 27 (1994)
15. M. Canton, S. Luvisetto, I. Schmehl, G. Azzone, The nature of mitochondrial respiration and discrimination between membrane and pump properties. *Biochem. J.* **310**, 477 (1995)
16. M. Banaji, P. Donnell, S. Baigent,  $P$  matrix properties, injectivity and stability in chemical reaction systems. *SIAM J. Appl. Math.* **67**(6), 1523 (2007)
17. R.B. Kellogg, On complex eigenvalues of  $M$  and  $P$  matrices. *Numer. Math.* **19**, 70 (1972)
18. D. Gale, H. Nikaïdo, The Jacobian matrix and global univalence of mappings. *Math. Ann.* **159**, 81 (1965)
19. R. Feßler, A proof of the two-dimensional Markus-Yamabe stability conjecture. *Ann. Polon. Math.* **62**, 45 (1995)
20. A.A. Glutsyuk, The complete solution of the Jacobian problem for vector fields on the plane. *Russ. Math. Surv.* **49**(3), 185 (1994)
21. C. Gutierrez, A solution to the bidimensional global asymptotic stability conjecture. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **12**, 627 (1995)
22. K. Ciesielski, On the Poincaré-Bendixson theorem. in *Lecture Notes in Nonlinear Analysis*, vol 3, eds. by W. Kryszewski, A. Nowakowski, *Proceedings of the 3rd Polish Symposium on Nonlinear Analysis* (2001)
23. J. Guckenheimer, P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields* (Springer-Verlag, 1983)
24. A. Cima, A.V. Essen, A. Gasull, E. Hubbers, F. Manosas, A polynomial counterexample to the Markus-Yamabe conjecture. *Adv. Math.* **131**(2), 453 (1997)
25. J.S. Muldowney, Compound matrices and ordinary differential equations. *Rocky Mt. J. Math.* **20**(4), 857 (1990)
26. M.Y. Li, L. Wang, A criterion for stability of matrices. *J. Math. Anal. Appl.* **225**, 249 (1998)
27. Scilab, A platform for numerical computation. Available at <http://www.scilab.org/>
28. W. Kafri, Robust  $D$ -stability. *Appl. Math. Lett.* **15**, 7 (2002)
29. M. Hirsch, H. Smith, Monotone dynamical systems. in *Handbook of Differential Equations: Ordinary Differential Equations*, vol 2, eds. by A. Canada, P. Drabek, A. Fonda (Elsevier, 2005)
30. D. Angeli, E. Sontag, Interconnections of monotone systems with steady-state characteristics. in *Optimal Control, Stabilization and Nonsmooth Analysis*, eds. by M.M.M. de Queiroz, P. Wolenski (Springer-Verlag, 2004)
31. D. Angeli, E. Sontag, Monotone control systems. *IEEE Trans. Automat. Contr.* **48**, 1684 (2003)
32. M.Y. Li, J.S. Muldowney, Dynamics of differential equations on invariant manifolds. *J. Differ. Equ.* **168**, 295 (2000)
33. L. Allen, T.J. Bridges, Numerical exterior algebra and the compound matrix method. Tech. rep. University of Surrey (2001)
34. T. Ström, On logarithmic norms. *SIAM J. Numer. Anal.* **12**(5), 741 (1975)
35. MAXIMA: A computer algebra system. Available at <http://maxima.sourceforge.net>