

8% inhibition of cholesterol synthesis. Activity testing continues.

Experimental Section⁶

7-Amino-3-(1,5-dimethylhexyl)dodecahydro-3a,6-dimethyl-1H-benz[e]indene-6-methylamine (II).—To a stirred solution of 4.00 g (9.52×10^{-3} mole) of I² in 35 ml of CHCl_3 was slowly added 12 ml of concd H_2SO_4 . To this was added very slowly 1.855 g (2.86×10^{-2} mole) of NaN_3 so that the temp of the solution did not exceed 45° . After the addition was complete, the mixture was warmed at $40\text{--}45^\circ$ for 15 min. The mixture was then cooled to $0\text{--}5^\circ$ and concd NH_4OH was slowly added to neutralize the acid. The resulting mixture was extracted 4 times with CHCl_3 . The extracts were evapd on a steam bath to give 3.00 g (86.5%) of the crude product. A sample of the highly hygroscopic product was recrystd using decolorizing charcoal in CHCl_3 ; mp $73\text{--}75^\circ$; $[\alpha]^{25\text{D}} + 10^\circ$ (CHCl_3).

7-Amino-3-(1,5-dimethylhexyl)dodecahydro-3a,6-dimethyl-1H-benz[e]indene-6-methylamine Dihydrochloride (IIa).—II (1 g, 2.77×10^{-3} mole) was dissolved in 15 ml of dry C_6H_6 . HCl gas was bubbled through the solu for 5 min. The white gelatinous mass was filtered and washed with C_6H_6 . The solvent was removed to give 1.05 g (87.6%) of the desired product. A sample was purified rigorously by dissolving some of the product in a minimum volume of hot H_2O , cooling the solu, and adding concd HCl . The resulting ppt was filtered, washed with dry C_6H_6 , and dried; mp $230\text{--}232^\circ$. *Anal.* ($\text{C}_{24}\text{H}_{48}\text{Cl}_2\text{N}_2$) C, H, Cl, N, neut equiv.

A series of derivatives of 7-amino-3-(1,5-dimethylhexyl)dodecahydro-3a,6-dimethyl-1H-benz[e]indene-6-methylamine (II) were prepared and characterized to confirm the functionality of the diamine. Among the derivatives prepared were the α -naphthylurea, the benzenesulfonamide, and the *p*-chlorobenzamide.

α -Naphthylurea.—II (0.2 g, 5.52×10^{-4} mole) was placed in a 25-ml flask and stoppered with a serum cap. α -Naphthylisocyanate (0.2 ml, 2.10×10^{-2} moles) was added to the diamine by injecting the sample through the serum cap with a syringe. The solution was heated at $40\text{--}50^\circ$ in a H_2O bath for 30 min. Absolute EtOH was added and the ppt filtered to give 0.300 g (77.6%) of product. A sample was recrystd from abs EtOH; mp $234\text{--}235.5^\circ$. *Anal.* ($\text{C}_{46}\text{H}_{60}\text{N}_4\text{O}_2$) C, H, N.

Benzenesulfonamide.—II (0.3 g, 8.28×10^{-4} mole) 10 ml of 10% aq NaOH and 0.50 ml (3.92×10^{-3} mole) of PhSO_2Cl were shaken vigorously, cooled, and aq HCl was added. The ppt was filtered, washed with ligroin, dried, and recrystd from EtOH to give 0.30 g (87%) of the product; mp $92\text{--}93.5^\circ$. *Anal.* ($\text{C}_{26}\text{H}_{34}\text{N}_2\text{S}_2\text{O}_4$) C, H, N, S.

***p*-Chlorobenzamide.**—To a solu of 0.50 g (1.38×10^{-3} mole) of II in 5 ml of dry $\text{C}_6\text{H}_5\text{N}$ and 10 ml of dry C_6H_6 was added a slight excess (0.60 ml, 4.75×10^{-3} mole) of *p*- $\text{ClC}_6\text{H}_4\text{COCl}$. The resulting mixture was heated on a H_2O bath at $60\text{--}70^\circ$ for 30 min, poured into 100 ml of H_2O , the C_6H_6 layer was separated and the aq layer washed with 10 ml of C_6H_6 . The combined C_6H_6 extracts were washed with H_2O and 5% aq Na_2CO_3 solu and dried (MgSO_4). The C_6H_6 solu was evapd to a small volume (3–4 ml), and hexane (20 ml) was stirred into the solu. This mixture was cooled. The solid substituted benzamide was filtered and washed with hexane. Recrystallization was effected from cyclohexane-hexane. The yield was 0.30 g (34%) mp $88\text{--}89^\circ$. *Anal.* ($\text{C}_{38}\text{H}_{52}\text{Cl}_2\text{N}_2\text{O}_2$) C, H, Cl, N.

[7-Amino-3-(1,5-dimethylhexyl)dodecahydro-3a,6-dimethyl-1H-benz[e]indene-6-methylamine]bis(ethylenediamine)cobalt(3+) Trichloride (III).—To a mixture of 1.00 g (3.51×10^{-3} mole) of *cis*-dichlorobisethylenediamine Co^{3+} chloride in 6 ml of MeOH was added a solu of 1.27 g (3.51×10^{-3} mole) of II in 10 ml of dry C_6H_6 . The mixture was stirred for 48 hr, filtered, and recrystd from H_2O –EtOH to yield 2.18 g (96%) of III; mp $240\text{--}242^\circ$ dec; λ_{max} 468 m μ ; $[\alpha]^{25\text{D}} + 2^\circ$ (H_2O). Cryoscopic particle number: Calcd, 4.00. Found, 4.06, 3.97. *Anal.* ($\text{C}_{28}\text{H}_{42}\text{Cl}_3\text{CoN}_6$) C, H, Cl, Co, N.

(6) Melting points were taken on a hot stage and are corrected. Infrared spectra were taken in KBr wafers on a Beckmann IR-12 spectrophotometer. Optical rotations were determined using a Rudolph polarimeter. Where analyses are indicated only by the symbols of the elements or functions, analytical data were within $\pm 0.3\%$ of the calculated values for those elements or functions.

Acknowledgments.—We are indebted to the National Science Foundation for support of this work under Traineeship Grant GE-7878, and we are indebted to Dr. K. L. Loening of the Chemical Abstracts Service for naming compounds II and III for us. Activity testing was done by Ayerst Laboratories.


Synthesis and Myotrophic-Androgenic Activity of 17 β -Hydroxy-5 α -androst-8(14)-en-3-one Derivatives¹

MANFRED E. WOLFF AND HONG-HSI CHANG

Department of Pharmaceutical Chemistry, School of Pharmacy, University of California, San Francisco, California 94122

Received May 21, 1970

In a previous study,² we described results which led us to suggest that the enhancement of myotrophic-androgenic activity by the 7 α -methyl group³ in steroids is due to flattening of the molecule towards the β face. An examination of molecular models revealed that a $\Delta^{8(14)}$ double bond would cause a similar effect, and the preparation of a number of 5 α -androst-8-(14)-ene derivatives (5–7) was undertaken on this basis. The compounds were prepared from 3 β ,17 β -dihydroxyandrost-8(14)-ene⁴ (1) by the methods described in the Experimental Section.



- | | |
|--|---------------------------|
| 1. $\text{R}_1 = \text{OH}; \text{R}_2 = \text{OH}$ | 8. $\text{R} = \text{H}$ |
| 2. $\text{R}_1 = \text{O}; \text{R}_2 = \text{O}$ | 9. $\text{R} = \text{Ac}$ |
| 3. $\text{R}_1 = (\text{MeO})_2; \text{R}_2 = \text{O}$ | |
| 4. $\text{R}_1 = (\text{MeO})_2; \text{R}_2 = \text{OH}$ | |
| 5. $\text{R}_1 = \text{O}; \text{R}_2 = \text{OH}$ | |
| 6. $\text{R}_1 = \text{O}; \text{R}_2 = \text{OAc}$ | |
| 7. $\text{R}_1 = \text{O}; \text{R}_2 = \text{OCOC}_2\text{H}_5$ | |

The data from the pharmacological testing⁵ are displayed in Table I. Since it appears likely that the active androgen is actually 5 α -dihydrotestosterone, 5 α -*H*- $\Delta^{8(14)}$ steroids were used in the present work. The enhancing effect of the 7 α -methyl substituent in the 5 α -*H* system was established by testing 8 and 9 which had been obtained in our previous study.² Both of these compounds were found to be far more active

(1) This investigation was supported in part by a Public Health Service Research Grant (AM-05016) from the National Institute of Arthritis and Metabolic Diseases, U. S. Public Health Service.

(2) M. E. Wolff, G. Zanati, G. Shanmugasundaram, S. Gupte, and G. Aadahl, *J. Med. Chem.*, **13**, 53 (1970).

(3) This enhancement has been reported, *inter alia* by (a) A. Segaloff, *Steroids*, **1**, 299 (1963), and J. A. Campbell, S. C. Lyster, G. W. Duncan, and J. C. Babcock, *ibid.*, **1**, 317 (1963), for testosterone derivatives; (b) G. Anner, J. Kalvoda, and P. Wieland, *Chimia*, **20**, 434 (1966), for 19-nortestosterone derivatives; and by (c) H. Kaneko, K. Nakamura, Y. Yamato, and M. Kurokawa, *Chem. Pharm. Bull. (Tokyo)*, **17**, 11 (1969), for 7 α -alkylthio and 7 α -arylthio derivatives.

(4) R. Antonucci, S. Bernstein, D. Giancola, and K. Sax, *J. Org. Chem.*, **16**, 1891 (1951).

(5) Pharmacological tests were performed at the Endocrine Laboratories, Madison, Wis., using essentially the method of L. G. Hershberger, E. G. Shipley, and R. K. Meyer, *Proc. Soc. Exp. Biol. Med.*, **83**, 175 (1953).

TABLE I
 ANDROGENIC-MYOTROPIC ASSAY

Compound (total dose, mg)	WG, mg ^a			Body wt, g	
	Ventral prostate	Seminal vesicle	Levator ani	Initial	Final
Castrate control	16.9 ± 1.07	12.0 ± 0.49	23.8 ± 2.05	54	92
Testosterone(0.5)	25.6 ± 2.47	15.1 ± 0.76	31.4 ± 2.50	55	96
<i>p</i>	<0.05	<0.05	<0.10-0.05		
Testosterone(0.6)	30.7 ± 3.50	17.2 ± 1.32	35.4 ± 1.71	55	94
<i>p</i>	<0.02	<0.05	<0.01		
Testosterone propionate(0.5)	55.5 ± 6.66	21.2 ± 1.25	34.8 ± 1.44	55	99
<i>p</i>	<0.01	<0.01	<0.02		
5 (3.0)	38.0 ± 8.62	12.5 ± 0.94	29.2 ± 2.65	54	91
<i>p</i>	<0.05	NS ^b	NS		
6 (3.0)	39.5 ± 6.38	13.1 ± 0.56	29.8 ± 3.31	54	92
<i>p</i>	<0.05	NS	NS		
7 (3.0)	39.0 ± 3.88	14.4 ± 0.59	27.6 ± 2.05	54	93
<i>p</i>	<0.01	<0.05	NS		
8 (0.3)	73.3 ± 4.64	26.5 ± 0.84	33.7 ± 1.93	54	98
<i>p</i>	<0.001	<0.001	<0.05		
9 (0.3)	111.8 ± 9.37	35.4 ± 2.84	41.4 ± 2.08	54	95
<i>p</i>	<0.001	<0.01	<0.01		

^a Mean ± standard error. ^b Not significant

than testosterone (Table I). On the other hand, the $\Delta^{8,14}$ compounds were only weakly active; in no case was potency higher than 0.2 of the corresponding testosterone activity observed. This could mean that the hypothesis of enhancement due to flattening toward the β face is incorrect. Alternatively, the presence of the double bond, or the absence of the $\delta\beta$ - or 14α -H may be responsible for the low order of activity.

Experimental Section⁶

5 α -Androsta-8(14)-3,17-dione (2).—A solution of 2 g of 1⁴ in 200 ml of Me₂CO was oxidized with Jones reagent at room temp. *i*-PrOH was added to destroy the excess Jones reagent, ice water was added, and the Me₂CO was removed under reduced pressure. The pptd powder was filtered to afford 1.8 g of product, mp 144–148°, which was recrystd from MeOH–H₂O to give a sample: mp 145–149°; nmr 0.93 (19 H₃), 1.10 (18 H₃), [α]_D²⁰ +347° (*c*, 1, CHCl₃). *Anal.* (C₁₇H₂₆O₂) C, H.

3,3-Dimethoxy-5 α -androst-8(14)-en-17-one (3).—A solution of 1.5 g of 2 and 1.5 g of SeO₂ in 60 ml of MeOH was heated at 50° for 15 min. It was cooled to room temp and a solution of 2.5 g of KOH in 20 ml of MeOH was added to make the solution alkaline. It was poured into ice water, and the pptd powder was filtered to afford 1.5 g of crude product, mp 103–104°. It was recrystd from MeOH containing 1 drop of methanolic KOH to give 3: mp 106–108°; nmr 0.70 (19 H₃), 1.07 (18 H₃), [α]_D²⁰ +222° (*c*, 1% CHCl₃). *Anal.* Calcd for C₂₁H₃₂O₃: C, 75.86; H, 9.70. Found: C, 75.10; H, 9.28.

17 β -Hydroxy-5 α -androst-8(14)-en-3-one (5).—To a solution of 1.2 g of 3 in 40 ml of MeOH was added slowly a solution of 1.2 g of NaBH₄ in 20 ml of MeOH. Ice water was added and the pptd powder was filtered to afford 1.0 g of crude 4, mp 184–187°. A solution of 1.0 g of this 4 in 5 ml of HOAc was warmed on a steam bath for 10 min and H₂O was added dropwise. The solution was cooled and the pptd powder was filtered to afford 0.7 g of 5, mp 180–183°. It was recrystd from MeOH–H₂O to give material, mp 182–183°, [α]_D²⁰ +65° (*c*, 1% CHCl₃). *Anal.* (C₁₉H₂₈O₂) C, H.

17 β -Hydroxy-5 α -androst-8(14)-en-3-one Acetate (6).—A solution of 0.1 g of 5 in 1 ml of C₆H₅N was added to 0.1 ml of Ac₂O and the mixture was kept at room temp for 24 hr. Ice water

was added and the pptd powder was filtered to afford 0.95 g of product, mp 145–147°, raised to mp 148–149° after recrystallization from MeOH, [α]_D²⁰ +41° (*c*, 1% CHCl₃). *Anal.* (C₂₁H₃₀O₃) C, H.

17 β -Hydroxy-5 α -androst-8(14)-en-3-one Propionate (7).—To a solution of 0.1 g of 5 in 1 ml of C₆H₅N was added 0.1 ml of (C₂H₅CO)₂O and the mixture was kept at room temp for 24 hr. Ice water was added and the pptd powder was filtered to afford 0.105 g of product, mp 158–163°. It was recrystd from MeOH–H₂O to give material: mp 160–162°; nmr 0.90 (19 H₃), 0.98 (18 H₃), [α]_D²⁰ +38° (*c*, 1% CHCl₃). *Anal.* (C₂₂H₃₂O₃) C, H.

Imidazole Derivatives.

Histidine Decarboxylase Inhibitors

LEO R. SWETT AND TOBIAS O. YELLIN¹

*Experimental Therapy Division, Abbott Laboratories,
North Chicago, Illinois*

Received March 13, 1970

Histamine has been implicated in a number of physiological processes,^{2,3} among them, the regulation of the microcirculation,⁴ gastric secretion,⁵ growth and repair processes,⁶ and certain hormone actions.⁷ Some clinical conditions in which histamine plays a role are anaphylaxis and allergy, wound healing, inflammation, and mastocytosis. The discovery of an inducible, specific histidine decarboxylase (HD) in mammalian tissues and the development of sensitive assays⁸ has opened up new approaches to the understanding of the physiological and pathological role of histamine. In recent years, interest in histidine decarboxylase inhibitors as tools in such studies and as agents for the treatment of disorders

(1) To whom inquiries should be directed.

(2) W. W. Douglas in the "Pharmacological Basis of Therapeutics," L. S. Goodman and A. Gilman, Ed., Macmillan Co., New York, N. Y., 1965, pp 615–627.

(3) D. M. Shepherd and D. Mackay, *Progr. Med. Chem.*, **5**, 199 (1967).

(4) R. W. Schayer, *Progr. Allergy*, **7**, 187 (1963).

(5) J. E. Fischer and S. H. Snyder, *Fed. Proc., Fed. Amer. Soc. Exp. Biol.*, **24**, 1334 (1965).

(6) G. Kahlson, *Perspect. Biol. Med.*, **5**, 179 (1962).

(7) C. M. Szego, *Fed. Proc., Fed. Amer. Soc. Exp. Biol.*, **24**, 1343 (1965).

(8) R. W. Schayer, *Methods Biochem. Anal.*, **16**, 273 (1968).

(6) Melting points were determined with a Thomas-Hoover apparatus equipped with a corrected thermometer. Microanalyses were performed by the Microanalytical Department, University of California, Berkeley, Calif. Nmr spectra were obtained at a field strength of 60 MHz on samples in CDCl₃ solution on a Varian A 60A instrument using TMS as internal standard. Optical rotations were obtained in a 0.5-dm tube with a Rudolph photoelectric polarimeter. Where analysis are indicated only by symbols of the elements or functions, analytical results obtained for those elements or functions were within ±0.4% of the theoretical values.