

was made to 0.5 ml with H₂O. The mixt was incubated for 5 min at 37°. Controls were included to correct for nonenzymatic decarboxylation.

Acknowledgment.—The authors gratefully acknowledge the support of this project by the National

Institutes of Health Grants GM-09254 and GM-01341. We wish to thank Dr. R. J. Taylor, Jr., and F. J. Leinweber, McNeil Laboratories, Fort Washington, Pa., for the histidine decarboxylase and dopa decarboxylase assays.

5-Homopyridoxals, 5-Thiopyridoxal, and Related Compounds. Synthesis, Tautomerism, and Biological Properties¹


W. KORYTNYK* AND H. AHRENS

Department of Experimental Therapeutics, Roswell Park Memorial Institute, Buffalo, New York 14203

Received January 19, 1971

Homopyridoxals with 1 or 2 additional CH₂ groups in the 5 position have been obtained by controlled oxidn of the corresponding homopyridoxols with MnO₂. More vigorous oxidn yielded the corresponding 4-homopyridoxic acids. 4-Deoxyhomopyridoxols have also been prepd from homopyridoxols by hydrogenolysis with hydrazine. Reaction of hydrazine-d₄ with pyridoxol gave 4-deoxyypyridoxol in which the α² and α⁴ Me groups and the 6 position were deuterated. These deuterations as well as the formation of 4-deoxyypyridoxol have been rationalized assuming the formation of quinone methide intermediates. 5-Thiopyridoxal was prepd and was found to be a hemiacetal in the narrow pH range in which it was stable. Likewise, the two homopyridoxals exist in a cyclic (hemiacetal) form in acid and neutral soln, whereas in an alkaline medium a marked tendency to revert to the aldehyde form, particularly in the 2 C homolog, has been observed. Derivatives of both the aldehyde and hemiacetal forms of these pyridoxal analogs have been obtained. Pyridoxal was found to undergo a Cannizzaro reaction when treated with alkali. The oximes of homopyridoxals and the 4-deoxyhomopyridoxols are inhibitors of pyridoxal phosphokinase. The effect of some of these compds on *Saccharomyces carlsbergensis*, tissue culture cells, and certain enzymes *in vivo* has also been determined.

In efforts to develop more selective antagonists of vitamin B₆ that might be active as anticancer agents² we previously synthesized a series of homologs of pyridoxol (I, R = CH₂OH) by extension and branching of the 4- and 5-hydroxymethyl side chains.^{3,4} Compds obtained by extension of the 5 position (I, R = CH₂OH; n = 2-4) were found to be inhibitors of *Saccharomyces carlsbergensis*⁴ but were ineffective in inhibiting mammalian systems.⁵



It was hoped that by modifying the 4-hydroxymethyl to a formyl (I, R = CHO; n = 2,3) or Me (I, R = CH₃; n = 2,3), improved inhibitors could be obtained, since they would more closely resemble the biologically more active form of vitamin B₆ or the well-known antimetabolite 4-deoxyypyridoxol (I, R = CH₃; n = 1), resp. (It has also been found that when 5-deoxyypyridoxol was converted to the corresponding 4-aldehyde, toxicity was increased markedly.⁶)

(1) (a) Pyridoxine Chemistry. 26. Preceding paper in this series: W. Korytnyk and B. Lachmann, *J. Med. Chem.*, **14**, 641 (1971). (b) A brief report of this study has appeared: Abstracts of the 158th National Meeting of the American Chemical Society, New York, N. Y., Sept 1969, MED148.

(2) For a review of the syn and biol activity of vitamin B₆ analogs see W. Korytnyk and M. Ikawa, *Methods Enzymol.*, **18A**, 524 (1970).

(3) W. Korytnyk and B. Paul, *J. Med. Chem.*, **13**, 187 (1970).

(4) W. Korytnyk, B. Paul, A. Bloch, and C. A. Nichol, *ibid.*, **10**, 345 (1967).

(5) Compd IX was tested against S-180 in Swiss mice fed complete or vitamin B₆ deficient diets. It was found to be inactive at doses up to 400 mg/kg per day x 7 ip or 0.025% in diets (Dr. E. Mihich, personal communication).

(6) F. Rosen, E. Mihich, and C. A. Nichol, *Vitam. Horm. (New York)*, **22**, 609 (1964).

In addn to compds of the types already mentioned, we have synthesized 5-thiopyridoxal (XVIII) and two 5-homopyridoxic acids (VIII and XIII). Chem properties of these compds, particularly ring-chain tautomerism, have been studied, and have been compared with those of pyridoxal. Some of these compds have been evaluated for their biol and enzymatic activity in several systems.

Chemistry. Synthesis.—Scheme I depicts the synthesis of the homopyridoxals (III and X), homopyridoxic acid (VIII and XIII), their derivs, and 4-deoxyhomopyridoxols (VI and XII); and Scheme II that of 5-thiopyridoxal (XVIII) and its ethyl acetal deriv.


Oxidn of the 4-CH₂OH group to the CHO and COOH groups has been carried out with MnO₂ as shown in Scheme I. Conditions for this oxidn had to be varied for each compd. Probably because of the ring-chain tautomerism of these compds (see below), the length of the side chain had a profound effect on the oxidizability of the 4-CH₂OH group. Thus conditions⁷ that had been worked out earlier for the oxidn of pyridoxol to pyridoxal and 4-pyridoxic acid could not be applied.

In the synthesis of 5-thiopyridoxal (XVIII, Scheme II), it was necessary to block the SH group of 5-thiopyridoxol by benzoylation, as in XVI. This was accomplished in a more direct manner and in better yield (from the blocked chloro derivative XIV) than has been reported previously.⁸ The most crucial step in this synthesis was the deblocking step to yield 5-thiopyridoxal from XVII. Both acid and alkaline hydrolysis of the thiobenzoate XVII gave a mixt of products, but a base-catalyzed transesterification with


(7) H. Ahrens and W. Korytnyk, *J. Heterocycl. Chem.*, **4**, 625 (1967).

(8) B. Paul and W. Korytnyk, *Tetrahedron*, **25**, 1071 (1969).

SCHEME I



SCHEME II



MeOH proved to be most satisfactory. The thiopyridoxal thus obtained was stable in a rather narrow pH range. Elemental anal. of this compd were not satisfactory, but the structure was indicated beyond any doubt by nmr spectroscopy, and by conversion to the ethyl acetal, which gave satisfactory anal. data.

Ring-Chain Tautomerism.— Ring-chain tautomerism of these compds has been studied by ir and nmr spectroscopy and by the prepn of appropriate derivs. In this regard, we have reexamined the ring-chain

tautomerism of pyridoxal in the light of more recent reports.⁹⁻¹¹

Pyridoxal has been found to exist in the hemiacetal form in both neutral and acid soln, as indicated by the

(9) W. Korytnyk and H. Ahrens, *Methods Enzymol.*, **18A**, 475 (1970).

(10) K. F. Turchin, V. F. Bystrov, M. Ya. Karpeisky, A. S. Olkhovoy, V. L. Florentiev, and Yu. N. Sheinker, "Pyridoxal Catalysis: Enzymes and Model Systems," E. R. Snell, A. E. Braunstein, E. S. Severin, and Yu. M. Torchinsky, Eds., Wiley, New York, N. Y., 1968, p 67.

(11) O. A. Gransow and R. H. Holm, *Tetrahedron*, **24**, 4477 (1968); *J. Amer. Chem. Soc.*, **91**, 5984 (1969).

presence of an AB quadruplet due to α^5 -CH₂ protons, one of which was found to be coupled to the α^4 (hemiacetal) proton.⁹ In alkaline soln the AB quadruplet collapsed to a singlet and the α^4 (hemiacetal) proton was shifted downfield. This has been explained by a fast equil between the aldehyde and hemiacetal forms by one group of investigators¹⁰ or by an equil between the hemiacetal and the hydrated form by another group.¹¹ The evidence obtained in the present study does not support the possibility of a hydrate \rightleftharpoons hemiacetal equil, but provides further arguments for the aldehyde \rightleftharpoons hemiacetal equil in addn to those adduced by us earlier.⁹ Thus pyridoxal was found to undergo a Cannizzaro reaction when heated with Ba(OH)₂, giving a mixt of pyridoxol (50%) and 4-pyridoxic acid, the latter isolated as the lactone (31%); this result is expected from the free aldehyde form. Addnl evidence of the existence of an aldehyde \rightleftharpoons hemiacetal equil was indicated by studies of the ring-chain tautomerism of the 2 homologs of pyridoxal, as about to be described.

5-Homopyridoxal (III) has also been shown to exist in the hemiacetal form in the solid state. (No C=O has been observed in its ir spectrum.) Its nmr spectrum is also consistent with the hemiacetal structure. In contrast to pyridoxal, homopyridoxal did not exhibit a shift of the hemiacetal proton in alkaline soln and hence the equil is entirely on the hemiacetal side.¹² This indication of the greater stability of the 6-membered hemiacetal ring in III as compared with the 5-membered ring in pyridoxal can be expected on the basis of Brown's I-strain theory.¹³ Homopyridoxal readily forms an ethyl acetal (IV); but a strong nucleophile, like NH₂OH, opens up the ring to give an oxime (V). Homopyridoxic acid (VIII) readily lactonizes in acid solution to VII, a behavior that is quite analogous to that of 4-pyridoxic acid.⁷

The next higher homolog of pyridoxal (X) forms a hemiacetal containing a 7-membered ring in the solid state. In D₂O, at acid and neutral pH, it also exists in the hemiacetal form; but in alkaline soln, it is converted entirely to the aldehyde form, as is shown by the aldehyde peak at -623 cps in its nmr spectrum. Here the equil between the aldehyde and hemiacetal forms as a function of pH is clearly indicated and cannot be misinterpreted as in the case of pyridoxal. Likewise, the stability of the hemiacetal rings with 5, 6, and 7 members is reflected by the equil in alkaline soln, and is in the expected order of 6 > 5 > 7.

The lactonization of homopyridoxic acids has also been compared. No tendency to form the 7-membered lactone ring in XIII could be observed, which is in sharp contrast to the 2 lower homologs (4-pyridoxic acid and VIII), which lactonize readily.

5-Thiopyridoxal was found to exist in the hemiacetal form as a solid, and in soln in the narrow pH range where it was stable. As was expected, it forms an ethyl acetal with EtOH.


4-Deoxyhomopyridoxols and Deuterations with Hydrazine-d₄.—Homopyridoxols (II and IX, resp) have also been converted to 4-deoxyhomopyridoxols (VI and XII, resp) by refluxing with anhyd N₂H₄

(12) The corresponding 5-amino analog of III has been prepared [T. L. Fisher and D. E. Metzler, *J. Amer. Chem. Soc.*, **91**, 5323 (1969)] and was also found to exist exclusively as the cyclic aldimine.

(13) H. C. Brown, R. S. Fletcher, and R. B. Johannesen, *J. Amer. Chem. Soc.*, **73**, 212 (1951).

(Scheme I). This method was introduced by Tabor-sky¹⁴ for the prepn of 4-deoxypyridoxol, and it seems to be general for this type of system.

In an effort to learn more about the nature of this reaction, we treated pyridoxol with hydrazine-d₄. Unexpectedly, we found that the resulting 4-deoxypyridoxol was almost fully deuterated in the 2- and 4-Me groups and in the 6 position, with only the 5-CH₂ protons left intact. Under the same conditions, 4-deoxypyridoxol was found to be deuterated in precisely the same manner. Thus deuteration is not introduced during hydrogenolysis of the 4-CH₂OH group. No exchange reactions were observed when γ -collidine was treated with hydrazine-d₄. Thus the phenolic OH ortho to the CH₃ and 4-CH₂OH groups activates those groups, presumably *via* quinone methide intermediates (XXa,b). Similarly, a quinonoid type of intermediate (XXc) may be invoked in explaining the proton exchange in the 6 position.¹⁵ The existence of these hypothetical quinone methide forms would also explain the selectivity of the hydrogenolysis of pyridoxol and its 5-homologs by hydrazine: refluxing with hydrazine produces an equilibrium mixture of XXa-XXc, of which only XXa is reduced to 4-deoxypyridoxol



(the diimide mechanism for reduction with hydrazine may be invoked)¹⁶ whereas the other two intermediates could readily revert to the starting pyridoxol. The process is continued until no starting material is left.

Deuteration of pyridoxol and its 4-deoxy analog provides for the first time experimental evidence of the relative lability of protons in these and related molecules. Earlier we observed facile deuteration of the 2-CH₃ group of pyridoxol *N*-methiodide in 1 *N* NaOH soln, but the 2-CH₃ group in the parent (non-quaternized) compd was found to be inert.^{17,18}

Quinone methide intermediates of the type XXb have been postulated in photophosphorylation reactions of pyridoxal Schiff bases.¹⁹ Quinone methides derived from pyridoxol, such as XXa, may explain the reactivity of its 4 position toward alcohols and in self-condensation reactions.²⁰ The same type of quinone methide (XXa) is most likely formed in the first fragmentation step from pyridoxol and pyridoxamine during mass spectrometry.²¹ Indeed, when pyridoxamine was re-

(14) R. Tabor-sky, *J. Org. Chem.*, **26**, 596 (1961).

(15) S. F. Contractor and B. Shane, *Biochemical Pharmacol.*, **19**, 1669 (1970), have studied the "metabolic" stability of ³H in labeled pyridoxol. From their experiments and ours, it would appear that the α^5 position would be the most stable for ³H labeling.

(16) (a) E. J. Corey, W. L. Mock, and D. J. Pardo, *Tetrahedron Lett.*, 347 (1961); (b) S. Hunig, H. R. Muller, and W. Thier, *ibid.*, 353 (1961).

(17) W. Korytnyk and R. P. Singh, *J. Amer. Chem. Soc.*, **85**, 2813 (1963).

(18) The reported deuteration of the 2-CH₃ of pyridoxol in D₂O [R. Hüttenrauch and K. Matthey, *Z. Chem.*, **6**, 421 (1966)] is in error.

(19) (a) K. Makino, Y. Murakami, and Y. Kobayashi, International Symposium on Pyridoxal Enzymes, Nagoya, 1967, Maruzen Co., Ltd., Tokyo, 1968, p 129; (b) Y. Kobayashi and K. Makino, *Jikeikai Med. J.*, **15**, 249 (1968).

(20) S. A. Harris, *J. Amer. Chem. Soc.*, **63**, 3363 (1941).

(21) D. C. DeJongh and W. Korytnyk, *Methods Enzymol.*, **18A**, 483 (1970), and ref cited therein.

fluxed with hydrazine, some 4-deoxypyridoxol was formed.

B. Biological Activity.—Selected compds described in this study were evaluated in various biol systems under the supervision of Drs. A. Bloch, M. Hakala, and F. Rosen of our department.

Antagonist Activity against *S. carlsbergensis*.²²—Although α^5 -homopyridoxol (II) was found⁴ to inhibit the growth of *S. carlsbergensis* by 50% at 5×10^{-8} M, the corresponding value for the 4-aldehyde (hemiacetal III) was 5×10^{-5} M; the corresponding ethyl acetal (IV) had a similar activity. Inhibitory activity was lost when this compd was converted to the carboxylic acid (VIII), which does not inhibit at 10^{-3} M. The lactone (VII) and oxime of the 3-C homolog XI also did not show any inhibitory activity at 10^{-3} M. By conversion of 5-thiobenzoylpyridoxol (XVI), which was found to be inactive at 10^{-3} M, to the corresponding 4-aldehyde XVII, the compd became active at 5×10^{-6} M. Thus, depending on the substituent in the 5 position, conversion of the 4-CH₂OH group to a related group increases or decreases the inhibitory activity of the resulting compd in relation to the parent compd.

Tissue-Culture Studies.²³—Compds VIII, XVII, and XVIII were ineffective as inhibitors of the mouse mammary adenocarcinoma cells grown in suspension in Eagle's vitamin B₆ free medium. The inactivity of the thio analog XVIII may be due to *in vivo* desulfuration, which was observed previously with 3-thiopyridoxol by Green and Montgomery.²⁴

In Vivo Systems.²⁵—No significant depression of liver L-tyrosine aminotransferase activity was seen in rats fed a vitamin B₆ deficient diet containing 20 mg% of the 4-DOP homolog XII for 5 days. Physical signs of vitamin B₆ deficiency were not aggravated, and the weight gain of the animals fed the diet containing the analog was slightly greater than that of those receiving the vitamin B₆ deficient diet only. There was no evidence of leukopenia, although 3 out of 4 animals showed a definite depression in neutrophil count. In the same experiment, homopyridoxol (II) was found to be without significant activity on lymphoid tissues, and had no effect on hepatic L-serine hydroxymethylase and L-serine dehydratase, but did depress L-tyrosine aminotransferase activity by 23%.

Inhibition of Pyridoxal Phosphokinase.—Some compds in this series were shown to be inhibitors of pyridoxal phosphokinase from rat liver.²⁶ With increasing length of the 5 side chain in 4-deoxyhomopyridoxols their binding to this enzyme progressively decreased as indicated by the following K_1 values: 4-deoxypyridoxol, 3.6×10^{-4} M; VI, 5.2×10^{-4} M, whereas XII did not bind. A conversion of the 4-CH₃ group to a 4-CH₂OH group also decreased the binding somewhat, as is shown by comparison of the K_1 value for homopyridoxol (II) (8.5×10^{-4} M) with that of the corresponding 4-deoxy compd, VI (5.2×10^{-4} M).

Oximes of homopyridoxals were shown to be more potent inhibitors of pyridoxal phosphokinase. The oxime of pyridoxal at 9.6×10^{-6} M inhibited the enzyme by 27%, the oxime of homopyridoxal (V) at 8.0×10^{-5} M by 31% (and at 8.0×10^{-4} M by 72%), and the oxime of the next higher homolog (XI) at 4.0×10^{-5} M by only 4% (and at 4×10^{-4} M by 65%). Thus a decrease of inhibitory activity with the lengthening of the 5 side chain is also indicated in the oxime series.

Experimental Section

Where analyses are indicated only by symbols of the elements, anal. results obtained for these elements were within +0.4% of the theoretical values.

Tlc was used routinely as described earlier.²⁷ Ir spectra were determined with a Perkin-Elmer 457 spectrometer, and nmr spectra with a Varian A 60A instrument, as 8-15% solutions in CDCl₃ or D₂O; positions of peaks are expressed in cps from TMS, or from sodium 3-(trimethylsilyl)-1-propanesulfonic acid as internal standards. Peaks were assigned on the basis of previous work.^{17,28}

α^5 -Homopyridoxal (III, 7-Methyl-3,4-dihydro-1H-pyrano-[4,3-c]pyridine-1,8-diol).— α^5 -Pyridoxylmethanol·HCl (II, 60 mg)⁴ was dissolved in H₂O (20 ml), to which concd H₂SO₄ (0.02 ml) was added. After addn of MnO₂ (500 mg),²⁹ the reaction mixt was shaken vigorously for 70 sec, then the MnO₂ was filtered off and washed with H₂O (25 ml). (The time for the reaction and washing should not exceed 6 min.) The filtrate was extd with CHCl₃ (3 × 50 ml), which removed α^5 -homo-4-pyridoxic acid lactone (VII), 9.8 mg (20%), mp 112-114°.

The aq layer was evapd *in vacuo* at 30°, dissolved in H₂O (1 ml), and carefully neutralized by the addn of solid NaHCO₃, when crystn occurred. The yield was 36.5 mg (74%) of material melting around 135°, which was recrystd from H₂O-Me₂CO: tlc, 1:1 MeOH-CHCl₃, R_f 0.77, retarded by boric acid²⁷ to R_f 0.46; nmr (in 1 N NaOD), (2-CH₃) -137.5, (α^5 -CH₂) -163.5 (tr J = 6 cps), (β^5 -CH₂) -192 (tr J = 6 cps), (α^4 -H) -381 (broad), (C₆-H) -422; at a neutral pH peaks appear broadened, and α^4 -H is at -364 cps; uv, $\lambda_{\max}^{0.1 N NaOH}$ 218.5 (ϵ 6700); 247 (ϵ 5800), 310 m μ (ϵ 5750); $\lambda_{\max}^{0.1 N HCl}$ 230 (ϵ 2500) shoulder, 292 m μ (ϵ 6750); $\lambda_{\max}^{pH 7}$ 255 (ϵ 4400), 312 (ϵ 7450); ir no C=O group (KBr pellet). Anal. (C₉H₁₁NO₃) C, H, N.

Ethyl Acetal of α^5 -Homopyridoxal (IV, 1-Ethoxy-7-methyl-3,4-dihydro-1H-pyrano-[4,3-c]pyridin-8-ol).— α^5 -Pyridoxylmethanol·HCl⁴ (II, 750 mg) in aq H₂SO₄ (30 ml of H₂O and 0.27 ml of concd H₂SO₄) was shaken with MnO₂¹⁵ (5.0 g) for 90 sec at room temp. The MnO₂ was filtered and washed with 40 ml of H₂O in small portions. The combined aq solns were extd with CHCl₃ (4 × 20 ml). The CHCl₃ exts yielded 133 mg of α^5 -homo-4-pyridoxic acid lactone, mp 114-116°.

The aq layer was evapd to dryness, dried at 0.1 mm, and shaken with EtOH (abs, 50 ml) for 24 hr. After filtration, the soln was evapd, and the oily material was purified on a silica gel column. The pure ethyl acetal was eluted with 1:1 CHCl₃-EtOH, giving 176 mg (25%) of the product, mp 135°. Further purification was accomplished by sublimation (0.15 Torr, 80-90°): mp 139-140°; nmr (in CDCl₃), (2-CH₃) -147, (α^4 -H) -337, (α^5 -CH₂) -166 (tr, J = 7.5 cps), (β^5 -CH₂) -233 (multiplet), (C₆-H) -472, (OEt) -80 (tr, J = 7.0 cps), -233 (multiplet); uv: $\lambda_{\max}^{0.1 N NaOH}$ 218.5 (ϵ 6600), 248 (ϵ 6400), 311 (ϵ 6650); $\lambda_{\max}^{0.1 N HCl}$ 293 (ϵ 7500), 230 (ϵ 3000 shoulder); $\lambda_{\max}^{pH 7.0}$ 254.5 (ϵ 5200), 323 (ϵ 8650). Anal. (C₁₁H₁₅NO₃) C, H, N.

The ethyl acetal (IV, 20.1 mg) was hydrolyzed by dissolving in H₂O (3 ml) and heating to 55° for 96 hr. An almost quant yield (17.3 mg) of pure hemiacetal (III) was obtained.

Oxime of 3-Hydroxy-5-(2-hydroxyethyl)-2-methylpyridine-4-carboxaldehyde (V).— α^5 -Pyridoxylmethanol·HCl (II, 210 mg) in dil acid (30 ml of H₂O plus 0.05 ml of concd H₂SO₄) was shaken with 2.0 g of MnO₂²⁹ for 90 sec at room temp. After filtration and washing of the ppt with H₂O (30 ml, in small portions), the filtrate was extd with CHCl₃ (3 × 25 ml) to remove

(22) Dr. A. Bloch, personal communication; testing procedure described in ref 4.

(23) Dr. M. Hakala, personal communication.

(24) J. L. Green and J. A. Montgomery, *J. Med. Chem.*, **7**, 17 (1964).

(25) Dr. F. Rosen, personal communication.

(26) Pyridoxal phosphokinase was obtd from rat liver as described by D. B. McCornick, M. E. Gregory, and E. E. Snell, *J. Biol. Chem.*, **236**, 2076 (1961).

(27) H. Ahrens and W. Korytnyk, *Anal. Biochem.*, **30**, 413 (1969).

(28) W. Korytnyk and B. Paul, *J. Heterocycl. Chem.*, **2**, 481 (1965).

(29) O. Mancera, G. Rosenkranz, and F. Sondheimer, *J. Chem. Soc.*, 2189 (1953).

VII. The H₂O phase was evapd to 3.5 ml, and NaOAc (1.1 g) and NH₂OH·HCl (150 mg) were added. After heating on a steam bath for 10 min, V crystd; it was kept at room temp overnight before filtration. The yield was 125 mg (67%), mp 197–198° dec. Recrystn from EtOH and from pyridine-H₂O raised the mp to 200–201° dec. Anal. (C₉H₁₂N₂O₂) C, H, N.

α⁴-Deoxy-α⁵-pyridoxylmethanol·HCl (VI).—α⁵-Pyridoxylmethanol·HCl (II, 257 mg) was refluxed with anhyd N₂H₄ (2.0 ml, purified according to the method of Taborsky¹⁴) for 18 hr, moisture being excluded. Excess N₂H₄ was distd off at 90° and 0.1 Torr. The cryst residue was taken up in EtOH (5 ml), and N₂H₄·HCl was filtered off. Addn of methanolic HCl and some Et₂O pptd additional amts of N₂H₄·HCl. The filtrate was evapd to approx 1.5 ml, and was treated with drops of Et₂O until turbid. On standing in a refrigerator for 12 hr, 122.5 mg of IV, mp 154–156°, was obtd. A further 14.1 mg was obtd from the mother liquors on addn of Et₂O, raising the yield to 137 mg (58%). Recrystn from EtOH and from MeCN raised the mp to 156–157°: nmr, (2-CH₃ and 4-CH₃) -149.5 and -160, (α⁵-CH₂) -184 (tr J = 7.0 cps), (β⁵-CH₂) -234.5 (tr J = 7.0 cps), (C₆-H) -482; uv, λ_{max}^{0.1 N HCl} 228 mμ (ε 2100), 286 (ε 7800); λ_{max}^{1 N NaOH} 245 (ε 6100), 302 (ε 6700); λ_{max}^{EtOH} 285 (ε 6950), 225 (shoulder). Anal. (C₉H₁₄ClNO₂) C, H, N.

3-Hydroxy-5-(2-hydroxyethyl)-2-methylpyridine-4-carboxylic Acid (4 → 5) Lactone (VII).—A soln of II (200 mg) in H₂O (5 ml) was added to a mixt of MnO₂ (1.65 g, prepd according to the method of Mancera, *et al.*²⁹), H₂O (15 ml), and concd H₂SO₄ (0.1 ml) and was stirred for 1 hr at room temp. MnO₂ was filtered off and washed with H₂O (30 ml). The filtrate was shaken with CHCl₃ (3 × 30 ml). The CHCl₃ ext was washed with H₂O (25 ml), dried (CaSO₄), and evapd, giving 125 mg (76%) of lactone VII, mp 119°. Recrystn from Et₂O-petr ether gave the anal. sample: mp 120°; nmr (CDCl₃); (2-CH₃) -151, (α⁵-CH₂) -183 (tr J = 6.0 cps, broad), (β⁵-CH₂) -279 (tr J = 6.0 cps, sharp), (C₆-H) -479 (broad), (3-OH) -634 (disappears on addn of D₂O); uv, λ_{max}^{0.1 N HCl} 247 (ε 2200, sh) 329 (ε 6550); ir, λ_{max}^{NaOH} 1700 cm⁻¹ (C=O). Anal. (C₉H₉NO₃) C, H, N.

3-Hydroxy-5-(2-hydroxyethyl)-2-methylpyridine-4-carboxylic Acid (VIII).—3-Hydroxy-5-(2-hydroxyethyl)-2-methylpyridine-4-carboxylic acid (4 → 5) lactone (VII, 25.0 mg) was dissd by gentle heating in 1 N NaOH (3 ml). After adding 0.1 N HCl till pH 8, the soln was evapd to 1.0 ml. More 0.1 N HCl was added, until, at pH 6, the acid crystd. After cooling in ice, the acid was filtered, yielding 16.5 mg (60%), mp 226°. Recrystn from EtOH did not raise the mp: nmr (in 1 N NaOD); (2-CH₃) -136, (α⁵-CH₂) -160 (tr J = 7.0 cps), (β⁵-CH₂) (tr J = 7.0 cps), (C₆-H) -440; λ_{max}^{0.1 N NaOH} 220 mμ (ε 10,500), 347 (ε 5500), 308 (ε 6400). Anal. (C₉H₁₁NO₄) C, H, N.

3-Hydroxy-5-(3-hydroxypropyl)-2-methylpyridine-4-carboxaldehyde (X).—(4-Hydroxymethyl)-5-(3-hydroxypropyl)-2-methyl-3-pyridinol·HCl (IX, 200 mg)³⁰ was dissolved in H₂O (25 ml). On addn of 1 N NaOH (4 ml) and MnO₂²⁹ (4.0 g), the mixt was stirred for 8 hr at room temp. The MnO₂ was filtered off and washed with H₂O (30 ml). The filtrates were evapd, giving a yellow oil. A very small amt of H₂O was added. The mixt was neutralized with 6 N HCl to pH 5.9, when crystals formed, which were filtered off and washed with H₂O. The yield was 89.1 mg, mp 124–126°. From the mother liquors, another 39.6 mg, mp 125–126°, was isolated. The total yield was 128.7 mg (76%). The first fraction of this material was converted to the hydrochloride by dissolving in dry Me₂CO (10 ml) and adding Et₂O (0.5 ml) satd with HCl gas. More Et₂O was added till cloudiness, and the product was allowed to cryst. The hydrochloride was then recrystd from Me₂CO: mp 132–135°; nmr (D₂O), pH 1–3.73 (2-CH₃) -160, (α⁴-H) -390 (C₆-H) -485; pH 8.45–11.0, (2-CH₃) -139, (C₆-H) -432 (very broad), (CHO) -623; the compd was not adequately sol in D₂O around pH 7 for an nmr study; uv, λ_{max}^{0.1 N HCl} 295 mμ (ε 8300); λ_{max}^{0.1 N NaOH} 307 (ε 1850), 394 (ε 6500), shoulder at 265 mμ; λ_{max}^{pH 7} 323 (ε 5200), 384 (ε 2750), shoulders at 249 and 266 mμ; ir, no C=O (KBr pellet). Anal. (C₁₀H₁₄ClNO₃) C, H, N, Cl.

Attempts to convert the aldehyde X to an ethyl acetal (ethanol and HCl) gave mixts of products, which were not further investigated.

Oxime of 3-Hydroxy-5-(3-hydroxypropyl)-2-methylpyridine-4-carboxaldehyde (XI).—To a soln of X·HCl (56.5 mg) in H₂O (5 ml), 1 N NaOH (1 ml) and MnO₂ (0.5 g, prepd according to the method of Mancera, *et al.*²⁹) were added, and the mixt was

stirred for 10 hr. Tlc (1:1 CHCl₃-MeOH) indicated the presence of an aldehyde (R_f 0.77) and a trace of fluorescent material (R_f 0.59). Excess MnO₂ was filtered off. NaOAc (0.2 g) and NH₂OH·HCl (50 mg) were added, and the soln was heated on a steam bath for 10 min, when the oxime pptd. After standing for 6 hr, the oxime was filtered. After drying, the yield was 32.9 mg (65%), mp 200–203°. Recrystn from pyridine raised the mp to 206–208°. Anal. (C₁₀H₁₄N₂O₃) C, H, N.

5-(3-Hydroxypropyl)-2,4-dimethyl-3-pyridinol (XII).—4-(Hydroxymethyl)-5-(3-hydroxypropyl)-2-methyl-3-pyridinol·HCl (IX, 250 mg) and N₂H₄ (95%+, 3 ml) were refluxed for 18 hr. Excess N₂H₄ was removed by distn under reduced pressure. The residue was extd with hot MeOH. The methanolic ext was concd to a small vol, when N₂H₄·HCl separated out; the latter was removed by filtration. The filtrate (MeOH ext) was evapd completely *in vacuo*, and the residue was extd several times with hot EtOAc. Combined EtOAc exts were evapd *in vacuo* to a small vol, when white cryst material septd out. The material was cooled, filtered, washed with anhyd Et₂O, and dried; yield, 155 mg (80%). The compd was crystd from EtOAc after treatment with charcoal, mp 135–136°. Anal. (C₁₀H₁₅NO₂) C, H, N.

3-Hydroxy-5-(3-hydroxypropyl)-2-methylpyridine-4-carboxylic Acid (XIII).—To 4-(hydroxymethyl)-5-(3-hydroxypropyl)-2-methyl-3-pyridinol (IX, 1.0 g) in 0.03 N HCl (70 ml), MnO₂²¹ (20 g) was added, and the reaction mixt was heated to 70°, with vigorous stirring, for 1.5 hr. The mixt was filtered (Celite Filter Aid), and the solid material was washed with H₂O and EtOH. After evapn to a small vol, the soln was applied to a Bio-Rad AG 1 × 8 anion-exchange column in the formate form (1.6 × 45 cm), and was eluted with 0.25 N HCO₂H. Fractions contg fluorescent material were combined, evapd to dryness, and crystd from EtOH. The yield was 146 mg plus 44 mg from the mother liquors (21% total); the compd decompd above 240°. Anal. (C₁₀H₁₃NO₄) C, H, N.

α⁵-S-Benzoylpyridoxthiol (XV).—Potassium thiobenzoate³² (5.0 g) in H₂O (30 ml) was added to XIV⁴ (7.92 g) in EtOH (180 ml) under N₂ for 30 min and was stirred for 1.5 hr. H₂O (150 ml) was added, and the soln was kept at 100° for 2 hr to hydrolyze the isopropylidene group. The reaction mixt was concd to 100 ml *in vacuo*, and was neutralized (NaHCO₃). The resulting ppt was collected and was washed with H₂O: yield 8.09 g (100%); mp 160–162°. The compd was identical with α⁵-S-benzoylpyridoxthiol prepared by a multistep procedure.⁸

α⁵-Thiobenzoylpyridoxal (XVII).—Compd XV (1.0 g) was dissolved in dry CHCl₃ (100 ml) and stirred for 2 hr with MnO₂²⁹ (5.0 g), moisture being excluded. The MnO₂ was filtered off and washed with CHCl₃ (100 ml), and the combined filtrates were evapd to about 10 ml, yielding 0.685 g of cryst material, plus 0.135 g from the mother liquors (81%). The anal. sample was obtained by filtration over a silica gel column (Woelm, activity grade I; EtOAc was used for elution): mp 116° dec; ir λ_{max}^{KBr} 1660 cm⁻¹ (C=O); nmr (CDCl₃), (2-CH₃) -151, (5-CH₂) -275.5, (Ph) -463 (multiplet), (C₆-H) -492.5, (α⁴-H) (aldehyde) -629, (phenolic OH) -693 cps; uv, λ_{max}^{EtOH} 271 (ε 9850), 354 mμ (ε 2460). Anal. (C₁₃H₁₃NO₃S) C, H, N.

α⁵-Thiopyridoxal (XVIII).—α⁵-Thiobenzoylpyridoxal (XVII, 200 mg) was added to 10 ml of dry MeOH containing 1% KOH, and the mixt was stirred for 2 hr. After thorough removal of MeOH *in vacuo*, the yellow residue was shaken with dry CHCl₃ (20 ml) to remove PhCO₂Me formed in the transesterification reaction. The yellow residue was dissolved in 2.3 ml of H₂O, and the pH was carefully adjusted to 9.0 with 6 N HCl (a pH meter was used), when crystn of the product began. At this point, more HCl was added, without any change in the pH, and the addn was interrupted once the pH started falling. The soln was chilled briefly in a freezer (*ca.* 10 min.), filtered, washed with ice-H₂O, and dried (P₂O₅, oil pump). The yield was 80 mg (63%). The compd decompd gradually between 150 and 170° (R_f 0.25 in EtOAc). The product was recrystd once from Me₂CO: ir spectrum, no C=O; nmr (DMSO-*d*₆) (2-CH₃) -143 (5-CH₂) -244, -264 (AB quadruplet, J = 15 cps), (α⁴-H, hemiacetal) -401.5, (C₆-H) -480 cps; uv, λ_{max}^{pH 7.0} 256 (ε 5850), 324 (ε 8950), λ_{max}^{0.1 N NaOH} 247.5 (ε 7400), 308 mμ (ε 7350). Anal. (C₈H₉NO₂S) C, H, S: calcd, 17.50; found, 15.29.

Various attempts to improve the anal. by using different sol-

(30) W. Korytnyk, *J. Med. Chem.*, **8**, 112 (1965).

(31) M. Harnfeist, A. Bavely, and W. A. Lazier, *J. Org. Chem.*, **19**, 1608 (1954).

(32) O. Kyn, *Ber.*, **32**, 3533 (1900).

vents were without success. The compd is quite unstable, and readily forms a black polymer. It was further characterized as the ethyl acetal (XIX).

α^5 -Thiopyridoxal Ethyl Acetal (XIX).— α^5 -Thiopyridoxal (XVIII, 50 mg) was suspended in EtOH (5 ml, dry), and 5 drops of Et₂O satd with HCl gas were added. The mixt was allowed to stand at room temp. The reaction was followed by tlc (EtOAc, *R_f* 0.25 starting material, 0.45 ethyl acetal). After standing for 7 days, only traces of starting material were left. The mixt was evapd to 1 ml *in vacuo*, and the product was sep'd by preparative tlc. The material was eluted with EtOAc and evapd to a small vol, and petr ether was added. Crystn yielded 6.8 mg of the ethyl acetal, hygroscopic crystals: mp 120–122° dec, nmr (CDCl₃) (CH₃CH₂) —77 (tr), (2-CH₃) —151, (CH₃CH₂) —218 (m), (5-CH₂) —257 (broad), (α^4 -H, hemiacetal) —405 (split singlet), (C₆-H) —487. *Anal.* (C₁₀H₁₃NSO₂) C, H, S.

Cannizzaro Reaction of Pyridoxal.—Pyridoxal·HCl (50 mmoles, 102 mg), dissolved in 10 ml of satd Ba(OH)₂ soln, was refluxed for 24 hr. Tlc of the reaction mixt indicated only spots due to pyridoxal and pyridoxic acid. (The identities of the products were confirmed by retardation by boric acid and a positive Gibbs test, as described earlier.²⁷) The mixt was then evapd to dryness, and was thoroughly dried *in vacuo*. The dry white powder was acetylated for gas chromatog (2.5 ml of pyridine and 2.5 ml of Ac₂O for 4 hr). Samples of this mixt were injected into a gas chromatograph operating under standard conditions.³³ Two peaks were observed, with retention times of 9.1 min (pyridoxal acetate) and 2.25 min (4-pyridoxic acid lactone acetate). Comparison of the areas under the curves with each other and with standards run separately showed that the total amts of the acetates were 24.9 mmoles for pyridoxal and 15.4 mmoles for 4-pyridoxic acid. The apparent loss of 4-pyridoxic acid during the reaction may be due to decarboxylation and degradation.

A similar mixt was obtained when pyridoxal·HCl (51 mg, 25 mmoles) was dissolved in strong NaOH soln (1 g of NaOH in 2.5 ml of water) and was heated to 110° for 24 hr.

Hydrazine-*d*₄ Experiments. (a) **4-Deoxyppyridoxol- α^2 -*d*₃, α^4 -*d*₃, α^6 -*d* from Pyridoxol.**—Pyridoxol·HCl (251 mg) and hydrazine-*d*₄ (2 ml, anhyd, supplied by Volk Radiochemical Co.) were refluxed for 16 hr, moisture being excluded. After evapd of

(33) W. Korytnyk, G. Fricke, and B. Paul, *Anal. Biochem.*, **17**, 66 (1966).

excess hydrazine [80° (0.1 mm)], the residue was extd with boiling EtOH (5 ml) for 10 min and cooled, and the hydrazine·2HCl that crystd was removed by filtration. To the filtrate, 1.5 ml of methanolic HCl (11.2% HCl) was added. On chilling, cryst 4-deoxyppyridoxol·HCl pptd. The yield was 178 mg (73%), mp 254° dec. On addn of Et₂O to the mother liquors, a further 40 mg of 4-deoxyppyridoxol could be obt'd; but it was contaminated. Recrystn of the main crop from boiling EtOH gave the pure product, mp 271° (lit.¹⁴ mp 273°), migrating as one spot on tlc (50:50 CHCl₃-MeOH; *R_f* 0.75, not retarded by boric acid). The nmr spectrum in 1 *N* D₂SO₄ shows only an α^4 -H₂ peak at —321 cps; α^2 -H₃, α^4 -H₃, and C₆-H appear as small bumps, indicating virtually complete deuteration.

(b) **4-Deoxyppyridoxol- α^2 -*d*₃, α^4 -*d*₃, α^6 -*d* from 4-Deoxyppyridoxol.**—4-Deoxyppyridoxol·HCl (251 mg) and hydrazine-*d*₄ (2.5 ml) were refluxed for 110 hr. The reaction mixt was worked up as in the preceding expt, yielding 139 mg (60%) of deuterated 4-deoxyppyridoxol·HCl. By using an internal standard and integration, it could be established that α^5 protons were not exchanged, but that α^2 -H₃, α^4 -H₃, and α^6 -H protons were exchanged to the extent of 94–95%.

(c).—Collidine (0.40 ml, pure by nmr spectroscopy) was heated with hydrazine-*d*₄ (2.0 ml) at 120° for 120 hr. On cooling, the reaction mixt sep'd into 2 layers, the upper one containing mostly γ -collidine. The nmr spectrum of this layer was exactly the same as that of the starting γ -collidine, indicating no D exchange.

Acknowledgments.—This study was supported in part by research grants (CA-08793 and CA-11047) from the National Cancer Institutes, U. S. Public Health Service. We wish to express our thanks to Drs. Harry B. Wood and Robert E. Engle of the Cancer Chemotherapy National Service Center for making available to us a quantity of α^5 -pyridoxylmethanol·HCl. Dr. B. Paul prep'd compd XII and Mr. N. Angelino compd XIII. We are also indebted to Drs. A. Bloch, M. Hakala, E. Mihich, and F. Rosen of our department for the biological evaluation of some of the compds reported, and to Mr. N. Angelino for the pyridoxal phosphokinase inhibition studies.

Antiestrogenic and Antifertility Compounds. 4. 1,1,2-Triarylalkan-1-ols and 1,1,2-Triarylalk-1-enes Containing Basic Ether Groups¹

D. J. COLLINS,* J. J. HOBBS, AND C. W. EMMENS

Department of Veterinary Physiology, University of Sydney, Sydney, New South Wales 2006, Australia

Received November 12, 1970

In an attempt to relate structure to antiestrogenic and antifertility activity, several 1,1,2-triarylalkan-1-ols and 1,1,2-triarylalk-1-enes containing a basic ether group have been synthesized, and their biological activities examined. Assignments of geometric isomerism in the triarylalkenes are made on the basis of nmr data.

The discoveries that the compds **3a**,^{2,3} **1**,^{4,5} and **2**⁶ are orally active antifertility agents, and that by sc administration they inhibit simultaneously applied estradiol,^{2,7–9} prompted us to undertake the synthesis

of **3b** and **3c**, which are previously unknown positional isomers of **3a**, and of compds **4**, which are open chain analogs of **1** and **2**. After we began this work, patents^{10,11} appeared describing some compds of the general type **4**, but these included only 2 of those described in this paper.

Chemistry.—Most of the compds were prepared by standard procedures described in the Experimental Section. Attempts to prepare 1-*p*-[2-(*N,N*-diethyl-

(1) For paper 3, see D. J. Collins and J. J. Hobbs, *Aust. J. Chem.*, **23**, 1605 (1970).

(2) L. J. Lerner, F. J. Holthaus, Jr., and C. R. Thompson, *Endocrinology*, **63**, 295 (1958).

(3) S. J. Segal and W. O. Nelson, *Proc. Soc. Exp. Biol. Med.*, **98**, 431 (1958).

(4) D. Lednicer, J. C. Babcock, S. C. Lyster, J. C. Stucki, and G. W. Duncan, *Chem. Ind. (London)*, 2098 (1961); D. Lednicer, J. C. Babcock, P. E. Marlatt, S. C. Lyster, and G. W. Duncan, *J. Med. Chem.*, **8**, 52 (1965).

(5) G. W. Duncan, J. C. Stucki, S. C. Lyster, and D. Lednicer, *Proc. Soc. Exp. Biol. Med.*, **109**, 163 (1962).

(6) G. W. Duncan, S. C. Lyster, J. J. Clark, and D. Lednicer, *ibid.*, **112**, 439 (1963).

(7) L. E. Barnes and R. K. Meyer, *Fert. Steril.*, **13**, 472 (1962).

(8) C. W. Emmens, R. I. Cox, and L. Martin, *Recent Progr. Hormone Res.*, **18**, 415 (1962).

(9) C. W. Emmens and L. Martin, *J. Reprod. Fert.*, **9**, 269 (1965).

(10) Imperial Chemical Industries Ltd., Belgian Patent 637389 (March 13, 1964); *Chem. Abstr.*, **62**, 10373 (1965).

(11) H. A. DeWald (Parke, Davis and Co.), U. S. Patent 3,288,806 (Nov 29, 1966); *Chem. Abstr.*, **66**, 37765 (1967).