

HCl. The product (0.17 g, 60%) was recrystd from *i*-PrOH, mp 300–304° dec. *Anal.* (C₇H₁₄N₅O₂Cl) C, H, N, Cl.

6-Chloro-9-(ethoxymethyl)purine.—To a suspension of 3.85 g (25.0 mmoles) of 6-chloropurine in 75 ml of DMF contg 8.0 ml (55.0 mmoles) of Et₃N was added 2.50 g (27.5 mmoles) of chloromethyl ethyl ether. After stirring the reaction mixt for 36 hr at 55–60°, it was cooled in ice and filtered. The filtrate was concd *in vacuo* to give a syrup which was chromatographed on 150.0 g of silica gel (200 mesh, column 3.0 cm i.d.) with a solvent of CHCl₃–MeOH (9.5:1) to give the crude product; yield, 3.12 g (56.3%); mp 72–75°. The analytical product was obtained by recrystg the crude material from Et₂O; yield, 2.26 g (40.8%); mp 79–81°. *Anal.* (C₈H₉ClN₄O) C, H, Cl, N.

9-Ethoxymethyladenine.—A soln of 0.64 g (3.0 mmoles) of 6-chloro-9-(ethoxymethyl)purine in 35.0 ml of 25% MeOH–NH₃ was heated at 60–70° for 18 hr in a bomb. The reaction mixt was evapd to dryness and the residue was extd with hot Me₂CO (2 × 15 ml). The Me₂CO ext was evapd *in vacuo* to give the crude product; yield 0.53 g (91.4%); mp 184–188°. One recrystn from Me₂CO and one from MeOH gave the pure product; yield, 0.49 g (84.5%); mp 190–192° (188° softens). *Anal.* (C₈H₁₁N₅O) C, H, N.

Reagents and Assay Procedures.—Adenosine deaminase (Type I, calf intestinal mucosa) was purchased from the Sigma Chemical Co. The enzyme experiments were performed at pH 7.6 in 0.05 M phosphate buffer at 25°. The *K_m* was determined by the procedure of Lineveaver and Burk.⁶ The assay for the study of reversible inhibitors has previously been described^{2a} and is a modification of the procedure of Kaplan⁷ based on the work of Kalckar.⁸

(6) H. Lineveaver and D. Burk, *J. Amer. Chem. Soc.*, **56**, 658 (1934).

(7) N. O. Kaplan, *Methods Enzymol.*, **2**, 473 (1955).

(8) H. M. Kalckar, *J. Biol. Chem.*, **167**, 461 (1947).

Xanthoquininic Acid Derivatives

LEONARD M. RICE,* BHAGVANDAS S. SHETH,
THEODORE B. ZALUCKY, AND J. W. WHEELER

College of Pharmacy and Department of Chemistry,
Howard University, Washington, D. C. 20001

Received October 24, 1970

Some 4-aminoquinolines have a high activity in the treatment of bronchial asthma^{1a–c} and malaria, as well as having antiarrhythmic^{2a–c} properties. Compound I, R = CH₃, X = 2, has antiasthmatic, antihistaminic, and antiarrhythmic activity but no antimalarial activity, whereas I, X = CCCC(C), R = C₂H₅, and chloroquine have good antimalarial activity. The latter also has antiasthmatic and antiarrhythmic activity. Thus as part of an earlier study we were interested in the modification of the quinoline nucleus to produce compds of type IIc and IIc.

The starting point in our synthesis was quininic acid (IIa) which was demethylated to xanthoquininic acid IIb³ and converted into its Et ester. This phenol was then subjected to the Mannich reaction to produce the 5-substituted derivs of type IIc. The 6-dialkylaminoalkoxy derivs IIc were prepared from the Na salt of the phenol with the appropriate dialkylaminoalkyl halides.

(1) (a) C. F. Geschickter, *Southern Med. J.*, **48**, 497 (1955). (b) P. Blanc, *Praxis*, **5**, 127 (1962). (c) R. C. Young, A. J. Murry, C. Carr, and K. A. Harden, *J. Nat. Med. Ass.*, **57**, 189 (1965).

(2) (a) D. M. Aviado, private communication, 1968. (b) M. A. Silver and D. M. Aviado, *Exp. Parasitol.*, **24**, 152 (1969). (c) J. W. Lawson, *J. Pharmacol. Exp. Ther.*, **160**, 22 (1968).


(3) H. John, *J. Prakt. Chem.*, **128**, 190 (1930).

During this work we treated 6-hydroxyquinoline-4-carboxylic acid Et ester with Br in AcOH. This readily



a, R = R'' = H; R' = CH₃

b, R = R' = H



II

c, R = secondary amino CH₂; R' = H; R'' = C₂H₅

d, R = H; R' = dialkylaminoalkyl; R'' = C₂H₅

introduced one Br atom and the product pptd out as the bromide·HBr. When boiled in H₂O this compd lost HBr and gave the free base. Introduction of the Br atom at the 5 position was confirmed by the pmr spectrum which exhibited two AB quartets in the aromatic region with *J*_{2,3} = 4.3 Hz and *J*_{7,8} = 9.2 Hz consistent with other quinoline derivatives.⁴

All compds prepared are listed in Table I. When evaluated for antiarrhythmic,^{2a–c} antimalarial,⁵ and antiasthmatic^{1a–c} activity none of the compds showed any appreciable activity.

Experimental Section

Elemental microanalyses were performed by Schwarzkopf Microanalytical Laboratory, Woodside, N. Y. Melting points of hydrates were taken on a Fisher-Johns block, those of non-hydrates in a Thomas-Hoover capillary type apparatus. Melting points are corrected. Where analyses are indicated only by symbols of the elements or functions, analytical results obtained for those elements or functions were within ±0.4% of the theoretical values.


6-Hydroxyquinoline-4-carboxylic Acid, Xanthoquininic Acid.—Quinic acid (6-methoxyquinoline-4-carboxylic acid, mp 275–278°, 101.6 g, 0.5 mole) was refluxed in 300 ml of HI, sp gr 1.7, for 12 hr. The reflux mixt was concd to 0.5 its vol, dild several times its vol with H₂O, made alk with 10% NaOH, treated with Norit A, and filtered. With stirring and cooling, the filtrate was made slightly acid with AcOH. Crystn started and was completed on standing in the refrigerator overnight. The pale yellow material was filtered, resuspended in H₂O, boiled for a few min, and filtered hot. The microcryst residue (88 g, 93%) was nearly colorless, mp 339–342° dec (lit. mp 320° dec.)³ On recrystn from DMF–H₂O, or redissolving in alk and repptg with AcOH, a colorless product was obtained which decompd at 340–342°. If placed in the oil bath at 300° decompn points as high as 347–348° were obtained. *Anal.* Calcd for C₁₀H₇NO₃: C, H, N.

6-Hydroxyquinoline-4-carboxylic Acid, Et Ester.—Xanthoquininic acid (94.5 g, 0.4 mole) was placed in a 1-l. flask, and a mixt of 450 ml of abs EtOH and 50 ml of concd H₂SO₄ was added. The whole was refluxed for 8 hr, most of the EtOH was removed at the aspirator, and the viscous dark red–brown mixt was poured with stirring into 1 l. of ice H₂O. With cooling and stirring, the mixt was neutralized with 20% NaOH and placed in the refrigerator overnight. The pale yellow ppt was filtered and washed co-

(4) L. M. Jackman and S. Sternhell, "Applications of Nuclear Magnetic Resonance Spectroscopy in Organic Chemistry," Pergamon Press, Oxford, 1969, pp 221 and 308.

(5) T. S. Osden, P. B. Russell, and L. Rane, *J. Med. Chem.*, **10**, 431 (1967).

TABLE I
XANTHOQUININIC ACID DERIVATIVES



Compd	R	R'	HCl, salt mp, °C	Formula	Analysis
1	H	H	186-187 ^a	C ₁₂ H ₁₁ NO ₃	C, H, N
2	H	H	198-200	C ₁₂ H ₁₃ ClNO ₃	C, H, N
3	Morpholinomethyl	H	153-155	C ₁₇ H ₂₂ Cl ₂ N ₂ O ₄ ·H ₂ O	C, H, Cl, N, O
4	Diethylaminomethyl	H	130-132	C ₁₇ H ₂₄ Cl ₂ N ₂ O ₃ ·H ₂ O	C, H, Cl, N, O
5	Piperidinomethyl	H	161-163	C ₁₈ H ₂₄ Cl ₂ N ₂ O ₃ ·H ₂ O	C, H, Cl, N, O ^b
6	H	2-Diethylaminoethyl	170-171.5	C ₁₈ H ₂₆ Cl ₂ N ₂ O ₃	C, H, Cl, N
7	H	2-Dimethylaminoethyl	202-203	C ₁₆ H ₂₂ Cl ₂ N ₂ O ₃ ·H ₂ O	C, H, N ^c
8	H	2-Morpholinoethyl	202.5-203.5	C ₁₈ H ₂₄ Cl ₂ N ₂ O ₄	C, H, N
9	H	2-Piperidinoethyl	181-183	C ₁₉ H ₂₆ Cl ₂ N ₂ O ₃	C, H, N
10	H	3-Diethylaminopropyl	172-175	C ₁₉ H ₂₅ Cl ₂ N ₂ O ₃	C, H, N
11	H	2-Diethylaminopropyl	160-162	C ₁₉ H ₂₅ Cl ₂ N ₂ O ₃	C, H, N
12	H	2-Dimethylaminopropyl	174-176	C ₁₇ H ₂₄ Cl ₂ N ₂ O ₃	C, H, N
13	Br	H	226-228 ^d	C ₁₂ H ₁₁ BrNO ₃	C, H, Br, N
14	Br	H	190-191 ^e	C ₁₂ H ₁₀ BrNO ₃	C, H, Br, N

^a Free base. ^b Weight loss *in vacuo*, 2 hr at 100° < 1%. Loss at 140°, 4.6%. Calcd for loss of H₂O, 4.4%. ^c Weight loss after 2 hr *in vacuo* at 100° < 1%. Loss at 140°, 4.8%. Calcd for loss of H₂O, 4.75%. ^d HBr salt.

iously with H₂O. There was obtained 82 g, 79% of product, mp 183-185°. On recrystn from EtOH-H₂O, the material had mp 186-187° (lit. mp 185.5°).³ The HCl salt of the ester was formed by bubbling dry HCl through a soln of the ester in EtOH that had been dild with Et₂O just to cloudiness and then cleared with a few drops of EtOH. It had mp 198-200° dec and was unchanged on recrystn from EtOH-Et₂O.

5-Morpholinomethyl-6-hydroxyquinoline-4-carboxylic Acid, Et Ester·2HCl (3). Experiment 1.—Xanthoquininic acid, Et ester (4.3 g, 0.02 mole) was dissolved in the minim of EtOH at room temp, 1.7 g of 40% formaldehyde soln and 2 g of morpholine were added, and the mixt was refluxed for 1 hr. The reaction mixt was evapd to dryness, the residue dissolved in abs EtOH, and an excess of alc HCl added. Addn of Et₂O to the soln gave a yellow cryst ppt, mp 153-155°. On recrystn of the material from EtOH-Et₂O, the mp spread rose to 151-157°. It was found finally that it was necessary to add a few drops of alc HCl to the recrystn solvents to narrow the mp range, mp 153-155°. The material was dried at 100° to yield 6 g (73%).

6-(2-Diethylaminoethoxy)quinoline-4-carboxylic Acid, Et Ester·2HCl (6).—Freshly cut Na (0.47 g, 0.02 g-atom) was dissolved in abs EtOH. To this soln was added 4.3 g, 0.02 mole, of Et xanthoquininate dissolved in the minim of abs EtOH. With stirring 2.7 g, 0.02 mole, of freshly distd 2-diethylaminoethyl chloride was added dropwise. The mixt was refluxed for 4 hr, cooled, and filtered to remove pptd NaCl. The EtOH was evapd and the residual oil dissolved in Et₂O. The Et₂O soln was extd repeatedly with aq 10% HCl. The aq soln of the dihydrochloride was made basic with 10% NaOH and extd 3 times with 50 ml of Et₂O. The Et₂O soln was dried (Na₂SO₄) and filtered, and the Et₂O stripped. The residual oil was dissolved in abs EtOH and an excess of alc HCl was added. The addn of Et₂O pptd the title compd, which was filtered and dried at 100°. There was obtained 6.3 g, 81% of product with mp 163-169°. Two recrystns from *i*-PrOH-Et₂O, to which 2 drops of alc HCl were added yielded anal. material, mp 170-171.5°.

5-Bromo-6-hydroxyquinoline-4-carboxylic Acid, Et Ester·HBr (13).—Ethyl xanthoquininate (5 g, 0.023 mole) was dissolved in glac AcOH and a soln of Br₂ in glac AcOH was added dropwise with stirring at room temp until the color of Br₂ persisted. The title comp pptd from the AcOH, was filtered off, washed with H₂O, and dried at 100°. There was obtained 7.4 g (85%) of compd, mp 215-220°. After 2 recrystns from EtOAc, the anal. material had mp 226-228° dec.

5-Bromo-6-hydroxyquinoline-4-carboxylic Acid, Et Ester (14).—During one attempt at the purification of **13**, 0.5 g of this material was suspended in 100 ml of H₂O, boiled for 20 min, and filtered hot. The residue, after drying at 100° had mp 185-190°. After 2 recrystns from EtOAc, it had mp 190-191°. This material gave no test for Br⁻.

Potential Psychotomimetics. Bromomethoxyamphetamines

CHARLES F. BARFKNECHT* AND DAVID E. NICHOLS¹

*Division of Medicinal Chemistry, College of Pharmacy,
The University of Iowa, Iowa City, Iowa 52240*

Received August 17, 1970

In the study of psychotomimetic amphetamines, 2,5-dimethoxy-4-methylamphetamine (DOM) is the most potent compound yet discovered (50-150 times mes-caline).² At least part of its potency is related to the nature of the para substituent. In light of Knoll's³ studies on the psychotomimetic effects of *p*-bromomethamphetamine and its cross-tolerance to LSD, the synthesis and evaluation of bromomethoxyamphetamines appeared to be a logical extension. Br has a comparable size, but different electronic character than Me. Kang and Green⁴ have recently demonstrated a correlation between the electronic character of the ring and hallucinogenic potency of methoxylated amphetamines. The substitution of Br into various ring positions of methoxylated amphetamines allows for several electronic arrangements.

Chemistry.—The general synthetic route involved preparation of the appropriately substituted benzaldehydes, condensation with EtNO₂, and reduction to the bromomethoxyamphetamines. Tables I and II summarize the compounds which have been prepared.

Attention is called to the report by Pandya and co-workers⁵ concerning the bromination of *m*-hydroxybenzaldehyde. The product of this reaction is claimed to be 3-hydroxy-4-bromobenzaldehyde; however, the

(1) NDEA Title IV fellow, 1969-present.

(2) A. T. Shulgin, T. Sargent, and C. Naranjo, *Nature (London)*, **221**, 537 (1969).

(3) J. Knoll in "Amphetamines and Related Compounds," E. Costa and S. Garattini, Ed., Raven Press, New York, N. Y., 1970, p 761.

(4) S. Kang and J. P. Green, *Nature (London)*, **226**, 645 (1970).

(5) K. C. Pandya, R. B. K. Pandya, and R. N. Singh, *J. Indian Chem. Soc.*, **29**, 363 (1952).