

(ED<sub>50</sub> > 125 mg/kg per day). The implied ability of cyclobut-A and cyclobut-G to cross the blood-brain barrier may be important in treating the neurological manifestations of AIDS. Despite its lower in vitro activity against HSV, cyclobut-A was as effective as cyclobut-G in protecting mice from a systemic challenge with 10 LD<sub>50</sub>'s of HSV-1 (Table IV). In this model, intraperitoneal (ip) therapy with either agent was initiated 7 h postinfection and then continued twice a day for 4 days. Both drugs produced 100% survival at 18.7 mg/kg per day and again appeared to be more potent than ara-A. In comparison, acyclovir produced only 40% survival at 200 mg/kg per day. Cyclobut-A and cyclobut-G were also effective in a murine CMV model (Table V). In this experiment, an ip dose of 16.7 mg/kg of either cyclobut-A, cyclobut-G, or ganciclovir twice daily for 5 days increased survival from 20% in the untreated controls to 93, 87, and 100% respectively.

The broad-spectrum activity of cyclobut-A and cyclobut-G against herpesviruses and HIV warrants further development of these compounds as possible agents for the treatment of AIDS. Preliminary studies have found neither cyclobut-A or cyclobut-G to be acutely toxic to mice at 1000 mg/kg ip. Ongoing investigations will address the mechanisms of action and pharmacodynamics of these promising agents, as well as the biological activities of the individual enantiomers.

**Acknowledgment.** The technical assistance of Bodil Lidin and Carroll Hartline is gratefully acknowledged. These studies were supported in part by Public Health Services Contract No. NO1-A1-82518 from the Antiviral Research Branch, NIAID, NIH.

† Southern Research Institute.

‡ National Cancer Institute.

§ University of Alabama at Birmingham.

**Daniel W. Norbeck,\* Earl Kern,§ Seiji Hayashi†  
William Rosenbrook, Hing Sham, Thomas Herrin  
Jacob J. Plattner, John Erickson, Jacob Clement  
Robert Swanson, Nathan Shipkowitz, Dwight Hardy  
Kennan Marsh, Gussie Arnett,† William Shannon†  
Samuel Broder,‡ Hiroaki Mitsuya†**

*Anti-infective Research Division  
Abbott Laboratories  
Abbott Park, Illinois 60064  
Southern Research Institute  
2000 Ninth Avenue, South  
Birmingham, Alabama 35255-5305  
The Clinical Oncology Program  
Division of Cancer Treatment  
Building 10, Room 13N248  
National Cancer Institute  
Bethesda, Maryland 20892  
Department of Pediatrics  
University of Alabama at Birmingham  
1670 University Boulevard  
University Station  
Birmingham, Alabama 35294*

*Received October 4, 1989*

### Hydroxyethylamine Analogues of the p17/p24 Substrate Cleavage Site Are Tight-Binding Inhibitors of HIV Protease


Sir:

A key step in the replication of the human immunodeficiency virus (HIV) occurs when HIV-1 protease, the

proteolytic enzyme encoded by the retrovirus, cleaves specific amide bonds in precursor *gag* and *pol* proteins to form the mature proteins needed for production of infectious viral particles.<sup>1</sup> Replacement of the catalytically active residues in protein precursors by site-directed mutagenesis techniques leads to the formation of noninfective virions,<sup>2</sup> and for this reason HIV protease is regarded as a potential target for developing agents for the treatment of acquired immunodeficiency syndrome (AIDS) and related diseases. Several X-ray crystal structures have established that the mature HIV protease is an aspartic proteinase that is formed from two identical 99 amino acid peptides, each subunit contributing one Asp-Thr-Gly unit.<sup>3-6</sup> The active site so formed closely resembles the active sites found in other well-characterized aspartic proteinases<sup>7-9</sup> and confirms some of the properties of HIV protease that were predicted on the basis of sequence homology.<sup>10,11</sup>

The discovery that mature HIV protease is an aspartic proteinase suggested to us and others<sup>12a-e</sup> that the general design strategy of replacing the P<sub>1</sub>-P<sub>1</sub>' cleavage point in substrates with transition-state analogues could be used to design tight-binding inhibitors of HIV protease, in the same way this approach was used to prepare inhibitors of other aspartic proteinases, e.g. pepsin, penicillopepsin, renin, cathepsin D.<sup>13</sup> We report herein the synthesis of hydroxyethylamine (HEA) dipeptidyl isosteres **1** and **2** that were designed to mimic the tetrahedral intermediate (**3**) for hydrolysis of Tyr-Pro, one of the partial substrate sequences cleaved by HIV protease (Figure 1). Incorporation of hydroxyethylamines **1** and **2** in peptides related

- (1) Krausslich, H.-G.; Wimmer, E. *Ann. Rev. Biochem.* **1988**, *57*, 701.
- (2) Seelmeier, S.; Schmidt, H.; Turk, V.; von der Helm, K. *Proc. Natl. Acad. Sci. U.S.A.* **1988**, *85*, 6612. Mous, J.; Heimer, E. P.; LeGrice, S. F. *J. Virol.* **1988**, *62*, 1433.
- (3) Wlodawer, A.; Miller, M.; Jaskolski, M.; Sathyanarayana, B. K.; Baldwin, E.; Weber, I. T.; Selk, L. M.; Clawson, L.; Schneider, J.; Kent, S. B. H. *Science* **1989**, *245*, 616.
- (4) Navia, M. A.; Fitzgerald, P.; McKeever, B.; Leu, C.-T.; Heimbach, J.; Herber, W.; Sigal, I.; Darke, P. *Nature* **1989**, *337*, 615.
- (5) Lapatto, R.; Blundell, T.; Hemmings, A.; Overington, J.; Wilderspin, A.; Wood, S.; Merson, J. R.; Whittle, P. J.; Danley, D. E.; Geoghegan, K. F.; Hawrylik, S. J.; Lee, S. E.; Scheid, K. G.; Hobart, P. M. *Nature (London)* **1989**, *342*, 299-302.
- (6) Miller, M.; Sathyanarayana, B. K.; Toth, M. V.; Marshall, G. R.; Clawson, L.; Selk, L.; Schneider, J.; Kent, S. B. H.; Wlodawer, A. *Science* **1989**, *246*, 1149.
- (7) Bott, R.; Subramanian, E.; Davies, D. R. *Biochemistry* **1982**, *21*, 6956.
- (8) James, M. N. G.; Sielecki, A. R.; Salituro, F.; Rich, D. H.; Hofmann, T. *Proc. Natl. Acad. Sci. U.S.A.* **1982**, *79*, 6137.
- (9) Blundell, T. L.; Cooper, J.; Foundling, S. I.; Jones, D. M.; Atrash, B.; Szelke, M. *Biochemistry* **1987**, *26*, 5585.
- (10) Toh, H.; Ono, M.; Saigo, K.; Miyata, T. *Nature* **1985**, *315*, 691.
- (11) Pearl, L. H.; Taylor, W. R. *Nature* **1987**, *328*, 482; *Ibid* **1987**, *329*, 351.
- (12) (a) Moore, M. L.; Bryan, W. M.; Fakhoury, S. A.; Magaard, V. W.; Huffman, W. F.; Dayton, B. D.; Meek, T. D.; Hyland, L.; Dreyer, G. B.; Metcalf, B. W.; Strickler, J. E.; Gorniak, J. G.; Debouck, C. *Biochem. Biophys. Res. Commun.* **1989**, *159*, 420. (b) Dreyer, G. B.; Metcalf, B. W.; Tomaszek, T. A., Jr.; Carr, T. J.; Chandler, A. C., III; Hyland, L.; Fakhoury, S. A.; Magaard, V. W.; Moore, M. L.; Strickler, J. E.; Debouck, C.; Meek, T. D. *Proc. Natl. Acad. Sci. U.S.A.* **1989**, *86*, 9752-9756. (c) Sigal, I. S.; Huff, J. R.; Darke, P. L.; Vacca, J. P.; Young, S. D.; Desolms, J. S.; Thompson, W. J.; Lyle, T. A.; Graham, S. L.; Ghosh, A. K., European Patent Appln. #0337714, 1988. (d) McQuade, T. J.; Tomasselli, A. G.; Lui, L.; Karacostas, V. G.; Moss, B.; Sawyer, T. K.; Heinrikson, R. L.; Tarpley, W. G. *Science* **1990**, *247*, 454-456. (e) Billich, S.; Knoop, M.-T.; Hansen, J.; Strop, P.; Sedlacek, J.; Mertz, R.; Moelling, K. *J. Biol. Chem.* **1988**, *263*, 17905-17908.




**Figure 1.** Structures of hydroxyethylamine analogues of p17/p27 Tyr-Pro substrate cleavage sites: 1, hydroxyethylamine corresponding to Phe-Pro sequence; 2, hydroxyethylamine corresponding to Leu-Pro sequence; 3, structure of a tetrahedral intermediate for the enzyme-catalyzed cleavage of substrate Tyr-Pro bonds.

to the p17/p24 substrate sequence<sup>14</sup> produces tight-binding inhibitors of HIV protease. HEA inhibitor 4 has a  $K_i = 0.66$  nM. A preliminary description of the X-ray crystal structure of inhibitor 4 bound to synthetic HIV protease has been presented.<sup>15</sup>

HEA inhibitor 4 was synthesized (Scheme I) by modification of the methods reported by Gordon and co-workers for the synthesis of other ketomethyleneamine and hydroxyethylamine inhibitors of ACE and renin.<sup>16</sup> Phenylalanine chloromethyl ketone (5)<sup>17</sup> was coupled stepwise with Boc-asparagine, Boc-leucine, Boc-(*O*-benzyl)serine, and acetyl chloride to give the corresponding protected tetrapeptide chloromethyl ketone 6. Leucine, which occupies the P<sub>3</sub> site in the p66/p51 substrate site, was placed in P<sub>3</sub> in place of glutamine for synthetic convenience. Right-hand fragment 6 was prepared from Boc-Ile-Val-OMe<sup>18</sup> by standard methods.<sup>19</sup> In the key reaction leading to 4, tetrapeptide chloromethyl ketone 6 was allowed to react with proline tripeptide 7 in DMF in the presence of NaI and NaHCO<sub>3</sub>. Ketomethyleneamine (KMA) 8 obtained by this procedure was reasonably pure, but is not stable to prolonged storage. Attempts to purify 8 and related ketones by chromatography led to significant ep-

### Scheme I<sup>a</sup>



<sup>a</sup>(a) 4 N HCl/dioxane; (b) *N*-methylmorpholine; (c) EDCl/HOBt;<sup>19</sup> (d) isobutyl chloroformate.

- (13) (a) For a discussion of the relationships between transition-state mimics and the structure of the tetrahedral intermediate, see: Rich, D. H.; Northrop, D. B. *Enzyme Kinetics in Drug Design: Implications of Multiple Forms of Enzyme on Substrate and Inhibitor Structure-Activity Correlations*. *Computer-Aided Drug Design*; Perun, T. J., Probst, C. L., Eds.; Marcel Dekker, Inc.: New York, 1989; pp 185-250. (b) Rich, D. H. *Peptidase Inhibitors*. *Comprehensive Medicinal Chemistry*; Sammes, P. G., Ed.; Pergamon Press: Oxford, Vol. 2, in press and references therein.
- (14) For a review of retroviral proteases including substrate sequences, see: Skalka, A. M. *Cell* 1989, 56, 911.
- (15) Miller, M.; Swain, A. L.; Jaskolski, M.; Sathyanarayana, B. K.; Marshall, G. R.; Rich, D. H.; Kent, S. B. H.; Wlodawer, A. In *Retroviral Proteases: Control of Maturation and Morphogenesis*; Pearl, L., Ed.; MacMillan Press: New York, in press.
- (16) (a) The hydroxyethylamine (HEA) and ketomethyleneamine (KMA) units were originally designated amino alcohols and amino ketones. Natarajan, S.; Gordon, E. M.; Sabo, E. F.; Godfrey, J. D.; Weller, H. N.; Puscec, J.; Rom, M. B.; Cushman, D. W. *Biochem. Biophys. Res. Commun.* 1984, 124, 141. Gordon, E. M.; Natarajan, S.; Pluscec, J.; Weller, H. N.; Godfrey, J. D.; Rom, M. B.; Sabo, E. F.; Engebrecht, J.; Cushman, D. W. *Biochem. Biophys. Res. Commun.* 1984, 124, 148. Gordon, E. M.; Godfrey, J. D.; Pluscec, J.; Von langen, D.; Natarajan, S. *Biochem. Biophys. Res. Commun.* 1985, 126, 419. We suggest the present nomenclature since it is derived from nomenclature currently used in the peptide mimetic field. (b) Ryono, D. E.; Free, C. A.; Neubeck, R.; Samaniego, S. G.; Godfrey, J. D.; Petrillo, E. W. In *Peptides: Structure and Function. Proceedings of the Ninth American Peptide Symposium*; Deber, C. M., Hrubby, V. J., Kopple, K. D., Eds.; Pierce Chemical Co., Rockford, IL, 1985; p 739.
- (17) Fittkau, S. J. *Prakt. Chem.* 1973, 315, 1037.
- (18) Lloyd, K.; Young, G. T. *J. Chem. Soc. C* 1971, 2890.
- (19) *The Peptides: Analysis, Synthesis, Biology*. Gross, E., Meinhofer, J., Eds.; Academic Press: New York, 1980; Vol. 1.

imerization of the carbon center  $\alpha$  to the ketone group. Consequently, crude KMA 8 was immediately reduced to hydroxyethylamines 4a,b by reaction with excess NaBH<sub>4</sub>. Purification of 4a,b by chromatography on silica gel, by eluting with methanol in chloroform, gave pure 4a,b in 46% yield as a mixture of diastereomers. The diastereomeric alcohols 4a,b did not separate upon chromatography, and their relative potencies have not yet been determined, although it is likely that the *S* diastereomer 4a is more active since this is the diastereomer present in the X-ray structure.<sup>15</sup> The other new inhibitors (9-18) listed in Table I were prepared by closely related procedures. Full details for the preparation of 4 are provided as supplementary material.<sup>32</sup>

Inhibition of either synthetic HIV protease<sup>20</sup> (in which the two Cys residues, Cys<sup>67</sup> and Cys<sup>95</sup>, have been replaced by the isosteric *L*- $\alpha$ -aminobutyric acid) or cloned material expressed in *E. coli*<sup>21</sup> was measured by using the HPLC<sup>22a</sup> and spectrofluorometric assays<sup>22b</sup> described by Toth et al. In all cases examined, both preparations gave identical results.<sup>22a</sup> Ac-Thr-Ile-Met-Met-Gln-Arg-NH<sub>2</sub> ( $K_m = 1.4$  mM,  $V_{max} = 725$  nmol/min per mg) was used as the substrate. For the inhibitor assays, a solution of the inhibitor in DMSO (final concentration 1-2%) was added to the

- (20) Schneider, J.; Kent, S. B. H. *Cell* 1988, 54, 363.
- (21) Recombinant enzyme provided by Dr. George Glover, Monsanto Co.
- (22) (a) Toth, M. V.; Chiu, F.; Glover, G.; Kent, S. B. H.; Ratner, L.; Heyden, N. V.; Green, J.; Rich, D. H.; Marshall, G. R. In *Proceedings of the 11th American Peptide Symposium*; Rivier, J., Ed.; ESCOM: Leiden, in press. (b) Toth, M. V.; Marshall, G. R., submitted for publication.

Table I. Inhibition Constants for HIV Protease Inhibitors<sup>a,b</sup>

no.	structure	$K_i$ , nM
4	Ac-Ser-Leu-Asn-[Phe-HEA-Pro]-Ile-Val-OMe	0.66 <sup>f</sup>
9	Ac-Ser-Leu-Asn-[Phe-HEA-Pro]-Ile-OMe	420
10	Ac-Leu-Asn-[Phe-HEA-Pro]-Ile-Val-OMe	21
11	Ac-Leu-Asn-[Phe-HEA-Pro]-Ile-OMe	4520
12	Ac-Ser-Leu-Asn-[Leu-HEA-Pro]-Ile-Val-OMe	11 <sup>g</sup>
13	Ac-Leu-Asn-[Leu-HEA-Pro]-Ile-Val-OMe	21
14	Ac-Ser-Leu-Asn-[Leu-HEA-Pro]-Ile-OMe	980
15	Ac-Leu-Asn-[Leu-HEA-Pro]-Ile-OMe	>10000
16	Ac-Ser-Leu-Asn-[Sta- ]-Ile-Val-OMe	1060
17	Boc-Ser-Leu-Asn-[AHPPA- ]-Ile-Val-OMe	2640
18	Ac-Ser-Leu-Asn-[AHPPA- ]-Ile-Val-OMe	8000
19	Ac-Ser-Gln-Asn-Phe[CH <sub>2</sub> N]Pro-Val-Val-NH <sub>2</sub>	13000
20	Ac-Thr-Ile-Nle[CH <sub>2</sub> NH]Nle-Gln-Arg-NH <sub>2</sub>	780
21	Ac-Ser-Gln-Asn-[AHPPA- ]-Val-Val-OMe	39000
22	Boc-Phe-[CHOHCH <sub>2</sub> ]Phe-Ile-Phe-OMe	0.6 <sup>f</sup>
23	Ala-Ala-Phe[CHOHCH <sub>2</sub> ]Gly-Val-Val-OMe	18 <sup>d</sup>
24	<sup>t</sup> BuAc-Cha[CHOHCH <sub>2</sub> ]Val-Ile-Amp	70 <sup>e</sup>

<sup>a</sup>Initial rates were determined at pH = 6.4, 37 °C, as described previously<sup>22</sup> and by using the fluorometric assay of Toth and Marshall.<sup>22b</sup> Error estimates are derived from multiple independent determinations. Each  $K_i$  represents the average of three or more independent determinations. The substrate Ac-Thr-Ile-Met-Gln-Arg-NH<sub>2</sub> was varied from 0.5 to 10 mM. Inhibitor assays contained 2–3% DMSO final concentration. <sup>b</sup>The structures of [Phe-HEA-Pro] and [Leu-HEA-Pro] are given in Figure 1. [HEA] indicates the hydroxyethylamine isostere reported by Gordon et al.<sup>16</sup>; [CH<sub>2</sub>NH] indicates the reduced amide isostere; Sta = statine, (3S,4S)-4-amino-3-hydroxy-6-methylheptanoic acid; AHPPA = (3S,4S)-4-amino-3-hydroxy-5-phenylpentanoic acid; Amp = 2-(aminomethyl)pyridine; Cha = cyclohexylalanine. <sup>c</sup>Taken from ref 12c. <sup>d</sup>Taken from ref 12b. <sup>e</sup>Taken from ref 12d. <sup>f</sup>Error = ±0.24 nM. <sup>g</sup>Error = ±5 nM.

protease, and the resulting solution was preincubated for 10 min prior to starting the reaction by the addition of substrate. Inhibition constants ( $K_i$ ; Table I) were calculated by fitting the initial velocity data to the equation for competitive inhibition as previously described for another aspartic protease.<sup>23</sup>

Systematic variation of the structure of inhibitor 4 (Table I) established that for the sequence related to the p17/p24 cleavage sequence, maximal binding was achieved with peptides which span subsites P<sub>4</sub>–P<sub>3</sub>. Replacement of the benzyl side chain in the P<sub>1</sub> position with an isobutyl group weakened binding to the enzyme. Incorporation of a P<sub>3</sub> substituent is important for tight binding to the enzyme for both series of analogues. However, binding of hydroxyethylamines derived from Leu-Pro to HIV protease was not significantly affected by incorporation of serine at P<sub>4</sub>. Other structure–activity results (data not shown) suggest that these related HEA derivatives probably bind to the HIV protease in slightly different modes in spite of their closely related structures. Ketomethyleneamine precursors of 4 and 9–15 (not shown) are weak inhibitors of HIV protease, suggesting that the hydroxyl group on the HEA unit, or a reasonable equivalent, will be needed for full inhibitory activity. Replacement of the HEA unit with statine<sup>24</sup> or the corresponding aromatic analogue of statine, AHPPA,<sup>25</sup> gave only moderately active HIV protease inhibitors (Table I). Analogues 16–18 and 21 are essentially equipotent with related inhibitors derived from other substrate sequences.<sup>12,22</sup> It is possible that any differences are due to slight variations in experimental procedures.<sup>26</sup>

Many potent inhibitors of aspartic proteinases<sup>27</sup> have been designed by application of the transition-state ana-

logue hypothesis<sup>28</sup> or by modification of the naturally occurring aspartic proteinase inhibitors<sup>29</sup> or by systematic structure–activity studies.<sup>13b</sup> Much of this work has been stimulated by the desire to develop novel inhibitors of renin for use as antihypertensive drugs.<sup>30</sup> In all cases, the most potent aspartic proteinase inhibitors contain a critical hydroxyl group (or its equivalent) that binds to the catalytic aspartic acid carboxyl groups in the enzyme–inhibitor complex. This hydroxyl group and adjacent tetrahedral carbon atom (or equivalent) are thought to mimic a reaction pathway intermediate for enzyme-catalyzed hydrolysis of amide bonds, and analogues that lack the critical hydroxyl group are orders of magnitude weaker inhibitors.<sup>13,24</sup> Our results with hydroxyethylamine inhibitor 4 are consistent with this pattern in that hydroxyethylamine inhibitor 4 is over 10 000-fold more potent than the closely related aminomethylene inhibitor 19, which lacks the hydroxyl group.<sup>20</sup> Furthermore, in the crystal structure of the HIV protease–inhibitor 4 complex described by Miller et al.<sup>15</sup> the hydroxyl group in 4 is hydrogen bonded to the two catalytically active aspartic acid residues, as is expected for an analogue that mimics a reaction pathway intermediate.<sup>13a,24</sup> We anticipate that the hydroxyl group in the hydroxyethylene isosteres<sup>31</sup> used to prepare HIV protease inhibitors 22–24<sup>12b,d,e</sup> will be bound in this enzyme's active site in a closely related fashion.

The remarkable binding of the Phe-Pro hydroxyethylamine moiety in peptide 4 to HIV protease provides an additional structural unit that can be used to generate tight-binding inhibitors of HIV protease. Hydroxyethylamine 4 is a versatile starting point for developing analogues with the pharmacodynamic properties that are required to obtain a clinically useful anti-HIV drug. The convergent route employed to synthesize 4 can be readily modified to generate analogues that probe the enzyme's binding pockets and that lead to metabolically stable analogues. Most importantly, the availability of the high-resolution crystal structure of inhibitor 4 complexed with HIV protease<sup>15</sup> provides medicinal chemists with the rare opportunity to design improved inhibitors of a therapeutically important enzyme by analyzing the crystal structure of a tight-binding inhibitor bound to the clinically relevant enzyme.

**Acknowledgment.** This work was supported by Grants AM20100 and AI27302 from the National Institutes of Health. High-resolution FAB mass spectral data were performed by the Midwest Center for Mass spectrometry, a National Science Foundation Regional Instrumentation Facility (Grant CHE 8620177).

- (23) Rich, D. H.; Sun, E. T. O. *Biochem. Pharmacol.* **1980**, *29*, 2205.  
 (24) Rich, D. H. *J. Med. Chem.* **1985**, *28*, 263.  
 (25) Rich, D. H.; Sun, E. T. O. *J. Med. Chem.* **1980**, *23*, 27.  
 (26) (a) Richards, A. D.; Broadhurst, A. V.; Ritchie, A. J.; Dunn, B. M.; Kay, J. *FEBS Lett.* **1989**, *253*, 214. (b) Richards, A. D.; Roberts, R. F.; Dunn, B. M.; Graves, M. C.; Kay, J. *FEBS Lett.* **1989**, *247*, 113.

- (27) Rich, D. H. In *Proteinase Inhibitors: Research Monographs in Cell and Tissue Physiology*; Barrett, A. J., Salvesen, G., Eds.; Elsevier Science Publishers: Amsterdam, 1986; Vol 12, pp 179–217.  
 (28) Wolfenden, R. *Acc. Chem. Res.* **1972**, *5*, 10–18. *Ibid. Ann. Rev. Biophys. Bioeng.* **1976**, *5*, 271–306.  
 (29) Boger, J.; Lohr, N. S.; Ulm, E. H.; Poe, M.; Blaine, E. H.; Fanelli, G. M.; Lin, T.-Y.; Payne, L. S.; Schorn, T. W.; Lamonte, B. I.; Vassil, T. C.; Stabilito, I. I.; Veber, D. F.; Rich, D. H.; Boparai, A. S. *Nature (London)* **1983**, *303*, 81.  
 (30) Greenlee, W. J. *Pharmaceutical Res.* **1987**, *4*, 364 and references therein.  
 (31) Szelke, M.; Leckie, B.; Hallett, A.; Jones, D. M.; Sueiras, J.; Atrash, B.; Lever, A. F. *Nature (London)* **1982**, *299*, 555–557.  
 (32) All new compounds reported here were homogeneous by TLC or HPLC and were characterized by their satisfactory IR, NMR, and high-resolution FAB mass spectral data.

**Supplementary Material Available:** Experimental procedures for the synthesis of hydroxyethylamine inhibitors **4a,b** (20 pages). Ordering information is given on any current masthead page.

\* To whom correspondence should be addressed.

† University of Wisconsin—Madison.

‡ Washington University School of Medicine.

§ Bond University.

Daniel H. Rich,<sup>\*,†</sup> Jeremy Green,<sup>†</sup> Mihaly V. Toth,<sup>‡</sup>  
Garland R. Marshall,<sup>‡</sup> Stephen B. H. Kent<sup>§</sup>

School of Pharmacy and Department of Chemistry  
University of Wisconsin—Madison  
Madison, Wisconsin 53706  
Department of Pharmacology  
Washington University School of Medicine  
St. Louis, Missouri 63110  
Graduate School of Science and Technology  
Bond University  
Queensland 4219, Australia

Received December 14, 1989

## Articles

### Hypoxia-Selective Antitumor Agents. 4. Relationships between Structure, Physicochemical Properties, and Hypoxia-Selective Cytotoxicity for Nitracrine Analogues with Varying Side Chains: The "Iminoacridan Hypothesis"

William A. Denny,<sup>\*,†</sup> Graham J. Atwell,<sup>†</sup> Robert F. Anderson,<sup>‡</sup> and William R. Wilson<sup>\*,§</sup>


Cancer Research Laboratory and Section of Oncology, Department of Pathology, University of Auckland School of Medicine, Private Bag, Auckland, New Zealand, and Cancer Research Campaign, Gray Laboratory, Mount Vernon Hospital, Northwood, Middlesex HA6 2RN, UK. Received July 5, 1989

The nitroacridine derivative nitracrine is a potent hypoxia-selective cytotoxin for mammalian cells in culture. In an attempt to modulate the degree of hypoxia selectivity among this class of compounds, we have studied a series of side-chain analogues of nitracrine. Both the electronic and steric properties of the side chain are shown to be important in determining the hypoxia selectivity of the compounds, by controlling the degree of aminoacridine/iminoacridan tautomerism. Studies with the repair-defective Chinese hamster cell line UV4 indicate that the cytotoxicity of all the compounds is due to nitro group reduction and subsequent macromolecular adduct formation. However, compounds such as the 9-amino derivative, which exist totally as the aminoacridine tautomer, form much less lethal lesions than the 9-alkylamino derivatives, which exist to varying degrees in the iminoacridan conformation. For the whole set of compounds, the degree of hypoxia-selective cytotoxicity correlates well with the proportion of iminoacridan tautomer present.

Nitracrine (**2**) is a nitroacridine DNA-intercalating ligand that has been used clinically as an anticancer drug.<sup>1,2</sup> We have previously shown<sup>3</sup> that nitracrine has potent hypoxia-selective cytotoxicity against tumor cells in culture. This property is of considerable interest, since hypoxic cells are relatively resistant to ionizing radiation and to most chemotherapeutic agents. Nitracrine represents a novel approach to the selective targeting of hypoxic tumor cells since it combines a readily reduced nitro group (the source of hypoxia-selective cytotoxicity via reductive bioactivation<sup>4</sup>) with a DNA-intercalating chromophore, which may enhance cytotoxicity by targeting reduced metabolites to DNA.

Although the net metabolic reduction of nitracrine is partially inhibited by oxygen (due to back-oxidation of the initial radical anion<sup>4</sup>), absolute rates of metabolism are high even in well-oxygenated tissue.<sup>4</sup> This leads to two problems which appear to limit activity against hypoxic cells in solid tumors. Firstly, metabolic activation in normal tissues with high nitroreductase content, such as liver,<sup>4</sup> probably contributes to host toxicity. Secondly, studies with multicellular spheroids indicate that metabolic consumption of drug in relatively well-oxygenated tissue is one

Scheme I



of the important determinants restricting diffusion to the target hypoxic microenvironments.<sup>4</sup> The development of nitroacridines with clinical utility against hypoxic tumor cells will require analogues with improved metabolic stability and/or more effective inhibition of metabolism (and cytotoxicity) by oxygen.

Improved metabolic stability has been sought by using a series of acridine-substituted analogues of nitracrine, where the nitro group reduction potentials are modulated

<sup>†</sup> Cancer Research Laboratory, University of Auckland School of Medicine.

<sup>‡</sup> Mount Vernon Hospital.

<sup>§</sup> Department of Pathology, University of Auckland School of Medicine.

(1) Radzikowski, C.; Ledochowski, A.; Hrabowska, M.; Stefanska, B.; Horowska, B.; Konopa, J. *Arch. Immunol. Ther. Exp.* **1969**, *17*, 99. *WHO Chron. (WHO, Geneva)* **1976**, *30*, No. 3 (Supplement), 11.

(2) Denny, W. A.; Baguley, B. C.; Cain, B. F.; Waring, M. J. In *Aspects of Anticancer Drug Action*; Neidle, S., Waring, M. J., Eds.; MacMillan: London, 1983; pp 1-34.

(3) Wilson, W. R.; Denny, W. A.; Twigden, S. J.; Baguley, B. C.; Probert, J. C. *B. J. Cancer* **1984**, *49*, 215.

(4) Wilson, W. R.; Denny, W. A.; Stewart, G. M.; Fenn, A.; Probert, J. C. *Int. J. Radiat. Oncol. Biol. Phys.* **1986**, *12*, 1235.