

M aqueous ammonium acetate (NH₄OAc) (pH 6.9) and CH₃CN and appropriate gradients. A flow rate of 7.0-8.0 mL/min was used. Elution of the peptides was monitored at 214 and/or 280 nm. Collected fractions were readily screened by analytical HPLC and pooled accordingly. The peptides thus obtained were subjected to rotary evaporation, in vacuo, to remove CH₃CN and then lyophilized twice. Purified peptides were analyzed for homogeneity by analytical HPLC on a μ Bondapak C₁₈ (0.39 × 15 cm, 10- μ m particle size) column using appropriate linear gradients of 0.01% aqueous TFA (pH 2.9) and 0.01% TFA-CH₃CN and of 0.01 M ammonium acetate (pH 6.9) and CH₃CN. Their amino acid composition and peptide content were assessed by quantitative amino acid analysis after acidic hydrolysis in vacuo (6 N HCl, 110 °C, 18 h), as we previously described.¹⁹ Individual amino acid recovery ranged from 0.83 to 1.08/residue, except for Cys and Trp. The molecular mass of peptides 6, 13, 19, 29, and 30 and therefore Cys and Trp integrity were assessed by FAB-MS on a Kratos MS-50 TATC instrument.

HSV-1 Ribonucleotide Reductase Assay. The inhibitory effect of the synthetic peptides, on HSV-1 RR activity, was determined as previously described¹⁹ using HSV-1 RR partially purified from quiescent BHK-21/C13 cells infected with strain F at 20 plaque-forming units per cell. The specific activity of the viral reductase preparation was 37 units/mg protein, one unit of RR being defined as the amount of enzyme generating 1 nmol of deoxycytidine 5'-diphosphate per hour under the assay conditions.

Acknowledgment. We gratefully acknowledge Claire Guilbault for preparing the RR extract and Marie-Françoise Marchal for secretarial assistance. This research was supported by the Medical Research Council of Canada and Bio-Méga Inc. Pierrette Gaudreau is the recipient of a scholarship from "Fonds de la Recherche en Santé du Québec".

Bis Basic Substituted Diaminobenzobisthiazoles as Potential Antiarthritic Agents

Ernest Cullen,*[†] Reinhold Becker,[†] Kurt Freter,[†] Thelma LeClerq,[‡] Genus Possanza,[‡] and Hin-Chor Wong[†]

Departments of Medicinal Chemistry and Pharmacology, Boehringer Ingelheim Pharmaceuticals Inc., 90 East Ridge, P.O. Box 368, Ridgefield, Connecticut 06877. Received November 2, 1990

A series of benzobisthiazoles were screened for antiinflammatory activity in the carrageenan paw edema and adjuvant arthritis tests. Compound 26, 2,6-bis(*N,N*-diethylamino)benzo[1,2-*d*:5,4-*d'*]bisthiazole, was found to inhibit the swelling of the uninjected paw in the prophylactic adjuvant arthritis model with an ED₅₀ of 2.3 mg/kg orally. As with most compounds of this series, 26 was inactive in acute model of inflammation, such as paw edema; like steroids, it showed activity in the granuloma pouch assay but did not inhibit cyclooxygenase, indicating a mode of action different from the classical nonsteroidal antiinflammatory drugs (NSAID's). At doses higher than those producing antiinflammatory activity, 26 had some immunoregulating properties.

Ever since it became clear that the classical nonsteroidal antiinflammatory drugs (NSAID's) produced gastrointestinal side effects by virtue of cyclooxygenase (CO) inhibition, research in medicinal laboratories has been directed at finding compounds which interfere with the underlying cause of the arthritic diseases. The name given to these elusive agents, "disease modifying agents" (DMA's), indicates the vagueness of the concept. One avenue of research was to concentrate on the aberrant autoimmune reaction believed to be the result of an inflammatory stimulus of unknown origin.¹

At the onset of the present work, very few chemicals were known to possess a selective mode of action on either the humoral or the cellular arms of the immune system.² One such experimental agent was tilorone (90a). Originally found to be an interferon inducer,³ it was later shown to suppress cell-mediated responses and, in contrast, to enhance antibody production in animals models.⁴ The compound was also reported to suppress adjuvant induced arthritis in rats⁴⁻⁶ and experimental allergic encephalomyelitis,^{4,6} two cell-mediated, delayed type reactions.

Tilorone has since generated an active search for bioisosteric analogues with an improved biological profile (for review see ref 7). In our laboratories, we have examined, as a potential source of antiarthritic drugs, the benzobisthiazole systems having as basic side chains (alkylamino)acetamido or related functions in lieu of the alkylamino ether moiety of tilorone. The compounds were screened in the acute paw edema and the chronic adjuvant arthritis assays. In the latter test, high activity in suppressing the secondary inflammation in the uninjected paw

was considered to be the result of interference with the immune response.⁸


Chemistry

Basic [(alkylamino)acyl]amino side chains were introduced into the diaminobenzobisthiazoles I, II, III, IV, and V (Y₁ = Y₂ = NH₂) (Tables I and II). Two general methods were used: (1) chloroacylation with excess chloroacyl chloride (or anhydride) followed by treatment of the resulting chloroacetylaminines with appropriate al-

- (1) (a) Zvaifler, N. *J. Immunol.* 1973, 16, 265. (b) Lloyd, T. M.; Panush, R. S. *J. Rheumatol.* 1977, 4, 231. (c) Fudenberg, H. H.; Wells, J. V. In *Infection and Immunity in the Rheumatic Diseases*; Dumonde, D. C., Ed.; Blackwell Scientific Publications: Oxford, 1976; p 549. (d) Zvaifler, N. J.; Silver, R. M. In *Immunology and Rheumatic Diseases*; Gupta, S., Talal, N., Eds.; Plenum Medical Books Co.: New York, 1985; p 517.
- (2) Hess, F. K.; Freter, K. R. In *Burger's Medicinal Chemistry*, 4th ed.; Wolff, M. E., Ed.; John Wiley and Sons: New York, 1979; Part II, p 671.
- (3) Krueger, R. F.; Mayer, G. D. *Science* 1970, 169, 1213.
- (4) (a) Megel, H.; Raychaudhuri, A.; Goldstein, S.; Kinsolving, C. R.; Shemano, I.; Michael, J. G. *Proc. Soc. Exp. Biol. Med.* 1974, 145, 513. (b) Megel, H.; Raychaudhuri, A.; Shemano, I.; Beaver, T. H.; Thomas, L. L. *Proc. Soc. Exp. Biol. Med.* 1975, 149, 89.
- (5) Chang, Y.-H.; Hoffman, W. W. *Pharmacologist* 1975, 17, 226.
- (6) Megel, H.; Raychaudhuri, A.; Shemano, I.; Gibson, J. P. In *Modulation of Host Immune Resistance in the Prevention or Treatment of Induced Neoplasias*; Chirigos, M. A., Ed.; U.S. Government Printing Office: Washington, DC, 1977; p 103.
- (7) (a) Radl, S.; Zikan, V. *Drug Future* 1985, 10, 215. (b) Levin, R. H.; Albrecht, W. L. In *Progress in Medicinal Chemistry*; Ellis, G. P., West, G. B., Eds.; Elsevier/North Holland Biomedical Press: Amsterdam, 1981; Vol. 18, p 135.
- (8) (a) Newbould, B. B. *Br. J. Pharmacol.* 1965, 24, 632. (b) Ward, J. R.; Cloud, R. S.; Kanitt, E. L.; Jones, R. S. *Arthr. Rheum.* 1964, 7, 654.

[†] Department of Medicinal Chemistry.

[‡] Department of Pharmacology.

Scheme I^a




	R ₁	R ₂	R ₃	R ₁	R ₂	R ₃
a	H	H	H	l	H	Cl
b	CH ₃	H	H	m	H	Br
c	C ₂ H ₅	H	H	n	H	Cl
d	C ₂ H ₅	Br	H	o	H	Cl
e	H	Br	H	p	H	Cl
f	H	Cl	H	q	H	Cl
g	H	CH ₃	H	r	H	CH ₃
h	H	OCH ₃	H	s	H	CH ₃
i	H	NO ₂	H	t	H	CH ₃
j	H	NH ₂	H	v	H	H
k	H	Cl	Cl	w	H	COOEt

^a (i) ClCO-A-Cl; (ii) H-N (method A); (iii) EtONa + EtOOC-A-N (method B). (*) See Table II reaction sequences. (A) = CH₂, CH₂CH₂, CH(CH₃). (-N) = aliphatic, heterocyclic, and carbocyclic amines. Schemes I and II illustrate routes for compounds of types I and V (Tables I and II). Types II, III, and IV (Table I) were prepared by similar routes from appropriate starting materials (Table III).

kylamines, heteroalkylamines, or heterocyclic amines (method A) and (2) aminolysis of ethyl aminoalkyl carboxylates with the diaminobenzobisthiazoles I-IV (Y₁ = Y₂ = NH₂) in the presence of sodium ethoxide⁹ (method B) (Schemes I and II). The later method was especially indicated when the competing transacylation reaction ii (method A) was favored over the desired alkylation of the amine, or when no reaction occurred. In several cases we could isolate the monoaminoacylamines which originated either from incomplete reaction (method B) or from cleavage of one of the chloroacetylamine bonds in method A. This cleavage was to some extent dependent on the solvent used. In most cases ethanol and DMF afforded

a mixture of products, while dioxane reduced the cleavage to a minimum. These monoaminoacetamides could be used to prepare a few unsymmetrical diaminoacylamides. Compound 89, for example, was prepared from 2-amino-6-[[[(diethylamino)acetyl]amino]benzo[1,2-d:5,4-d']bis-thiazole (87) by method A with piperidine. Direct monoacylation was never observed. Even when a large excess of diaminobenzobisthiazole was used, only mixtures of mono- and diacylated products resulted. Nuclear substituted products (Table II) were prepared through acylation, by method A or B, of their corresponding diaminobenzobisthiazoles (Table III), or through introduction of substituents in the proper preacylated compounds by standard methods (see reaction sequences in Table II). Starting materials listed in Table III were prepared by standard procedures^{10,11} from the appropriate phenylene-


(9) De Feoand, R. J.; Strickler, P. D. *J. Org. Chem.* 1963, 28, 2915.

Scheme II^a

	R ₁	R ₂	R ₃
X	H	Cl	H
Y	H	H	OCH ₃
Z	H	Cl	OCH ₃

^a (i) ClCO-A-Cl; (ii) H-N (method A); (iii) EtONa + EtOOC-A-N (method B). (*) See Table II reaction sequences. (A) = CH₂, CH₂CH₂, CH(CH₃). (-N) = aliphatic, heterocyclic, and carbocyclic amines. Schemes I and II illustrate routes for compounds of types I and V (Tables I and II). Types II, III, and IV (Table I) were prepared by similar routes from appropriate starting materials (Table III).


diamines. Where necessary, the linear versus angular nature of the products was determined by UV¹¹ or NMR¹² analysis. The bis[(alkylamino)alkyl]amines 85 and 86 were prepared via the corresponding thioureas 21a and 21b (Scheme III). In the same fashion the bisacetic acid 83 was obtained from ester 22c via 21c. These derivatives, however, proved to be unsuitable precursors for the synthesis of the carbamoylmethyl derivative 84, in which carbonyl and methylene were reversed as compared with the amide 26. Instead 84 was prepared by the reaction of 2,6-dichlorobenzo[1,2-*d*:5,4-*d'*]bisthiazole 23 and glycine diethylamide¹⁴ (Scheme IV). The product could also be obtained in small yield through oxidative cyclization of 1,3-bis[*N'*-(*N,N*-diethylacetamido)thioureido]benzene (21c). Attempts to obtain 85 directly by reduction of 26 were unsuccessful.

Scheme III^a

a	CH ₂ CH ₂ N(CH ₂ CH ₃) ₂	85*
b	CH(CH ₃)CH ₂ CH ₂ CH ₂ N(CH ₂ CH ₃) ₂	86*
c	CH ₂ COOCH ₂ CH ₃	
d	CH ₂ COOH	83*
e	CH ₂ CON(CH ₂ CH ₃) ₂	84*

^a (*) Compound number from Table I.

Scheme IV



Biological Results and Discussions

Tables I–III show the activities in the acute carrageenin paw edema (CPE) and in the chronic rat adjuvant arthritis (AA) tests. A prophylactic treatment protocol, as described in the Experimental Section, was used in the latter case. The percent change in body weight, in the AA test, as compared to the weight of the untreated animals, was considered a preliminary indication of the compound's toxicity.

Few compounds showed significant activity in the CPE assay even at the high screening dose of 200 mg/kg and, for these, there was no correlation with the inhibition of edema observed in the injected paw in the D-AA screen. It is well accepted that in D-AA the injected paw swelling is related to the acute phase of inflammation. Moreover when retested at 50 mg/kg, these compounds, except 40, showed no activity (compounds of Table II were not retested). The high activity of the bisamines 111 and 131 in CPE may be attributed to the stimulation of the adrenal cortex. Indeed 111 was found negative in adrenalectomized animals. In the AA screen, for the straight-chain alkyl-amino derivatives, high activity resulted with the methyl (25) and the ethyl (26, 34) while the higher homologues (27, 28, 29, 35) were inactive.

Elongation (30), branching (31), or shortening (79) of the acylamino linkage resulted in loss of activity. It was established also that the nitrogen of the amido group could not bear groups larger than methyl without compromising the activity: see 26, 32 versus 33 (Table I) and 94 versus 107 (Table II).

Bis(monoalkylamino) derivatives would present greater activity than their corresponding dialkyl analogues. See 34 versus 26; 37 and 40 versus 43 and 44, respectively. However, the monoethyl 34, at the screening dose of 50 mg/kg, caused the death of all animals at day 4 of the

- (10) Barnikow, G.; Kunzek, H.; Hofmann, M. *J. Prakt. Chem.* 1965, 27(4), 271.
 (11) Landquist, J. K. *J. Chem. Soc. C* 1967, 2212.
 (12) Solar, S. L.; Cox, R. J.; Clecak, N. J. *J. Org. Chem.* 1968, 33, 2132.
 (13) Clecak, N. J.; Cox, R. J.; Solar, S. L.; Wurster, H. K. U.S. Patent 3,489,558, 1970; *Chem. Abstr.* 1970, 73, 4974a.
 (14) Haworth, R. D.; Peacock, D. H.; Smith, W. R.; MacGillivray, R. *J. Chem. Soc.* 1952, 2972.

experiment. This is in sharp contrast with the monocycloalkylamino derivatives which induced, at the same dose, a body weight gain in the animals.

In the heterocyclic amine series significant activity was observed for quite a few piperidino (51–63) and piperazino (64–77) derivatives. With respect to ring substitution only the monomethyl- or monoethylpiperidino compounds presented activity. Rings having two methyl groups or larger, bulkier substituents had no activity with the exception of some *N*-phenyl analogues, e.g. 73.

Substitution at the tricyclic nucleus in the 4-position (Table II) in general had a negative influence; either the activity was retained but with a corresponding increase in the toxicity of the compound, as for the methyl (91), 4-chloro (93), and 4-bromo (94) derivatives, or the activity was completely abolished, as for the 4-methoxy (92), 4-nitro (95), and 4-amino (96) derivatives. No clear SAR was recognized for substitution in the 8-position nor in the 4-plus 8-positions.

Linear and angular anellation isomers of benzo[1,2-*d*:5,4-*d'*]bisthiazole (I) with the basic [(diethylamino)acetyl]amino side chains of the prototype 26 had no activity in AA; see 80, 81, 82 (Table I) and 108 (Table II). The borderline inhibition observed with 108 was probably a reflection of the severe loss in body weight induced by the compound.

The mono[(dialkylamino)acyl]amide 87 showed only weak activity in comparison with 26. This seems surprising in view of the precursor activity, discussed below.

The precursor 111 had been reported to be active in some inflammatory tests,¹¹ and we also observed potent activity in AA (Table III). However, it seems unlikely that the antiarthritic activities of the bis[(alkylamino)acyl]amines are due to *in vivo* cleavage to the diamine 111 for the following reasons: (a) It would be unusual that some (aminoacyl)amides are metabolized to the diamine and others are not. (b) The diamine 127, isomeric to 111, is highly active while the bis[(aminoacyl)amide] 80, isomeric to the prototype 26 is not. (c) Conversely, the 2,6-diamino ring-substituted analogues 113, 114, 115 (Table III) were all found inactive as such but were quite active after introduction of the [(diethylamino)acetyl]amino side chains 94, 99, and 100 (Table II). Lastly, it would appear that the presence of a carbonyl function in the side chains is a requisite for activity. Thus, the reduced analogue 85 turned out to be completely inactive. Shifting the carbonyl function as in the carbamyl derivative 84 reduced the activity considerably as compared with 26. A few selected members of the series were further evaluated in a secondary screening program comprising established adjuvant arthritis (E-AA) (with a therapeutic dosing regimen), granuloma pouch, and analgesic assays.

Interestingly, all compounds found active in the E-AA assay significantly restored the body weight of the arthritic animals. In particular, with compound 26 the increase was greater than in the polyarthritic controls. The oral ED₅₀ for this product was found to be 41 mg/kg for the injected paw and 5 mg/kg for the uninjected paw.

We could not detect any significant analgesic activity for 26 by the acetic acid writhing test. The granuloma pouch assay primarily detects steroidal compounds and as a rule gives negative results for the NSAID's. In that assay 26 significantly inhibited the exudate volume (ED₅₀ 35.7 mg/kg) and the granuloma weight (ED₅₀ 46.7 mg/kg).

Unlike typical NSAID's 26 did not induce ulceration in rats up to 200 mg/kg, the highest dose tested. As expected, the compound did not inhibit human platelet cyclooxygenase *in vitro* at concentration from 0.1 to 30 μM.¹⁵

This product inhibited the T-cell-dependent antibody production in the Jerne plaque forming cell (PFC) assay (Table IV). However, this was observed at doses at least 10 times in excess of those necessary for anti-AA activity. It is also to be noted that, in our hands, tilorone failed to enhance, in the same assay, antibody production.

Preliminary acute toxicity studies in mice, rats, rabbits, and guinea pigs established oral LD₅₀'s of 2000 mg/kg. Further studies to determine the pharmacological and immunological profile will be published elsewhere.

Conclusion

Bis basic side chains were introduced in each one of the isosteric structures of diaminobenzobisthiazoles I–V, and the compounds were evaluated for antiinflammatory activity. Only the linear structure I was found active. This activity was restricted to the bis(lower alkylamino) (e.g. 26), the bis(cycloalkylamino) (e.g. 37), and some of the bis(heterocyclic amino) derivatives (e.g. 53). Furthermore an acetamido linkage to the nucleus was a requisite for activity. These products had no significant activity in a model of acute inflammation but were orally active in model of chronic inflammation. In particular compound 26 had an oral ED₅₀ of 2.3 mg/kg in D-AA and 5.0 mg/kg in E-AA for the uninjected paw. Unlike NSAID's 26 showed activity in granuloma pouch assay, which suggested some analogy with steroids. The product had no effect on CO activity *in vitro* and as expected induced no ulceration *in vivo* in rats. Immunosuppression could be involved although inhibitory effect in Jerne plaque cells assay was seen at doses 10× the dose required for AI activity.

Experimental Section


Infrared spectra were recorded with a Perkin-Elmer Model 237B or 267 spectrophotometer using KBr disks. ¹H NMR spectra were taken in the indicated solvent with a Varian Model T-60 NMR spectrometer. UV spectra were recorded in 0.1 N HCl with a Perkin-Elmer Model 320 instrument. Melting points were determined with an electrothermal melting point apparatus and are uncorrected. Elemental analyses were performed by Micro Tech Laboratories Inc., Skokie, IL and were within ±0.4% of the calculated values except as indicated.

2,6-Bis[(*N,N*-diethylamino)acetyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (26) and 2-Amino-6-[(*N,N*-diethylamino)acetyl]amino]benzo[1,2-*d*:4,5-*d'*]bisthiazole (87). **Method A. A stirred mixture of 2,6-bis[(chloroacetyl)amino]benzo[1,2-*d*:4,5-*d'*]bisthiazole (132) (15.2 g, 0.0405 mol) and diethylamine (16 g, 0.22 mol) in dioxane (120 mL) was heated at 100 °C overnight in a pressure vessel. After cooling, the solvent was evaporated *in vacuo*, the residue was taken up in chloroform (500 mL), and the solution was washed with water and brine and dried over MgSO₄. Evaporation of this solution left 15 g of a crude solid, which was applied to a 450-g dry silica gel column and eluted with chloroform/methanol, 97/3. Fractions were pooled according to TLC. Compound 26 (9.0 g, 49.5%) eluted first. It crystallized from ethanol: mp 225–227 °C; IR 3400, 1700, 1660 sh, 1640 sh, 1615, 1550 cm⁻¹; UV λ_{max} (log ε) 214 (4.46), 277 (4.68), 319 (sh) (4.36), 332 nm (4.46); NMR (CDCl₃) δ 1.1 (t, 12 H), 2.7 (q, 8 H), 3.3 (s, 4 H), 8.15 (s, 2 H), 10.5 (s, br, 2 H ex). Anal. (C₂₀H₂₂N₆O₂S₂) C, H, N, S. Following 26, the monoacylated product 87 was collected. It melted at 203–205 °C after one crystallization from ethanol/DMF (0.30 g, 2.2%): IR 3300 sh, 3190, 3060, 1710, 1680, 1670, 1650, 1575 sh, 1555 cm⁻¹; NMR (Me₂SO-*d*₆) δ 1.0 (t, 6 H), 2.6 (q, 4 H), 3.4 (s, 2 H), 7.45 (s, br, 2 H ex), 7.7 (s, 1 H), 8.2 (s, 1 H), 8–10 (1 H, ex). Anal. (C₁₄H₁₇N₅O₂S₂) C, H, N, S.**


2,6-Bis[(chloroacetyl)amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (132). Chloroacetyl chloride (20 mL, 0.294 mol) was added portionwise, at 5–10 °C, to a vigorously stirred suspension of 2,6-diaminobenzobisthiazole¹⁰ (111) (18.5 g, 0.0832 mol) in DMF (130 mL). The mixture was then heated on a water

(15) Farina, P.; Graham, A. Personal communication.


Table I. Chemical and Biological Screening Data for *N*-Aminoacyl (25–82, 87–89), *N*-Carboxy and *N*-Carbamoylalkyl (83–84), and *N*-Aminoalkyl (85–86) Derivatives of Diaminobenzobisthiazoles^a of Structures I, II, III, and IV




I



II



III



IV



















compd	type	Y ₁ = Y ₂	method ^b	mp, °C ^c	crystn ^d solv	formula ^e	CPE/ ^f (mg/kg) % inhibn	D-adjuvant arthritis/ ^g		change body wt
								inj/ uninj	% inhibn	
25	I	NHCOCH ₂ N(CH ₃) ₂	B	288–290	I	C ₁₆ H ₂₀ N ₆ O ₂ S ₂	8 (200)	1/81 (50)*		-37
26	I	NHCOCH ₂ N(C ₂ H ₅) ₂	A, B	225–227	II	C ₂₀ H ₂₈ N ₆ O ₂ S ₂	9 (200)	76/98 (50)* ^h		-22
27	I	NHCOCH ₂ N(C ₃ H ₇) ₂	A	≥190 dec	III	C ₂₄ H ₃₆ N ₆ O ₂ S ₂ ·2HCl	46 (200)*	29/47 (50)		-11
28	I	NHCOCH ₂ N(i-C ₃ H ₇) ₂	B	208–210	II	C ₂₄ H ₃₆ N ₆ O ₂ S ₂	0 (50)	0/0 (50)		-5
29	I	NHCOCH ₂ N- (CH ₂ CH ₂ OC ₂ H ₅) ₂	A	82–84	II	C ₂₈ H ₄₄ N ₆ O ₆ S ₂	0 (50)	16/21 (50)		0
30	I	NHCOCH ₂ CH ₂ N(C ₂ H ₅) ₂	B	>300 dec	I	C ₂₂ H ₃₂ N ₆ O ₂ S ₂ ·0.1H ₂ O	16 (50)	29/48 (50)		-5
31	I	NHCOC(CH ₃)HN(C ₂ H ₅) ₂	A	230–235 dec	IV	C ₂₂ H ₃₂ N ₆ O ₂ S ₂ ·2HCl	0 (50)	0/0 (50)		-19
32	I	N(CH ₃)COCH ₂ N(C ₂ H ₅) ₂	A	244–246 dec	V	C ₂₂ H ₃₂ N ₆ O ₂ S ₂ ·2HCl	16 (200)	0/16 (50)		-16
33	I	N(C ₂ H ₅)COCH ₂ N(C ₂ H ₅) ₂	A	133–135	II	C ₂₄ H ₃₆ N ₆ O ₂ S ₂	18 (200)	14/0 (50) ⁱ		-4
34	I	NHCOCH ₂ NHC ₂ H ₅	B	230–250 dec	VI	C ₁₆ H ₂₀ N ₆ O ₂ S ₂ ·2HCl	21 (50)	79/100 (25)* ^j		-28
35	I	NHCOCH ₂ NHCH(CH ₃) ₂	B	255–257	VII	C ₁₈ H ₂₄ N ₆ O ₂ S ₂ · 0.5H ₂ O ^{k,l}	28 (50)	9/11 (50)		+7
36	I	NHCOCH ₂ NH- 	B	203–205	VII	C ₁₈ H ₂₀ N ₆ O ₂ S ₂ ·H ₂ O	18 (200)	0/16 (50)		+14
37	I	NHCOCH ₂ NH- 	B	258–260	VII	C ₂₂ H ₂₈ N ₆ O ₂ S ₂ · 0.75H ₂ O	13 (200) 9 (50)	43/70 (50)*		+10
38	I	NHCOCH ₂ NH- 	B	220–222	VII	C ₂₄ H ₃₂ N ₆ O ₂ S ₂ ^{k2}	11 (200)	11/30 (50)		-7
39	I	NHCOCH ₂ NH- 	B	238–240	VII	C ₂₄ H ₃₂ N ₆ O ₂ S ₂	36 (200)* 18 (50)	0/0 (50)		-22
40	I	NHCOCH ₂ NH- 	B	257–259 dec	VII	C ₂₄ H ₃₂ N ₆ O ₂ S ₂	20 (200) 15 (50)	3/59 (50)*		+8
41	I	NHCOCH ₂ NH- 	B	250–252 dec	VII	C ₂₆ H ₃₆ N ₆ O ₂ S ₂ ^{k3}	3 (50)	21/54 (50)		-4
42	I	NHCOCH ₂ NH- 	B	240–242	VII	C ₂₆ H ₃₆ N ₆ O ₂ S ₂ ^{k4}	0 (50)	6/59 (50)*		+1
43	I	NHCOCH ₂ N- 	A	189–191	II	C ₂₄ H ₃₂ N ₆ O ₂ S ₂	6 (50)	21/3 (50)		-18
44	I	NHCOCH ₂ N- 	A	235–237 dec	V	C ₂₆ H ₃₄ N ₆ O ₂ S ₂ ·2HCl· 1.5H ₂ O	6 (50)	32/27 (100)		-4
45	I	NHCOCH ₂ N- 	A	137–139	II	C ₃₀ H ₄₄ N ₆ O ₂ S ₂	0 (50)	22/29 (50)		-3
46	I	NHCOCH ₂ NH- 	B	295–297 dec	VIII	C ₂₄ H ₂₀ N ₆ O ₂ S ₂ ^{k5}	0 (50)	0/0 (50)		-11
47	I	NHCOCH ₂ N- 	A	290–292 dec	VII	C ₂₀ H ₂₄ N ₆ O ₂ S ₂	17 (50)	18/21 (50)		-21
48	I	NHCOCH ₂ N- 	A	263–265	VII	C ₁₈ H ₂₀ N ₆ O ₂ S ₄ ·0.5H ₂ O	11 (50)	26/27 (50)		+11
49	I	NHCOCH ₂ N- 	A	280–295 dec	VI	C ₂₀ H ₂₄ N ₆ O ₂ S ₂ ·2HCl	17 (50)	18/21 (50)		-21
50	I	NHCOCH ₂ N- 	A	303–305 dec	VII	C ₂₀ H ₂₄ N ₆ O ₂ S ₄	7 (50)	0/43 (200)		+6
51	I	NHCOCH ₂ N- 	A	293–295 dec	VI	C ₂₂ H ₂₈ N ₆ O ₂ S ₂ ·2HCl· 0.5H ₂ O	0.8 (100)	56/79 (200)*		-6
52	I	NHCOCH ₂ N- 	A	245–247	VIII	C ₂₄ H ₃₂ N ₆ O ₂ S ₂	20 (200)	70/100 (50)*		-26
53	I	NHCOCH ₂ N- 	A	235 dec	II	C ₂₄ H ₃₂ N ₆ O ₂ S ₂	0 (50)	41/82 (50)*		+6

Table I (Continued)
























compd	type	Y ₁ = Y ₂	method ^b	mp, °C ^c	crystn ^d solv	formula ^e	CPE/ ^f (mg/kg) % inhibn	D-adjuvant arthritis/ ^g	
								inj/ uninj paw (mg/kg) % inhibn	% change body wt
54	I		A	240-242	I	C ₂₄ H ₃₂ N ₆ O ₂ S ₂	19 (200)	34/63 (200)*	+6
55	I		A	201-203	II	C ₂₆ H ₃₆ N ₆ O ₂ S ₂	12 (200)	34/65 (200)*	-1
56	I		A	262-265	VII	C ₂₆ H ₃₆ N ₆ O ₂ S ₂	15 (200)	29/11 (50)	-3
57	I		A	185-187	II	C ₂₆ H ₃₆ N ₆ O ₄ S ₂ ·0.5H ₂ O	30 (200)* 6 (50)	4/5 (50)	-9
58	I		A	138-140	II	C ₂₈ H ₃₆ N ₆ O ₆ S ₂	20 (200)	6/0 (50)	-16
59	I		A	226-228	VII	C ₂₄ H ₃₂ N ₆ O ₄ S ₂	16 (200)	25/47 (50)	-4
60	I		A	110	IV	C ₃₆ H ₅₂ N ₆ O ₂ S ₂ ·0.75H ₂ O	0 (200)	0/0 (50)	+1
61	I		A	233-235	VII	C ₃₆ H ₄₀ N ₆ O ₂ S ₂	9 (200)	15/13 (50)	-16
62	I		A	288-290 dec	VII	C ₃₄ H ₃₆ N ₆ O ₂ S ₂	1 (50)	40/12 (50)	-12
63	I		A	272-274	VII	C ₃₂ N ₄₆ N ₈ O ₂ S ₂ ·0.33H ₂ O	0 (200)	46/38 (50)	-3
64	I		A	230-250 dec	VII	C ₂₀ H ₂₆ N ₈ O ₂ S ₂ ·H ₂ O	19 (50)	31/27 (50)	-9
65	I		A	256-258	II	C ₂₂ H ₃₀ N ₈ O ₂ S ₂ ·0.33H ₂ O	28 (50)	0/0 (50)	-25
66	I		B	260 dec	II	C ₂₄ H ₃₄ N ₈ O ₂ S ₂ ·4HCl	0 (200)	0/0 (50)	-6
67	I		A	238-240	VII	C ₂₄ H ₃₄ N ₈ O ₄ S ₂ ·H ₂ O	24 (200)	16/72 (50)*	0
68	I		A	225 dec	X	C ₂₆ H ₃₄ N ₈ O ₆ S ₂ ·2HCl	17 (200)	26/60 (50)*	-17
69	I		A	158-160	II	C ₂₈ H ₃₈ N ₈ O ₆ S ₂ ·H ₂ O	14 (200)	0/0 (50)	-11
70	I		B	270-275 dec	I	C ₂₄ H ₂₈ F ₆ N ₈ O ₂ S ₂	25 (200)	0/10 (50)	-13
71	I		A	247-249	VII	C ₃₄ H ₃₈ N ₈ O ₂ S ₂ ·H ₂ O	26 (200) 4 (50)	5/6 (50)	-6
72	I		A	293-295	VII	C ₃₂ H ₃₄ N ₈ O ₂ S ₂ ·0.33H ₂ O	0 (200)	40/56 (50) ⁱ	+1
73	I		A	268-270	VII	C ₃₂ H ₃₂ F ₂ N ₈ O ₂ S ₂	11 (200)	41/79 (50)*	+12
74	I		A	225-227	VII	C ₃₂ H ₃₂ Cl ₂ N ₈ O ₂ S ₂ ·H ₂ O	12 (200)	21/41 (50)	-2
75	I		A	>330	VII	C ₃₆ H ₃₈ N ₈ O ₄ S ₂ ·2H ₂ O	40 (200)* 1 (50)	0/0 (50)	-7
76	I		A	232-234	I	C ₃₄ H ₃₂ F ₆ N ₈ O ₂ S ₂	0 (200)	39/73 (50)*	+4

Table I (Continued)

compd	type	Y ₁ = Y ₂	method ^b	mp, °C ^c	crystn ^d solv	formula ^e	CPE/ ^f (mg/kg) % inhibn	D-adjvant arthritis ^{f,g}	
								inj/d	uninj/d
77	I		A	285 dec	VII	C ₃₀ H ₃₂ N ₁₀ O ₂ S ₂ ·H ₂ O	9 (200)	0/16 (50)	-2
78	I		A	245-247	II	C ₂₄ H ₃₂ N ₆ O ₂ S ₂	0 (50)	26/27 (50)	
79	I	NHCONHC ₂ H ₅	E	>330 dec	VII	C ₁₄ H ₁₆ N ₆ O ₂ S ₂	6 (200)	0/15 (50)	-1
80	II	NHCOCH ₂ N(C ₂ H ₅) ₂	A	280-315 dec	V	C ₂₀ H ₂₈ N ₆ O ₂ S ₂ ·2HCl	9 (50)	11/0 (5) ^m	-33
81	III	NHCOCH ₂ N(C ₂ H ₅) ₂	A	283-283.5	X	C ₂₀ H ₂₈ N ₆ O ₂ S ₂ ·2HCl· 0.5H ₂ O	0 (50)	2/36 (50)	-20
82	IV	N(CH ₃)COCH ₂ N(C ₂ H ₅) ₂	A	165-167	II	C ₂₂ H ₃₂ N ₆ O ₂ S ₂ · 0.25H ₂ O ^h	21 (200)	0/34 (50)	+4
83	I	NHCH ₂ COOH	C	310 dec	VII	C ₁₂ H ₁₀ N ₄ O ₄ S ₂ ^{h7}	2 (200)	0/2 (200)	-25
84	I	NHCH ₂ CON(C ₂ H ₅) ₂	D	235-237	VII	C ₂₀ H ₂₈ N ₆ O ₂ S ₂	5 (200)	58/55 (50)* ⁿ	-18
85	I	NHCH ₂ CH ₂ N(C ₂ H ₅) ₂	C	oil	IX	C ₂₀ H ₃₂ N ₆ S ₂ ·0.5H ₂ O	14 (100)	33/0 (100)	+3
86	I		C	145-150 d	XII	C ₂₈ H ₄₄ N ₆ S ₂ ·H ₂ O	0 (200)	0/0 (50)	-17
87	I	Y ₁ = NH ₂ Y ₂ = NHCOCH ₂ N(C ₂ H ₅) ₂	A	203-205	I	C ₁₄ H ₁₇ N ₅ OS ₂ ^{h8}	13 (50)	34/61 (50)*	-20
88	I	Y ₁ = NHCOCH ₂ Cl Y ₂ = NHCOCH ₂ N(C ₂ H ₅) ₂	A	280-282	II	C ₁₆ H ₁₈ ClN ₅ O ₂ S ₂	25 (200) 14 (50)	38/0 (50)	-16
89	I	Y ₁ = NHCOCH ₂ N(C ₂ H ₅) ₂ 	A	221-223	XI	C ₂₁ H ₂₈ N ₆ O ₂ S ₂	20 (50)	50/58 (50)*	-14
90a	tilorone						o	59/45 (100)* ^p 53/0 (50)	+10 -8
90b	cyclophosphamide						18 (100) ^q	86/100 (25)* ^d	-19
90c	indomethacin						63 (10)*	71/83 (2.5)*	+18
							34 (3)*	61/69 (1.0)*	+28

^a [I, [1,2-*d*:5,4-*d'*]; type II, [1,2-*d*:4,5-*d'*]; type III, [1,2-*d*:4,3-*d'*]; type IV, [2,1-*d*:3,4-*d'*]; type V (see Table II), [1,2-*d*:3,4-*d'*]. ^b A and B, Schemes I and II; C, Scheme III; D, Scheme IV; E, see Experimental Section. ^c Uncorrected. ^d I, EtOH-DMF; II, EtOH; III, ethanol-ether; IV, free base crystallized from ethanol and treated with ethereal HCl; V, the free base as eluted from a SiO₂ column was treated with HCl; VI, EtOH-H₂O; VII, DMF; VIII, EtOH-CHCl₃; IX, CH₃OH; X, H₂O; XI, CH₃CN; XII, see Experimental Section. ^e All analyses were within ±0.4% of theoretical value except where indicated. ^f Animals were dosed orally. See Experimental Section for description of screening tests. ^g Statistically significant results with two-tail student's tests (*p* < 0.05) are marked with an asterisk (*). ^h The oral ED₅₀ was established for the uninjected paw at 2.3 (0.8-6.9) mg/kg, *p* = 0.05. ⁱ 2/6 animals died. ^j At 50 mg/kg, all animals died. ^{k1} H: calcd, 5.86; found, 5.43. ^{k2} H: calcd, 6.44; found, 6.03. ^{k3} S: calcd, 12.13; found, 12.55. ^{k4} S: calcd, 12.13; found, 11.70. ^{k5} N: calcd, 17.20; found, 16.72. ^{k6} S: calcd, 13.32; found, 13.77. ^{k7} S: calcd, 18.95; found, 18.53. ^{k8} N: calcd, 20.88; found, 21.35. ^l 1/6 animals died. ^m Compound was screened at 1/5 the LD₅₀. ⁿ At 200 mg/kg, all rats died between day 8 and day 12. ^o Mengel et al., see ref 4b, reported the ED₇₀ at 82 mg/kg, the compound being administered 24 and 1 h prior to the carrageenan challenge. ^p At 100 mg/kg p.o., significant inhibition in the injected paw and complete inhibition in the noninjected paw was reported by Mengel et al., see refs 4a and 7b. ^q Doses were administered subcutaneously.

bath for 1 h. A solid crystallized and was collected after cooling. It was treated with sodium bicarbonate solution, washed well with water, and dried. The product (23 g, 73.6%) was sufficiently pure (by TLC) to be used as such. It crystallized from DMF/H₂O: mp 320 °C; IR 3220, 1685, 1620, 1560 cm⁻¹; NMR (Me₂SO-*d*₆) δ 4.4 (s, 4 H), 8.0 (s, 1 H), 8.4 (s, 1 H), 12.9 (s, br, 2 H ex). Anal. (C₁₂H₈Cl₂N₄O₂S₂) C, H, Cl, N, S.

4-Bromo-8-chloro-2,6-bis[[(*N,N*-diethylamino)acetyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (104). Method B. A mixture of crude 2,6-diamino-4-bromo-8-chlorobenzo[1,2-*d*:5,4-*d'*]bisthiazole (124) (2.6 g, 0.01 mol), 30 mL of 1 M sodium ethoxide solution in ethanol (0.03 mol), and ethyl *N,N*-diethylglycinate (4.5 g, 0.028 mol) was stirred at room temperature overnight. The residue obtained on evaporation of the resulting solution was dissolved in water (200 mL), neutralized with 2 N HCl, and extracted with chloroform. The extract was washed, dried, and taken to near dryness in vacuo. The concentrated solution was then chromatographed on a 60-g dry silica gel column, using chloroform as eluant. The first eluates (TLC) contained the title product (2.1 g, 37%). It crystallized from DMF/chloroform: mp 265-267 °C; IR 3430, 3270, 1700, 1670 sh, 1600, 1580 cm⁻¹; NMR (CDCl₃) δ 1.2 (t, 6 H), 2.7 (q, 4 H), 10.3 (s, br, 2 H ex); UV λ_{max} (log ε) 220 (4.34), 278 (sh) (4.60), 286 (4.72), 323 (sh) (4.28), 338 nm (4.20). Anal. (C₂₀H₁₆BrClN₆O₂S₂) C, H, S, Br.

4,8-Dichloro-2,6-bis[[(*N,N*-diethylamino)acetyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (98). Chlorine (6.0 g, 0.085 mol) was absorbed into trimethyl phosphate (TMP) (50 mL) at -10 °C, and the solution was added to a stirred suspension of 2,6-bis[[(*N,N*-diethylamino)acetyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (26) (7.0 g, 0.0156 mol) in TMP (50 mL) at -10 °C. The resulting solution was kept at 0-5 °C for 8 h, poured into ice/water (150 mL), and neutralized with ammonium hydroxide. The pale off-white solid was collected, washed with water, dried, dissolved in methylene chloride, and treated with charcoal. The colorless solution was evaporated to dryness, and the residue was crystallized twice from DMF to yield 3.5 g (44%) of compound 98: mp 278-280 °C; IR 3450, 3220, 1710 sh, 1700, 1610, 1550 cm⁻¹. NMR (CDCl₃) δ 1.1 (t, 12 H), 2.7 (q, 8 H), 3.4 (s, 4 H), 10.4 (s, br, 2 H ex). Anal. (C₂₀H₂₆Cl₂N₆O₂S₂) C, H, Cl, N, S.

2,6-Bis[(chloroacetyl)amino]-4-nitrobenzo[1,2-*d*:5,4-*d'*]bisthiazole (137). Fuming nitric acid (10 mL) was added dropwise to a stirred suspension of 2,6-bis[(chloroacetyl)amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (132) (1.8 g, 4.8 mmol) in acetic acid (50 mL) with heating on a water bath. The reaction mixture was then heated to reflux for 0.5 h, cooled, and diluted with ice water. The precipitated solid was collected and crystallized twice from DMF to give 137: 1.2 g (60%); mp 300 °C; IR 3310, 3210, 1700, 1515 cm⁻¹; NMR (Me₂SO-*d*₆) δ 4.5 (s, 4 H),

Table II. Chemical and Biological Screening Data for Substituted 2,6-[[Diethylamino)acetyl]amino]benzo[1,2-d:5,4-d']bisthiazoles (90-107) and 2,7-[[Diethylamino)acetyl]amino]benzo[1,2-d:3,4-d']bisthiazoles (108-110)

I

V

compd	type	R ₁	R ₂	R ₃	method (Scheme I) reaction sequences ^a	mp, °C ^b	crystn ^c solvent	formula ^d	CPE ^{e,f} (200 mg/kg) % inhibn	D-adjvant arthritis ^{e,f}	
										inj/d	uninj/d paw (mg/kg)
91	I	H	CH ₃	H	4g \xrightarrow{i} 5g \xrightarrow{ii} 6g	225 dec	V	C ₂₁ H ₃₀ N ₆ O ₂ S ₂ ·2HCl·H ₂ O	8	59/100 (50)*	-37
92	I	H	OCH ₃	H	4h \xrightarrow{iii} 6h	175-177	II	C ₂₁ H ₃₀ N ₆ O ₃ S ₂ ·H ₂ O	17	19/0 (50)	+10
93	I	H	Cl	H	4f \xrightarrow{iii} 6f	225-226	II	C ₂₀ H ₂₇ ClN ₆ O ₂ S ₂	8	94/100 (50)* ^g	-46
94	I	H	Br	H	1a \xrightarrow{iv} 4e \xrightarrow{iii} 6e \xleftarrow{v} 3a	196-198	II	C ₂₀ H ₂₇ BrN ₆ O ₂ S ₂	0	80/100 (50)*	-45
95	I	H	NO ₂	H	1a \xrightarrow{i} 2a \xrightarrow{vi} 5i \xrightarrow{ii} 6i	276-278	I	C ₂₀ H ₂₇ N ₇ O ₄ S ₂	23	11/12 (50) ^h	-16
96	I	H	NH ₂	H	4j \xrightarrow{iii} 6j	160-162	II	C ₂₀ H ₂₉ N ₇ O ₂ S ₂	53*	26/38 (50)	-35
97	I	H	H	COOEt	10w \xrightarrow{i} 11w \xrightarrow{ii} 12w	215-217	II	C ₂₃ H ₃₂ N ₆ O ₄ S ₂	32*	7/0 (50)	-9
98	I	H	Cl	Cl	1a \xrightarrow{iii} 3a \xrightarrow{vii} 9k	278-280	VII	C ₂₀ H ₂₆ Cl ₂ N ₆ O ₂ S ₂	19	30/71 (50)* ^j	-33
99	I	H	Cl	CF ₃	7n \xrightarrow{i} 8n \xrightarrow{ii} 9n	240-245	IV	C ₂₁ H ₂₆ ClF ₃ N ₆ O ₂ S ₂ · 2HCl·0.2H ₂ O	18	57/100 (25)* ^j	-22
100	I	H	Cl	CN	7o \xrightarrow{i} 8o \xrightarrow{ii} 9o	260-265	IV	C ₂₁ H ₂₆ ClN ₇ O ₂ S ₂ ·2HCl	19	80/100 (50)*	-35
101	I	H	Cl	C ₆ H ₅	7q \xrightarrow{iii} 9q	247-249	VIII	C ₂₆ H ₃₁ ClN ₆ O ₂ S ₂	35*	13/7 (50)	-6
102	I	H	Cl	COOEt	10w \xrightarrow{i} 11w \xrightarrow{vi} 8p \xrightarrow{ii} 9p	260-262	VIII	C ₂₃ H ₃₁ ClN ₆ O ₄ S ₂	5	0/0 (50)	-16
103	I	H	Cl	Br	4f \xrightarrow{i} 5f \xrightarrow{v} 8m \xrightarrow{ii} 9m	278-280	VII	C ₂₀ H ₂₆ BrClN ₆ O ₂ S ₂ ·0.5H ₂ O	24	55/72 (50)*	-33
104	I	H	Br	Cl	10v \xrightarrow{v} 7l \xrightarrow{iii} 9l	265-267	XII	C ₂₀ H ₂₆ BrClN ₆ O ₂ S ₂	8	68/100 (50)*	-49
105	I	H	CH ₃	Cl	4g \xrightarrow{i} 5g \xrightarrow{ii} 6g \xrightarrow{vii} 9s	245-247	VII	C ₂₁ H ₂₉ ClN ₆ O ₂ S ₂ ·0.75H ₂ O	18	42/16 (50)	-37
106	I	H	CH ₃	COOEt	7r \xrightarrow{iii} 9r	256-257	VII	C ₂₄ H ₃₄ N ₆ O ₄ S ₂	35*	15/50 (50)*	-12
107	I	C ₂ H ₅	Br	H	1c \xrightarrow{iv} 4d \xrightarrow{i} 5d \xrightarrow{ii} 6d	158-160	II	C ₂₄ H ₃₅ BrN ₆ O ₂ S ₂	16	0/0 (50)	-16
(Scheme II)											
108	V	H	Cl	H	13x \xrightarrow{i} 14x \xrightarrow{ii} 15x	220-225	V	C ₂₀ H ₂₇ ClN ₆ O ₂ S ₂	35*	0/62 (100)*	-58
109	V	H	H	OCH ₃	16y \xrightarrow{iii} 17y	176-178	II	C ₂₁ H ₃₀ N ₆ O ₃ S ₂	32*	47/51 (50)*	+7
110	V	H	Cl	OCH ₃	17y \xrightarrow{vii} 18z	172-174	II	C ₂₁ H ₂₉ ClN ₆ O ₃ S ₂ ·0.25H ₂ O	17	38/8 (50)	-11

^aNumbers are those assigned to structures in Scheme I and II; (i) ClCOCH₂Cl; (ii) HNEt₂ (method A); (iii) NaOEt + EtOOCCH₂NEt₂ (method B); (iv) in situ bromination of the starting material (e.g. 1a, 1c, or 7l) formed by oxidative cyclization of the appropriate 1,3-phenylenebisthiourea with >3 Br₂ equivalents. See Experimental Section (method E) 115, 124, and 126; (v) Br₂; (vi) HNO₃; (vii) Cl₂. ^bUncorrected. ^cSee Table I, footnote d. ^dAll compounds had elemental analyses within ±0.4% of theoretical value. ^eAnimals were dosed orally. See Experimental Section for tests description. ^fStatistically significant results with two-tail student's tests (*p* < 0.05) results are marked with an asterisk (*). ^g3/6 animals died. ^h2/6 animals died. ⁱAt 50 mg/kg, 3/6 animals died. ^jAt 50 mg/kg 2/6 animals died.

8.9 (s, 1 H), 13.3 (s, 2 H ex). Anal. (C₁₂H₇Cl₂N₅O₄S₂) C, H, Cl.

8-Bromo-4-chloro-2,6-bis[(chloroacetyl)amino]benzo[1,2-d:5,4-d']bisthiazole (139). Bromine (2.85 mL, 0.056 mol) was added portionwise to a stirred suspension of 4-chloro-2,6-bis[(chloroacetyl)amino]benzo[1,2-d:5,4-d']bisthiazole (138) (7.0 g, 0.0171 mol) in TMP (80 mL), and the mixture was stirred at room temperature overnight. The solution was then poured on ice/water, and the resulting solid was collected, washed, and crystallized three times from DMF to yield 5.0 g (77%) of pure 139: mp 320 °C; IR 3280, 1700, 1610, 1550 cm⁻¹. Anal. (C₁₂H₆BrCl₂N₄O₂S₂) H, N; C: calcd, 29.49; found, 29.93.

2,6-Diamino-4-chloro-8-(trifluoromethyl)benzo[1,2-d:5,4-d']bisthiazole (113). Method F. A bromine (3 mL, 0.058 mol) solution in acetic acid (30 mL) was added over a period of 20 min to an ice-cold stirred mixture of 2-chloro-5-(trifluoromethyl)-1,3-phenylenediamine¹⁶ (5.3 g, 0.025 mol) and potassium thiocyanate (19.4 g, 0.20 mol) in acetic acid (150 mL) containing methanol (20 mL). The reaction mixture was stirred at room temperature for 1 h, poured into water (200 mL), and basified with ammonium hydroxide to precipitate the free base. It crystallized from DMF: 6.0 g (73%); mp 320 °C; IR 3495, 3280, 1640, 1530 cm⁻¹. Anal. (C₉H₄ClF₃N₄S₂) C, H, Cl, F, N, S.

2,6-Diamino-4-chloro-8-cyanobenzo[1,2-d:5,4-d']bisthiazole (114). The title product was prepared from 2-chloro-5-cyano-1,3-phenylenediamine¹⁶ by method F—see above—in 66% yield. It crystallized from methanol: mp 350 °C; IR 3400, 3275, 3070, 2225, 1630, 1540 cm⁻¹. Anal. (C₉H₄ClN₅S₂·1/4DMF) C, H, N, Cl.

2,6-Diamino-8-carboxy-4-methylbenzo[1,2-d:5,4-d']bisthiazole Sodium Salt (116). The title product was prepared

from 3,5-diamino-4-methylbenzoic acid¹⁷ in 46% yield by method F, see above. The crude product was crystallized from 1 N sodium hydroxide solution to yield the sodium salt: mp 350 °C; IR 3600-2700, 1670, 1635, 1615, 1540, 1375 cm⁻¹; NMR (Me₂SO-*d*₆) δ 2.4 (s, 3 H), 7.0 (s, 4 H ex). Anal. (C₁₀H₇N₄O₂S₂Na·H₂O) C, H, N, S.

2,6-Diamino-8-carbomethoxy-4-methylbenzo[1,2-d:5,4-d']bisthiazole (121). The title product was prepared from ethyl 3,5-diamino-4-methylbenzoate¹⁸ in 80% yield (crude) by method F, see above. The product crystallized from DMF: mp 350 °C; IR 3430, 3300, 3150, 1660, 1640 sh, 1630, 1530 cm⁻¹; NMR (Me₂SO-*d*₆) δ 1.4 (t, 3 H), 2.7 (s, 3 H), 4.2 (q, 2 H), 7.5 (s, br, 4 H ex). Anal. (C₁₂H₁₂N₄O₂S₂) C, H, N, S.

2,4,6-Triaminobenzo[1,2-d:5,4-d']bisthiazole (122). The title product was prepared from 2-amino-1,3-phenylenediamine¹⁹ by method F, see above. The crude product (47%) obtained after treatment with ammonia was collected, washed, and dried: mp 350 °C; IR 3410, 3320, 3275, 3140, 2250 w, 1620 s, 1530 cm⁻¹. The weak band at 2250 cm⁻¹ would indicate some uncyclized product in this sample. The product was used as such to prepare 96.

2,6-Diamino-4-chloro-8-phenylbenzo[1,2-d:5,4-d']bisthiazole (123). The title product was prepared from 3,5-diamino-4-chlorobiphenyl²⁰ in 63% yield by method F, see above. It crystallized from DMF: mp 350 °C; IR 3400, 3350, 3100, 1615, 1515 cm⁻¹; NMR (Me₂SO-*d*₆) δ 7.6 (m, 9 H, 4 ex). Anal. (C₁₄H₉ClN₄S₂·1/3H₂O) C, H, S.


(17) Claus, A.; Joachim, J. *Liebigs Ann. Chem.* 1891, 266, 209.

(18) Barben, I. K.; Suschitzky, H. *J. Chem. Soc.* 1960, 672.

(19) von Auwers, K.; Frese, E. *Ber.* 1926, 59, 539.

(20) Wright, J. B. Ger. Offen. 2,525,249, 1976; *Chem. Abstr.* 1976, 84, 135315m.

(16) Blahak, J.; Meckel, W.; Mueller, E. Ger. Offen. 2,025,896, 1971; *Chem. Abstr.* 1971, 75, 141742d.

Table III. Chemical and Biological Screening Data for Diaminobenzobisthiazoles,^a Types I-V, Used as Starting Materials in Tables I and II

compd (Schemes I and II)	type	R ₁	R ₂	R ₃	method ^b (lit.)	crystn ^c solvent	mp, °C ^d	formula ^e	CPE/ ^f (mg/kg) % inhibn	D-adjuvant arthritis/ ^g	
										injnd/uninjnd paw (mg/kg) % inhibn	% change body wt
111 (1a)	I	H	H	H	(10), (11)	XIII	>400	C ₈ H ₆ N ₄ S ₂	60 (200)* ^h	79/92 (100)*	-20
112 (4b)	I	CH ₃	H	H	(13)	VII	318-319	C ₈ H ₁₀ N ₄ S ₂ ·2HCl	24 (200)	71/100 (25)* ⁱ	-30
113 (7n)	I	H	Cl	CF ₃	F	VII	>320	C ₈ H ₄ ClF ₃ N ₄ S ₂	13 (200)	0/12 (200)	+3
114 (7o)	I	H	Cl	CN	F	XIV	>320	C ₈ H ₄ ClN ₅ S ₂ ·0.25H ₂ O	7 (200)	47/37 (200)	+3
115 (4e)	I	H	Br	H	G	XIII	>320	C ₈ H ₅ BrN ₄ S ₂	33 (200)*	47/37 (200)	+10
116 (7t)	I	H	CH ₃	COONa	F	XV	>320	C ₁₀ H ₇ N ₄ O ₂ S ₂ ·Na·H ₂ O	22 (200)	7/0 (50)	-15
117 (7w)	I	H	H	COOEt	H	II	>320	C ₁₁ H ₁₀ N ₄ O ₂ S ₂ ·1.25H ₂ O	2 (200)	23/0 (50)	-14
118 (4g)	I	H	CH ₃	H	(13)	VII	>320	C ₈ H ₈ N ₄ S ₂			
119 (4h)	I	H	OCH ₃	H	(13)	VII	305 dec	C ₈ H ₈ N ₄ OS ₂			
120 (4k)	I	H	Cl	H	(13)	XIII	>350	C ₈ H ₅ ClN ₄ S ₂			
121 (7p)	I	H	CH ₃	COOEt	F	VII	>320	C ₁₂ H ₁₂ N ₄ O ₂ S ₂			
122 (4j)	I	H	NH ₂	H	F	XVI	>320	C ₈ H ₇ N ₅ S ₂ ^j			
123 (7g)	I	H	Cl	C ₆ H ₅	F	VII	>320	C ₁₄ H ₉ ClN ₄ S ₂ ·0.33H ₂ O			
124 (7l)	I	H	Br	Cl	G	XVI	>320	C ₈ H ₄ BrClN ₄ S ₂ ^j			
125 (1c)	I	C ₂ H ₅	H	H	(11)	VII	306-308	C ₁₂ H ₁₄ N ₄ S ₂			
126 (4d)	I	C ₂ H ₅	Br	H	G	VII	273-275	C ₁₂ H ₁₃ BrN ₄ S ₂ ^j			
127	II	H	H	H	(11)	XIV	350	C ₈ H ₆ N ₄ S ₂	4 (100)	88/100 (10)* ^h	-30
128 ⁱ	III	H	H	H	(11)	XVII	320	C ₈ H ₆ N ₄ S ₂			
129	IV	CH ₃	H	H	(11)	XIII	315-316	C ₁₀ H ₁₀ N ₄ S ₂			
130 (13x)	V	H	Cl	H	(11)	XIII	360	C ₈ H ₅ ClN ₄ S ₂			
131 (16y)	V	H	H	OCH ₃	H	II	320	C ₈ H ₈ N ₄ OS ₂ ·1/4H ₂ O	43 (200)*	34/35 (50) ^m	-29

^a See Table I, note a. ^b As described in literature (lit.) or in the Experimental Section, methods F-H. ^c II, ethanol; VII, DMF; XIII, pyridine; XIV, DMF-H₂O; XV, H₂O; XVI, product was not purified and used as such; XVII, product was purified via the hydrochloride and then neutralized with ammonia. ^d Uncorrected. ^e The elemental analysis of new compounds within ±0.4% of the theoretical value. ^f Animals were dosed orally. For description of screening tests see Experimental Section. ^g Statistically significant results with two-tail student's tests ($p < 0.05$) are marked with an asterisk (*). ^h Product showed no activity when tested with adrenalectomized animals. ⁱ 50 mg/kg for 4 days and 25 mg/kg for 10 days; 2/6 animals died. At 10 mg/kg, compound was inactive. ^j Product was not analyzed, see Experimental Section. ^k At 50 mg/kg, all animals died by day 7. ^l Prepared from 1,4-phenylenediamine; product contained 25% of the corresponding linear isomer, 127, by NMR analysis. ^m 2/6 animals died on days 2 and 3.

Table IV. Effect of 26 in Plaque-Forming Cell (PFC) in AKR Mice^a

compd	dose, ^b mg/kg po	N ^c	total PFC/spleen % suppression ^d
26	10	6	11
	50	6	67*
	100	6	82*
	200	6	83*
90a	125	6	28

^a See experimental. ^b Day 1-3. ^c Number of animals. ^d Results with $p < 0.05$ are marked with an asterisk (*).

2,6-Diamino-8-carbomethoxybenzo[1,2-*d*:5,4-*d'*]bisthiazole (117). **Method H.** A solution of bromine (2.7 g, 17 mmol) in chloroform (10 mL) was added dropwise to 5-carbomethoxy-1,3-phenylenebisthiourea (150) (2.4 g, 8.05 mmol) in chloroform (100 mL). When the addition was completed, the mixture was refluxed for 2 h, and the solid was collected, resuspended in water, and neutralized with ammonium hydroxide. The base was collected, washed, dried, and crystallized from ethanol, yielding 117 (0.90 g, 38%): mp 350 °C; IR 3320, 3180, 1650 cm⁻¹; NMR (Me₂SO-*d*₆) δ 1.2 (t, 3 H), 4.4 (q, 2 H), 7.4 (4 H ex), 7.55 (s, 1 H); UV λ_{max} (log ε) 215 (3.79), 238 (4.15), 254 (sh) (3.96), 346 nm (3.61). Anal. (C₁₁H₁₀N₄O₂S₂·1.25H₂O) C, H, N, S.

2,7-Diamino-4-methoxybenzo[1,2-*d*:3,4-*d'*]bisthiazole (131). The title compound was prepared in 48% yield from 5-methoxy-1,3-phenylenebisthiourea by method H, see above: mp 255 °C; IR 3340, 3200, 1640 cm⁻¹; NMR (Me₂SO-*d*₆) δ 3.85 (s, 3 H), 6.8 (s, 1 H), 7.68 (s, br, 4 H ex); UV λ_{max} (log ε) 210 (4.30), 221 (sh) (4.28), 252 (4.30), 306 nm (sh) (3.63). Anal. (C₉H₈N₄OS₂·3/4H₂O) C, H, N, S.

2,6-Diamino-4-bromobenzo[1,2-*d*:5,4-*d'*]bisthiazole (115). **Method G.** Bromine (9 mL, 0.175 mol) was added dropwise to a solution of 1,3-phenylenebisthiourea¹⁰ (8.5 g, 0.0377 mol) in chloroform (150 mL). The mixture was stirred at room temperature overnight and then refluxed for 2 h. After cooling, a solid precipitated, which was collected, suspended in water, and treated with ammonium hydroxide. The free base was collected, washed with water, and crystallized from pyridine to yield 2.8 g (27.5%) of a light pink crystalline solid: mp 320 °C; IR 3450, 3280, 1630, 1535 cm⁻¹; NMR (Me₂SO-*d*₆) δ 7.6 (s, br, 4 H ex), 7.8 (s, 1 H); UV λ_{max} (log ε) 210 (3.98), 253 (4.21), 304 (sh) (3.72), 315 nm (3.87). Anal. (C₈H₅BrN₄S₂) C, H, N.

2,6-Diamino-4-bromo-8-chlorobenzo[1,2-*d*:5,4-*d'*]bisthiazole (124). The title product was prepared from 5-chloro-1,3-phenylenebisthiourea and excess bromine according to method G, see above. The crude product was treated with boiling DMF (25 mL/g), the insoluble material was removed by filtration, and the filtrate was rotoevaporated to dryness. The residue (mp 300 °C; IR 3470, 3270, 3100, 1630, 1540 cm⁻¹) was used as such for the preparation of 104.

4-Bromo-2,6-bis(ethylamino)benzo[1,2-*d*:5,4-*d'*]bisthiazole (126). The title product was prepared from *N,N'*-diethyl-1,3-phenylenebisthiourea¹¹ and excess bromine in 49% yield according to method G, see above. The product crystallized from DMF: mp 273-275 °C; IR 3200, 2975, 1675 sh, 1625, 1570 cm⁻¹; NMR (Me₂SO-*d*₆) δ 1.2 (t, 6 H), 3.3 (m, 4 H), 7.8 (s, 1 H), 8.1 (t, 2 H ex). The product was used as such for the preparation of 107.

5-Carbomethoxy-1,3-phenylenebisthiourea (149). A mixture of ethyl 3,5-diaminobenzoate²¹ (10 g, 0.0395 mol) and ammonium

(21) Brill, H. C. *J. Am. Chem. Soc.* 1921, 43, 1320.

Table V. Intermediate Acyl Chlorides of Diaminobenzobisthiazoles^a

compd	type	X ₁ = X ₂	R ₂	R ₃	mp, °C ^b	formula ^c
132	I	NHCOCH ₂ Cl	H	H	>300	C ₁₂ H ₈ Cl ₂ N ₄ O ₂ S ₂
133	I	NH(CH ₃)COCH ₂ Cl	H	H	>250 dec	C ₁₄ H ₁₂ Cl ₂ N ₄ O ₂ S ₂
134	I	NH(C ₂ H ₅)COCH ₂ Cl	H	H	>220 dec	C ₁₆ H ₁₆ Cl ₂ N ₄ O ₂ S ₂
135	I	NHCOCH(CH ₃)Cl	H	H	265–270	C ₁₄ H ₁₂ Cl ₂ N ₄ O ₂ S ₂ ·1.5DMF
136	I	NHCOCH ₂ Cl	CH ₃	H	>300	C ₁₃ H ₁₀ Cl ₂ N ₄ O ₂ S ₂
137	I	NHCOCH ₂ Cl	NO ₂	H	>300	C ₁₂ H ₇ Cl ₂ N ₄ O ₂ S ₂
138	I	NHCOCH ₂ Cl	Cl	H	>300 ^d	C ₁₂ H ₇ Cl ₃ N ₄ O ₂ S ₂
139	I	NHCOCH ₂ Cl	Cl	Br	>300	C ₁₂ H ₆ BrCl ₂ N ₄ O ₂ S ₂ ^e
140	I	NHCOCH ₂ Cl	Cl	CF ₃	>300	C ₁₃ H ₆ Cl ₃ F ₃ N ₄ O ₂ S ₂
141	I	NHCOCH ₂ Cl	Cl	CN	>300	C ₁₃ H ₆ Cl ₃ N ₅ O ₂ S ₂ ·0.33H ₂ O
142	I	NHCOCH ₂ Cl	H	COOEt	>300 dec	C ₁₅ H ₁₂ Cl ₂ N ₄ O ₄ S ₂ ·0.5H ₂ O
143	I	NH(C ₂ H ₅)COCH ₂ Cl	Br	H	>300 dec	C ₁₆ H ₁₅ BrCl ₂ N ₄ O ₂ S ₂
144	II	NHCOCH ₂ Cl	H	H	>300 dec	C ₁₂ H ₈ Cl ₂ N ₄ O ₂ S ₂
145	III	NHCOCH ₂ Cl	H	H	>300 ^f	C ₁₂ H ₈ Cl ₂ N ₄ O ₂ S ₂ ·0.33H ₂ O
146	IV	N(CH ₃)COCH ₂ Cl	H	H	280–285	C ₁₄ H ₁₂ Cl ₂ N ₄ O ₂ S ₂
147	V	NHCOCH ₂ Cl	Cl	H	>300	C ₁₂ H ₇ Cl ₃ N ₄ O ₂ S ₂ ·0.67DMF
148	I	X ₁ , X ₂ X ₁ = NHCOCH ₂ N(C ₂ H ₅) ₂ X ₂ = NHCOCH ₂ Cl	H	H	280–282	C ₁₆ H ₁₈ ClN ₅ O ₂ S ₂

^a See Table I, note a. ^b Products were crystallized from DMF except as indicated. ^c Analyses were within ±0.4% of the theoretical value except where indicated. ^d Product could not be crystallized. It was used crude and not analyzed. ^e C: calcd, 29.50; found, 29.93. ^f DMF-CH₃COOH. ^g Product contained 25% of isomer 144 (NMR).

thiocyanate (6.5 g, 0.0854 mol) in water (30 mL) was heated to reflux for 3 h. The reaction product was cooled, the solid was collected and crystallized from water and a trace amount of DMF to yield 2.4 g (20.4%) of the title product: mp 210–212 °C; NMR (Me₂SO-*d*₆) δ 1.2 (t, 3 H), 4.4 (q, 2 H), 7.58 (s, 4 H ex), 7.8 (s, 2 H), 7.9 (s, 1 H), 10.0 (s, 2 H ex). Anal. (C₁₁H₁₄N₄O₂S₂·1/4H₂O) C, H, N: calcd, 18.50; found, 18.09.

5-Chloro-1,3-phenylenebisthiourea (150). 5-Chloro-1,3-phenylenediamine²² (15.5 g, 0.109 mol), carbon disulfide (53 mL, 0.109 mol), and triethylamine (53 g, 0.524 mol) were mixed together in acetone (450 mL) and stirred at room temperature for 2 days. A solid was collected which was added to methanol (100 mL) and treated with iodomethane (25 g, 0.173 mol) for 2 h at room temperature. Dimethyl 5-chloro-1,3-phenylenebis(dithiocarbamate) precipitated. It was collected and dissolved in methanol (200 mL), and the solution was treated with concentrated ammonia (75 mL). The mixture was heated to reflux for 2 h, cooled to room temperature, and the resulting suspension was filtered. The solid was washed with methanol and dried. Another crop was obtained by evaporation of the mother liquor and treatment of the residue with ethanol: yield 11.4 g (38%); mp 185–187 °C; NMR (Me₂SO-*d*₆) δ 7.3 (s, 2 H), 7.5 (s, 1 H), 7.5 (s, br, 4 H ex), 9.7 (s, 2 H ex). The product was used as such.

5-Methoxy-1,3-phenylenebisthiourea (151). This product was obtained from 5-methoxy-1,3-phenylenediamine²³ in three steps as described above for 5-chloro-1,3-phenylenebisthiourea: yield 53%; mp 193–195 °C; IR 3380, 3250, 3150, 1615 cm⁻¹; NMR (Me₂SO-*d*₆) δ 3.7 (s, 3 H), 6.8 (d, 2 H), 7.0 (t, 1 H), 7.5 (s, 4 H ex), 9.5 (s, 2 H ex). The product was used as such.

2,6-Bis[(carboxymethyl)amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (83). 2,6-Bis[(carboxymethyl)amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (22c) (18 g, 0.0545 mol) in a mixture of ethanol (300 mL) and 1 N NaOH (150 mL) was heated in a water bath for 30 min. Water (300 mL) was added, and the resulting solution was concentrated in vacuo to about 300 mL and then neutralized with 2 N HCl. The solid was collected, washed with water, dried, and crystallized from DMF to yield 13 g (84%) of the title product melting at 310 °C: IR 3500–2600 with max. at 2900, 1640, 1620

sh, 1565, 1420, 1300 cm⁻¹; NMR (TFA-*d*) δ 4.6 (s, 4 H), 8.0 (s, 1 H), 8.3 (s, 1 H). Anal. (C₁₂H₁₀N₄O₄S₂) C, H, N, S: calcd, 18.95; found, 18.53.

2,6-Bis[(carboxymethyl)amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (22c). Bromine (26 g, 0.163 mol) in chloroform (100 mL) was added dropwise at room temperature to a stirred solution of 1,3-bis[*N'*-(carboxymethyl)thioureido]benzene (21c) (31.5 g, 0.079 mol) in chloroform (500 mL). The resulting mixture was heated to reflux for 1 h. A solid was collected, suspended in water, treated with ammonia, and again collected by filtration. It was washed well with water and then crystallized from ethanol/DMF, 2/1, to yield 20 g (65%) of the title compound: mp 215–217 °C; IR 3340, 3300, 1735 sh, 1725, 1600, 1550 cm⁻¹; NMR (Me₂SO-*d*) δ 1.2 (t, 6 H), 4.2 (m, 8 H), 7.4 (s, 1 H), 8.0 (s, 1 H), 8.4 (t, br, 2 H ex). Anal. (C₁₆H₁₈N₄O₄S₂) C, H, N, S.

1,3-Bis[*N'*-(carboxymethyl)thioureido]benzene (21c). A stirred mixture of 1,3-phenylenediamine (2.5 g, 0.023 mol) and ethyl isothiocyanatoacetate²⁴ (8.0 g, 0.055 mol) in dioxane (30 mL) was heated on a water bath for 1 h. The resulting solution was evaporated, and the residue was crystallized from ethanol to give 8.0 g (87%) of the title product: mp 122–124 °C; IR 3340, 1740, 1620 m, 1550, 1530 cm⁻¹; NMR (Me₂SO-*d*₆) δ 1.2 (t, 6 H), 4.15 (m, 8 H), 7.2 (s, br, 3 H), 7.4 (s, br, 1 H), 8.0 (t, 2 H ex), 9.6 (s, br, 2 H ex). Anal. (C₁₆H₂₂N₄O₄S₂) C, H, N, S.

2,6-Bis[[(*N,N*-diethylcarbamoyl)methyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (84). Method D. A stirred mixture of 2-chloro-6-[[*(N,N*-diethylcarbamoyl)methyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (24) (4.0 g, 0.0113 mol) and glycine-diethylamide¹⁴ (1.7 g, 0.0135 mol) was heated neat at 170 °C for 30 min, allowed to cool to room temperature, and treated with ethanol. The resulting solid was collected and crystallized from DMF: yield 2.3 g (45%); mp 235–237 °C; IR 3250, 1640, 1600 w, 1540 cm⁻¹; NMR (CDCl₃) δ 1.2 (m, 12 H), 3.4 (m, 8 H), 4.3 (s, br, 4 H), 6.6 (s, br, 2 H ex), 7.8 (s, 1 H), 7.9 (s, 1 H). Anal. (C₂₀H₂₈N₆O₂S₂) C, H, N, S.

2-Chloro-6-[[*(N,N*-diethylcarbamoyl)methyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole (24). A stirred mixture of 2,6-dichlorobenzo[1,2-*d*:5,4-*d'*]bisthiazole¹⁰ (9.4 g, 0.036 mol), gly-

(22) Cohn, P. *Monatsh. Chem.* 1909, 22, 118.(23) Zemplen, G.; Bognar, R.; Thiele, K. *Ber.* 1944, 77, 446.(24) Johnson, T. B.; Renfrew, A. G. *J. Am. Chem. Soc.* 1925, 47, 240.

cine-diethylamide¹⁴ (9.4 g, 0.072 mol), and triethylamine (8.0 g, 0.079 mol) in dioxane (50 mL) was heated to reflux for 1 h. The reaction mixture was evaporated, the residue was dissolved in chloroform, and the solution was washed with H₂O and dried. It was then concentrated to a few milliliters and applied to a 200-g silica gel column. The reaction products were eluted with chloroform/methanol, 95/5. The first fraction (8.5 g, 66%) crystallized from ethanol/DMF and corresponded to the title product: mp 178–180 °C; IR 3220, 1640 sh, 1630, 1600, 1560 cm⁻¹; NMR (CDCl₃) δ 1.1 (m, 6 H), 3.4 (m, 4 H), 4.25 (d, 2 H), 6.9 (s, br, 1 H ex), 7.8 (s, 1 H), 8.0 (s, 1 H). Anal. (C₁₄H₁₅ClN₂OS₂) C, H, Cl, N. A second fraction yielded 2.0 g of the bis-adduct (identical IR to 84 above), which was contaminated with an impurity of near identical R_f, and which could not be removed by crystallization.

2,6-Bis[[2-(*N,N*-diethylamino)ethyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole Tetrahydrochloride (85). Method C. 2-(Diethylamino)ethyl isothiocyanate²⁵ (8.0 g, 0.051 mol) was added to a solution of 1,3-phenylenediamine (2.7 g, 0.025 mol) in dioxane (20 mL), and the mixture was heated to reflux for 3 h. The product, 1,3-bis[*N'*-(*N,N*-diethylamino)ethyl]thioureido]benzene (21a) [11 g, 99%; NMR (CDCl₃) δ 0.9 (m, 12 H), 2.5 (q, 12 H), 3.6 (q, br, 6 H, 2 H ex), 7.2 (m, 6 H, 2 H ex)], obtained on evaporation, was used as such for the oxidative cyclization of the next step. The compound (7.0 g, 0.0165 mol) was dissolved in chloroform (100 mL), and excess HCl gas was bubbled through the solution. Bromine (4.5 g, 0.028 mol) in chloroform (70 mL) was then added dropwise, at room temperature, and under strong stirring to the resulting suspension of the hydrochloride. The mixture was then heated to reflux for 1 h and cooled, and the resulting solid was collected. It was dissolved in water, and the solution was treated with ammonia and extracted with chloroform (3 × 100 mL). The combined extracts were washed with water and brine and dried over sodium sulfate. The residue obtained on evaporation (6.2 g) was passed through a column of silica gel using chloroform/methanol, 9/1, as eluant. The title product (2.5 g, 36%) was obtained as thick oil: IR 3360, 2970, 2830, 1620, 1550 cm⁻¹; NMR (CDCl₃) δ 1.0 (t, 12 H), 2.5 (m, 12 H), 3.4 (t, 4 H), 6.2 (s, br, 2 H ex), 7.6 (s, 1 H), 7.65 (s, 1 H). Anal. (C₂₀H₃₂N₆S₂·0.5H₂O) C, H, N, S. The oil was dissolved in chloroform, and the hydrochloride was precipitated from ethereal HCl. It was recrystallized from methanol, mp 195–200 °C. Anal. (C₂₀H₃₂N₆S₂·4HCl·2CH₃OH) C, H, N.

2,6-Bis[[5-(diethylamino)-2-pentyl]amino]benzo[1,2-*d*:5,4-*d'*]bisthiazole Tetrahydrochloride (86). The title product was prepared as described above for compound 85 from 1,3-phenylenediamine and 1-(diethylamino)-4-isothiocyanatopentane. The latter product was prepared according to Schmidt et al.²⁶ It had a bp of 68–72 °C (0.1 mmHg): IR 2950, 2800, 2100, 1475, 1450 cm⁻¹. The hydrochloride was obtained from an ethanolic solution of the free base with excess ethereal HCl. It became a crystalline solid after treatment with acetone, ether, and ethyl acetate: yield 57%; mp 145–150 °C dec; NMR (Me₂SO-*d*₆) δ 1.2 (t, 21 H), 1.8 (m, br, 8 H), 3.1 (m, br, 14 H), 4.3 (m, br, 2 H), 8.0 (s, 1 H), 8.4 (s, 1 H), 10.7–11.0 (m, br, 7 H ex). Anal. (C₂₆H₄₄N₆S₂·4HCl·EtOH) C, Cl, S, H: calcd, 7.80; found, 7.32; N: calcd, 12.07; found, 12.54. Neutralization of an aqueous solution of this salt with 2 N NaOH afforded the pure free base as a solid, mp 85–86 °C. Anal. (C₂₆H₄₄N₆S₂·H₂O) C, H, N, S.

2,6-Bis(*N*-ethylureido)benzo[1,2-*d*:5,4-*d'*]bisthiazole (79). Method E. A stirred mixture of 2,6-diaminobenzo[1,2-*d*:5,4-*d'*]bisthiazole¹⁰ (4.45 g, 0.020 mol) and ethyl isocyanate (3.0 g, 0.042 mol) in dioxane (100 mL) was heated to reflux for 2 h. The resulting suspension was left at room temperature overnight. A beige solid (5.6 g, 77%) was collected, washed, dried, and crystallized from DMF: mp 330 °C; IR 3400, 3260, 3200 sh, 3080, 1680, 1660, 1615 br cm⁻¹; NMR (CF₃COOD) δ 1.3 (t, 6 H), 3.5 (q, 4 H), 8.4 (s, 1 H), 8.5 (s, 1 H), 12 (s, br, 4 H). Anal. (C₁₄H₁₆N₆O₂S₂) C, H, N.

Biological Methods. Carrageenan Paw Edema. Edema was produced in the right hind paw of male Sprague–Dawley rats

(150 ± 10 g) by the subplantar injection of 0.1 mL of a 1% carrageenan suspension in saline.²⁷ Test compounds or vehicle was administered orally in 0.5% methylcellulose 1 h prior to carrageenan injection. Five rats were used in each group. The paw volume was determined by measuring the amount of mercury displaced after immersing the paw to the level of the lateral malleolus. Paw volumes were measured just prior to test compound administration and again 3 h after carrageenan injection and the difference was designated as edema volume. Statistics were performed using Analysis of Variance (ANOVA) and Dunnett's *t* test.

Ulcerogenicity Test. Ulcerogenic activity was evaluated by the method described by Wong et al.³⁰

Acetic Acid Writhing. The analgesic activity of 26 was determined by using the procedure of Hendershot and Forsaith,³¹ but with acetic acid³² in place of phenylquinone.

Plaque Forming Cell (PFC) Test. The PFC assay was performed according to the method of Jerne et al.³³ For each test three groups consisting of six AKR mice were set up. One control group was neither immunized nor treated and was used to establish the number on nonspecific PFC/spleen. Mice in another control group were immunized on day 0 with 0.2 mL of a 15% suspension of sheep erythrocytes and the test groups were treated with test drug starting 1 day after immunization and continued until day 3. All groups were killed on day 4, and the spleens were removed. The spleen cells were teased out into a culture medium, washed, counted, and diluted to a suitable concentration. Aliquots were then mixed with sheep erythrocytes and agarose and plated in triplicate on Petri dishes. These were incubated for 1 h at 37 °C, and then fresh guinea pig serum was added to provide complement. After incubation for a further 30 min, the plates were stained with benzidine, and the number of plaques were counted. The total PFC/spleen in the test group was expressed as a percentage of those in the control group. The data were analyzed using the Student's *t* test or Analysis of Variance.

Acknowledgment. We are indebted to Anne Marie Thomas for recording the NMR and UV spectra and to Mary Feron and Kay Gurry for their able assistance in the preparation of the manuscript.

Registry No. 19, 108-45-2; 20a, 32813-52-8; 20b, 104093-88-1; 20c, 24066-82-8; 20d, 4385-41-5; 20e, 137697-50-8; 21a, 137697-51-9; 21b, 137697-52-0; 21c, 137697-55-3; 21d, 137697-53-1; 21e, 137697-54-2; 22c, 137697-56-4; 24, 137697-58-6; 25, 70175-42-7; 26, 70175-10-9; 27, 74071-27-5; 27-2HCl, 70175-45-0; 28, 70175-69-8; 29, 70175-70-1; 30, 70175-73-4; 31, 137698-01-2; 31-2HCl, 137697-59-7; 32, 70175-14-3; 32-2HCl, 137697-60-0; 33, 137698-02-3; 33-2HCl, 74071-07-1; 34, 70175-44-9; 35, 74071-42-4; 36, 74071-28-6; 37, 70175-51-8; 38, 74071-29-7; 39, 74071-30-0; 40, 70175-50-7; 41, 70175-48-3; 42, 70175-49-4; 43, 70175-24-5; 44, 70175-21-2; 44-2HCl, 70175-22-3; 45, 70175-23-4; 46, 70175-71-2; 47, 70175-17-6; 48, 70175-18-7; 49, 74083-79-7; 49-2HCl, 137697-61-1; 50, 70175-19-8; 51, 74071-25-3; 52, 70175-64-3; 53, 70175-66-5; 54, 70175-57-4; 55, 70175-59-6; 56, 74071-09-3; 57, 74071-35-5; 58, 74071-11-7; 59, 137697-62-2; 60, 74071-36-6; 61, 74071-10-6; 62, 70175-20-1; 63, 137697-63-3; 64, 70175-68-7; 65, 70175-67-6; 66, 137698-03-4; 66-4HCl, 74071-38-8; 67, 70175-55-2; 68, 70175-52-9; 68-2HCl, 74071-31-1; 69, 74071-32-2; 70, 74071-33-3; 71, 74071-41-3; 72, 74083-81-1; 73, 70175-61-0; 74, 70175-63-2; 75, 137697-64-4; 76, 74071-37-7; 77, 74071-40-2; 78, 70175-47-2; 79, 137697-65-5; 80, 137698-04-5; 80-2HCl, 137697-66-6; 81, 137698-07-8; 81-2HCl,

- (25) Schmidt, E.; Ross, D.; Kittl, J.; von Duesel, H. H.; Wamsler, K. *Liebigs Ann. Chem.* 1958, 612, 11.
 (26) Schmidt, E.; Zaller, F.; Moosmueller, F.; Kammerl, E. *Liebigs Ann. Chem.* 1954, 585, 230.

- (27) Winter, C. A.; Risley, E. A. *J. Pharmacol. Exp. Ther.* 1963, 141, 369.
 (28) Newbould, B. B. *Br. J. Pharmacol.* 1963, 21, 127.
 (29) Robert, A.; Nezamis, J. E. *Acta Endocrinol.* 1957, 25, 105.
 (30) Wong, S.; Garducki, J. F.; Pruss, T. P. *J. Pharmacol. Exp. Ther.* 1973, 185, 127.
 (31) Hendershot, L. C.; Forsaith, J. *J. Pharmacol. Exp. Ther.* 1959, 125, 237.
 (32) Koster, R.; Anderson, W.; DeBeer, E. *J. Fed. Proc. Fed. Am. Soc. Exp. Biol.* 1959, 18, 412.
 (33) Jerne, N. K.; Nordin, A. A.; Henz, C. In *Cell Bound Antibodies*; Amos, B., Koprowski, H., Eds.; Wistar Institute Press: Philadelphia, 1963; p 107.

137697-67-7; 82, 137697-68-8; 83, 137697-69-9; 84, 137697-70-2; 85, 137697-71-3; 86, 137698-00-1; 86·4HCl, 137697-72-4; 87, 70175-11-0; 88, 70175-39-2; 89, 70175-40-5; 91, 70175-36-9; 91·2HCl, 70175-37-0; 92, 74071-15-1; 93, 74071-13-9; 94, 70175-33-6; 95, 70175-32-5; 96, 74071-23-1; 97, 74071-18-4; 98, 70175-31-4; 99, 74071-43-5; 99·2HCl, 70175-26-7; 100, 137698-05-6; 100·2HCl, 70175-29-0; 101, 74071-20-8; 102, 74071-21-9; 103, 74071-24-2; 104, 74071-22-0; 105, 70215-11-1; 106, 74071-19-5; 107, 74071-08-2; 108, 137697-74-6; 109, 137697-75-7; 110, 137697-76-8; 111, 2789-24-4; 112, 137698-06-7; 112·2HCl, 16351-82-9; 113, 70175-27-8; 114, 70175-30-3; 115, 137697-77-9; 116, 137697-78-0; 117, 137697-79-1; 118, 27052-41-1; 119, 27052-39-7; 120, 27052-36-4; 121, 137697-80-4; 122, 137697-81-5; 123, 137697-82-6; 124, 137697-83-7; 125, 16351-85-2; 126, 137697-84-8; 127, 16162-28-0; 128, 16203-56-8; 129, 137697-85-9; 130, 16351-89-6; 131, 137697-86-0; 132, 70175-09-6; 133, 70175-13-2; 134, 137697-87-1; 135, 137697-88-2; 136, 70175-35-8; 137, 137697-89-3; 138, 137697-90-6; 139, 137697-91-7; 140, 70175-25-6; 141, 70175-28-9; 142, 137697-92-8; 143, 137697-93-9; 144, 137697-94-0; 145, 137697-95-1; 146, 137697-96-2; 147,

137697-97-3; 148, 70175-39-2; 150, 137697-99-5; diethylamine, 109-89-7; chloroacetyl chloride, 79-04-9; *N,N*-diethylglycinate, 2644-21-5; 2-chloro-5-(trifluoromethyl)-1,3-phenylenediamine, 34207-44-8; potassium thiocyanate, 333-20-0; 2-chloro-5-cyano-1,3-phenylenediamine, 34207-46-0; 3,5-diamino-4-methylbenzoic acid, 6633-36-9; ethyl 3,5-diamino-4-methylbenzoate, 42908-12-3; 2-amino-1,3-phenylenediamine, 608-32-2; 3,5-diamino-4-chlorobiphenyl, 58495-18-4; 5-carbomethoxy-1,3-phenylenebisthiourea, 137697-98-4; 5-methoxy-1,3-phenylenebisthiourea, 137697-57-5; 1,3-phenylenebisthiourea, 2591-01-7; *N,N'*-diethyl-1,3-phenylenebisthiourea, 16349-50-1; 3,5-diaminobenzoic acid ethyl ester, 1949-51-5; ammonium thiocyanate, 1762-95-4; 5-chloro-1,3-phenylenediamine, 33786-89-9; dimethyl 5-chloro-1,3-phenylenebisthiocarbamate, 137697-73-5; 5-chloro-1,3-phenylenebisthiourea, 100-96-9; 2,6-dichlorobenzo[1,2-*d*:5,4-*d'*]-bisthiazole, 2591-03-9; 2-(diethylamino)ethyl isothiocyanate, 32813-52-8; 1-(diethylamino)-4-isothiocyanatopentane, 104093-88-1; 2,6-diaminobenzo[1,2-*d*:5,4-*d'*]bisthiazole, 2789-24-4; ethyl isocyanate, 109-90-0.

New 8-(Trifluoromethyl)-Substituted Quinolones. The Benefits of the 8-Fluoro Group with Reduced Phototoxic Risk

J. P. Sanchez,* A. J. Bridges, R. Bucsh, J. M. Domagala, R. D. Gogliotti, S. E. Hagen, C. L. Heifetz, E. T. Joannides, J. C. Sesnie, M. A. Shapiro, and D. L. Szotek

Parke-Davis Pharmaceutical Research Division, Warner-Lambert Company, Ann Arbor, Michigan 48106.

Received July 29, 1991

A series of 8-(trifluoromethyl)-substituted quinolones has been prepared and evaluated for in vitro and in vivo antibacterial activity, and phototolerance in a mouse phototolerance assay. These analogues were compared to the corresponding series of 6,8-difluoro- and 6-fluoro-8*H*-quinolones (ciprofloxacin type). Although their in vitro antibacterial activities are less than the 6,8-difluoro analogues, the 8-(trifluoromethyl)quinolones are generally equivalent to their 8*H* analogues. In vivo, they are comparable to the 6,8-difluoro series and show up to 10-fold improvement in efficacy when compared to their ciprofloxacin counterparts vs *Streptococcus pyogenes* and *Streptococcus pneumoniae*. In the phototolerance model, the 8-(trifluoromethyl)quinolones are comparable to the 8*H*-quinolones. Both of these series display much higher no effect doses (greater tolerance) than the corresponding 6,8-difluoroquinolones.

The quinolone antibacterials have emerged as an area of intense interest because of their broad spectrum of activity in vitro and their in vivo chemotherapeutic efficacy.¹ Several quinolones are already being marketed, e.g., norfloxacin 1,² ofloxacin 2,³ enoxacin (Flumark) 3,⁴ and

ciprofloxacin 4.⁵ The success of these compounds has caused an increase in efforts to produce even more efficacious agents, leading to the current list of compounds with exciting clinical potential. These candidates include lomefloxacin (5),⁶ tosufloxacin (6),⁷ sparfloxacin (AT-4140/CI-978) (7),⁸ WIN 57273 (8),⁹ and PD 127391 (9)¹⁰

- (1) For a comprehensive review through 1976, see: (a) Albrecht, R. Development of Antibacterial Agents to the Nalidixic Acid Type. *Prog. Drug. Res.* 1977, 21, 9-104. More recent review include: (b) White, D. R.; Davenport, L. C. Antibacterial Agents. *Annu. Rep. Med. Chem.* 1990, 25, 109-118. (c) Heck, J. V. Antibacterial Agents. *Annu. Rep. Med. Chem.* 1989, 24, 101-110. (d) Jack, D. B. Recent Advances in Pharmaceutical Chemistry. The 4-Quinolone Antibiotics. *J. Clin. Hosp. Pharm.* 1986, 11, 75-93. (e) Bergan, T. Quinolones. In *The Antimicrobial Agents Annual/1*; Peterson, P. K., Verhoef, J., Eds.; Elsevier: New York, 1986; Vol. 12, pp 164-178. (f) Vergin, H.; Metz, R. Review of Developments in Fluoroquinolones. *Drugs Today* 1991, 27, 177-192.
- (2) Koga, H.; Itoh, A.; Murayama, S.; Suzue, S.; Irikura, T. Structure-Activity Relationships of Antibacterial 6,7- and 7,8-Disubstituted 1-Alkyl-1,4-dihydro-4-oxoquinoline-3-carboxylic Acids. *J. Med. Chem.* 1980, 23, 1358-1363.
- (3) Hayakawa, I. Eur. Pat. 47005; *Chem. Abstr.* 1982, 97, 55821b.
- (4) Matsumoto, J.; Miyamoto, T.; Minamida, A.; Nishimura, Y.; Egawa, H.; Nishimura, H. Pyridinecarboxylic Acids as Antibacterial Agents. 2. Synthesis and Structure-Activity Relationships of 1,6,7-Trisubstituted-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic Acids, Including Enoxacin, a New Antibacterial Agents. *J. Med. Chem.* 1984, 27, 292-301.

- (5) Wise, R.; Andrews, J. M.; Edwards, L. J. In Vitro Activity of Bay 09867, a New Quinoline Derivative, Compared with Those of Other Antimicrobial Agents. *Antimicrob. Agents Chemother.* 1983, 23, 559-564.
- (6) Chin, N. X.; Novelli, A.; Neu, H. C. In Vitro Activity of Lomefloxacin (SC-4711; NY-198), a Difluoroquinoline-3-carboxylic Acid, Compared with Those of Other Quinolones. *Antimicrob. Agents Chemother.* 1988, 32, 656-662. Wise, R.; Andrews, J. M.; Ashby, J. P.; Matthews, R. S. In Vitro Activity of Lomefloxacin, a New Quinolone Antimicrobial Agent, in Comparison with Those of Other Agents. *Antimicrob. Agents Chemother.* 1988, 32, 617-622.
- (7) Chu, D. T. W.; Prabhavathi, B. F.; Akiyo, K. C.; Pihuleac, E.; Nordeen, C. W.; Maleczka, R. E.; Pernet, A. G. Synthesis and Structure-Activity Relationships of Novel Arylfluoroquinolone Antibacterial Agents. *J. Med. Chem.* 1985, 28, 1558-1564.
- (8) Miyamoto, T.; Matsumoto, J.; Chiba, K.; Egawa, H.; Shibamori, K.; Minamida, A.; Nishimura, Y.; Okada, H.; Kataoka, M.; Fujita, M.; Hirose, T.; Nakano, J. Synthesis and Structure-Activity Relationships of 5-Substituted 6,8-Difluoroquinolones, Including Sparfloxacin, a New Quinolone Antibacterial Agent with Improved Potency. *J. Med. Chem.* 1990, 33, 1645-1656.