

Derivatives of 5-[[1-(4'-Carboxybenzyl)imidazolyl]methylidene]hydantoin as Orally Active Angiotensin II Receptor Antagonists^{1,2}

Jeremy J. Edmunds,* Sylvester Klutchko, James M. Hamby, Amy M. Bunker, Cleo J. C. Connolly, R. Thomas Winters, John Quin III, Ila Sircar, John C. Hodges, Robert L. Panek, Joan A. Keiser, and Annette M. Doherty

Department of Chemistry and Pharmacology, Parke-Davis Pharmaceutical Research, Division of Warner-Lambert Company, 2800 Plymouth Road, Ann Arbor, Michigan 48106-1047

Received May 8, 1995[®]

A series of 5-[[1-(4'-carboxybenzyl)imidazolyl]methylidene]hydantoin have been prepared and evaluated as *in vitro* and *in vivo* angiotensin II (Ang II) antagonists. Variation of substituents on the hydantoin ring leads to potent and selective Ang II antagonists with nanomolar IC₅₀ values at the AT₁ receptor and negligible affinity for the AT₂ receptor. Preferred substituents include an *n*-butyl at R₁ and an alkyl or heteroarylmethyl substituent at R₂. The selection of the R₂ substituent was guided in part by the calculation of its log *P* since a significant correlation was observed between CLOGP and AT₁ binding affinity. The biphenyl tetrazole pharmacophore, common to a number of AT₁ antagonists, could be replaced by, for example, a 4-carbomethoxyphenyl substituent resulting in potent Ang II antagonists both *in vitro* and *in vivo*. A representative compound of this series is **57**, which reduced the mean arterial blood pressure of renal hypertensive rats by 40% at 30 mg/kg *po* and by 25% at 10 mg/kg *po*. In addition this compound was efficacious in the salt-deplete normotensive monkey model maximally decreasing blood pressure 27% at 10 mg/kg *po*. In summary, these compounds belong to a novel class of Ang II antagonists that lack the biphenyl tetrazole moiety yet display appreciable and long lasting oral activity.

Introduction

The renin angiotensin system plays an important role in the regulation of blood pressure in addition to electrolyte homeostasis.^{3,4} The kidney responds to a reduction of blood pressure by the release of renin from the juxtaglomerular cells surrounding the afferent arterioles.⁵ This enzyme cleaves angiotensinogen to angiotensin I which is in turn converted to the potent vasoconstrictor angiotensin II (Ang II) by the nonspecific carboxydipeptidase angiotensin-converting enzyme (ACE).⁶ As a vasoconstrictor and stimulator of aldosterone release, Ang II increases peripheral resistance and expands blood volume as dual mechanisms for restoring blood pressure. In renin dependent hypertension, this blood pressure-regulating mechanism is compromised resulting in abnormal cardiac output and peripheral resistance.⁷

The renin angiotensin system offers several pharmacologically distinct approaches for antihypertensive therapy. These approaches include inhibitors of renin release⁸ as well as those of the enzyme renin.⁹ Although a number of renin inhibitors have entered clinical trials to date, their poor bioavailability has thus far precluded further development.¹⁰ A particularly successful pharmacologic approach, however, has involved the inhibition of the ACE.^{11,12} Inhibitors such as captopril and enalapril have been shown to be clinically efficacious in the treatment of hypertension and, more recently, congestive heart failure.¹³ These ACE inhibitors are not without side effects,^{14–16} and as a result, recent efforts have focused upon the discovery of Ang II receptor antagonists.¹⁷

Saralasin, a specific Ang II antagonist, lowered blood pressure in patients with elevated renin levels but, in

addition, exhibited partial agonist effects *in vivo* and poor bioavailability.^{18,19} This peptide, however, has served to demonstrate the usefulness of Ang II blockade and hence established several research programs directed toward the discovery of nonpeptide Ang II antagonists.^{20–25} The first orally active, nonpeptide Ang II receptor antagonist, DuP 753 (Cozaar), has recently been marketed for the clinical treatment of hypertension.²⁶

We have previously described aspects of our research programs relating to both AT₁²⁷ and AT₂²⁸ specific ligands and now describe a novel series of potent AT₁ ligands. One of our strategies for developing an alternate series of Ang II antagonists required an investigation of the notion that the biphenyl tetrazole moiety of most Ang II antagonists is a requirement for oral activity.²⁹ In fact recent reports have demonstrated that the biphenyl tetrazole moiety of L-158,809³⁰ could be replaced with acylsulfonamides³¹ or (dialkylphenoxy)phenylacetic acids³² while still retaining excellent oral activity. Furthermore, several reports from the SKF group^{33–36} describing the structure–activity relationship (SAR) of a series of 1-(carboxybenzyl)imidazole-5-acrylic acids, exemplified by SKF 108566 (**1**), was particularly interesting since this series lacked the biphenyl tetrazole moiety. We considered that a suitably substituted hydantoin could replace the acrylic acid of SKF 108566 and allow the appropriate positioning of hydrogen bond donor and hydrophobic functionality as previously dictated by SAR studies.^{27,37–39} The present paper describes the synthesis and SAR of a series of 5-[[1-(4'-carboxybenzyl)imidazolyl]methylidene]hydantoin **2** that resulted from this analysis (Figure 1).

Chemistry

Our initial synthetic route to prepare the requisite alkylidenehydantoin involved a base-catalyzed conden-

[®] Abstract published in *Advance ACS Abstracts*, August 15, 1995.





Figure 1. SKF 108566 (**1**) and 5-[[1-(4'-carboxybenzyl)-imidazolyl]methylidene]hydantoin **2**.

Scheme 1.^a Synthesis of Isomeric Mixture of (*Z*)- and (*E*)-Methylidenehydantoin^a




^a Reagents: (a) 1.1 equiv of HCl, KNCO, H₂O; (b) concentrated HCl or NaOMe, MeOH; (c) **4**, (Me)₄NOH, MeOH, H₂O, 100–120 °C.

sation of an alkyl-substituted hydantoin with methyl 4-[(2-butyl-5-formyl-1*H*-imidazol-1-yl)methyl]benzoate (**5**). The intermediate hydantoin **4** were prepared in a two-step process, starting from the appropriately *N*-substituted-glycine ester (Scheme 1).⁴⁰ For example, treatment of **3** with dilute aqueous HCl and KNCO provided an intermediate urea. Cyclization to 1-butyl-2,4-imidazolidinedione (1-butylhydantoin,⁴¹ **4**) was affected either with hot concentrated HCl or by treatment with NaOMe in MeOH. The condensation of **4** with **5** was initially performed under strongly basic conditions using (Me)₄NOH in alcohol–water mixtures at 100–120 °C. These conditions led to a mixture of regioisomeric products with the *E* isomer predominating and caused hydrolysis of the methyl ester to afford the corresponding acid. These *Z* and *E* regioisomeric products **6** and **7**, respectively (**97**–**105**), were then separated with some difficulty by crystallization or chromatography. The *Z* isomer was distinguished from the *E* isomer via a nuclear Overhauser effect. Particularly characteristic of the *E* isomer was the cross enhancement that was observed upon irradiating the methylene of the NCH₂R substituent and the vinylic proton of the methylidenehydantoin. In addition it was routinely observed that the imidazole proton of the *E* isomer exhibited a downfield shift attributable to the anisotropic deshielding by the π electrons in the carbonyl group.


An alternative method of condensation that offered greater control of regiochemistry about the double bond involved the use of diethyl (2,5-dioxo-3-butyl-4-imidazolyl)phosphonate (**8**) (Scheme 2).^{42,43} Thus **8** was condensed with the aldehyde **5** in the presence of 1,8-diazabicyclo[4.4.0]undec-7-ene (DBU) in methylene chloride to afford predominantly the required *Z* isomer **9**. The small amount of the corresponding *E* isomer **10** was separated through chromatography. These alkylidenehydantoin were then alkylated in excellent yield by

Scheme 2.^a Procedure for the Regioselective Synthesis of Methylidenehydantoin and Subsequent Alkylation^a



^a Reagents: (a) DBU, CH₂Cl₂; (b) R₁-X, K₂CO₃, DMF.

Scheme 3.^a Synthetic Route Allowing Functionalization of Hydantoin N₁ and N₃^a




^a Reagents: (a) BuI, K₂CO₃, DMF, room temperature; (b) **5**, DBU, CH₂Cl₂; (c) 2-methyl-4-(chloromethyl)thiazole, K₂CO₃, DMF.

treatment with an alkyl iodide and potassium carbonate in DMF to afford, for example, **11**.

Once preliminary SAR studies established that the preferred substituent at R₁ was butyl, we modified the synthetic route to allow alkylation of R₂ as the final synthetic step. As outlined in Scheme 3 the intermediate aldehyde **5** was condensed with the hydantoin phosphonate **13** to afford the useful intermediate **14** in good yield. Although this condensation afforded some corresponding *E* isomer, typically ca. 10%, as evident by NMR of the crude reaction mixture, the *Z* isomer could be isolated directly by crystallization of the reaction mixture after workup.

The phosphonate **13** was prepared, in excellent yield, by the treatment of **12** with butyl iodide and potassium carbonate in DMF. In fact this was a general procedure with the reaction tolerant of a number of electrophiles. In the case of less reactive electrophiles such as SEM-Cl, MOM-Cl, and MEM-Cl and some alkyl bromides, it was necessary to employ NaH in THF or DMF to afford good yields of the required products **80**–**88**.

This intermediate **14** was then reacted with a variety of electrophiles to afford the majority of compounds used to explore the SAR described herein. Again K₂CO₃ or NaH was employed as base, and only 1 equiv of the

Scheme 4. Synthesis of 5-[[1-(4'-Carboxybenzyl)imidazolyl]methylidene]hydantoin^s

^a Reagents: (a) 13, DBU, CH₂Cl₂; (b) BuI, K₂CO₃, DMF, room temperature; (c) TFA, CH₂Cl₂.

electrophile was added to prevent quarternization of the substituted imidazole. The final products **38–79** were simply isolated as their hydrochloride salts which allowed direct crystallization from the reaction mixture (Scheme 3).


During the attempted saponification of the esters, we noted that only partial hydrolysis was affected by dilute aqueous hydrochloric acid and that with concentrated acid substantial *Z* to *E* isomerization occurred. Basic saponification was equally fruitless due to competing urea formation as a result of hydantoin ring cleavage.

The problem of saponification was overcome by the use of *tert*-butyl benzoate esters (Scheme 4). These esters behaved similarly to the corresponding methyl esters during the reaction with the hydantoin phosphonates and the alkylation procedure but, in addition, allowed facile deesterification with trifluoroacetic acid in methylene chloride without concomitant *Z* to *E* isomerization to afford the required products **89–96** and **107–110**.


Scheme 5 outlines the chemistry used to prepare a number of alkylidenehydantoin that incorporate a variety of benzoate ester surrogates. The aldehydes **19**,^{21,44} **23**,³³ **27**,⁴⁵ **30**,⁴⁶ and **35**⁴⁷ underwent condensation with the hydantoin phosphonates in the expected fashion. In these cases the *Z* isomers were typically separated by chromatography in good yield. Fortunately detritylation of the protected tetrazoles, with citric or acetic acid, occurred without significant *Z* to *E* isomerization to afford **20**, **21** and **33**, **34**.

Biological Assays


The alkylidenehydantoin were evaluated as Ang II antagonists by a combination of *in vitro* AT₁ receptor binding and isolated vascular ring contraction assays as previously described.⁴⁸ Selected compounds were also evaluated in an AT₂ binding assay, using membranes isolated from the rabbit uterus, and were consistently found to lack significant binding activity at this receptor subtype. The AT₁ binding assay, which evaluated the ability of the test compounds to displace [¹²⁵I]Ang II in a rat liver membrane preparation, was used to select compounds for a functional assay. This functional assay studied the ability of the test compound to inhibit Ang II-induced vasoconstriction in isolated

Scheme 5. Analogs Incorporating Benzoate Ester Surrogates


Reagents: (a) 8, NaH, MeOH; CH₃CO₂H, MeOH (b) Trityl-20, H₂, Pd/C; CH₃CO₂H, MeOH



Reagents: (a) 8, NaH, MeOH; (b) BuI, K₂CO₃, DMF




Reagents: (a) 8, NaH, MeOH; (b) Citric acid, MeOH, reflux



Reagents: (a) 13, NaH, MeOH; 2-methyl-4-(chloromethyl)thiazole, K₂CO₃, DMF; (b) 2N HCl, reflux

rabbit aorta rings (RAR). Compounds which displayed good activity in these *in vitro* screens were then evaluated in two *in vivo* screens. The majority of the potent compounds were examined in conscious, renal hypertensive rats, and the decrease in blood pressure was monitored over a period of 20 h.²⁷ A compilation of the activities of these hydantoin is given in Table 1. Finally selected compounds were tested in conscious, high-renin, normotensive male cynomolgus monkey model.⁴⁹

Table 1. Compilation of Rat Liver Membrane Binding (AT₁), Inhibition of Ang-II Induced Rabbit Aortic Ring Contraction (RAR), and Oral Antihypertensive Activity in Renal Hypertensive Rats (RHR) of Substituted Alkylidenehydantoin^a

compd	R ₁	R ₂	R ₃	isomer	IC ₅₀ (nM)		RHR % max	
					AT ₁	RAR	30 mg/kg	10 mg/kg
9	H	butyl	CO ₂ Me	Z	3.5	30	-11	NT
20	H	butyl	2-(1 <i>H</i> -tetrazol-5-yl)phenyl	Z	1.4	40	IA	NT
21	H	butyl	2-(1 <i>H</i> -tetrazol-5-yl)phenyl	E	2.7	NT	NT	NT
24	H	butyl	2-carbomethoxyphenyl	Z	15.9	NT	NT	NT
25	H	butyl	2-carbomethoxyphenyl	E	9.9	NT	NT	NT
26	Butyl	butyl	2-carbomethoxyphenyl	Z	29.6	NT	NT	NT
28	H	butyl	trifluoromethanesulfonamido	Z	3.3	120	-11	NT
29	H	butyl	trifluoromethanesulfonamido	E	4.2	NT	NT	NT
33	H	butyl	1 <i>H</i> -tetrazol-5-yl	Z	7.1	40	IA	NT
34	H	butyl	1 <i>H</i> -tetrazol-5-yl	E	2.5	NT	NT	NT
36	butyl	(2-methyl-4-thiazolyl)methyl	1-(methoxycarbonyl)cyclopent-1-yl	Z	8.0	NT	NT	NT
37	butyl	(2-methyl-4-thiazolyl)methyl	1-carboxycyclopent-1-yl	Z	1.9	NT	-16	NT
38	butyl	(4-aminophenyl)methyl	CO ₂ Me	Z	3.0	NT	NT	NT
39	butyl	(3,4-dihydroxyphenyl)methyl	CO ₂ Me	Z	4.9	NT	NT	NT
40	butyl	(4-methoxyphenyl)methyl	CO ₂ Me	Z	9.8	NT	NT	NT
41	butyl	(4-cyanophenyl)methyl	CO ₂ Me	Z	11.1	NT	NT	NT
42	butyl	phenylmethyl	CO ₂ Me	Z	12.4	NT	NT	NT
43	butyl	[4-methoxycarbonyl]phenyl]methyl	CO ₂ Me	Z	14.3	NT	NT	NT
44	butyl	(4-nitrophenyl)methyl	CO ₂ Me	Z	16.9	NT	NT	NT
45	butyl	(4-chlorophenyl)methyl	CO ₂ Me	Z	17.6	500	-15	NT
46	butyl	(4-methylphenyl)methyl	CO ₂ Me	Z	34.6	NT	NT	NT
47	butyl	[4-(trifluoromethyl)phenyl]methyl	CO ₂ Me	Z	159.0	NT	NT	NT
48	butyl	butyl	CO ₂ Me	Z	8.5	460	-27	NT
49	butyl	hexyl	CO ₂ Me	Z	40.2	NT	NT	NT
50	butyl	(2-naphthalenyl)methyl	CO ₂ Me	Z	104.0	NT	NT	NT
51	butyl	diphenylmethyl	CO ₂ Me	Z	884.0	NT	NT	NT
52	butyl	c-hexylmethyl	CO ₂ Me	Z	3.3	NT	NT	NT
53	butyl	pentyl	CO ₂ Me	Z	6.1	NT	NT	NT
54	butyl	4,4,4-trifluorobutyl	CO ₂ Me	Z	6.4	NT	NT	NT
55	butyl	CH=CMe ₂	CO ₂ Me	Z	11.1	NT	NT	NT
56	butyl	(2-thienyl)methyl	CO ₂ Me	Z	3.4	370	-28	NT
57	butyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	3.8	350	-40	-25
58	butyl	(3-thienyl)methyl	CO ₂ Me	Z	4.8	NT	-36	NT
59	butyl	phenethyl	CO ₂ Me	Z	31.0	NT	NT	NT
60	butyl	(5-bromo-2-thienyl)methyl	CO ₂ Me	Z	17.8	NT	NT	NT
61	butyl	(3-methyl-2-thienyl)methyl	CO ₂ Me	Z	23.9	NT	NT	NT
62	butyl	(4-bromo-2-thienyl)methyl	CO ₂ Me	Z	21.9	NT	NT	NT
63	butyl	(2-amino-4-thiazolyl)methyl	CO ₂ Me	Z	1.2	290	-37	-39
64	butyl	(3,5-dimethyl-4-isoxazolyl)methyl	CO ₂ Me	Z	198.0	NT	NT	NT
65	butyl	(3-thienyl)ethyl	CO ₂ Me	Z	14.6	NT	NT	NT
66	butyl	(2-thienyl)ethyl	CO ₂ Me	Z	14.5	NT	NT	NT
67	butyl	(2-pyridinyl <i>N</i> -oxide)methyl	CO ₂ Me	Z	260.0	NT	NT	NT
68	butyl	(2-pyridinyl)methyl	CO ₂ Me	Z	5.8	490	-20	NT
69	butyl	(5-bromo-3-thienyl)methyl	CO ₂ Me	Z	25.4	NT	NT	NT
70	butyl	(5-isoxazolyl)methyl	CO ₂ Me	Z	10.1	>100	-23	NT
71	butyl	(4-thiazolyl)methyl	CO ₂ Me	Z	8.0	NT	NT	NT
72	butyl	(1 <i>H</i> -tetrazol-5-yl)methyl	CO ₂ Me	Z	41.7	NT	NT	NT
73	butyl	(1 <i>H</i> -imidazol-5-yl)methyl	CO ₂ Me	Z	54.5	NT	NT	NT
74	butyl	(5-methyl-3-oxazolyl)methyl	CO ₂ Me	Z	8.5	NT	NT	NT
75	butyl	(3-chlorophenyl)methyl	CO ₂ Me	Z	117.0	NT	NT	NT
76	butyl	(2-methyl-4-imidazolyl)methyl	CO ₂ Me	Z	245.0	NT	NT	NT
77	butyl	(3-methylphenyl)methyl	CO ₂ Me	Z	18.5	NT	NT	NT
78	butyl	(3-furanyl)methyl	CO ₂ Me	Z	8.3	>100	-22	NT
79	butyl	[5-methoxycarbonyl]-2-thienyl]methyl	CO ₂ Me	Z	4.2	NT	NT	NT
80	methyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	16.6	NT	NT	NT
81	ethyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	11.1	NT	NT	NT
82	propyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	9.0	NT	NT	NT
83	hexyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	5.3	>100	-34	NT
84	hydroxymethyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	35.7	NT	NT	NT
85	3-hydroxypropyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	47.9	NT	NT	NT
86	methoxymethyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	47.8	NT	NT	NT
87	(thiomethyl)ethyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	9.6	NT	NT	NT
88	trifluorobutyl	(2-methyl-4-thiazolyl)methyl	CO ₂ Me	Z	70.0	NT	NT	NT
89	butyl	butyl	CO ₂ H	Z	1.3	14	-45	-21
90	butyl	butyl	CO ₂ H	E	0.5	NT	NT	IA
91	butyl	(2-thienyl)methyl	CO ₂ H	Z	0.7	13	-28	NT
92	butyl	(2-thienyl)methyl	CO ₂ H	E	0.6	NT	NT	NT

Table 1 (Continued)

compd	R ₁	R ₂	R ₃	isomer	IC ₅₀ (nM)		RHR % max	
					AT ₁	RAR	30 mg/kg	10 mg/kg
93	butyl	(2-amino-4-thiazolyl)methyl	CO ₂ H	Z	0.6	2.9	IA	NT
94	butyl	(2-amino-4-thiazolyl)methyl	CO ₂ H	E	0.5	NT	NT	NT
95	butyl	(2-methyl-4-thiazolyl)methyl	CO ₂ H	Z	1.2	2.6	-34	NT
96	butyl	(2-methyl-4-thiazolyl)methyl	CO ₂ H	E	1.7	NT	NT	NT
97	H	butyl	CO ₂ H	Z	2.8	28	-17	NT
98	H	butyl	CO ₂ H	E	2.4	44	IA	NT
99	H	3-methylbutyl	CO ₂ H	Z	6.6	32	NT	NT
100	H	3-methylbutyl	CO ₂ H	E	4.9	74	NT	NT
101	H	phenylmethyl	CO ₂ H	Z	4.0	35	NT	NT
102	H	phenylmethyl	CO ₂ H	E	3.4	>100	NT	NT
103	H	(2-thienyl)methyl	CO ₂ H	Z	2.3	18	-21	NT
104	H	(2-thienyl)methyl	CO ₂ H	E	2.6	NT	IA	NT
105	H	(4-chlorophenyl)methyl	CO ₂ H	Z	3.0	28	-17	NT
106	H	(2-thienyl)methyl	CO ₂ Me	Z	7.1	28	-19	NT
107	butyl	(5-methyl-2-thienyl)methyl	CO ₂ H	Z	0.6	5.3	-22	NT
108	butyl	(4-methyl-2-thienyl)methyl	CO ₂ H	Z	1.5	21	-16	NT
109	butyl	(5-methyl-3-thienyl)methyl	CO ₂ H	Z	0.7	9.2	-23	-28
110	butyl	(3-thienyl)methyl	CO ₂ H	Z	0.9	10	-42	-28

^a NT, not tested; IA, inactive (% max < 10%).




Figure 2. Initial chemical lead.

Results and Discussion

With the discovery of our initial lead, (*E*)-4-[[2-butyl-5-[(2,5-dioxo-3-phenyl-4-imidazolidinylidene)methyl]-1*H*-imidazol-1-yl]methyl]benzoic acid (AT₁ IC₅₀ = 0.5 μM) (Figure 2) we chose to optimize in vitro activity by a three-stage approach. Our first step was to vary the substituents at R₁ and R₂ guided in part by the physicochemical characteristics of the substituents, SAR of SKF 108566, and peptide analogs of Ang II.^{50,51} Secondly, we investigated the in vitro activity of the *Z* and *E* geometrical isomers of the more potent analogs. Finally we replaced the 4-carboxybenzyl substituent with a number of substituents of varying p*K*_a and steric requirements.⁵²

The selection of an optimal R₂ hydantoin substituent was pursued by a combination of three approaches. One of these involved the incorporation of a number of substituted aromatics at R₂ while targeting the *Z* isomer with an *n*-butyl substituent at R₁ (38–47). The *Z* isomer was pursued initially since it corresponded to the orally active geometrical isomer from the SKF 108566 series, and an *n*-butyl substituent was incorporated at R₁ for synthetic simplicity. Our selection of aromatic substituents was guided by correlating the σ (electronic) and Π (hydrophobic) values of the substituent with in vitro activity.⁵³ As an initial approach we attempted to correlate the in vitro activity of the derivatives via the Topliss^{54,55} approach and by employing Craig plots.⁵⁶ Both the Topliss and Craig approaches indicated that substituents with both a negative σ and Π were optimal for in vitro activity (38, 39). However, at this point we could not determine whether the in vitro activity was a result of the electronic or lipophilic nature of the substituent. We therefore compared the in vitro activity via regression analysis




Figure 3. Comparison of the in vitro AT₁ receptor binding affinity of compounds 38–79 with calculated log *P* of the substituent R₂.

with the σ, Π, MR (molecular refractivity), *F* (field inductive), and *R* (resonance) values for the aromatic substituents. Significant correlation was only readily apparent with Π, indicating that the lipophilicity of the substituent was crucial in determining activity.⁵⁷ Furthermore by measuring the HPLC retention times of each of the derivatives (38–47), under isocratic conditions, we observed the excellent correlation⁵⁸ between experimentally measured HPLC retention time and Π.⁵⁹ This data prompted the interesting conclusion that in vitro binding activity could be predicted by simply observing the HPLC retention time of the derivative.

Concomitantly to the synthesis of benzyl substituents at R₂, we also investigated the binding affinity of analogs which were alkyl substituted at R₂ (48, 49, 52–55). Interestingly these analogs also demonstrated excellent in vitro activity, and in the light of the results of the above analysis, this was not surprising due to the similarity in the CLOGP⁶⁰ values for the alkyl substituents and benzyl substituents (42, 48). However, although in vitro activity compared favorably with the lipophilicity of the substituent, the unexpected potent in vitro activity of compound 52 demonstrates a limitation of simply relying on the CLOGP values to predict activity (Figure 3).

As an extension of this correlation of in vitro activity with lipophilicity of the R₂ substituent, we then investigated a variety of heterocycles^{61,63} (56–79) which were calculated to have CLOGP values that expanded upon the range of the alkyl and substituted benzylys previously

examined. Although we realized that many problems exist in calculating log *P* values for heterocycles due to hidden steric and electronic properties, we were simply looking for a semiquantitative guide to aid in the selection of appropriate heterocycles.

In fact as can be seen from Table 1 and Figure 3, a remarkable number of heterocycles afforded potent in vitro compounds, the most potent of which tended to be substituted thiazole and thiophene heterocycles (57, 58). Furthermore Figure 3 demonstrates a parabolic dependence ($P < 0.005$) between the in vitro activity and the lipophilic nature of the substituent, although it should be pointed out that calculating an optimal log *P* is hindered by the apparent spread ($R^2 = 0.35$) of compounds. Two compounds, 64 and 76, in particular showed surprisingly poor IC₅₀'s based on their calculated log *P*. This may be the result of unusual electronic or steric effects of the substituents and hence an unfavorable association with the receptor. Of particular interest was the excellent activity of 63. This unexpected level of activity, based upon its corresponding log *P*, may be the result of the known inherent problems associated with calculating the log *P* of substituents that contain basic residues. However, in order to address the question of whether the calculated log *P* values are truly reflective of the measured log *P* we selected several representative compounds (42, 48, 49, 52, 56–58, 60, 61) and measured their log *P* by the octanol/water partition method. Gratifyingly the regression analysis indicated an excellent correlation⁶⁴ between the calculated and measured log *P* and furthermore demonstrated the usefulness of the CLOGP algorithm in calculating the log *P* of at least some of the heteroaryl substituents studied here.

Having studied the SAR's for the R₂ position, we then systematically studied the effect of the R₁ substituent to investigate whether a butyl substituent was optimal for activity. We uniformly increased the alkyl chain length at R₁, in addition to introducing a number of alkyl chains containing heteroatoms, while incorporating one of the more potent heterocycles in the form of (2-methyl-4-thiazolyl)methyl at the R₂ position (57, 80–88). From this analysis we determined that optimal in vitro binding affinity was indeed afforded by a four-carbon alkyl chain at R₁.

At this point then, we felt assured that the R₁ and R₂ substituents had been adequately explored, and hence we turned our attention to some secondary considerations. The first of these was the excellent in vitro activity of the corresponding *E* isomers. As can be seen in Table 1, the in vitro activity was maintained in both the *E* and *Z* isomers 89–104. Furthermore although only a limited number of these *E* isomers (98, 100, 102) were evaluated in the RAR assay, it was apparent that they inhibited the contraction of isolated rabbit aorta rings but to a lesser extent than the corresponding *Z* isomers (97, 99, 101). It soon became apparent that despite such good in vitro activity, the *E* isomers 90, 98, and 104 were devoid of oral activity in the renal hypertensive rat (RHR) despite the fact that the corresponding *Z* isomers (89, 97, 103) displayed reasonable oral activity in the same assay. We were further surprised to find that the saturated analog of 97 or 98 had relatively poor in vitro activity.⁶⁵ One possible explanation for the lack of oral activity of the *E* isomer may be a preferential metabolic deactivation by Michael

addition to the *E* isomer.⁶⁶ Clearly it is difficult to draw definitive conclusions concerning the oral activity of the *E* isomers due to the lack of metabolic studies and the relatively few *E* isomers that were tested orally.

In our final approach to optimizing in vitro activity, we examined a variety of benzoic acid replacements (20, 21, 24–26, 28, 29, 33, 34, 36, 37). Although these compounds displayed excellent in vitro activity, their lack of in vivo activity and lengthy synthesis precluded further study.

Compounds which showed appreciable inhibition of [¹²⁵I]Ang II in rat liver membrane preparations (AT₁) were generally also evaluated for antagonism of Ang-II induced contraction of rabbit aortic rings (RAR). The results of this assay clearly demonstrated that the acids were more potent antagonists than the corresponding esters and usually confirmed the potency of the most active analogs. However, the critical test was the determination of oral activity in renal hypertensive rats. Due to the intensive nature of this assay, we limited the test to compounds that displayed in vitro activity (AT₁) better than a threshold value of 20 nM and/or showed significant activity in RAR. Initially compounds were tested at a single dose to demonstrate oral activity, and then the most active analogs were tested at multiple doses to demonstrate a dose–response relationship. Compounds 9 and 48, 56 and 106, 89 and 97, and 91 and 103 demonstrate the trend toward enhanced in vivo activity afforded by an alkyl chain at R₁. Furthermore, although the in vitro activity was generally slightly superior for the analogs incorporating an acid at R₃ (9 and 97, 48 and 89, 56 and 91, 57 and 95), the in vivo activity of the acids and esters were similar, yet there was a notable exception (63, 93). Due to the inherent variability associated with measuring blood pressure responses after oral dosing, it is difficult to speculate as to why certain analogs show relatively poor oral activity in light of potent in vitro activity. Obviously such factors as absorption, distribution, metabolism, and elimination are important, and the correlation of these parameters with oral activities is better addressed by studies which determine the pharmacokinetics and pharmacodynamics of the compounds under study, which is beyond the scope of this report.

Although all of the compounds previously described were competitive antagonists, we discovered that slight modification of the benzoic acid pharmacophore by the introduction of a 2-chloro substituent⁶⁷ at R₄ afforded an insurmountable antagonist,⁶⁸ 111 (Figure 4), as has been previously seen in the SKF 108566 series of compounds. The increased lipophilicity afforded by the chloro substitution presumably causes a receptor–ligand interaction that leads to a partial but constant suppression of the maximal contraction of the rabbit aortic ring induced by Ang II.⁶⁹ However, when dosed orally, 111 demonstrated a sustained and rapid reduction in blood pressure at doses of 3 and 10 mg/kg, which was similar to the effect seen with 57 (Figures 5 and 6).

To this point all of the alkylidenehydantoin had been tested in tissues isolated from rat and rabbit and the oral activity of the compounds assayed in the renal





Figure 4. Surmountable (**57**) and insurmountable (**111**) AT₁ receptor antagonists in rabbit aorta.




Figure 5. Effect of compound **57** at 3 mg/kg po ($n = 7$) and 30 mg/kg po ($n = 9$) on mean arterial blood pressure in renal hypertensive rats.




Figure 6. Effect of compound **111** at 10 mg/kg po ($n = 4$) and 3 mg/kg po ($n = 5$) on mean arterial blood pressure in renal hypertensive rats.

hypertensive rat. We therefore considered that a primate model of hypertension would be a reasonable extension of the previous studies and hence provide some insight to the suitability of these compounds for

Table 2. Effect of Compounds **57** and **111** at 10 mg/kg on Salt-Deplete Normotensive Monkeys

compd	blood pressure (mm/Hg)		
	baseline	min.	at 4.5 h
57	101 ± 8	74 ± 6	88 ± 5
111	100 ± 4	80 ± 2	85 ± 6

Table 3. Comparison of AT₁ and AT₂ Binding Affinities of Representative [(Benzylimidazolyl)alkylidene]hydantoin

compd	R ₁	R ₂	R ₃	IC ₅₀ (nM)	
				AT ₁	AT ₂
20	H	butyl	2-(1 <i>H</i> -tetrazol-5-yl)phenyl	1.4	789
37	butyl	(2-methyl-4-thiazolyl)methyl	1-carboxycyclopent-1-yl	1.9	549
48	butyl	butyl	methoxycarbonyl	8.5	5870

clinical studies. Gratifyingly compounds **57** and **111**, when dosed orally in a primate model of renin dependent hypertension, caused a reduction in blood pressure that was evident up to 4.5 h (Table 2).

Finally the selectivity of these Ang II antagonists for the AT₁ receptor subtype was evaluated (Table 3). From this analysis it was determined that modification of the benzyl 4-substituent afforded compounds with modest AT₂ binding affinity. In fact this trend has been observed previously⁷⁰⁻⁷² and suggests that further modification of the R₃ substituent would also be a fruitful avenue of exploration in the design of dual AT₁/AT₂ ligands for this series of compounds.

Conclusion

This report describes a series of novel substituted 5-[[1-(4'-carboxybenzyl)imidazolyl]methylidene]hydantoin that are potent orally active angiotensin II (AT₁) receptor antagonists. The in vitro potency was demonstrated to closely correlate with the lipophilicity of the R₂ substituent. A good agreement was observed between the calculated log *P*, the log *P* measured by the octanol/water partition protocol, and the log *P* calculated by the HPLC for a variety of R₂ substituents, which further indicated that the CLOGP algorithm is a useful tool for predicting in vitro activity. A butyl substituent at R₁ was optimal for in vitro activity and furthermore enhanced the antihypertensive activity in renal hypertensive rats. A 4-carboxybenzyl substituent generally afforded the most potent compounds, although the corresponding esters were also orally active. Rather surprisingly replacement of the carboxylic acid with a tetrazole, (trifluoromethyl)sulfonamide, or tetrazolylphenyl substituent, although displaying excellent in vitro activity, afforded orally inactive compounds when tested in the renal hypertensive rat.

This class of compounds is represented by compound **57** which displayed a dose-related lowering of blood pressure, and the effect (~25% at 10 mg/kg po) was sustained for at least a period of 24 h in the renal hypertensive rat. This compound also caused a maximal blood pressure reduction of 27% (at 10 mg/kg po)

and a 13% reduction at 4.5 h in salt-deplete normotensive monkeys. This profile of activity and duration of action makes these compounds suitable candidates for antihypertensive therapy.

Experimental Section

Melting points are uncorrected and were determined on a Thomas-Hoover capillary melting point apparatus. Infrared (IR) spectra were determined in KBr on a Mattson Cygnus 100 or a Nicolet MX1 FTIR spectrophotometer. Proton magnetic resonance (NMR) were recorded on either a Varian XL-200 or XL-300, Unity-400, or Bruker AM250 instrument; shifts are reported in δ units relative to internal tetramethylsilane. Mass spectra were obtained on a Finnigan 4500 or a VG analytical 7070 E/HF spectrometer. Elemental analyses were performed on a CEC Model 240 elemental analyzer, and all compounds prepared had analytical results within 0.4% of the theoretical values, unless otherwise noted. Column chromatography was performed using E. Merck silica gel 60 (230–400 mesh). Thin layer chromatography was performed with glass-backed silica gel (60 F 254) plates employing either UV light or iodine vapor for visualization. Purity of compounds was assayed by analytical HPLC employing a Beckman Ultrasphere C18 column (0.46 cm \times 25 cm) with a mobile phase consisting of MeCN/H₂O and a flow rate of 1.5 mL/min detecting at 214 or 254 nm; in all cases the purity exceeded 98%.

1-Butyl-2,4-imidazolidinedione (1-Butylhydantoin, 4). To a solution of ethyl *N*-butylglycinate (31.80 g, 0.20 mol) in 1 N HCl (220 mL) was added potassium cyanate (24.4 g, 0.30 mol) in portions. The resulting solution was warmed to 90 °C and held at this temperature for 10 min. Concentrated HCl (250 mL) was added cautiously and the mixture heated at 90–100 °C for 30 min. After concentrating to dryness, the residue was extracted with 1:1 ethanol/methylene chloride (400 mL). The extract was evaporated in vacuo and then triturated with water (100 mL). Filtration afforded pure product (21.22 g, 68%): ¹H NMR (CDCl₃) 0.95 (t, *J* = 7.24 Hz, 3H), 1.33 (m, 2H), 1.52 (m, 2H), 3.37 (t, *J* = 7.35 Hz, 2H), 3.91 (s, 2H), 8.73 (brs, 1H). Anal. (C₇H₁₂N₂O₂) C, H, N.

(Z)-4-[[2-Butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoic Acid (6). A mixture of methyl 4-[[2-butyl-5-formyl-1H-imidazol-1-yl]methyl]benzoate (5) (3.44 g, 0.012 mol), 1-butylhydantoin (4) (2.40 g, 0.016 mol), 25% tetramethylammonium hydroxide in methanol (25.92 g, 0.071 mol), and water (12 mL) was heated in an open flask at an oil bath temperature of 135–150 °C which caused the methanol to distill over. A yellow solution formed during this process, and the mixture was maintained at an oil bath temperature of 135–150 °C for a total of 45 min. The solution was cooled, and glacial acetic acid (5.04 g, 0.084 mol) and water (100 mL) were added. The precipitate was collected and washed with water to afford the title compound as a mixture of *Z* and *E* isomers (3.60 g, 70%). Crystallization from methanol and dichloromethane readily afforded the *E* isomer (1.16 g, 23%). The pure *Z* isomer (0.300 g, 6%) was obtained by repeated crystallization from ethyl acetate and hexane: ¹H NMR (DMSO) 0.63 (m, 3H), 0.84 (m, 7H), 1.28 (m, 2H), 1.60 (m, 2H), 2.68 (t, *J* = 7.33 Hz, 2H), 3.65 (brt, 2H), 5.38 (s, 2H), 6.25 (s, 1H) 7.10 (d, *J* = 8.55 Hz, 2H), 7.12 (s, 1H), 7.89 (d, *J* = 8.55 Hz, 2H), 11.42 (s, 1H), 13.00 (brs, 1H). Anal. (C₂₃H₂₈N₄O₄) C, H, N.

(E)-4-[[2-Butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoic acid (7): ¹H NMR (DMSO) 0.64 (m, 3H), 0.84 (t, *J* = 7.31 Hz, 3H), 0.98 (m, 4H), 1.33 (m, 2H), 1.60 (m, 2H), 2.72 (t, *J* = 7.36 Hz, 2H), 3.45 (brs, 2H), 5.54 (s, 2H), 5.86 (s, 1H), 7.07 (d, *J* = 8.25 Hz, 2H), 7.91 (d, *J* = 8.25 Hz, 2H), 8.24 (s, 1H), 11.22 (s, 1H), 13.00 (brs, 1H). Anal. (C₂₃H₂₈N₄O₄·0.25H₂O) C, H, N.

Diethyl (2,5-Dioxo-3-butyl-4-imidazolyl)phosphonate (8). To a solution of 1-butylhydantoin (4) (5.25 g, 0.034 mol) in acetic acid (45 mL), under nitrogen, at 90–100 °C was added a solution of bromine (1.86 mL) in acetic acid (15 mL) dropwise over 5 min. The mixture was maintained at this temperature for 90 min. At this time an aliquot revealed the reaction to be complete as judged by NMR. The solution was cooled,

concentrated, and then evaporated again from toluene (50 mL). This oil was dissolved in ether (70 mL) and treated with triethyl phosphite (5.0 mL, 0.029 mol). After stirring for 16 h the mixture was concentrated in vacuo. The oil was dissolved in chloroform and filtered through a pad of silica gel. Concentration in vacuo afforded the product as a viscous, colorless oil (7.69 g, 80%): ¹H NMR (CDCl₃) 0.93 (t, *J* = 7.00 Hz, 3H), 1.26–1.43 (m, 8H), 1.55 (m, 2H), 3.39 (m, 1H), 3.78 (m, 1H), 4.26 (m, 4H), 4.35 (d, *J* = 15.00 Hz, 1H), 8.40 (brs, 1H). Anal. (C₁₁H₂₁N₂O₅P) C, H, N.

Methyl (Z)-4-[[2-Butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoate (9). To a suspension of diethyl (3-butyl-2,5-dioxo-4-imidazolyl)phosphonate (8) (0.59 g, 1.85 mmol) and LiCl (0.160 g, 3.72 mmol) in acetonitrile (5 mL) was added DBU (0.554 mL, 3.70 mmol). After stirring for 5 min, methyl 4-[[2-butyl-5-formyl-1H-imidazol-1-yl]methyl]benzoate (5) (0.461 g, 1.54 mmol) was added in one portion. This wine-red solution was stirred for 16 h and evaporated in vacuo. Silica gel chromatography, eluting with chloroform, afforded the *Z* isomer (0.441 g, 65%): ¹H NMR (CDCl₃) 0.74 (t, *J* = 6.83 Hz, 3H), 0.89 (t, *J* = 7.27 Hz, 3H), 1.11 (m, 4H), 1.36 (m, 2H), 1.73 (m, 2H), 2.64 (t, *J* = 7.78 Hz, 2H), 3.78 (m, 2H), 3.90 (s, 3H), 5.19 (s, 2H), 6.31 (s, 1H) 7.03 (d, *J* = 8.12 Hz, 2H), 7.16 (s, 1H), 7.98 (d, *J* = 8.18 Hz, 2H), 8.60 (bs, 1H). Anal. (C₂₄H₃₀N₄O₄) C, H, N.

The more polar *E* isomer was then eluted from the column (0.096 g, 14%). **Methyl (E)-4-[[2-butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoate (10):** ¹H NMR (CDCl₃) 0.75 (t, *J* = 6.83 Hz, 3H), 0.90 (t, *J* = 7.27 Hz, 3H), 1.14 (m, 4H), 1.40 (m, 2H), 1.72 (m, 2H), 2.74 (t, *J* = 7.78 Hz, 2H), 3.38 (m, 2H), 3.91 (s, 3H), 5.25 (s, 2H), 5.61 (s, 1H), 7.06 (d, *J* = 8.12 Hz, 2H), 8.04 (d, *J* = 8.18 Hz, 2H), 8.45 (s, 1H) (NH absent). Anal. (C₂₄H₃₀N₄O₄) C, H, N.

Diethyl (1-Butyl-2,5-dioxo-4-imidazolyl)phosphonate (13). To a solution of diethyl (2,5-dioxo-4-imidazolyl)phosphonate (12) (6.00 g, 0.025 mol) in DMF (15 mL), under N₂, were added K₂CO₃ (17.5 g, 0.127 mol) and 1-iodobutane (2.9 mL, 0.025 mol). The mixture was stirred at room temperature for 16 h and then filtered and the solid washed with ethyl acetate. The filtrate was concentrated in vacuo to afford a yellow oil which was redissolved in ethyl acetate, washed with water and brine, and then dried over MgSO₄. Evaporation of solvents under high vacuum afforded the required product as a viscous yellow oil which solidified on standing (6.45 g, 88%): ¹H NMR (CDCl₃) 0.92 (t, *J* = 7.26 Hz, 3H), 1.29–1.42 (m, 8H), 1.59 (m, 2H), 3.51 (t, *J* = 7.17 Hz, 2H), 4.26 (m, 4H), 4.38 (d, *J* = 15.00 Hz, 1H) 6.80 (brs, 1H). Anal. (C₁₁H₂₁N₂O₅P) C, H, N.

Methyl (Z)-4-[[2-Butyl-5-[(1-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoate (14). To a solution of diethyl (1-butyl-2,5-dioxo-4-imidazolyl)phosphonate (13) (0.37 g, 1.28 mmol) in methylene chloride (5 mL) at room temperature were added DBU (0.38 mL, 2.54 mmol) and then methyl 4-[[2-butyl-5-formyl-1H-imidazol-1-yl]methyl]benzoate (5) (0.3 g, 1.00 mmol). This mixture was stirred for 16 h, diluted with ethyl acetate, and washed with brine. After drying over MgSO₄, the product was isolated, by chromatography eluting with 1% methanol in chloroform, as a yellow solid (0.305 g, 70%). A sample was recrystallized from ethyl acetate: ¹H NMR (CDCl₃) 0.83 (t, *J* = 7.07 Hz, 3H), 0.90 (t, *J* = 7.07 Hz, 3H), 1.22–1.40 (m, 4H), 1.62 (m, 4H), 2.67 (t, *J* = 7.36 Hz, 2H), 3.58 (t, *J* = 7.17 Hz, 2H), 3.91 (s, 3H), 5.23 (s, 2H), 6.29 (s, 1H), 7.00 (d, *J* = 8.12 Hz, 2H), 7.56 (s, 1H), 8.00 (d, *J* = 8.12 Hz, 2H), 9.28 (brs, 1H). Anal. (C₂₄H₃₀N₄O₄) C, H, N.

A small amount (10% yield) of the *E* isomer was isolated as well. **Methyl (E)-4-[[2-butyl-5-[(1-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoate (15):** ¹H NMR (CDCl₃) 0.86 (t, *J* = 7.27 Hz, 3H), 0.90 (t, *J* = 7.27 Hz, 3H), 1.28–1.38 (m, 4H), 1.62–1.80 (m, 4H), 2.65 (t, *J* = 7.26 Hz, 2H), 3.58 (t, *J* = 7.08 Hz, 2H), 3.89 (s, 3H), 5.21 (s, 2H), 5.90 (s, 1H), 7.00 (d, *J* = 8.12 Hz, 2H), 8.00 (d, *J* = 8.12 Hz, 2H), 8.10 (brs, 1H), 8.45 (s, 1H). Anal. (C₂₄H₃₀N₄O₄·0.33H₂O) C, H, N.

1,1-Dimethylethyl (Z)-4-[[2-Butyl-5-[(1-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoate (17). To a solution of diethyl (1-butyl-2,5-dioxo-

4-imidazolyl)phosphonate (**13**) (1.42 g, 4.86 mmol) in dichloromethane (10 mL), at -78°C , was added DBU (0.731 mL, 4.88 mmol). After stirring for 15 min, a solution of 1,1-dimethylethyl 4-[(2-butyl-5-formyl-1*H*-imidazol-1-yl)methyl]benzoate (**16**) (1.39 g, 4.06 mmol) in dichloromethane (5 mL) was added dropwise. This mixture was allowed to warm to room temperature and stirred for 16 h. The mixture was diluted with dichloromethane (30 mL) and washed with brine. After drying over MgSO_4 , the solvents were removed in vacuo to afford a yellow solid which was collected and washed with ether to afford the title compound (1.307 g, 67%). ^1H NMR revealed 8% of the corresponding *E* isomer (characteristic ^1H signals at δ 5.38, 6.07, 8.24) which could not be separated by chromatography: ^1H NMR (DMSO) 0.83 (t, $J = 7.38$ Hz, 3H), 0.86 (t, $J = 7.41$ Hz, 3H), 1.27 (m, 4H), 1.52 (s, 9H), 1.54 (m, 4H), 2.65 (t, $J = 7.27$ Hz, 2H), 3.42 (t, $J = 6.62$ Hz, 2H), 5.46 (s, 2H), 6.22 (s, 1H), 7.05 (d, $J = 8.16$ Hz, 2H), 7.75 (s, 1H), 7.87 (d, $J = 8.22$ Hz, 2H), 10.40 (s, 1H). Anal. ($\text{C}_{27}\text{H}_{36}\text{N}_4\text{O}_4 \cdot 0.2\text{H}_2\text{O}$) C, H, N.

2-Butyl-1-[[2'-(1*H*-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-5-formylimidazole (19**)**. A solution of 2-butyl-1-[[2'-(1*H*-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-4-chloro-5-formylimidazole (3.90 g, 5.89 mmol) in THF (100 mL), potassium acetate (0.6 g, 6.12 mmol), and Pd/C (10%) (0.5 g) was hydrogenated for 19 h. The solvents were removed in vacuo, and the product was purified by chromatography eluting with 30–50% ethyl acetate in hexanes. Unreacted starting material was recovered (0.700 g, 18%) and then the title compound eluted (2.43 g, 66%): ^1H NMR (CDCl_3) 0.87 (t, $J = 7.26$ Hz, 3H), 1.32 (m, 2H), 1.70 (m, 2H), 2.57 (t, $J = 7.27$ Hz, 2H), 5.49 (s, 2H), 6.80–7.50 (m, 22H), 7.79 (s, 1H), 7.95 (m, 1H), 9.67 (s, 1H). Anal. ($\text{C}_{41}\text{H}_{56}\text{N}_6\text{O}$) C, H, N.

(*Z*)-1-Butyl-5-[[2-butyl-3-[[2'-(1*H*-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methylene]-3*H*-imidazol-4-yl]methylene]-2,4-imidazolidinedione (20**)**. To a solution of diethyl (3-butyl-2,5-dioxo-4-imidazolyl)phosphonate (**8**) (0.770 g, 2.73 mmol) in methanol (20 mL) was added sodium hydride (0.16 g, 5.33 mmol), and the mixture was stirred for 5 min. A solution of 2-butyl-1-[[2'-(1*H*-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-5-formylimidazole (**19**) (1.15 g, 1.83 mmol) in methylene chloride (100 mL) was then added. The mixture was stirred for 16 h. The solvents were removed in vacuo and ethyl acetate (100 mL) and methylene chloride (10 mL) added. The solution was washed with saturated NH_4Cl and brine. After drying over Na_2SO_4 , the *Z* isomer (0.815 g, 58%) and *E* isomer (0.550 g, 39%) were purified by chromatography eluting with 1% methanol in chloroform. The *Z* isomer (0.800 g, 1.04 mmol) was added to methanol (100 mL) and 5% aqueous citric acid (5 mL). This white heterogenous mixture was refluxed for 3 h in which time a solution formed. The solution was cooled to room temperature and diluted with water (10 mL) and hexanes (100 mL). The aqueous methanol fraction was evaporated in vacuo and then dissolved in a mixture of ethyl acetate and water. After drying the organic fraction over Na_2SO_4 , the product was precipitated by the addition of ether to a methylene chloride solution of the product (0.440 g, 46% overall): ^1H NMR (DMSO) 0.61 (m, 3H), 0.85 (m, 7H), 1.18 (m, 2H), 1.59 (m, 2H), 2.76 (t, $J = 7.94$ Hz, 2H), 3.60 (s, 2H), 5.35 (s, 2H), 6.26 (s, 1H), 6.99 (d, $J = 7.73$ Hz, 2H), 7.07 (d, $J = 8.07$ Hz, 2H), 7.35 (s, 1H), 7.47 (d, $J = 7.94$ Hz, 2H), 7.6–7.70 (m, 3H), 11.55 (s, 1H); HPLC eluant 55% 0.1% TFA in H_2O :45% MeCN, t_R 2.98 (99%); MS (CI) 525 (M + 1).

1-Butyl-5-[[2-butyl-3-[[2'-(1*H*-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-3*H*-imidazol-4-yl]methyl]-2,4-imidazolidinedione (22**)**. To 1-butyl-5-[[2-butyl-3-[[2'-*N*-(triphenylmethyl)tetrazol-5-yl][1,1'-biphenyl]-4-yl]methyl]-3*H*-imidazol-4-yl]methyl]-2,4-imidazolidinedione (0.080 g, 0.10 mmol) were added methanol (30 mL) and glacial acetic acid (0.06 mL). This white heterogenous mixture was refluxed for 5 h in which time a solution formed. The solution was cooled to room temperature and diluted with water (10 mL) and hexanes (50 mL). The aqueous methanol fraction was evaporated in vacuo and then dissolved in a minimal amount of methanol. Ether was added to precipitate a white solid: ^1H NMR (DMSO) 0.83 (m, 6H), 1.17–1.52 (m, 8H), 2.51 (m, 1H), 2.92 (m, 4H), 3.40 (m, 1H), 4.34 (s, 1H), 5.21 (s, 2H), 6.68 (s, 1H), 6.89 (d, $J = 7.80$ Hz, 2H), 7.08 (d, $J = 7.80$ Hz, 2H), 7.47–7.63 (m, 4H), 10.90

(s, 1H) (NH absent); HPLC eluant 67% 0.1% TFA in H_2O :33% MeCN, t_R 5.4 (98%). Anal. ($\text{C}_{25}\text{H}_{34}\text{N}_8\text{O}_2 \cdot \text{H}_2\text{O}$) C, H, N: calcd, 20.57; found, 20.07.

Methyl (*E*)- and (*Z*)-4'-[[2-butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1*H*-imidazol-1-yl]methyl]-[1,1'-biphenyl]-2-carboxylate (24** and **25**)**. To a solution of diethyl (3-butyl-2,5-dioxo-4-imidazolyl)phosphonate (**8**) (0.540 g, 1.91 mmol) in methanol (15 mL) was added NaH (80% in oil, 0.115 g, 3.83 mmol). The solution was stirred for 5 min, and then 2-butyl-1-[(2'-carbomethoxy[1,1'-biphenyl]-4-yl)methyl]-1*H*-imidazole-5-carboxaldehyde (**23**) (0.515 g, 1.37 mmol) was added. The yellow solution was stirred for 16 h at room temperature and then diluted with ethyl acetate and saturated ammonium chloride. The organic fraction was washed with brine and then dried over Na_2SO_4 . Chromatography separated the *Z* (0.304 g, 43%) and *E* (0.280 g, 40%) isomers, eluant 0–2% methanol in chloroform.

Methyl (*Z*)-4'-[[2-butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1*H*-imidazol-1-yl]methyl]-[1,1'-biphenyl]-2-carboxylate (24**)**: ^1H NMR (CDCl_3) 0.71 (t, $J = 6.50$ Hz, 3H), 0.89 (t, $J = 7.17$ Hz, 3H), 1.18 (m, 4H), 1.34 (m, 2H), 1.73 (m, 2H), 2.71 (t, $J = 7.60$ Hz, 2H), 3.61 (s, 3H), 3.77 (m, 2H), 5.17 (s, 2H), 6.38 (s, 1H), 6.99 (d, $J = 8.12$ Hz, 2H), 7.20–7.52 (m, 5H), 7.30 (s, 1H), 7.80 (d, $J = 7.60$ Hz, 1H), 10.50 (brs, 1H). Anal. ($\text{C}_{30}\text{H}_{34}\text{N}_4\text{O}_4$) H, N; C: calcd, 70.02; found, 69.58.

Methyl (*E*)-4'-[[2-butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1*H*-imidazol-1-yl]methyl]-[1,1'-biphenyl]-2-carboxylate (25**)**: ^1H NMR (CDCl_3) 0.81 (t, $J = 6.93$ Hz, 3H), 0.92 (t, $J = 7.17$ Hz, 3H), 1.20–1.45 (m, 6H), 1.76 (m, 2H), 2.79 (t, $J = 7.27$ Hz, 2H), 3.43 (m, 2H), 3.66 (m, 3H), 5.23 (s, 2H), 5.78 (s, 1H), 7.02 (d, $J = 8.11$ Hz, 2H), 7.25–7.52 (m, 5H), 7.84 (d, $J = 7.60$ Hz, 1H), 8.35 (brs, 1H), 8.45 (s, 1H). Anal. ($\text{C}_{30}\text{H}_{34}\text{N}_4\text{O}_4$) C, H, N.

(*Z*)-*N*-[4-[[2-Butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1*H*-imidazol-1-yl]methyl]phenyl]-1,1,1-trifluoromethanesulfonamide (28**)**. To a solution of diethyl (3-butyl-2,5-dioxo-4-imidazolyl)phosphonate (**8**) (1.48 g, 5.06 mmol) in ethanol (10 mL) was cautiously added sodium hydride batchwise (1.0 g, 25 mmol). The cloudy solution was stirred for 30 min, and then 2-butyl-1-[[4-(1,1,1-trifluoromethanesulfonamido)phenyl]methyl]-5-formylimidazole (**27**) (0.68 g, 1.75 mmol) in ethanol (10 mL) was added over 30 min. This solution was heated at reflux for 2 h. The reaction mixture was cooled to room temperature and treated with glacial acetic acid (2 mL). Evaporation in vacuo afforded a thick gel. Chromatography eluting with 0–4% methanol in chloroform afforded the required *Z* isomer which was triturated with ether/hexanes to afford the title compound as an amorphous solid (0.565 g, 61%). The more polar *E* isomer was then eluted: ^1H NMR (DMSO) 0.62 (t, $J = 6.5$ Hz, 3H), 0.84 (m, 7H), 1.30 (m, 2H), 1.58 (m, 2H), 2.88 (t, $J = 7.4$ Hz, 2H), 3.53 (brs, 2H), 5.31 (s, 2H), 6.26 (s, 1H), 6.95 (d, $J = 8.4$ Hz, 2H), 7.04 (d, $J = 8.4$ Hz, 2H), 7.51 (s, 1H), 11.75 (s, 1H) (NH absent). Anal. ($\text{C}_{23}\text{H}_{28}\text{N}_5\text{O}_4\text{F}_3\text{S}$) C, H, N.

(*E*)-*N*-[4-[[2-Butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1*H*-imidazol-1-yl]methyl]phenyl]-1,1,1-trifluoromethanesulfonamide (29**)**: 0.11 g, 12%; ^1H NMR (DMSO) 0.80 (t, $J = 7.0$ Hz, 3H), 0.86 (t, $J = 7.2$ Hz, 3H), 0.95 (m, 2H), 1.19 (m, 2H), 1.31 (m, 2H), 1.61 (m, 2H), 2.96 (t, $J = 8.0$ Hz, 2H), 3.51 (brs, 2H), 5.48 (s, 2H), 5.98 (s, 1H), 6.91 (d, $J = 8.5$ Hz, 2H), 7.04 (d, $J = 8.5$ Hz, 2H), 8.32 (s, 1H), 11.45 (s, 1H) (NH absent). Anal. ($\text{C}_{23}\text{H}_{28}\text{N}_5\text{O}_4\text{F}_3\text{S} \cdot 0.25\text{H}_2\text{O}$) C, H, N.

(*Z*)- and (*E*)-1-Butyl-5-[[2-butyl-3-[[4-(1*H*-tetrazol-5-yl)phenyl]methyl]-3*H*-imidazol-4-yl]methylene]-2,4-imidazolidinedione (33** and **34**)**. To a solution of diethyl (3-butyl-2,5-dioxo-4-imidazolyl)phosphonate (**8**) (0.240 g, 0.86 mmol) in ethanol (10 mL) was added sodium hydride (0.072 g, 1.80 mmol), and the mixture was stirred for 15 min. A solution of 2-butyl-1-[[4-(1-(triphenylmethyl)tetrazol-5-yl)phenyl]methyl]-5-formylimidazole (**30**) (0.400 g, 0.72 mmol) in methylene chloride (5 mL) was then added. The mixture was stirred for 16 h. The solvents were removed in vacuo and the *Z* and *E* isomers purified by chromatography eluting with 20% ethyl acetate in methylene chloride. The *Z* isomer **31** (0.36 g, 72%) and the *E* isomer **32** (0.09 g, 18%) were each separately

dissolved in methanol (5 mL) containing a 10% aqueous solution of citric acid (0.5 mL). The reaction mixture was refluxed overnight. Water (20 mL) was added and the mixture extracted with hexane. The aqueous fraction was then extracted with ethyl acetate, washed with brine, and dried over MgSO_4 . After evaporating in vacuo, the residue was crystallized from a combination of hexane, ethyl acetate, and methanol. This procedure afforded **33** (0.06 g, 17%): $^1\text{H NMR}$ (DMSO) 0.61 (m, 3H), 0.88 (m, 7H), 1.34 (m, 2H), 1.62 (m, 2H), 2.74 (t, $J = 7.3$ Hz, 2H), 3.64 (brs, 2H), 5.41 (s, 2H), 6.28 (s, 1H), 7.18 (s, 1H), 7.23 (d, $J = 8.23$ Hz, 2H), 8.00 (d, $J = 8.23$ Hz, 2H), 11.44 (s, 1H) (NH absent). Anal. ($\text{C}_{23}\text{H}_{28}\text{N}_8\text{O}_2\cdot\text{S}\cdot 0.9\text{H}_2\text{O}$) C, H, N.

34: $^1\text{H NMR}$ (DMSO) 0.61 (m, 3H), 0.86 (t, $J = 7.24$ Hz, 3H), 1.05 (m, 4H), 1.35 (m, 2H), 1.62 (m, 2H), 2.76 (t, $J = 7.3$ Hz, 2H), 3.47 (brs, 2H), 5.49 (s, 2H), 5.96 (s, 1H), 7.09 (d, $J = 8.20$ Hz, 2H), 7.98 (d, $J = 8.20$ Hz, 2H), 8.24 (s, 1H), 11.24 (s, 1H) (NH absent). Anal. ($\text{C}_{23}\text{H}_{28}\text{N}_8\text{O}_2\cdot\text{S}\cdot 1.67\text{H}_2\text{O}$) C, H, N.

Methyl (Z)-1-[4-[[2-Butyl-5-[[1-butyl-3-[(2-methyl-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]phenyl]cyclopentanecarboxylate Monohydrochloride (36). Methyl (Z)-1-[4-[[2-butyl-5-[[1-butyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]phenyl]cyclopentanecarboxylate (1.00 g, 1.97 mmol) and 4-(chloromethyl)-2-methylthiazole monohydrochloride (0.363 g, 1.97 mmol) were dissolved in DMF (10 mL). K_2CO_3 (3.0 g, 21.7 mmol) was added and the mixture stirred for 16 h. The mixture was filtered and evaporated. The residue was added to water, and the gum that formed was dissolved in ether. After washing the ethereal fraction with water and drying over K_2CO_3 , the product was purified by chromatography: eluant 0.5–3% methanol in chloroform. The appropriate fraction was dissolved in ether and treated with ethereal HCl to afford the required product (0.64 g, 48%). A small amount of the corresponding *E* isomer (10%) was also collected: $^1\text{H NMR}$ (DMSO) 0.77 (t, $J = 7.10$ Hz, 3H), 0.90 (t, $J = 7.25$ Hz, 3H), 1.15–1.85 (m, 14H), 2.51 (m, 2H), 2.56 (s, 3H), 2.93 (t, $J = 7.23$ Hz, 2H), 3.54 (brt, $J = 7.0$ Hz, 2H), 3.51 (s, 3H), 4.82 (s, 2H), 5.20 (s, 2H), 6.08 (s, 1H), 7.00 (s, 1H), 7.02 (d, $J = 8.33$ Hz, 2H), 7.29 (d, $J = 8.33$ Hz, 2H), 7.78 (s, 1H). Anal. ($\text{C}_{34}\text{H}_{43}\text{N}_6\text{O}_4\cdot\text{S}\cdot\text{HCl}\cdot\text{H}_2\text{O}$) C, H, N.

(Z)-1-[4-[[2-Butyl-5-[[1-butyl-3-[(2-methyl-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]phenyl]cyclopentanecarboxylic Acid (37). Methyl (Z)-1-[4-[[2-butyl-5-[[1-butyl-2,5-dioxo-3-[(2-methyl-4-thiazolyl)methyl]-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]phenyl]cyclopentanecarboxylate monohydrochloride (**36**) (0.340 g, 0.506 mmol) and 2 N HCl (15 mL) were heated at reflux for 2 h. After evaporation in vacuo, the residue was dissolved in methanol and treated with saturated aqueous NaOAc to precipitate a gum. This gum crystallized from ethyl acetate and was then recrystallized from methanol/ether to afford the title compound (0.153 g, 50%): $^1\text{H NMR}$ (DMSO) 0.79 (t, $J = 7.34$ Hz, 3H), 0.88 (t, $J = 7.39$ Hz, 3H), 1.15–1.77 (m, 14H), 2.50 (m, 2H), 2.55 (s, 3H), 2.56 (t, $J = 6.15$ Hz, 2H), 3.51 (brt, $J = 6.83$ Hz, 2H), 4.96 (s, 2H), 5.01 (s, 2H), 6.27 (s, 1H), 6.73 (s, 1H), 6.80 (d, $J = 8.29$ Hz, 2H), 7.12 (s, 1H), 7.24 (d, $J = 8.29$ Hz, 2H) (CO_2H absent). Anal. ($\text{C}_{33}\text{H}_{41}\text{N}_6\text{O}_4\cdot\text{S}$) C, H, N.

Methyl (Z)-4-[[2-Butyl-5-[[1-butyl-2,5-dioxo-3-(phenylmethyl)-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate Monohydrochloride (42). To a solution of methyl (Z)-4-[[2-butyl-5-[[1-butyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate (**14**) (0.509 g, 1.16 mmol) in DMF (6 mL) was added K_2CO_3 (0.800 g, 5.80 mmol). After stirring for 5 min, benzyl bromide (0.150 mL, 1.26 mmol) was added. The mixture was stirred for 16 h, diluted with ethyl acetate, washed with water and brine, and then dried over MgSO_4 . A viscous oil (0.525 g, 86%) was obtained upon chromatography on silica gel (eluant 1% methanol in chloroform) and evaporation of the solvents in vacuo. This yellow oil was dissolved in ether and treated with ethereal HCl to afford a foamy solid upon evaporation in vacuo. This foam was crystallized from 2-propanol and ether: $^1\text{H NMR}$ (CDCl_3) 0.84 (t, $J = 7.2$ Hz, 3H), 0.96 (t, $J = 7.2$ Hz, 3H), 1.35 (m, 4H), 1.65 (m, 4H), 2.76 (t, $J = 7.7$ Hz, 2H), 3.68 (t, $J = 7.2$ Hz, 2H), 3.91 (s, 3H), 4.32 (s, 2H), 4.90 (s, 2H), 5.84

(s, 1H), 6.70 (m, 2H), 6.82 (d, $J = 7.92$ Hz, 2H), 7.28 (m, 3H), 7.30 (s, 1H), 7.98 (d, $J = 7.92$, 2H). Anal. ($\text{C}_{31}\text{H}_{38}\text{N}_4\text{O}_4\cdot\text{HCl}\cdot 1.25\text{H}_2\text{O}$) C, H, N.

Methyl (Z)-4-[[2-Butyl-5-[[1,3-dibutyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate Monohydrochloride (48). To a solution of methyl (Z)-4-[[2-butyl-5-[[1-butyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate (**14**) (0.140 g, 0.32 mmol) in DMF (2 mL) was added K_2CO_3 (0.200 g, 1.44 mmol). After stirring for 5 min, 1-iodobutane (0.050 mL, 0.44 mmol) was added. The mixture was stirred for 16 h, diluted with ethyl acetate, washed with water and brine, and then dried over MgSO_4 . A viscous oil was obtained by evaporation of the solvents in vacuo. This yellow oil was dissolved in ether and treated with ethereal HCl to precipitate a gummy solid. This gum was crystallized from 2-propanol/ether to afford the title compound (0.130 g, 82%): $^1\text{H NMR}$ (DMSO) 0.62 (t, $J = 6.8$ Hz, 3H), 0.86 (m, 8H), 1.23–1.75 (m, 10H), 2.50 (brs, 2H), 3.11 (t, $J = 7$ Hz, 2H), 3.49 (m, 2H), 3.83 (s, 3H), 5.66 (s, 2H), 6.26 (s, 1H), 7.34 (d, $J = 8.21$ Hz, 2H), 7.92 (d, $J = 8.21$ Hz, 2H), 7.95 (s, 1H). Anal. ($\text{C}_{28}\text{H}_{38}\text{N}_4\text{O}_4\cdot\text{HCl}$) C, H, N.

Methyl (Z)-4-[[2-Butyl-5-[[1-butyl-3-[(2-methyl-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate Dihydrochloride (57). To a solution of methyl (Z)-4-[[2-butyl-5-[[1-butyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate hydrochloride (prepared by treating **14** with ethereal HCl and crystallization from 2-propanol/ether) (3.00 g, 6.3 mmol) in DMF (20 mL) was added K_2CO_3 (10.0 g, 72 mmol). After stirring for 5 min, 2-methyl-4-(chloromethyl)thiazole hydrochloride (1.33 g, 7.2 mmol) was added. An additional amount of DMF (10 mL) was added after a period of 4 h to aid stirring. The mixture was stirred for 3 days and then filtered. The filtrate was concentrated in vacuo and added to water (20 mL). The gum was separated, dissolved in ether (50 mL), washed with water, and dried over K_2CO_3 and charcoal. A viscous oil was obtained by evaporation of the solvents in vacuo which was dissolved in ether, treated with ethereal HCl, and then evaporated in vacuo to afford a solid foam. Recrystallization from 2-propanol/ether afforded the title compound (2.90 g, 73%): $^1\text{H NMR}$ (DMSO) 0.80 (t, $J = 7.34$ Hz, 3H), 0.88 (t, $J = 7.37$ Hz, 3H), 1.25 (m, 4H), 1.54 (m, 4H), 2.56 (s, 3H), 2.96 (brt, $J = 7.65$ Hz, 2H), 3.52 (brt, $J = 6.84$ Hz, 2H), 3.86 (s, 3H), 4.84 (s, 2H), 5.36 (s, 2H), 5.98 (s, 1H), 7.05 (s, 1H), 7.20 (d, $J = 8.24$ Hz, 2H), 7.84 (s, 1H), 7.92 (d, $J = 8.26$ Hz, 2H). Anal. ($\text{C}_{29}\text{H}_{35}\text{N}_5\text{O}_4\cdot\text{S}\cdot 2\text{HCl}\cdot 0.5\text{H}_2\text{O}$) C, H, N.

Methyl (Z)-4-[[2-Butyl-5-[[1-butyl-3-[(2-amino-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate Dihydrochloride (63). To a solution of methyl (Z)-4-[[2-butyl-5-[[1-butyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate hydrochloride (**14**) (3.70 g, 7.8 mmol) in DMF (30 mL) was added K_2CO_3 (15.0 g, 0.1 mol). After stirring for 5 min, 2-amino-4-(chloromethyl)thiazole hydrochloride (1.85 g, 10.0 mmol) was added. The mixture was stirred for 72 h and then diluted with ethyl acetate and filtered. The filtrate was concentrated in vacuo and added to water (50 mL). The brown solid that formed was dissolved in methylene chloride (100 mL), washed with water, and dried over K_2CO_3 and charcoal. A viscous oil (4.5 g) was obtained by evaporation of the solvents in vacuo. This oil was dissolved in ether and treated with ethereal HCl to precipitate a gum which was collected by decantation. This gum was recrystallized from 2-propanol/ether to afford the title compound as a white solid (3.20 g, 62%): $^1\text{H NMR}$ (DMSO) 0.82 (t, $J = 7.20$ Hz, 3H), 0.88 (t, $J = 7.30$ Hz, 3H), 1.28 (m, 4H), 1.55 (m, 4H), 3.00 (brt, $J = 7.4$ Hz, 2H), 3.50 (brt, $J = 6.6$ Hz, 2H), 3.86 (s, 3H), 4.67 (s, 2H), 5.47 (s, 2H), 6.00 (s, 1H), 6.38 (s, 1H), 7.27 (d, $J = 8.02$ Hz, 2H), 7.77 (s, 1H), 7.96 (d, $J = 8.02$ Hz, 2H) (NH_2 absent). Anal. ($\text{C}_{28}\text{H}_{34}\text{N}_6\text{O}_4\cdot\text{S}\cdot 2\text{HCl}\cdot 2\text{H}_2\text{O}$) C, H, N.

(Z)-4-[[2-Butyl-5-[[1,3-dibutyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoic Acid (89). To a solution of 1,1-dimethylethyl (Z)-4-[[2-butyl-5-[[1-butyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate (**17**) (2.04 g, 4.25 mmol) in DMF (20 mL) were added K_2CO_3 (2.93 g, 5 equiv) and 1-iodobutane (1.70 g, 2 equiv). The mixture was stirred for 16 h and then evapo-

rated in vacuo. The ethyl acetate extract was washed with brine, dried over $MgSO_4$, and then evaporated in vacuo. The yellow oil that resulted (2.33 g) was used directly in the subsequent reaction. 1,1-Dimethylethyl (Z)-4-[[2-butyl-5-[(1,3-dibutyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoate (18) (2.33 g, assume 4.25 mmol) was dissolved in methylene chloride (20 mL) and trifluoroacetic acid (15 mL). After stirring for 4 h, the reaction mixture was evaporated in vacuo. The yellow residue was dissolved in dichloromethane (50 mL) and washed with aqueous NaOAc (10%, 60 mL) and brine. A yellow foam was obtained upon evaporation of the organic extract after drying over $MgSO_4$. This foam solidified upon trituration with ether. The required Z isomer (1.085 g, 52%) was separated from the E isomer 90 (12%) by chromatography, eluting with 2–3% methanol in dichloromethane: 1H NMR (DMSO) 0.63 (m, 3H), 0.86 (m, 8H), 1.22 (m, 2H), 1.30 (m, 2H), 1.50 (m, 2H), 1.61 (m, 2H), 2.69 (t, $J = 7.60$ Hz, 2H), 3.33 (m, 2H), 3.44 (t, $J = 7.2$ Hz, 2H), 3.70 (m, 2H), 5.40 (s, 2H), 6.37 (s, 1H), 7.10 (d, $J = 8.40$ Hz, 2H), 7.15 (s, 1H), 7.88 (d, $J = 8.40$ Hz, 2H), 13.00 (brs, 1H). Anal. ($C_{27}H_{36}N_4O_4$) C, H, N.

(Z)-4-[[2-Butyl-5-[[1-butyl-3-[(2-methyl-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoic Acid (95). A mixture of methyl (Z)-4-[[2-butyl-5-[[1-butyl-3-[(2-methyl-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate (57) (0.428 g, 0.65 mmol) and 2 N HCl (12 mL) was refluxed for 1 h in which time all the solid dissolved. The solvents were removed in vacuo, and the mixture was evaporated from ethanol (5 mL) three times. The solid that resulted was treated with 5% aqueous sodium acetate (10 mL) and water (10 mL) to precipitate a gummy solid which was dissolved in methylene chloride (30 mL). The organic layer was dried over Na_2SO_4 and then evaporated in vacuo. The residue was recrystallized from hot ethyl acetate/hexane to afford the title compound (0.279 g, 80%): 1H NMR (DMSO) 0.80 (t, $J = 7.27$ Hz, 3H), 0.88 (t, $J = 7.29$ Hz, 3H), 1.27 (m, 4H), 1.53 (m, 4H), 2.54 (s, 3H), 2.55 (t, $J = 7.35$ Hz, 2H), 3.50 (t, $J = 6.75$ Hz, 2H), 4.95 (s, 2H), 5.15 (s, 2H), 6.22 (s, 1H), 6.73 (s, 1H), 6.94 (d, $J = 8.13$ Hz, 2H), 7.15 (s, 1H), 7.84 (d, $J = 8.11$ Hz, 2H), 13.00 (brs, 1H). Anal. ($C_{28}H_{33}N_5O_4S$) C, H, N.

Isomerization of (Z)-4-[[2-Butyl-5-[[1-butyl-3-[(2-methyl-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoic Acid (95) to (E)-4-[[2-Butyl-5-[[1-butyl-3-[(2-methyl-4-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoic Acid (96). Methyl (Z)-4-[[2-butyl-5-[[1-butyl-3-[(4-methyl-2-thiazolyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoic acid (95) (0.25 g, 0.46 mmol) was dissolved in concentrated HCl (15 mL) and the mixture refluxed for 4 h. The mixture was concentrated in vacuo and aqueous sodium acetate added. The precipitate was collected and washed with water. This solid was dissolved in methanol (2 mL), treated with charcoal, and then filtered and concentrated in vacuo. The residue was recrystallized from hot ethyl acetate to afford the title compound (0.04 g, 16%): 1H NMR (DMSO) 0.80 (t, $J = 7.27$ Hz, 3H), 0.89 (t, $J = 7.29$ Hz, 3H), 1.27 (m, 4H), 1.54 (m, 4H), 2.47 (s, 3H), 2.62 (t, $J = 7.31$ Hz, 2H), 3.52 (t, $J = 6.85$ Hz, 2H), 4.85 (s, 2H), 5.42 (s, 2H), 6.17 (s, 1H), 6.92 (d, $J = 8.20$ Hz, 2H), 6.95 (s, 1H), 7.82 (d, $J = 8.21$ Hz, 2H), 8.29 (s, 1H) (CO_2H absent). Anal. ($C_{28}H_{33}N_5O_4S \cdot 0.25H_2O$) C, H, N.

Hydrogenation of 98 To Afford 4-[[2-Butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinyl)methyl]-1H-imidazol-1-yl]methyl]benzoic Acid. To a solution of (E)-4-[[2-butyl-5-[(3-butyl-2,5-dioxo-4-imidazolidinylidene)methyl]-1H-imidazol-1-yl]methyl]benzoic acid (98) (0.84 g, 2.00 mmol) in 1 N NaOH (2.04 mL, 2.04 mmol) and water (500 mL) was added PtO_2 (0.200 g), and then a hydrogen balloon was attached. The mixture was stirred for 3 h, 1 N HCl (2.04 mL, 2.04 mmol) was added, and then the mixture was concentrated at reduced pressure. Ethanol (50 mL) was added and the mixture filtered. The filtrate was evaporated in vacuo, and the residue (0.700 g, 90% pure) was purified by crystallization and chromatography eluting with 5–15% methanol in chloroform: 1H NMR

(DMSO) 0.78 (t, $J = 7.39$ Hz, 3H), 0.81 (t, $J = 7.20$ Hz, 3H), 1.25–1.48 (m, 8H), 2.50 (brt, 2H), 2.86 (m, 2H), 3.40 (m, 2H), 4.31 (brt, 1H), 5.25 (s, 2H), 6.58 (s, 1H), 6.98 (d, $J = 8.21$ Hz, 2H), 7.89 (d, $J = 8.21$ Hz, 2H) (CO_2H and NH absent). Anal. ($C_{23}H_{30}N_4O_4S \cdot H_2O$) C, H, N.

(Z)-4-[[2-Butyl-5-[[1-butyl-3-[(5-methyl-3-thienyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoic Acid (109). To a solution of 1,1-dimethylethyl (Z)-4-[[2-butyl-5-[[1-butyl-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate (17) (1.00 g, 2.08 mmol) in THF (15 mL) was added NaH (0.11 g, 2.75 mmol). This mixture was stirred for 15 min to afford a yellow solution. 4-(Bromomethyl)-2-methylthiophene (0.65 g, 3.39 mmol) in THF (5 mL) was added dropwise over 30 min. After 3 h potassium iodide (0.5 g, 3 mmol) was added, and then the mixture was stirred for 16 h. The mixture was filtered, and the solvents were removed in vacuo. The product was isolated, by chromatography eluting with 0–8% acetone in dichloromethane (0.78 g, 64%). The 1,1-dimethylethyl (Z)-4-[[2-butyl-5-[[1-butyl-3-[(5-methyl-3-thienyl)methyl]-2,5-dioxo-4-imidazolidinylidene]methyl]-1H-imidazol-1-yl]methyl]benzoate (0.78 g, 1.32 mmol) was dissolved in methylene chloride (14 mL) and trifluoroacetic acid (10 mL). After stirring for 2 h, the reaction mixture was evaporated in vacuo. After chromatography, eluant 0–6% methanol in dichloromethane, the foam was dissolved in ethyl acetate, washed with 10% aqueous NaOAc and brine, and then dried over $MgSO_4$. Crystallization from ether afforded 109 (0.319 g, 45%): 1H NMR (DMSO) 0.81 (t, $J = 7.31$ Hz, 3H), 0.87 (t, $J = 7.18$ Hz, 3H), 1.26 (m, 4H), 1.53 (m, 4H), 2.34 (s, 3H), 2.58 (t, $J = 7.33$ Hz, 2H), 3.49 (t, $J = 6.75$ Hz, 2H), 4.84 (s, 2H), 5.13 (s, 2H), 6.21 (s, 1H), 6.31 (s, 1H), 6.44 (s, 1H), 6.96 (d, $J = 8.11$ Hz, 2H), 7.16 (s, 1H), 7.84 (d, $J = 8.02$ Hz, 2H) (CO_2H absent). Anal. ($C_{29}H_{34}N_4O_4S$) C, H, N.

Biological Assays. Angiotensin Receptor Binding Assay (AT₁). Methods for the preparation of rat liver membranes have been reported previously in detail.¹⁷ Solutions (10 mM) of the test compounds were diluted to the appropriate concentration and added to 1 mg of rat liver membrane homogenate, 1 mg of wheat germ agglutinin (WGA)-coated scintillation proximity assay (SPA) beads, and 50 pM [¹²⁵I]angiotensin II. After an incubation of 180 min at 25 °C, the light emitted by the stimulated SPA bead was measured. Nonspecific binding was defined as radioactivity retained on the beads in the presence of 10 μ M saralasin. The inhibitory concentration (IC_{50}) of an inhibitor that gave 50% displacement of the specifically bound [¹²⁵I]angiotensin II was then calculated. Concentrations of the test compound were tailored to adequately span the concentration inhibiting binding by 50%. Typically IC_{50} 's were determined via nonlinear regression curve fitting employing seven points and in duplicate. Individual points were determined with an average error of $\pm 5\%$. For comparison, using this protocol, reference values for antagonists were DuP 753,²¹ 9.2 nM; L-158809,³⁰ 0.8 nM; and SKF 108566,³⁶ 2.8 nM.

For assessing AT₂ receptor binding, a similar protocol was employed except rabbit uterine membranes were used.

Angiotensin II-Induced Contraction in Isolated Vascular Tissue (RAR). Details for the rabbit thoracic aorta ring preparation have been reported previously.¹⁷ Briefly the aortic rings were connected to a pressor transducer, and the tissue preparation was allowed to equilibrate at a mechanically induced tension of 4 g for 1.5 h before use. A control–response curve to exogenous Ang II was recorded. The test compound in DMSO was added to the bath (Krebs-bicarbonate solution gassed continuously with 5% CO_2 in oxygen) at several concentrations that were expected to span the IC_{50} , as predicted from receptor binding studies. The tissue preparation was allowed to equilibrate for 10 min. For each concentration of antagonist, the pressor response to 10 nM Ang II was recorded. The preparation was then allowed to equilibrate for 1 h between Ang II challenges. The IC_{50} of the test compounds represents the mean concentration of antagonist that inhibited by 50% the contractile response to Ang II for two rings within the same bath.

Evaluation of the Test Compound in Conscious Renal Hypertensive Rats (RHR). The protocol was similar to that

previously described.¹⁷ Briefly young male Sprague–Dawley rats underwent bilateral renal artery clipping to afford rats with mean blood pressure of greater than 150 mmHg. The test compounds were administered in 0.5% methocel in water by oral gavage in a volume of 2 mL/kg, and the blood pressure was measured by a cannula inserted in the abdominal aorta. Data reported represent the maximum percent reduction in blood pressure relative to base line and were averaged over three to four animals. Standard error mean was typically <5% in control and <8% in drug-treated animals. Heart rate did not vary significantly from base line.

Evaluation of the Test Compound in Sodium-Deplete Normotensive Monkeys. The protocol was similar to that previously described.⁴⁹ Briefly animals were rendered renin dependent by a combination of a sodium-restricted diet and pretreatment with the diuretic furosemide. In this model saralasin (20 µg/kg/min) infusion reduced mean arterial blood pressure by 23 ± 2 mmHg. The test compounds were administered in 0.5% methocel in water by oral gavage in a volume of 2 mL/kg, and the blood pressure was measured via an arterial vascular access port. Data reported represent the maximum percent reduction in blood pressure relative to base line and were averaged over two animals. Heart rate did not vary significantly from base line.

Supporting Information Available: Tables of HPLC retention times, log *P* measured by octanol/water partition method, analytical data, blood pressure effect of compounds on renal hypertensive rats, and inhibition of Ang II-induced pressor responses in normotensive rats (9 pages). Ordering information is given on any current masthead page.

References

- Klutchko, S.; Edmunds, J. J.; Hodges, J. C.; Hamby, J. M.; Sircar, I.; Bunker, A. M.; Connolly, C. J. C.; Winters, R. T.; Quin, J., III; Lu, G. H.; Panek, R. L.; Doherty, A. M. Derivatives of 1-Benzylimidazole-5-Methylidenehydantoins as Potent, Nonpeptide Angiotensin II (AT₁) Receptor Antagonists. Presented at the 206th National Meeting of the ACS, Chicago, IL, August 1993; Abstr. Medi 71.
- Sircar, I.; Keiser, J. A.; Panek, R. L.; Major, T. C.; Taylor, D. G.; Klutchko, S.; Edmunds, J. J.; Hamby, J. M.; Winters, R. T.; Bunker, A. M.; Quin, J., III; Connolly, C. J. C.; Doherty, A. M. Substituted 1-Benzylimidazole-5-Methylidene Hydantoins As Orally Active Angiotensin II (AT₁) Receptor Antagonists. Presented at the 206th National Meeting of the ACS, Chicago, IL, August 1993; Abstr. Medi 73.
- Ferrario, C. M. The Renin-Angiotensin System: Importance in Physiology and Pathology. *J. Cardiovasc. Pharmacol.* **1990**, *15* (Suppl. 3), S1–S5.
- Greenlee, W. J.; Siegl, P. K. S. Angiotensin/Renin Modulators. *Annu. Rep. Med. Chem.* **1992**, *27*, 59–68.
- Valloton, M. B. The renin-angiotensin system. *Trends Pharmacol. Sci.* **1987**, *8*, 69–74.
- Martinez Bru, C. Dipeptidyl carboxypeptidases (angiotensin converting enzyme). *Rev. Soc. Esp. Quim. Clin.* **1992**, *11* (1), 8–16.
- Brunner, H. R.; Nussberger, J.; Waeber, B. Control of vascular tone by renin and angiotensin in cardiovascular disease. *Eur. Heart J.* **1993**, *14* (Suppl. I), 149–53.
- Stella, A.; Zanchetti, A. Control of renal renin release. *Kidney Int., Suppl.* **1987**, *31* (20), S89–S94.
- Ocain, T. D.; Abou-Gharbia, M. Renin-angiotensin system (RAS) dependent antihypertensive agents: 1. Renin inhibitors. *Drugs Future* **1991**, *16* (1), 37–51.
- Wood, J. M.; Cumin, F.; Maibaum, J. Pharmacology of renin inhibitors and their application to the treatment of hypertension. *Pharmacol. Ther.* **1994**, *61* (3), 325–44.
- Ondetti, M. A.; Cushman, D. W. Inhibition of the Renin-Angiotensin System. A New Approach to the Therapy of Hypertension. *J. Med. Chem.* **1981**, *24*, 355–61.
- Patchett, A. A.; Harris, E.; Tristram, E. W.; Wyvratt, M. J.; Wu, M. T.; Taub, D.; Peterson, E. R.; Ikeler, T. J.; ten Broeke, J.; Payne, L. G.; Ondeyka, D. L.; Thorsett, E. D.; Greenlee, W. J.; Lohr, N. S.; Hoffsommer, R. D.; Joshua, H.; Ruyle, W. V.; Rothrock, J. W.; Aster, S. D.; Maycock, A. L.; Robinson, F. M.; Hirschmann, R.; Sweet, C. S.; Ulm, E. H.; Gross, D. M.; Vassil, T. C.; Stone, C. A. A new class of angiotensin-converting enzyme inhibitors. *Nature* **1980**, *288*, 280–3.
- Dzau, V. J.; Pratt, R.; Gibbons, G. H. Angiotensin as local modulating factor in ventricular dysfunction and failure due to coronary artery disease. *Drugs* **1994**, *47* (Suppl. 4), 1–13.
- Coulter, D. M.; Edwards I. R. Cough associated with captopril and enalapril. *Brit. Med. J.* **1987**, *294*, 1521–3.
- McEwan, J. R.; Fuller, R. W. Angiotensin Converting Enzyme Inhibitors and Cough. *J. Cardiovasc. Pharmacol.* **1989**, *13* (Suppl. 3), S67–S69.
- Chin, H. L.; Buchan, D. A. Severe Angioedema after Long-Term Use of an Angiotensin-Converting Enzyme Inhibitor. *Ann. Intern. Med.* **1990**, *112*, 312–3.
- Hodges, J. C.; Hamby, J. M.; Blankley, C. J. Angiotensin II receptor binding inhibitors. *Drugs Future* **1992**, *17* (7), 575–93.
- Moore, A. F.; Fulton, R. W. Angiotensin II Antagonists-Saralasin. *Drug Dev. Res.* **1984**, *4*, 331–49.
- Pals, D. T.; Denning, G. S., Jr.; Keenan, R. E. Historical development of saralasin. *Kidney Int.* **1979**, *15* (9), 7–10.
- Duncia, J. V.; Chiu, A. T.; Carini, D. J.; Gregory, G. B.; Johnson, A. L.; Price, W. A.; Wells, G. J.; Wong, P. C.; Calabrese, J. C.; Timmermans, P. B. M. W. M. The Discovery of Potent Nonpeptide Angiotensin II Receptor Antagonists: A New Class of Potent Antihypertensives. *J. Med. Chem.* **1990**, *33*, 1312–29.
- Furukawa, Y.; Kishimoto, S.; Nishikawa, K. Hypotensive Imidazole Derivatives. U.S. Patent 4,340,598, 1982.
- Furukawa, Y.; Kishimoto, S.; Nishikawa, K. Hypotensive Imidazole-5-Acetic Acid Derivatives. U.S. Patent 4,355,040, 1982.
- Furukawa, Y.; Kishimoto, S.; Nishikawa, K. 4-Chloro-2-Phenylimidazole-5-Acetic Acid Derivatives. Eur. Pat. Appl. 103,647, 1984.
- Middlemiss, D.; Watson, S. P.; Ross, B. C.; Dowle, M. D.; Scopes, D. I. C.; Montana, J. G.; Hirst, G. C.; Panchal, T. A.; Paton, J. M. S.; Hubbard, T.; Stuart, G.; Drew, G. M.; Hilditch, A.; Travers, A.; Robertson, M. J.; Hunt, A. A. E.; Palmer, E.; Manchee, G. R. Benzofuran Based Angiotensin II Antagonists Related to GR117289: Part II; Amino Acid Amides. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 2043–6.
- Ashton, W. T.; Cantone, C. L.; Chang, L. L.; Hutchins, S. M.; Strelitz, R. A.; MacCoss, M.; Chang, R. S. L.; Lotti, V. J.; Faust, K. A.; Chen, T.-B.; Bunting, P.; Schorn, T. W.; Kivlighn, S. D.; Siegl, P. K. S. Nonpeptide Angiotensin II Antagonists Derived from 4*H*-1,2,4-Triazoles and 3*H*-Imidazo[1,2-*b*]1,2,4-triazoles. *J. Med. Chem.* **1993**, *36*, 591–609.
- Timmermans, P. B. M. W. M.; Carini, D. J.; Chiu, A. T.; Duncia, J. V.; Price, W. A., Jr.; Wells, G. J.; Wong, P. C.; Johnson, A. L.; Wexler, R. R. The discovery of a new class of highly specific nonpeptide angiotensin II receptor antagonists. *Am. J. Hypertens.* **1991**, *4*, 275S–281S.
- Sircar, I.; Hodges, J. C.; Quin, J., III; Bunker, A.; Winters, R. T.; Edmunds, J. J.; Kostlan, C. R.; Connolly, C.; Kesten, S. J.; Hamby, J. M.; Topliss, J. G.; Keiser, J. A.; Panek, R. L. Nonpeptide Angiotensin II Receptor Antagonists. 2. Design, Synthesis, and Structure-Activity Relationships of 2-Alkyl-4-(1*H*-pyrrol-1-yl)-1*H*-imidazole Derivatives: Profile of 2-Propyl-1-[[2-(1*H*-tetrazol-5-yl)-[1,1'-biphenyl]-4-yl]-methyl]-4-[2-(trifluoroacetyl)-1*H*-pyrrol-1-yl]-1*H*-imidazole-5-carboxylic Acid (CI-996). *J. Med. Chem.* **1993**, *36*, 2253–65.
- Blankley, C. J.; Hodges, J. C.; Klutchko, S. R.; Himmelsbach, R. J.; Chucholowski, A.; Connolly, C. J.; Neergaard, S. J.; Van Nieuwenhze, M. S.; Sebastian, A.; Quin, J., III; Essenburg, A. D.; Cohen, D. M. Synthesis and Structure-Activity Relationships of a Novel Series of Non-Peptide Angiotensin II Receptor Binding Inhibitors Specific for the AT₂ Subtype. *J. Med. Chem.* **1991**, *34*, 3248–60.
- Ellingboe, J. W.; Antane, M.; Nguyen, T. T.; Collini, M. D.; Antane, S.; Bender, R.; Hartupce, D.; White, V.; McCallum, J.; Park, C. H.; Russo, A.; Osler, M. B.; Wojdan, A.; Dinsh, J.; Ho, D. M.; Bagli, J. F. Pyrido[2,3-*d*]pyrimidine Angiotensin II Antagonists. *J. Med. Chem.* **1994**, *37*, 542–50.
- Mantlo, N. B.; Chakravarty, P. K.; Ondeyka, D. L.; Siegl, P. K. S.; Chang, R. S.; Lotti, V. J.; Faust, K. A.; Schorn, T. W.; Chen, T.-B.; Schorn, T. W.; Sweet, C. S.; Emmert, S. E.; Patchett, A. A.; Greenlee, W. J. Potent, orally active imidazo[4,5-*b*]pyridine-based angiotensin II receptor antagonists. *J. Med. Chem.* **1991**, *34*, 2919–22.
- Chakravarty, P. K.; Naylor, E. M.; Chen, A.; Chang, R. S. L.; Chen, T.-B.; Faust, K. A.; Lotti, V. J.; Kivlighn, S. D.; Gable, R. A.; Zingaro, G. J.; Schorn, T. W.; Schaffer, L. W.; Broten, T. P.; Siegl, P. K. S.; Patchett, A. A.; Greenlee, W. J. A Highly Potent, Orally Active Imidazo[4,5-*b*]pyridine Biphenyl Acylsulfonamide (MK-996; L-159,282): A New AT₁-Selective Angiotensin II Receptor Antagonist. *J. Med. Chem.* **1994**, *37*, 4068–72.
- Dhanoa, D. S.; Bagley, S. W.; Chang, R. S. L.; Lotti, V. K. S.; Chakravarty, P. K.; Patchett, A. A.; Greenlee, W. J. (Dipropylphenoxy)phenylacetic acids: a new generation of nonpeptide angiotensin II receptor antagonists. *J. Med. Chem.* **1993**, *36*, 3738–42.
- Keenan, R. M.; Weinstock, J.; Finkelstein, J. A.; Franz, R. G.; Gaitanopoulos, D. E.; Girard, G. R.; Hill, D. T.; Morgan, T. M.; Samanen, J. M.; Peishoff, C. E.; Tucker, L. M.; Aiyar, N.; Griffin, E.; Ohlstein, E. H.; Stack, E. J.; Weidley, E. F.; Edwards, R. M. Potent Nonpeptide Angiotensin II Receptor Antagonists. 2. 1-(Carboxybenzyl)imidazole-5-acrylic Acids. *J. Med. Chem.* **1993**, *36*, 1880–92.
- Keenan, R. M.; Weinstock, J.; Finkelstein, J. A.; Franz, R. G.; Gaitanopoulos, D. E.; Girard, G. R.; Hill, D. T.; Morgan, T. M.; Samanen, J. M.; Hempel, J.; Eggleston, D. S.; Aiyar, N.; Griffin,

- E.; Ohlstein, E. H.; Stack, E. J.; Weidley, E. F.; Edwards, R. Imidazole-5-acrylic Acids: Potent Nonpeptide Angiotensin II Receptor Antagonists Designed Using a Novel Peptide Pharmacophore Model. *J. Med. Chem.* **1992**, *35*, 3858-72.
- (35) Weinstock, J.; Hill, D. T.; Keenan, R. M.; Franz, R. G.; Gaitanopoulos, D. E.; Girard, G. R.; Eggleston, D. R.; Aiyar, N.; Griffin, E.; Ohlstein, E.; Brooks, D. P.; Gellai, M.; Frederickson, T. A.; Weidley, E. F.; Edwards, R. M. A Potent Long-Acting Imidazole-5-Acrylic Acid Angiotensin II AT-1 Receptor Antagonist. *Bioorg. Med. Chem. Lett.* **1994**, *4* (1), 23-8.
- (36) Weinstock, J.; Keenan, R. M.; Samanen, J.; Hempel, J.; Finkelshtein, J. A.; Franz, R. G.; Gaitanopoulos, D. E.; Girard, G. R.; Gleason, J. G.; Hill, D. T.; Morgan, T. M.; Peishoff, C. E.; Aiyar, N.; Brooks, D. P.; Fredrickson, T. A.; Ohlstein, E. H.; Ruffolo, R. R., Jr.; Stack, E. J.; Sulpizio, A. C.; Weidley, E. F.; Edwards, R. M. 1-(Carboxybenzyl)imidazole-5-acrylic Acids: Potent and Selective Angiotensin Receptor Antagonists. *J. Med. Chem.* **1991**, *34*, 1514-17.
- (37) Matsoukas, J. M.; Agelis, G.; Hondrelis, J.; Yamdagni, R.; Wu, Q.; Ganter, R.; Smith, J. R.; Moore, D.; Moore, G. J. Synthesis and Biological Activities of Angiotensin II, Sarileisin, and Sarmesin Analogues Containing Aze or Pip at Position 7. *J. Med. Chem.* **1993**, *36*, 904-11.
- (38) Plucinska, K.; Kataoka, T.; Yodo, M.; Cody, W. L.; He, J. X.; Humblet, C.; Lu, G. H.; Lunney, E.; Major, T. C.; Panek, R. L.; Schelkun, P.; Skean, R.; Marshall, G. R. Multiple Binding Modes for the Receptor-Bound Conformations of Cyclic AII Agonists. *J. Med. Chem.* **1993**, *36*, 1902-13.
- (39) Masek, B. B.; Merchant, A.; Matthew, J. B. Molecular Shape Comparison of Angiotensin II Receptor Antagonists. *J. Med. Chem.* **1993**, *36*, 1230-38.
- (40) Kavalek, J.; Machacek, V.; Svobodova, G.; Sterba, V. Base Catalysed Cyclization of Substituted Esters of Hydantoic and Thiohydantoic Acids. *Collect. Czech. Chem. Commun.* **1986**, *51*, 375-90.
- (41) Grande, M. T.; Trigo, G. C.; Soellhuber, M. M. Synthesis of 2- and 3-substituted-1,2,3,4-tetrahydrodibenzof[h]isoquinolines. *J. Heterocycl. Chem.* **1986**, *23* (3), 929-33.
- (42) Meanwell, N. A.; Roth, H. R.; Smith, E. C. R.; Wedding, D. L.; Wright, J. J. K. Diethyl 2,4-Dioxoimidazolidine-5-phosphonates: Horner-Wadsworth-Emmons Reagents for the Mild and Efficient Preparation of C-5 Unsaturated Hydantoin Derivatives. *J. Org. Chem.* **1991**, *56*, 6897-904.
- (43) Khokhlov, P. S.; Kashemirov, B. A.; Mikityuk, A. D.; Chimishkyan, A. L.; Strepikheev, Y. A. Phosgenation of α -amino-phosphonoacetates. *Zh. Obshch. Khim.* **1984**, *54* (10), 2237-41.
- (44) **19** was prepared by benzylation of 5-bromo-2-butyl-3H-imidazole-4-carbaldehyde with 4-(bromomethyl)-2-[1-(triphenylmethyl)tetrazol-5-yl]biphenyl followed by selective dehalogenation with H₂ and Pd/C.
- (45) **27** was prepared by benzylation of 5-bromo-2-butyl-3H-imidazole-4-carboxaldehyde with 4-nitrobenzyl bromide followed by reduction with Ra Ni/H₂ and treatment with trifluoromethanesulfonic anhydride.
- (46) **30** was prepared by benzylation of 5-chloro-2-butyl-3H-imidazole-4-carbaldehyde with 5-[4-(bromomethyl)phenyl]-1-(triphenylmethyl)-1H-tetrazole followed by dehalogenation with H₂ and Pd/C.
- (47) Hodges, J. C.; Klutchko, S. (Phenylmethyl)imidazole angiotensin II receptor antagonists. PCT Int. Appl. WO 9313077, 1993.
- (48) Dudley, D. T.; Panek, R. L.; Major, T. C.; Lu, G. H.; Bruns, R. F.; Klinkefus, B. A.; Hodges, J. C.; Weishaar, R. E. Subclasses of Angiotensin II Binding Sites and Their Functional Significance. *Mol. Pharmacol.* **1990**, *38*, 370-7.
- (49) Ryan, M. J.; Hicks, G. W.; Batley, B. L.; Rapundalo, S. T.; Patt, W. C.; Taylor, D. G.; Keiser, J. A. Effect of an Orally Active Renin Inhibitor CI-992 on Blood Pressure in Normotensive and Hypertensive Monkeys. *J. Pharmacol. Exp. Ther.* **1994**, *268* (1), 372-9.
- (50) Juliano, L.; Paiva, A. C. M. Conformation of angiotensin II in aqueous solutions. Titration of several peptide analogs and homologs. *Biochemistry* **1974**, *13* (11), 2445-50.
- (51) Khosla, M. C.; Leese, R. A.; Maloy, W. L.; Ferreira, A. T.; Smeby, R. R.; Bumpus, F. M. Synthesis of some analogs of angiotensin II as specific antagonists of the parent hormone. *J. Med. Chem.* **1972**, *15* (8), 792-5.
- (52) *A Textbook of drug design and development*; Krogsgaard-Larsen, P., Bundgaard, H., Eds.; Harwood Academic Publishers GmbH: Switzerland, 1994; ISBN 3718650991.
- (53) *Quantitative drug design: a critical introduction*; Martin, Y. C., Ed.; Marcel Dekker Inc.: New York, 1978; ISBN 0824765745.
- (54) Topliss, J. G. Utilization of operational schemes for analog synthesis in drug design. *J. Med. Chem.* **1971**, *15*, 1006-11.
- (55) Topliss, J. G. A manual method for applying the Hansch approach to drug design. *J. Med. Chem.* **1977**, *20*, 463-9.
- (56) Craig, P. N. Interdependence between physical parameters and selection of substituent groups for correlation studies. *J. Med. Chem.* **1971**, *14*, 680-4.
- (57) Regression plot: $\log(\text{IC}_{50})$ vs CLOGP(R₂), $Y = -0.11 + 0.502X$, $R^2 = 0.721$, $P < 0.005$; $\log(1/\text{IC}_{50})$ vs CLOGP(R₂), $Y = -0.458 + 0.012X - 0.105 \times 2$, $R^2 = 0.737$, $P < 0.01$. The log P was calculated for the substituent R₂ since the CLOGP algorithm was unable to calculate the log P of the hydantoin. This does not affect the correlation since only the R₂ substituent is varied in the comparison of the analogs prepared.
- (58) Regression plot: HPLC(t_R) vs CLOGP(NR), $Y = -1.294 + 3.479X$, $R^2 = 0.912$, $P < 0.0001$. See supporting information for data.
- (59) Kaliszan, R. High performance liquid chromatographic methods and procedures of hydrophobicity determination. *Quant. Struct.-Act. Relat.* **1990**, *9* (2), 83-87.
- (60) Leo, A.; Weininger, D.; Weininger, A. CLOGP. CMR Medicinal Chemistry Project, Pomona College, Claremont, CA. Release 3.54 distributed by Daylight Chemical Information Systems, 1989.
- (61) Taylor, P. J.; Wait, A. R. Values of σ_I for heterocycles. *J. Chem. Soc., Perkin Trans. II* **1986**, *11*, 1765-70.
- (62) Lewis, S. J.; Mirrless, M. S.; Taylor, P. J. Rationalizations among heterocyclic partition coefficients. Part 2: The azines. *Quant. Struct.-Act. Relat.* **1983**, *2* (3), 100-11.
- (63) Bradshaw, J.; Taylor, P. J. Rationalizations among heterocyclic partition coefficients. Part 3: Π -Excessive heterocycles. Some comments on the CLOGP algorithm. *Quant. Struct.-Act. Relat.* **1989**, *8* (4), 279-87.
- (64) Regression plot: $\text{obsd log } P$ vs CLOGP(NR), $Y = 2.671 + 0.539X$, $R^2 = 0.931$, $P < 0.0001$. See supporting information for data.
- (65) The racemic saturated analog of compound **97** or **98** has an IC₅₀ of 161 nM (AT₁).
- (66) Ma, Z. Application of Michael addition method on metabolites of chemical substance chemo-synthesis. *Weisheng Yanjiu* **1994**, *23* (6), 333-4.
- (67) Kraemer, T.; Dressel, J. D.; Fey, P.; Hanco, R.; Huebsch, W.; Mueller, U.; Mueller-Gliemann, M.; Beuck, M.; Kazda, S.; Knorr, A.; Stasch, J.-P.; Wohlfeil, S. Preparation of cycloalkyl- and heterocyclic substituted imidazolylpropenic acid derivatives. *Eur. Pat. Appl. EP 563705*, 1993.
- (68) Reitz, D. B.; Garland, D. J.; Norton, M. B.; Collins, J. T.; Reinhard, E. J.; Manning, R. E.; Olin, G. M.; Chen, S. T.; Palomo, M. A.; McMahon, E. G.; Koehler, K. F. N1- Sterically Hindered 2H-Imidazol-2-one Angiotensin II Receptor Antagonists: The Conversion of Surmountable Antagonists to Insurmountable Antagonists. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 1055-60.
- (69) Schambye, H. T.; Hjorth, S. A.; Bergsma, D. J.; Sathe, G.; Schwartz, T. W. Differentiation between binding sites for angiotensin II and nonpeptide antagonists on the angiotensin II type 1 receptors. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91* (15), 7046-50.
- (70) Ashton, W. T.; Chang, L. L.; Flanagan, K. L.; Hutchins, S. M.; Naylor, E. M.; Chakravarty, P. K.; Patchett, A. A.; Greenlee, W. J.; Chen, T.-B.; Faust, K. A.; Chang, R. S. L.; Lotti, V. J.; Zingaro, G. J.; Schorn, T. W.; Siegl, P. K. S.; Kivlighn, S. D. Triazolone Biphenylsulfonamide Derivatives as Orally Active Angiotensin II Antagonists with Potent AT₁ Receptor Affinity and Enhanced AT₂ Affinity. *J. Med. Chem.* **1994**, *37* (17), 2808-24.
- (71) Wong, P. C.; Duncia, J. V.; Santella, J. B., III; Smith, R. D.; Wexler, R. R.; Timmermans, P. B. M. W. M.; Chiu, A. T. EXP597, a nonpeptide angiotensin II receptor antagonist with high affinities for the angiotensin AT₁ and AT₂ receptor subtypes. *Eur. J. Pharmacol.* **1994**, *260* (2-3), 261-4.
- (72) Chang, L. L.; Ashton, W. T.; Flanagan, K. L.; Chen, T.-B.; O'Malley, S. S.; Zingaro, G. J.; Siegl, P. K. S.; Kivlighn, S. D.; Lotti, V. J.; Chang, R. S. L.; Greenlee, W. J. Triazolone Biphenylsulfonamides as Angiotensin II Receptor Antagonists with High Affinity for Both the AT₁ and AT₂ Subtypes. *J. Med. Chem.* **1994**, *37* (26), 4464-78.