

Design, Synthesis, and Structure-Activity Relationships of a New Series of α -Adrenergic Agonists: Spiro[(1,3-diazacyclopent-1-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)]

Alex A. Cordi,* Jean-Michel Lacoste, Jean-Jacques Descombes, Christine Courchay, Paul M. Vanhoutte, Michel Laubie, and Tony J. Verbeuren

Institut de Recherches Servier, 11, rue des Moulineaux, 92150 Suresnes, France

Received March 20, 1995[⊗]

The contractions induced by a partial α_1 -adrenoceptor agonist in cutaneous veins, such as the saphenous vein, show a particular sensitivity to changes in local temperature: the contractility to a partial α_1 -adrenoceptor agonist increases when the temperature is raised, a response that contrasts to that noted with full α_1 - and α_2 -adrenoceptor agonists. This observation may be of importance for the treatment of the symptoms of venous insufficiency, favored during warm summer days. A new series of full and partial α -adrenergic agonists was designed and synthesized, the spiro[(1,3-diazacyclopent-1-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)] **7a-7k** or spiro-imidazolines. Using *in vitro* (femoral artery and saphenous vein) and *in vivo* (pithed rat) biological evaluations, structure-activity relationships could be defined which allowed the discovery of a full α_2 -agonist (**34b**), a full α_1 -agonist (**7s**), and a nonselective partial α_1/α_2 -agonist (**7aa**) endowed with an outstanding veinotonic selectivity as compared to its effect on mean arterial pressure. The latter compound is presently undergoing extensive pharmacological and toxicological evaluations, as a clinical candidate.

Introduction

α -Adrenoceptors play a key role in the control of the vascular tone. Thus, the neurotransmitter noradrenaline, by interacting with α -adrenoceptors present on the vasculature smooth muscle cells, contracts blood vessels thereby increasing pressure. Moreover, the release of noradrenaline from peripheral and cerebral nerve endings is subject to feedback regulation via α -adrenoceptors of the α_2 -subtype. The importance of the α -adrenoceptors is emphasized¹ by the existence of two classes of antihypertensive drugs: the peripheral α_1 -adrenoceptor antagonists and the centrally acting α_2 -adrenoceptor agonists. The field has been recently reviewed,²⁻⁴ and the number, as well as the diversity of the molecular entities which have been evaluated in this area, is breath taking.

Cutaneous veins are very sensitive to changes in environmental physical conditions such as temperature. Using the dog saphenous vein as a model, it was previously reported that contractions to α -adrenoceptor agonists were differentially altered by local temperature changes; thus contractions to full α_1 and α_2 -adrenoceptor agonists increase, while those to partial α_1 -adrenoceptor agonists decrease when the temperature is decreased.⁵ The reverse is true when the temperature is raised. A deficit in venous blood return, leading to venous stasis, is particularly favored during the hot days of summer. Since these symptoms can be, at least in part, counteracted by vasoconstrictor agents, we hypothesized that treatment with a partial adrenoceptor agonist, which will increase venous tone specifically as temperature rises, might be preferred.

Partial and full α_1 and α_2 -adrenoceptor agonists belong to two well-defined classes of chemical structures: phenylethylamines and imidazolines with exclusive structure-activity relationships^{3,4} (SAR). In our




Figure 1.


present design, the imidazoline moiety was used since this class of molecules can lead to very potent ligands endowed with good pharmacokinetic characteristics, good metabolic stability, and limited toxicological liabilities.⁶

A prototypical partial α_1 -adrenoceptor agonist is St 587 (**2**) (Figure 1). This compound has been characterized in many different isolated organ preparations.⁷ The close structural resemblance of this molecule with the partial α_2 -adrenoceptor selective agonist clonidine (**1**) (Figure 1), a central antihypertensive drug,⁸ indicates that the SARs dictating the α_1/α_2 -adrenoceptor selectivity are very narrow.

A common feature of most of these drugs is the connectivity between the aromatic ring and one of the basic, potentially sp^2 nitrogens of the imidazoline: two atoms, a nitrogen and an sp^2 carbon. In addition, many authors⁹⁻¹¹ have speculated about the conformation the molecule has to adopt to interact with the receptor: the basic nitrogen must lie at a short distance above the plane of the aromatic ring with the plane of the imidazoline perpendicular to the plane of the aromatic ring, thus preventing resonance interaction between the aromatic portion and the imidazoline ring. From in-

[⊗] Abstract published in *Advance ACS Abstracts*, August 15, 1995.

Scheme 1



vestigations related to the phenethylamines,¹² it was found that optimum activity resides in compounds where the basic nitrogen is exocyclic to a saturated six-membered ring such as in compound 3. By combining all these elements, our design target could therefore be stipulated as an aromatic ring linked by two atoms of a saturated six-membered ring to a basic sp^2 nitrogen, the latter included in an imidazoline moiety in such a way that the imidazole plane lies perpendicular to the aromatic ring.

To fulfill these requirements, the idea of linking the imidazoline moiety to the tetraline nucleus through a spiranic carbon was investigated. Due to the tetrasubstituted nature of the carbon, it may no longer be sp^2 hybridized, and in order to keep the basic nitrogen atom sp^2 hybridized, the imidazoline ring has to be substituted in the 4(5) position instead of the classical 2 position. To our knowledge, the corresponding open form (4(5)-benzylimidazoline) has never been evaluated as a potential α -adrenergic ligand. In these structures, the two nitrogens are no longer equivalent, which in turn implies that the two tautomeric forms of the amidine are not equalized by symmetry which could potentially lead to a decrease of affinity due to disfavored entropy change. Consideration of Dreiding models confirmed that some of the conformations of 3 could be shared by the rigid spiro-imidazoline 7. Therefore, it was hoped that the rigidification of what was supposed to be the active conformation of agonist ligands at α -adrenoceptors would compensate for the unfavorable entropy contribution built in the 4(5)-substituted imidazolines.

The spiro-imidazolines 7a-kk as well as the close analogues 25-28 and 32-34 were synthesized with the


aim of exploring these hypotheses. The compounds were evaluated *in vitro* and *in vivo*, and the SARs were defined with the hope that an increase in venous tone could be achieved without much influence on the mean blood pressure. These efforts led to the selection of a potent partial adrenoceptor agonist which is presently in development as a potential human venotonic agent.

Synthesis

The spiro-imidazolines of interest, 7a-kk, were synthesized initially through a three-step procedure starting from the tetralones 4a-kk (Scheme 1, route A). The first step was the synthesis of the corresponding amino nitriles 5a-kk by the Strecker reaction.¹³⁻¹⁵ Reduction of the amino nitriles with lithium aluminum hydride in diethyl ether or tetrahydrofuran led to the ethylenediamine derivatives 6a-kk, which in turn were cyclized by mild reaction with formamidine acetate in ethanol at room temperature to provide 7a-kk.

In many cases, the Strecker reaction was sluggish, supposedly due to either the insolubility of the tetralones or their instability. Therefore, alternate routes were devised in order to improve the yield of this critical step. In a first variant (route B), the tetralones 4a-kk were transformed into cyanohydrins 8a-kk by the action of hydrocyanic acid in a two-phase (ether/water) system; treatment of these compounds with a methanolic solution of ammonia led to the corresponding amino nitriles 5a-kk which were transformed into the spiro-imidazolines according to the procedure of method A.

In some cases, a more elaborated approach was used (method C) in which the tetralones were transformed into the enamines 9a-kk by reaction with benzylamine

Scheme 2^a

^a Reagents and conditions: (a) SOCl_2 , toluene or SOCl_2 , DMF, CHCl_3 ; (b) AlCl_3 , CH_2Cl_2 , $\text{CH}_2=\text{CH}_2$; (c) PPA; (d) $\text{Ph}_3\text{PCH}_3^+\text{Br}^-$, KOtBu , ether; (e) $\text{Ti}(\text{NO}_3)_3$, MeOH, CHCl_3 ; (f) $(\text{COCl})_2$, DMF, CHCl_3 ; (g) CH_2N_2 ; (h) $\text{Rh}(\text{AcO})_2$, CH_2Cl_2 ; (i) $\text{CF}_3\text{CO}_2\text{H}$; (j) NaBH_4 , EtOH; (k) TsOH, toluene; (l) mCPBA, CH_2Cl_2 ; (m) $\text{BF}_3\cdot\text{Et}_2\text{O}$, ether; (n) ethylene glycol, TsOH, Δ ; (o) I_2 , $\text{Hg}(\text{OAc})_2$, AcOH; (p) CH_3SLi , Cu_2O , DMSO; (q) AcOH, H_2O .


in the presence of a catalytic amount of *p*-toluenesulfonic acid and azeotropic removal of water by refluxing toluene. Unexpectedly, these enamines reacted with trimethylsilyl cyanide in the presence of zinc iodide to give the benzylamino nitriles in the presence of zinc iodide. Indeed, this route was inspired by the work of Ojima¹⁶ who showed that the benzylimines reacted smoothly with trimethylsilyl cyanide to afford the benzylamino nitriles in the presence of zinc iodide. Therefore, we were surprised to realize that the condensation of the tetralones with benzylamine led, in all cases, to the enamines in place of the imines. Apparently the reaction of enamines with trimethylsilyl cyanide has not been reported in the literature. The favorable course of reaction is perhaps linked to the presence of a very low concentration of the imine (undetectable in the NMR spectra) due to a putative equilibrium between the imine and the enamine allowing the reaction to go slowly to completion.

After reduction of the nitrile function with lithium aluminum hydride in diethyl ether or tetrahydrofuran, the benzyl-protected ethylenediamines **11a–kk** were produced and then transformed into **6a–kk** by debenylation with ammonium formate in the presence of palladium on carbon. The correct choice of synthetic method for a specific ethylenediamine was achieved through trial and error; today we are still unable to deduce rules pertaining to the choice of one method over another. However, experimenting with these three methods, we were always able to reach our target spiroimidazolines.

Very few of the starting tetralones **4a–kk** were commercially available; some of them were not even described. Therefore we investigated the different known routes reported to lead to 2-tetralones (Scheme 2). From these experiences, we came up with the conclusion that depending on the substitution patterns, routes A and B are the more efficient and reliable ways


to attain the different 2-tetralones **4a–kk**. Indeed, while trying to synthesize 2-tetralones with substituents in the 6 or 8 position, route A led unambiguously to the required compounds starting from the corresponding para- or ortho-substituted phenylacetic acids, respectively; whereas starting from para- or ortho-substituted phenylpropionic acids, 2-tetralones bearing substituents in the 7 or 5 position were obtained, respectively, following the procedure of route B. Route C which relied on the rhodium-catalyzed decomposition of diazo ketones led to conflicting results regarding the exact position of the substituents.¹⁷ Finally, route D, which involves the transposition of 1-tetralones into 2-tetralones, is lengthy and limited by the availability of 4-phenylbutyric acid starting materials.

To reach the different methyl-substituted spiroimidazolines **25**, **27**, and **28**, small variances of this synthetic scheme were elaborated. Indeed, when ammonium chloride was replaced with methylamine hydrochloride in the Strecker reaction (Scheme 3), the (methylamino)(aminomethylene)tetrahydronaphthalene derivatives **24** were obtained after the reduction step. Through condensation with formamidine acetate or acetamidine hydrochloride, the spiroimidazolines **25** and **26** were produced, respectively. The reaction of the ethylenediamine **6a** with either acetamidine hydrochloride or cyanogen bromide (Scheme 3) led to the 2-substituted spiroimidazolines **27** and **33**, respectively. Finally, the third methyl-substituted spiroimidazoline isomer **28** was prepared through nonselective alkylation of the parent imidazoline **7a** with dimethyl sulfate in the presence of triethylamine followed by chromatographic separation of the two isomers (Scheme 3). The structure of **28** was unambiguously assigned by NMR spectroscopy through comparison with an authentic sample of **25** synthesized by the Strecker reaction with methylamine as described above. As expected¹⁸ from

Scheme 3^a

^a Reagents and conditions: (a) $\text{CH}_3\text{NH}_2\cdot\text{HCl}$, KCN, AcOH, MeOH; (b) LAH, ether; (c) $\text{HN}=\text{C}(\text{R}_5)\text{NH}_2\cdot\text{AcOH}$, EtOH; (d) $\text{HN}=\text{C}(\text{CH}_3)\text{NH}_2\cdot\text{HCl}$, EtOH; (e) BrCN, CH_2Cl_2 , 0 °C; (f) $(\text{CH}_3\text{O})_2\text{SO}_2$, Et_3N , CH_2Cl_2 .

Scheme 4




Reagents and conditions: a) HCO_2H , HCl; b) HCl, H_2O , Δ ; c) LAH, ether; d) HCO_2Me ; e) DAST, CH_2Cl_2 ; f) BrCN, Et_3N , CH_2Cl_2 .

the relative steric crowding of the two nitrogen reactive centers, the yield of **28** in the alkylation reaction was higher than the yield of **25**.

To reach spiro-oxazolines **32** and **34a,b**, a stepwise hydrolysis of the amino nitrile (Scheme 4) was necessary to prevent degradation of the amino nitrile, probably from a retro-Strecker reaction leading to the starting tetralone. Therefore, the amino amide was obtained by hydration of the nitriles **5a,b** in formic acid saturated with dry hydrochloric acid, and the amides **29a,b** were then safely hydrolyzed into the acids **30a,b**; the latter were reduced with lithium aluminum hydride into the amino alcohols **31a,b** which in turn were cyclized by either diethylaminosulfur trifluoride (DAST) or cyanogen bromide to give the spiro-oxazolines **32** and **34a,b**, respectively.

It is worth noting that attempts to synthesize the isomeric spiro-oxazoline **37** failed; this is due to the high reactivity of the quaternary hydroxyl group toward an elimination reaction leading to a compound bearing a conjugated trisubstituted double bond, **38** (Scheme 5).

Scheme 5^a

^a Reagents and conditions: (a) TMSCN, ZnI_2 ; (b) LAH, ether; (c) HCO_2Me .

Biology

The choice of the different tests used for screening was made after the evaluation in terms of reliability, time efficiency, and product throughput, of many biological assays relying to various organs from different animal species. Since the aim of our research was the discovery of new venotonic agents possessing α -adrenergic agonist properties, particular attention was paid to a possible hypertensive effect which could be a potential major side effect.

Therefore, the compounds described above were tested for their biological activity in the pithed rat (*in vivo*) and on segments of isolated canine saphenous vein and femoral artery (*in vitro*). The pithed rat assay allows the clean and quantitative determination of the impact of the tested compounds on the vascular tone defined by the dose (in $\mu\text{g}/\text{kg}$ iv) which increases the arterial blood pressure by 20 mmHg (C_{20} in Tables 1 and 2). Additionally, the maximum pressure response caused by the agonist (max), expressed in mmHg, is an indication of the efficacy of the tested compounds. In the same assay, the use of selective α_1 - and α_2 -adrenoceptor antagonists (prazosin and yohimbine) in the presence of the tested compounds allows the definition of a C_{20} ratio (praz and yohim, respectively) which is indicative of the contribution of α_1 - and α_2 -adrenergic mechanisms

Table 1. *In Vitro* and *in Vivo* α -Adrenergic Activity of Compounds 7a–kk

compd	structure				pithed rat				isolated blood vessel			
	R1	R2	R3	R4	control		$C_{20}(\text{antag})^c / C_{20}(\text{control})$		femoral artery		saphenous vein	
					C_{20}^a	max ^b	praz	yohim	EC_{50}^d (μM)	E_{max}^e	EC_{50}^d (μM)	E_{max}^e
2				ST 587	10 ± 3	73 ± 6	33	2.2	4.7 ± 0.7	22 ± 3	5.1 ± 1.1	81 ± 3
42				SDV NVI 085	23 ± 7	123 ± 3	5.1	2.4	38 ± 53	28 ± 11	0.023 ± 0.07	38 ± 2
7a	H	H	H	H	4 ± 0.4	120 ± 1	2	5.3	7.5 ± 2	19 ± 4	1.3 ± 0.2	99 ± 3
7b	Cl	H	H	H	0.3 ± 0.06	133 ± 3	1.9	4.7	1.6 ± 0.4	83 ± 7	0.24 ± 0.07	95 ± 2
7c	F	H	H	H	4.6 ± 1.7	131 ± 14	0.6	1	57 ± 21	75 ± 7	2.3 ± 0.9	99 ± 9
7d	H	Cl	H	H	6.3 ± 0.8	116 ± 5	2.6	5.8	ND	0	5.5 ± 1.1	72 ± 1
7e	H	F	H	H	7.8 ± 3	108 ± 6	5	7.7	ND	8 ± 0.7	11 ± 1	67 ± 7
7f	H	H	Cl	H	6.3 ± 0.8	120 ± 7	2.8	12.8	ND	0	6 ± 1	66 ± 6
7g	H	H	H	Cl	7 ± 1.4	125 ± 8	1.3	2.8	NT	0	NT	0
7h	Cl	H	Cl	H	4.2 ± 0.5	137 ± 7	NT	NT	8 ± 2	75 ± 1	6 ± 0.7	100 ± 2
7i	Cl	H	H	CH ₃ O	NT	NT	NT	NT	0.27 ± 0.16	102 ± 7	0.027 ± 0.008	106 ± 3
7j	H	Cl	Cl	H	33 ± 5	101 ± 5	7	3.1	ND	0	44 ± 36	36 ± 3
7k	H	Cl	H	Cl	19 ± 5	91 ± 10	9.5	3.1	21 ± 19	10 ± 3	2 ± 0.5	45 ± 3
7l	H	F	H	F	31 ± 10	84 ± 7	2.4	5.7	ND	0	6.3 ± 2.9	35 ± 8
7m	CH ₃ O	H	H	H	0.64 ± 0.2	139 ± 3	2	4.6	7.2 ± 5	97 ± 3	0.5 ± 0.1	100 ± 3
7n	OH	H	H	H	4.3 ± 0.4	119 ± 3	0.9	12	39 ± 3	22 ± 1	10 ± 4	66 ± 2
7o	H	CH ₃ O	H	H	1.5 ± 0.4	106 ± 6	3.9	6.9	ND	0	7.1 ± 0.3	49 ± 5
7p	H	H	CH ₃ O	H	42 ± 6	105 ± 7	2.6	1.5	ND	3	8.8 ± 2	60 ± 4
7q	H	H	iPrO	H	74 ± 27	62 ± 8	11	66	ND	0	3.9 ± 4	15 ± 3
7r	H	H	H	CH ₃ O	0.85 ± 0.2	139 ± 4	5.4	4.7	2 ± 0.7	93 ± 4	0.8 ± 0.2	101 ± 2
7s	CH ₃ O	H	H	CH ₃ O	0.05 ± 0.02	128 ± 7	21	2	0.12 ± 0.04	77 ± 4	0.3 ± 0.07	106 ± 2
7t	CH ₃ O	H	H	CH ₃ S	0.003 ± 0.01	136 ± 3	75	45	0.29 ± 0.2	96 ± 2	0.18 ± 0.13	98 ± 1
7u	CH ₃ O	H	H	C ₂ H ₅ O	0.06 ± 0.004	147 ± 15	21.5	4.8	0.08 ± 0.05	85 ± 5	0.8 ± 0.2	98 ± 0.5
7v	C ₂ H ₅ O	H	H	C ₂ H ₅ O	0.6 ± 0.1	157 ± 10	24	7.7	0.85 ± 0.5	55 ± 3	2.7 ± 1.3	68 ± 14
7w	H	CH ₃ O	CH ₃ O	H	98 ± 18	129 ± 10	1.8	NT	0	0	NT	0
7x	H	CH ₃ O	OH	H	1.2 ± 0.3	134 ± 7	9.2	2.8	1.4 ± 0.5	69 ± 8	19 ± 11	97 ± 1
7y	H	CH ₃ O	H	CH ₃ O	7.9 ± 1.9	109 ± 9	2.9	2.6	0.7 ± 0.7	25 ± 10	7.3 ± 3	44 ± 8
7z	H	H	CH ₃ O	CH ₃ O	142 ± 26	107 ± 4	2.9	2.4	50 ± 20	21 ± 5	52 ± 21	67 ± 13
7aa	H	CH ₃	H	H	8 ± 1.5	111 ± 2	6.1	7.9	74 ± 56	5 ± 0.5	0.26 ± 0.01	64 ± 5
7bb	H	Et	H	H	209 ± 119	61 ± 13	2.5	2.6	ND	0	75 ± 57	50 ± 8
7cc	H	iPr	H	H	245 ± 44	66 ± 13	2.9	2.9	ND	0	4.3 ± 3	53 ± 20
7dd	H	tBut	H	H	465 ± 54	45 ± 1	1.7	1.6	ND	0	40 ± 2	35 ± 3
7ee	H	CF ₃	H	H	320 ± 55	55 ± 8	10	2.7	ND	0	ND	7
7ff	H	H	CH ₃	H	5.4 ± 1.3	114 ± 1	1.3	6.1	14 ± 10	12 ± 3	2.1 ± 1.1	44 ± 10
7gg	CH ₃	H	H	CH ₃	0.5 ± 0.09	140 ± 5	2.1	4.8	3 ± 1.7	67 ± 3	2.4 ± 1.8	84 ± 7
7hh	H	CH ₃	CH ₃	H	37 ± 4	72 ± 8	0.5	2.5	ND	0	2.7 ± 3.1	64 ± 13
7ii	H	.CH ₂ -CH ₂ -CH ₂ .	H	H	74 ± 11	68 ± 10	1.1	5	ND	0	0.07 ± 0.01	17 ± 2
7jj	H	CH ₃	H	CH ₃	20 ± 3.5	91 ± 10	2.9	2.8	0.8 ± 0.2	5 ± 0.5	2.9 ± 0.8	66 ± 9
7kk	H	CH ₃	H	CH ₃ O	10 ± 6	136 ± 3	1.6	1	0.8 ± 0.5	60 ± 9	3.9 ± 1.9	81 ± 1

^a Concentration which increases the arterial pressure by 20 mmHg in the pithed rat, expressed in $\mu\text{g}/\text{kg}$ iv (mean ± SD). NT, not tested; ND, not determined due to the very low activity of the compound. ^b Maximal pressure response (mean ± SD) caused by the agonist, expressed in mmHg (the maximum obtained with phenylephrine is 150 mmHg). ^c Ratio of C_{20} obtained in the absence or presence of 0.1 mg of prazosin or 1 mg of yohimbine. ^d EC_{50} value for the agonist in the isolated blood vessel shown in μM (mean ± SD). ^e E_{max} = maximal contractile response obtained with the agonist expressed as the percentage (mean ± SD) of the maximal contraction to phenylephrine in the blood vessel indicated.

to the pressure response. The *in vitro* evaluation allows the comparison in terms of affinity (EC_{50}) and efficacy (E_{max}) on two types of vascular beds which might be responsible for either the expected therapeutic efficacy (saphenous vein) or the potential side effect (femoral artery).

Results and Discussion


The results of the biological evaluation of the spiro-imidazolines are presented in Tables 1 and 2. A first conclusion to be drawn from these data is that a large number of active compounds was detected in this series. Moreover, the unsubstituted spiro-imidazoline **7a** is a very potent compound as compared to the 2-(phenyl-amino)imidazoline series (e.g., ST 587 and clonidine) where a 2,6 or 2,5 substitution pattern is requisite for

high activity. A possible explanation is that the relative geometry of the nitrogen and the aromatic ring in the spiro-imidazolines fits particularly well the pharmacophore geometry of the adrenergic receptor agonist conformation.

A further interpretation of these tables in terms of SAR illustrated that the set of compounds encompasses the whole spectra of α -adrenergic activities; indeed, the substances cover the whole range from full to partial α_1/α_2 -adrenoceptor agonists.

From the hypothesis stated above, it is suspected that the nitrogen which mimicks the basic function of norepinephrine is N1 of the spiro-imidazoline. In addition, it might be that this nitrogen needs to be sp^2 hybridized.

In the spiro-imidazolines, there are two tautomeric forms: one with N1 sp^2 hybridized and the other with

Table 2. *In Vitro* and *In Vivo* α -Adrenergic Activity of Compounds 25–34b


compd	structure				pithed rat				isolated blood vessel			
	R	X	Y	Z	control		$C_{20}(\text{antag}^r)/C_{20}(\text{control})$		femoral artery		saphenous vein	
					C_{20}^a	max ^b	praz	yohim	EC ₅₀ ^d (μM)	E_{max}^e	EC ₅₀ ^d (μM)	E_{max}^e
7a	H	N	NH	H	4 ± 0.4	120 ± 3	2	5.3	7.5 ± 2	19 ± 4	1.3 ± 0.2	99 ± 3
25	H	NCH ₃	N	H	675 ± 138	73 ± 28	1.55	1.9	ND	0	ND	0
26	H	NCH ₃	N	CH ₃	1059 ± 237	73 ± 27	1	1.1	ND	0	ND	0
27	H	N	NH	CH ₃	166 ± 32	113 ± 7	1.8	5.3	ND	0	400 ± 27	23 ± 6
28	H	N	NCH ₃	H	150 ± 27	77 ± 8	2.1	2.5	ND	0	25 ± 10	31 ± 6
32	H	N	O	H	>10,000	38	ND	ND	ND	0	ND	0
33	H	N	NH	NH ₂	57 ± 4	53 ± 6	2.5	26	50 (2)	2	52 ± 3	36 ± 9
34a	H	N	O	NH ₂	5.2 ± 0.9	95 ± 11	3.5	19.6	26 ± 23	18 ± 7	2.3 ± 0.3	76 ± 6
34b	8-Cl	N	O	NH ₂	0.2 ± 0.02	140 ± 5	2	3	0.36 ± 0.09	68 ± 1	0.014 ± 0.04	117 ± 7

^a Concentration which increases the arterial pressure by 20 mmHg in the pithed rat, expressed in $\mu\text{g}/\text{kg}$ iv (mean \pm SD). NT, not tested; ND, not determined due to the very low activity of the compound. ^b Maximal pressure response (mean \pm SD) caused by the agonist, expressed in mmHg (the maximum obtained with phenylephrine is 150 mmHg). ^c Ratio of C_{20} obtained in the absence or presence of 0.1 mg of prazosin or 1 mg of yohimbine. ^d EC₅₀ value for the agonist in the isolated blood vessel shown in μM (mean \pm SD). ^e E_{max} = maximal contractile response obtained with the agonist expressed as the percentage (mean \pm SD) of the maximal contraction to phenylephrine in the blood vessel indicated.

N3 sp² hybridized. To check which one of the two tautomers is actually present as the active ligand, the methyl-substituted spiro-imidazolines **25** and **28** were synthesized. It is clear from the activity of compound **27** that the steric crowding caused by a methyl in this area of the molecule has a dramatic negative influence. In terms of basicity, the influence of the methyl group on the delocalized amidine system must be the same for the two compounds. Therefore, the small difference in activity between the two compounds might correspond to a relatively better fit of the 3-substituted imidazoline (N1 being sp² hybridized) **28** than the 1-substituted imidazoline (N3 being sp² hybridized) **25**, which is in agreement with the hypothesis. Due to the lack of a strong discriminative activity with these compounds, the isomeric spiro-oxazolines **32** and **37** were synthesized with the aim of corroborating the hypothesis. As mentioned earlier, **37** is not stable and isomerized into **38**; in contrast, **32** was stable enough to be evaluated but was almost completely inactive. This might be the consequence of the very low basicity of oxazolines as compared to imidazolines. To increase the basicity of the nitrogen atom, the 2-aminoimidazoline **33** and 2-aminoxazoline **34a** were synthesized and provided much more active compounds. A delocalized basic function was restored which, when protonated, bears only one sp²-hybridized atom: the C2.

Full adrenoceptor agonists in Table 1 and 2 are identified through the comparison of the E_{max} values in the femoral artery and saphenous vein preparations. When the values of the E_{max} are high and almost identical to each other whether the receptor reserve is small (femoral artery) or large (saphenous vein),^{19,20} the compounds behave as full agonists as in cases **7b,c,h,i,m,r-u**. This is further emphasized by the observation that these compounds produce maximal pressor responses in the pithed rat comparable to those produced by the full α_1 -adrenoceptor agonist phenylephrine.²¹ The use of the latter criterion alone would have led to the selection of many misleading compounds such as **7a,d,f,g,n,v,w,x,gg,kk**. All the structures mentioned above but one, **7r**, share a common structural

feature, that is, a substituent in the 8 position (R1 \neq H) of the tetrahydronaphthalene portion of the molecule.

The α_1/α_2 -adrenoceptor selectivity of the compounds was tested by comparing of the C_{20} values in the presence of prazosin, a selective α_1 -adrenoceptor antagonist,²² or yohimbine, a selective α_2 -adrenoceptor antagonist.²³ The sensitivity to the antagonists is expressed by the ratio of the C_{20} in their presence or absence: $r(\text{praz})$ and $r(\text{yohim})$ in Tables 1 and 2. This analysis indicates that selective α_2 -adrenoceptor agonists bear a substituent only in position 8 (R1 \neq H) such as **7b,m**, while selective α_1 -adrenoceptor agonists possess substituents in positions 5 and 8 (R1 and R4 \neq H) such as **7s-u**. Similar relationships existed for the partial agonists such as **7n** and **34b** for the selective α_2 compounds and **7v** for the selective α_1 compounds, **7gg** being an exception to this rule.

In addition, it appears from Table 2 that successful modifications of the heterocycle spiro-imidazolines always lead to selective α_2 compounds, for example, compare **7a** with 2-amino-spiro-imidazoline **33** and 2-amino-spiro-oxazoline **34a**. However, the effect of substitution of the tetrahydronaphthalene and the effect of the modification of the heterocycle is not cumulative as **34b** shows only limited selectivity for the α_2 -adrenoceptor.

Graphically, the conclusions could be represented by the three Markus formulae of Figure 2, where structure I represents mainly the selective α_2 -adrenoceptor agonists, structure II the selective α_1 -adrenoceptor agonists, and structure III the nonselective partial α -adrenoceptor agonists of various efficacy. It is clear from this picture that the SARs so far described are in agreement with previous data from the literature since structure I can mimic the substitution pattern of clonidine (**1**), a partial α_2 -adrenoceptor agonist, and structure II represents the kind of substitution prevailing in ST 587 (**2**) and SDV NVI 085 (**42**),²⁴ two partial α_1 -adrenoceptor agonists.²⁵

A major goal of the present project was to detect compounds acting specifically on the veins while minimally affecting mean blood pressure. From the results of the biological evaluations presented in Tables 1 and




Figure 2.

Table 3. Ratio C_{20}/EC_{50} of Selected Compounds 7

compd	C_{20} ($\mu\text{g}/\text{kg}$) pithed rat ^a	EC_{50} (μM) saphenous vein ^b	ratio C_{20}/EC_{50}
7d	6.3 \pm 0.8	5.5 \pm 1.1	1.15
7e	7.8 \pm 3	11 \pm 1	0.71
7f	6.3 \pm 0.8	6 \pm 1	1.05
7o	1.5 \pm 0.4	7.1 \pm 0.3	0.21
7p	42 \pm 6	8.8 \pm 2	4.77
7aa	8 \pm 1.5	0.26 \pm 0.01	31
7cc	245 \pm 44	4.3 \pm 3	57
7hh	37 \pm 4	2.7 \pm 3	14
7ii	74 \pm 11	0.07 \pm 0.01	1,057
7jj	20 \pm 3.5	0.8 \pm 0.5	25

^a Concentration which increases the arterial pressure by 20 mmHg in the pithed rat, expressed in $\mu\text{g}/\text{kg}$ iv (mean \pm SD). ^b EC_{50} value for the agonist in the isolated blood vessel shown in μM (mean \pm SD).

2, it is obvious that such compounds have been detected. For instance, by comparing the EC_{50} and E_{max} values on veins and arteries for compounds **7d–f, o, p, aa, c, hh–jj**, a high level of selectivity for venous tissues could be detected since all these compounds powerfully contracted the veins but possessed very limited activity on arteries with EC_{50} values too low to be defined. Another way of specifying the venous selectivity was by calculating the empirical ratio between the C_{20} value in the pithed rat and the EC_{50} value obtained in the vein. The use of this ratio proved to be more discriminating than the use of the ratio of EC_{50} arteries versus veins since many compounds were only poorly active on the arteries *in vitro*.

Using the ratio reported in Table 3, it is possible to rank the molecules starting from the most selective compound to the least selective one: **7ii** > **7cc** > **7aa** > **7ii** > **7hh** > **7p** \approx **7o** \approx **7d** \approx **7f** \approx **7e**. The compounds of Table 3 present a very high structural homogeneity, in the sense that they all bear a substituent in the 6 and/or 7 position of the tetrahydronaphthalene (R2 and/or R3 \neq H) and only in these positions. In addition, regarding the best profile, one can deduce that the better substituent is an alkyl (methylene, methyl, or isopropyl) in position 7; indeed, the first five of the eight compounds which have been ranked contain such a substituent. In terms of pharmacological characterization, the results of Table 1 indicate that these compounds obviously behave as partial adrenergic agonists with a low selectivity for the α -adrenoceptor subtypes. Despite this lack of selectivity, compound **7aa** proved to be the most interesting venotonic agent detected in

Table 4. Binding Affinities (K_i , μM) of Selected Spiro-imidazolines for the α_1 , α_{2a} , and α_{2b} -Adrenergic Sites

compd	α_1 rat cortex ^a	α_{2a} human platelets ^a	α_{2b} neonatal rat lung ^a
2	65.8	1.56	0.179
7a	14.4	0.034	0.275
7c	2.5	0.012	0.152
7k	0.7	0.003	0.095
7m	16	0.051	0.099
7o	16	0.018	0.193
7p	29.4	0.087	1.2
7r	3.4	0.012	0.159
7s	2		
7t	0.6		
7aa	1	0.007	0.05
7kk	0.6		
34b	0.8	0.0016	0.02

^a Mean of at least three independent determinations made in duplicate. The standard deviations were always less than 45% of the geometric mean.

this series. The design of the molecule was built from known compounds having known pharmacology, but the desired compound was a molecule endowed with a unique pharmacological profile. Therefore, the spiro-imidazolines were investigated in the search for the optimum compound, not knowing if this goal was realistic or not. At the start of the program, neither the exact selectivity between the α_1 - and α_2 -adrenoceptors nor the efficacy at each adrenoceptor site, necessary to fulfill our target, could be anticipated. Today, it is our goal to define the exact pharmacology of our lead compound and the mechanism of the physiological effects measured.

Extensive binding evaluation of the compounds described has not been performed because the pharmacological results obtained with several of the initial molecules synthesized were disappointing. Table 4 shows the results of this evaluation regarding the α_1 -site and two subtypes of the α_2 -adrenergic receptor, the α_{2a} and the α_{2b} . The affinities detected in the α_2 binding tests are well in line with the data collected in the isolated organs and the *in vivo* assays; the affinities detected for the α_1 binding site are much weaker than could be expected from the result of the saphenous vein assays for instance. Indeed, for a full agonist, the EC_{50} values should be in the same range as the K_i , and for a partial agonist, the EC_{50} values should be larger than the K_i .²⁶ Comparing the results of Table 4 with the results of Tables 1 and 2, it is illustrated that in most cases the K_i is larger than the EC_{50} , even when the compound clearly behaves as a partial agonist (**7aa**). We suspect that α_1 -agonist binding is not correctly measured by the displacement of a tritiated antagonist such as prazosin.²⁷ Indeed, for some receptors, agonist binding needs to be defined by the displacement of a labeled agonist, while affinities of antagonists can only be correctly measured by the displacement of labeled antagonists.²⁸

In conclusion, by developing a new family of adrenoceptor ligands derived from a spiro-imidazoline skeleton, we were able to define SARs leading to the discovery of a full α_2 -adrenoceptor agonist (**34b**, S 17824-1), a full α_1 -adrenoceptor agonist (**7s**, S 17265-1), and a nonselective partial α_1/α_2 -adrenoceptor agonist (**7aa**, S 17089-1) endowed with an outstanding venous selectivity. The latter compound is presently undergoing extensive pharmacological and toxicological evaluation as a clini-

cal candidate. Reports dealing with the resolution of the two enantiomers of **7aa** as well as with its complete pharmacological characterization are currently in preparation.

Experimental Section

Biology. Pithed Rat. Male Sprague-Dawley rats (300–400 g) were anesthetized with ether. After cannulation of the trachea, the animals were pithed (to destroy the spinal cord) and immediately placed under artificial ventilation. The vagus nerves were sectioned, and the carotid arteries were ligated. One of the carotid arteries was cannulated for monitoring of the arterial blood pressure. Three more catheters were inserted into both jugular veins and the penile vein; these catheters were used for injection of compounds. The body temperature of the animals was maintained at 36 °C. The animals were pretreated with a dose of the β -adrenoceptor blocker tertatolol (100 mg/kg) and, in some cases, 10 min later with either the α_1 -adrenoceptor antagonist prazosin (100 μ g/kg) or the α_2 -adrenoceptor antagonist yohimbine (1 mg/kg). Another 10 min later, cumulative doses of the compounds to be tested were injected at 20 s intervals. The changes in arterial pressure were recorded with a pressure transducer (Statham P23XL) and expressed in mmHg. The experimental setup allows the calculation of the dose which augments the arterial pressure with 20 mmHg (C_{20}) as well as the maximal effect of the compound which is expressed in mmHg. The reference α_1 -adrenoceptor agonist phenylephrine augments arterial pressure maximally by 150 mmHg. The results are expressed as the mean (\pm SD) of three to six independent determinations. The α_1 - or α_2 -adrenergic properties of the compounds are evaluated by comparing the C_{20} values in the presence and absence of either prazosin or yohimbine by calculating the ratio $C_{20}(\text{antagonist})/C_{20}(\text{control})$.

Isolated Blood Vessels. These experiments were performed on blood vessels obtained from mongrel dogs (weight 15–25 kg) anesthetized with sodium pentobarbital (30 mg/kg iv) using a technique previously described.¹⁹ After anesthesia, saphenous vein and femoral artery segments were prepared and immediately placed in physiological saline solution (composition in mM: NaCl, 118; KCl, 4.7; NaHCO₃, 25; CaCl₂, 1.25; MgSO₄, 1.2; KH₂PO₄, 1.1; and glucose, 10). Rings were prepared from the blood vessels, and they were mounted under a tension of 4 g (femoral artery) or 1 g (saphenous vein) in organ chambers filled with the salt solution at 37 °C and oxygenated with 95% O₂/5% CO₂. The isometric tension development was continuously monitored via a force transducer (Statham) coupled to a computerized measuring system. After mounting, the tissues were allowed to equilibrate for 90 min prior to experimentation; the organ bath solution was replaced every 30 min. The blood vessels were then incubated, throughout the duration of the experiments, in physiological saline solution containing rauwolscine (10⁻⁷ M) in order to block the α_2 -adrenoceptors. After readjustment of the basal tension, a contraction was evoked in each tissue with KCl (100 mM). After rinsing and return to base line conditions, a contraction was induced with a submaximal concentration of phenylephrine (1 μ M). After rinsing and return to basal conditions, a concentration–response curve was performed with one of the putative α -adrenoceptor agonists using increasing concentrations separated by 1/2 log. This allowed determination of the maximal contraction to the agonist and calculation of the EC₅₀ value (concentration that caused half-maximal contraction) using standard regression analysis techniques. Those values are the mean (\pm SD) of three to six independent determinations, unless otherwise indicated (2), and are used to compare the activities of the different substances in arterial and venous tissues.

Radioligand-Binding Assays. Binding to rat (α_1 ²⁹ and α_2 ³⁰) and human (α_{2a} ³¹) adrenergic receptors was measured using [³H]prazosin (0.3 nM) in rat cortex and [³H]RX821002 (0.5 and 1.5 nM) in neonatal rat lung and human platelets, respectively. For each receptor, data were collected for a key reference ligand, prazosin ($K_i = 0.14$ nM) or phentolamine ($K_i = 11$ and 14 nM), respectively. Nonspecific binding was

defined in the presence of cold prazosin (10 μ M) and phentolamine (10 μ M); the percentage of specific binding was 87%, 90%, and 95%, respectively. Drugs were tested on the specific binding of the three receptors. IC₅₀ values for the displacement of the [³H]radioligands were determined by log-probit analysis of data from inhibition experiments in which nine concentrations of drugs spanning 3 orders of magnitude were used. K_i values were calculated according to the equation:³² $K_i = IC_{50}/(1 + L/K_d)$ with L the concentration and K_d the apparent dissociation constant of the [³H]ligand obtained from Scatchard analysis of saturation experiment data. Each K_i value was determined in at least three independent experiments made in duplicate.

Chemistry. Reagents were commercially available and of synthetic grade. ¹H NMR spectra relative to TMS were recorded on Bruker 200 or 400 MHz spectrometers. Infrared spectra were obtained as Nujol emulsion, on a Bruker Fourier transform spectrometer. All new substances were homogeneous in TLC and exhibited spectroscopic data consistent with the assigned structures. Elemental analyses (C, H, N) were performed on a Carlo Erba 1108 instrument and agree with the calculated values within the $\pm 0.4\%$ range. Melting points were obtained on a Reichert hot stage microscope and are uncorrected. Silica gel 60, Merck 230–400 mesh, was used for both flash and medium pressure chromatography. TLC was performed on precoated 5 \times 10 cm, Merck silica gel 60 F254 plates (layer thickness 0.25 mm).

Synthesis of Tetralones. Route B. Synthesis of Precursors (Substituted Phenylpropionic Acids). Ethyl 3-(2-Ethoxy-5-methoxyphenyl)prop-2-enoate. Maintaining a reaction temperature of 5–10 °C, a solution of ethyl diethylphosphonoacetate (6.30 g, 280 mmol) in toluene (230 mL) was added dropwise, under N₂, to a cooled (5 °C), stirred suspension of NaH (6.70 g, 280 mmol) in toluene (200 mL). The reaction mixture was allowed to warm to room temperature and stirred for 1 h. A solution of 2-ethoxy-5-methoxybenzaldehyde³³ (50.5 g, 280 mmol) in toluene (200 mL) was added dropwise and the medium stirred at 20 °C overnight. The mixture was poured onto ice/water (2 L) and extracted with Et₂O (2 \times 500 mL). The organic layers were dried and concentrated under vacuum to give the crude cinnamate as an oil (52.5 g, 75%), used without further purification for the next step. ¹H NMR (CDCl₃): δ 8.00 (d, 1H), 6.50 (d, 1H), 7.05 (d, 1H), 6.90–6.75 (m, 2H), 4.25 (q, 2H), 4.05 (q, 2H), 3.75 (s, 3H), 1.40 (t, 3H), 1.30 (t, 3H).

Ethyl 3-(2,5-Diethoxyphenyl)prop-2-enoate. Using the above procedure, but starting from 2,5-diethoxybenzaldehyde,³⁴ ethyl (2,5-diethoxyphenyl)propenoate was obtained as an oil. ¹H NMR (CDCl₃): δ 8.00 (d, 1H), 7.05 (d, 1H), 6.90 (dd, 1H), 6.85 (d, 1H), 6.85 (d, 1H), 6.50 (d, 1H), 4.25 (q, 2H), 4.15–3.90 (2q, 4H), 1.50–1.20 (3t, 9H).

Ethyl 3-(2-Ethoxy-5-methoxyphenyl)propanoate. Reduction of the above cinnamate (50 g, 200 mmol) under 1 atm of H₂ and in the presence of a catalytic amount of 10% Pd/C in EtOH (700 mL), followed by filtration and evaporation of the solvent, yielded the propanoate as a white solid (49.5 g, 98%). ¹H NMR (CDCl₃): δ (m, 3H), 4.15 (q, 2H), 4.00 (q, 2H), 3.75 (s, 3H), 2.90 (m, 2H), 2.60 (m, 2H), 1.35 (t, 3H).

Ethyl 3-(2,5-Diethoxyphenyl)propanoate. The above procedure applied to ethyl 3-(2,5-diethoxyphenyl)prop-2-enoate provided a solid, mp 125–126 °C. ¹H NMR (CDCl₃): δ 6.80–6.60 (m, 3H), 4.10 (q, 2H), 4.00 (m, 4H), 2.90 (m, 2H), 2.60 (m, 2H), 1.35 (2t, 6H), 1.25 (t, 3H).

3-(2-Ethoxy-5-methoxyphenyl)propanoic Acid (14u). Saponification of the corresponding ethyl ester (48 g, 190 mmol) with NaOH (2 N, 200 mL) in EtOH (350 mL) and acid workup gave the desired acid (41.50 g, 97%), mp 160–162 °C. ¹H NMR (DMSO-*d*₆): δ 12.10 (br d, 1H), 6.85 (d, 1H), 6.70 (dd, 1H), 4.00 (q, 2H), 3.65 (s, 3H), 2.75 (m, 2H), 2.50 (m, 2H), 1.30 (t, 3H).

3-(2,5-Diethoxyphenyl)propanoic Acid (14v). 14v was prepared in a similar fashion from the corresponding ester, mp 173–174 °C. ¹H NMR (DMSO-*d*₆): δ 12.00 (br d, 1H), 6.80 (d, 1H), 6.70 (d, 1H), 6.65 (dd, 1H), 4.00–3.80 (2q, 4H), 2.75 (m, 2H), 2.45 (m, 2H), 1.45–1.20 (2t, 6H).

Step a: 6-Ethyl-1-indanone (15bb). 3-(4-Ethylphenyl)propanoic acid³⁵ (30 g, 168 mmol) was added portionwise to 500 g of polyphosphoric acid (PPA), warmed up to 60 °C. The mixture was stirred at 80 °C for 1 h, the reaction quenched with an ice/water mixture, and then the mixture extracted with CH₂Cl₂. Combined extracts were washed with water, dried (MgSO₄), and concentrated under vacuum to yield the indanone **15bb** (26 g, 97%) as a yellow solid. A sample was recrystallized to analytical purity from hexane, mp 55–56 °C. ¹H NMR (CDCl₃): δ 7.60 (d, 1H), 7.45 (dd, 1H), 7.40 (d, 1H), 3.10 (m, 2H), 2.70 (q, 2H), 2.65 (m, 2H), 1.25 (t, 3H).

The following compounds were obtained in the same manner.

6-Isopropyl-1-indanone (15cc): mp 73–74 °C. ¹H NMR (CDCl₃): δ 7.65 (m, 1H), 7.50 (dd, 1H), 7.45 (d, 1H), 3.10 (m, 2H), 3.00 (m, 1H), 2.70 (m, 2H), 1.25 (d, 6H).

6-tert-Butyl-1-indanone (15dd): mp 94–95 °C. ¹H NMR (CDCl₃): δ 7.80 (d, 1H), 7.70 (dd, 1H), 7.45 (d, 1H), 3.10 (m, 2H), 2.70 (m, 2H), 1.35 (s, 9H).

4-Ethoxy-7-methoxy-1-indanone (15u): oil. ¹H NMR (CDCl₃): δ 7.00 (d, 1H), 6.70 (d, 1H), 4.05 (q, 2H), 3.90 (s, 3H), 3.10–2.60 (m, 4H), 1.40 (t, 3H).

4,7-Diethoxy-1-indanone (15v): oil. ¹H NMR (CDCl₃): δ 6.65 (AB system, 2H), 4.05 (q, 4H), 3.10–2.60 (m, 4H), 1.40 (2t, 6H).

Step b: 6-Ethyl-1-methyleneindan (16bb). Following the procedure of Cannon et al.,³⁶ methyltriphenylphosphonium bromide (103 g, 284 mmol) and KOtBu (31.90 g, 284 mmol) were added successively to a solution of **15bb** (23 g, 143 mmol) in Et₂O (500 mL) and CH₂Cl₂ (160 mL). After stirring at 20 °C under N₂ for 1 h, the mixture was diluted with water (800 mL) and extracted with EtOAc (3 × 300 mL). The combined organic extracts were washed with water, dried (MgSO₄), and evaporated under reduced pressure. The oily brown residue was purified by chromatography (SiO₂, cyclohexane as eluent) to give a pale yellow solid (12.40 g, 55%), mp 72–73 °C. ¹H NMR (CDCl₃): δ 7.30 (d, 1H), 7.20 (d, 1H), 7.05 (dd, 1H), 5.45 (m, 1H), 5.00 (m, 1H), 3.00–2.70 (m, 4H), 2.60 (q, 2H), 1.25 (t, 3H).

The following compounds were prepared by the above procedure.

6-Isopropyl-1-methyleneindan (16cc): mp 78–79 °C. ¹H NMR (CDCl₃): δ 7.35 (m, 1H), 7.25–7.05 (m, 2H), 5.45 (m, 1H), 5.00 (m, 1H), 5.00 (m, 1H), 3.00–2.70 (m, 5H), 1.35 (d, 6H).

6-tert-Butyl-1-methyleneindan (16dd): mp 87–88 °C. ¹H NMR (CDCl₃): δ 7.55 (d, 1H), 7.30 (dd, 1H), 7.20 (d, 1H), 5.45 (m, 1H), 5.00 (m, 1H), 2.95 (m, 2H), 2.80 (m, 2H), 1.35 (s, 9H).

5,6-Dimethyl-1-methyleneindan (16hh): starting from the known 5,6-dimethyl-1-indanone,³⁷ oil. ¹H NMR (CDCl₃): δ 7.25 (s, 1H), 7.00 (s, 1H), 5.35 (m, 1H), 4.95 (m, 1H), 3.00–2.70 (m, 4H), 2.25 (s, 6H).

3,5,6,7-Tetrahydro-1-methylene-1H-s-indacene (16ii): starting from the corresponding known indanone,³⁸ mp 104–105 °C. ¹H NMR (CDCl₃): δ 7.35 (s, 1H), 7.10 (s, 1H), 5.40 (m, 1H), 4.95 (br s, 1H), 3.05–2.75 (m, 8H), 2.25–2.00 (m, 2H).

4-Ethoxy-7-methoxy-1-methyleneindan (16u): oil. ¹H NMR (CDCl₃): δ 6.90 (d, 1H), 6.70 (d, 1H), 5.90 (m, 1H), 5.15 (m, 1H), 4.05 (q, 2H), 3.90 (s, 3H), 3.00–2.70 (m, 4H), 1.40 (t, 3H).

4,7-Diethoxy-1-methyleneindan (16v): mp 90–91 °C. ¹H NMR (CDCl₃): δ 6.60 (s, 2H), 5.90 (m, 1H), 5.15 (m, 1H), 4.15–3.90 (m, 4H), 3.00–2.70 (m, 4H), 1.55–1.30 (2t, 6H).

Step c: 1,2,3,4-Tetrahydro-7-ethyl-2(1H)-naphthalenone (4bb). Using the procedure of Cannon et al.,³⁴ a freshly prepared solution of Ti(NO₃)₃ (25.30 g, 57 mmol) in MeOH (75 mL) was added in one portion to a solution of **16bb** (9.0 g, 57 mmol) in MeOH (180 mL). The mixture was stirred for 10 min and then diluted with CH₂Cl₂ (300 mL). The resulting precipitate was removed by filtration and the filtrate washed with saturated aqueous NaHCO₃ and water before drying (MgSO₄) and concentrating under reduced pressure. The residue was purified by chromatography (SiO₂, CH₂Cl₂) to afford the tetralone **4bb** as a pale yellow solid (6.50 g, 66%), mp 85–86 °C. ¹H NMR (CDCl₃): δ 7.15 (d, 1H), 7.05 (dd, 1H),

6.95 (d, 1H), 3.60 (s, 2H), 3.00 (t, 2H), 2.65 (q, 2H), 2.55 (t, 2H), 1.25 (t, 3H).

The following compounds were prepared by the above procedure.

1,2,3,4-Tetrahydro-7-isopropyl-2(1H)-naphthalenone (4cc): mp 67–68 °C. ¹H NMR (CDCl₃): δ 7.20–6.90 (m, 3H), 3.55 (s, 2H), 3.05 (m, 2H), 3.20–2.75 (m, 1H), 2.55 (m, 2H), 1.25 (d, 6H).

1,2,3,4-Tetrahydro-7-tert-butyl-2(1H)-naphthalenone (4cd): mp 79–80 °C. ¹H NMR (CDCl₃): δ 7.30–7.10 (m, 3H), 3.60 (s, 2H), 3.05 (m, 2H), 2.55 (m, 2H), 1.30 (s, 9H).

1,2,3,4-Tetrahydro-5-ethoxy-8-methoxy-2(1H)-naphthalenone (4u): oil. ¹H NMR (CDCl₃): δ 6.70 (AB system, 2H), 4.00 (q, 2H), 3.80 (s, 3H), 3.50 (s, 2H), 3.10 (m, 2H), 2.55 (m, 2H), 1.45 (t, 3H).

1,2,3,4-Tetrahydro-5,8-diethoxy-2(1H)-naphthalenone (4v): mp 88–89 °C. ¹H NMR (CDCl₃): δ 6.70 (AB system, 2H), 4.00 (m, 4H), 3.50 (s, 2H), 3.10 (m, 2H), 2.55 (m, 2H), 1.40 (m, 6H).

Route C. Step a: 3-[4-(Trifluoromethyl)phenyl]propanoyl Chloride. A mixture of 3-[4-(trifluoromethyl)phenyl]propanoic acid³⁸ (5.50 g, 25 mmol) and thionyl chloride (10 mL) in toluene (100 mL) was refluxed for 2 h. After cooling followed by evaporation under reduced pressure, the acid chloride was obtained as an oil. ¹H NMR (CDCl₃): δ 7.50 (m, 2H), 7.30 (m, 2H), 3.20 (m, 2H), 3.00 (m, 2H).

Step b: 1-Diazo-4-[4-(trifluoromethyl)phenyl]butan-2-one (17ee). A solution of the above acid chloride (5.20 g, 22 mmol) in ether (50 mL) was added to a cooled (–10 °C) solution of diazomethane (72 mmol) in ether (180 mL) at such a rate that the reaction temperature did not exceed 0 °C. After total addition, the temperature was allowed to rise to 20 °C and stirring was continued overnight. After evaporation of the solvent, **17ee** was obtained as a crystalline solid (5.20 g, 98%), mp 45–46 °C.

Step c: 6-(Trifluoromethyl)-3,4-dihydroazulen-1(2H)-one (18ee). A solution of crude **17ee** (4.90 g, 20 mmol) in CH₂Cl₂ (150 mL) was added dropwise to a boiling solution of rhodium acetate (4 mg, 0.4 mmol) in CH₂Cl₂ (200 mL). After evolution of gas had ceased, the solution was cooled and washed with water (2 × 150 mL) and saturated aqueous NaHCO₃ (150 mL), dried (MgSO₄), and concentrated. The brownish oily residue (4 g, 94%) was used for the next step without further purification. ¹H NMR (400 MHz, CDCl₃): δ 6.95 (d, 1H), 6.62 (d, 1H), 6.02 (d, 1H), 6.02 (t, 1H), 2.95 (d, 2H), 2.70 (dd, 2H), 2.66 (dd, 2H).

Step d: 1,2,3,4-Tetrahydro-7-(trifluoromethyl)-2(1H)-naphthalenone (4ee). A stirred solution of crude **18ee** (3.86 g, 18 mmol) in CH₂Cl₂ (80 mL) was acidified by addition of trifluoroacetic acid (TFA) (1.50 mL). After stirring at ambient temperature for 30 min, the mixture was washed with saturated aqueous NaHCO₃ (2 × 70 mL), dried (MgSO₄), and concentrated. The residue was purified by chromatography (SiO₂) eluting with hexane/EtOAc (9/1) to afford the desired tetralone **4ee** (2.5 g, 65%) as an oil. ¹H NMR (400 MHz, CDCl₃): δ 7.55 (d, 1H), 7.42 (s, 1H), 7.40 (d, 1H), 3.70 (s, 2H), 3.10 (t, 2H), 2.45 (t, 2H).

1,2,3,4-Tetrahydro-5,7-dimethyl-2(1H)-naphthalenone (4jj). **4jj** was prepared as described above for **4ee** but using 3-(3,5-dimethylphenyl)propanoic acid³⁹ as starting material, mp 52–53 °C. ¹H NMR (CDCl₃): δ 6.90 (s, 1H), 6.80 (s, 1H), 3.55 (s, 2H), 3.00 (m, 2H), 2.55 (m, 2H), 2.30 (2s, 6H).

1,2,3,4-Tetrahydro-5-methoxy-7-methyl-2(1H)-naphthalenone (4kk). **4kk** was prepared as described for **4ee**, starting from 3-(3-methoxy-5-methylphenyl)propanoic acid.⁴⁰ ¹H NMR (CDCl₃): δ 6.60–6.50 (d, 2H), 3.80 (s, 3H), 3.55 (s, 2H), 3.05 (m, 2H), 2.50 (m, 2H), 2.30 (s, 3H).

1,2,3,4-Tetrahydro-5-methoxy-8-chloro-2(1H)-naphthalenone (4i). The procedure described for **4ee** was employed. Starting from 3-(2-chloro-5-methoxyphenyl)propanoic acid,⁴¹ **4i** was obtained as a pale yellow solid, mp 74–75 °C. ¹H NMR (CDCl₃): δ 7.15 (d, 1H), 6.65 (d, 1H), 6.15 (s, br d, 1H), 5.25 (s, br d, 1H), 3.80 (s, 3H), 3.00–2.70 (m, 4H).

Route E. Step a: Spiro[(1,3-dioxocyclopentane)-2,2'-(8'-methoxy-1',2',3',4'-tetrahydronaphthalene)] (39). A mixture of 1,2,3,4-tetrahydro-8-methoxy-2(1H)-naphthalen-

none⁴² (**4m**) (10 g, 56 mmol), ethylene glycol (3.47 g, 56 mmol), and a catalytic amount of *p*-toluenesulfonic acid in toluene (500 mL) was heated under reflux in a Dean–Stark apparatus. After 3 h of heating, another batch of ethylene glycol (3.40 g) was added and azeotropic distillation continued for a further 3 h. After cooling, the mixture was filtered, washed with water, dried (MgSO₄), and concentrated under vacuum to give a solid (10 g, 81%), mp 124–125 °C. ¹H NMR (CDCl₃): δ 7.10 (t, 1H), 6.75 (d, 1H), 6.65 (q, 1H), 4.05 (m, 4H), 3.80 (s, 3H), 3.00 (t, 2H), 2.90 (s, 2H), 1.90 (t, 2H).

Step b: Spiro[(1,3-dioxocyclopentane)-2,2'-(5'-iodo-8'-methoxy-1',2',3',4'-tetrahydronaphthalene)] (40). To a warm (50 °C) solution of the above product (4.40 g, 20 mmol) in acetic acid (100 mL), solutions of I₂ (15.80 g, 62 mmol) in acetic acid (620 mL) and Hg(OAc)₂ (10.20 g, 32 mmol) in acetic acid (500 mL) were added dropwise, separately and simultaneously, while maintaining the reaction temperature between 45 and 55 °C. The mixture was stirred for 15 min at 50 °C, cooled to room temperature, and stirred for an additional 2 h. The acetic acid was evaporated off, and the crystalline residue was dissolved in concentrated NH₃ and extracted with CH₂Cl₂/H₂O. The organic fraction was dried (MgSO₄) and concentrated to leave a solid (6.0 g, 86%), mp 145–147 °C. ¹H NMR (CDCl₃): δ 7.55 (d, 1H), 6.40 (d, 1H), 4.00 (m, 4H), 3.70 (s, 3H), 2.85 (m, 2H), 2.80 (s, 2H), 1.85 (m, 2H).

Step c: Spiro[(1,3-dioxocyclopentane)-2,2'-(5'-(thiomethyl)-8'-methoxy-1',2',3',4'-tetrahydronaphthalene)] (41). To a suspension of CH₃SLi (8.10 g, 150 mmol) in DMSO (120 mL) were added a solution of the above iodo derivative (6.0 g, 17 mmol) in DMSO (70 mL) and Cu₂O (33.50 g, 230 mmol). After stirring for 5 h at 80 °C under an Ar atmosphere, the mixture was filtered through Celite and washed with CH₂Cl₂; the filtrate was concentrated under vacuum. The residue was taken up in CH₂Cl₂/ice, the organic layer was separated, and the aqueous fraction was extracted with CH₂Cl₂. The combined organic fractions were dried (MgSO₄) and concentrated. The crude product was purified by column chromatography (SiO₂, cyclohexane/Et₂O = 80/20) to yield the thio ether as a yellow solid (3.40 g, 75%), mp 125–127 °C. ¹H NMR (CDCl₃): δ 7.05 (d, 1H), 6.60 (d, 1H), 4.00 (m, 4H), 3.70 (s, 3H), 2.95 (t, 2H), 2.30 (s, 3H), 1.90 (t, 2H).

Step d: 1,2,3,4-Tetrahydro-5-(thiomethyl)-8-methoxy-2(1H)-naphthalenone (4t). A suspension of the above dioxolane (3 g, 11 mmol) in aqueous acetic acid (50 mL, v/v = 30%) was heated overnight at 90 °C. The mixture was diluted with water and extracted with CH₂Cl₂; the organic phase was washed with water (until neutral pH), dried, and concentrated under vacuum to afford the tetralone **4t** as a yellow solid (2.32 g, 95%), mp 110–112 °C. ¹H NMR (CDCl₃): δ 7.20 (d, 1H), 6.70 (d, 1H), 3.75 (s, 3H), 3.50 (s, 2H), 3.20 (m, 2H), 2.50 (m, 2H), 2.30 (s, 3H).

Typical Procedures for the Synthesis of Spiro-imidazolines. Method A. Step a: 2-Amino-2-cyano-8-chloro-1,2,3,4-tetrahydronaphthalene (5b). To a vigorously stirred solution, maintained under N₂ atmosphere and containing 8-chloro-1,2,3,4-tetrahydronaphthalenone⁴³ (**4b**) (7.77 g, 43 mmol), EtOH (60 mL), and water (30 mL) were added KCN (2.87 g, 44 mmol) and NH₄Cl (2.35 g, 44 mmol) successively. After stirring for 20 h at 20 °C, the mixture was concentrated under vacuum and the residue was taken up in EtOAc. The organic phase was washed with water and then extracted with 1 N HCl. Rendering of the aqueous phase alkaline with 35% aqueous NaOH followed by extraction with EtOAc, drying, and evaporation under reduced pressure provided the desired amino nitrile **5b** (5.25 g, 58%) as a solid, mp 67–69 °C. ¹H NMR (CDCl₃): δ 7.30–7.00 (m, 3H), 3.40 (d, 1H), 3.05 (m, 2H), 2.90 (d, 1H), 2.15 (m, 1H), 2.00 (m, 1H).

Step b: 2-Amino-2-(aminomethyl)-8-chloro-1,2,3,4-tetrahydronaphthalene (6b). A solution of **5b** (5.25 g, 25 mmol) in anhydrous THF (40 mL) was added dropwise to a suspension of LiAlH₄ (1.63 g, 43 mmol) in anhydrous THF (50 mL) while maintaining the reaction temperature below 20 °C. The mixture was stirred for 20 min, cooled to 0 °C, and then hydrolyzed by addition of H₂O (1.6 mL), 2 N NaOH (1.6 mL), and H₂O (3.5 mL); the resulting suspension was filtered and the filtrate evaporated to give an oily residue. The diamine

6b was obtained, after purification by column chromatography (SiO₂, CH₂Cl₂/CH₃OH/NH₄OH = 90/10/1), as an oil (4.05 g, 77%). ¹H NMR (CDCl₃): δ 7.20–6.90 (m, 3H), 3.20–2.60 (m, 6H), 1.70 (m, 2H), 1.70–1.20 (m, br d, 4H).

Step c: Spiro[(1,3-diazacyclopent-1-ene)-5,2'-(8'-chloro-1',2',3',4'-tetrahydronaphthalene)], Fumarate (7b). A mixture of the above diamine **6b** (2.10 g, 10 mmol) and formamidine acetate (1.18 g, 11.5 mmol) in EtOH (60 mL) was stirred at 20 °C under an N₂ atmosphere for 10 h. The solvent was evaporated and the residue taken up in 1 N HCl. The acidic phase was washed with EtOAc and rendered basic with 35% aqueous NaOH; the mixture was extracted with EtOAc and the organic layer washed with saturated aqueous NaCl and evaporated. The solid residue was dissolved in EtOH (20 mL) and treated with a solution of fumaric acid (0.81 g, 7 mmol) in EtOH (20 mL). After evaporation of the solvent and recrystallization of the residue in EtOH, **7b** was isolated as a white solid (2.28 g, 68%), mp 213–214 °C. ¹H NMR (DMSO-*d*₆): δ 8.15 (s, 1H), 7.30 (dd, 1H), 7.15 (m, 2H), 6.45 (s, 2H), 4.70 (br d, 2H), 3.65 (AB system, 2H), 2.95 (m, 4H), 1.95 (m, 2H). Anal. (C₁₂H₁₃ClN₂C₄H₄O₄) C, H, N, Cl.

Method B. Step a: 2-Amino-2-cyano-5-(methylthio)-8-methoxy-1,2,3,4-tetrahydronaphthalene (5t). A stirred mixture of tetralone **4t** (2 g, 9 mmol) and NaCN (1.42 g, 29 mmol) in water (50 mL) and Et₂O (40 mL) was cautiously treated with concentrated HCl (1 mL). After stirring for 1 h at 20 °C, the organic phase was separated, washed with water, dried, and concentrated. The oily residue was then treated with 3.5 N methanolic NH₃ solution (10 mL) and stirred in a closed vessel for 6 h at 20 °C. After evaporation of the solvent, the oil was taken up in Et₂O and extracted with 1 N HCl. The aqueous phases were rendered alkaline with 35% aqueous NaOH and then extracted with Et₂O. The amino nitrile **5t** was obtained as a solid after drying and evaporation of the organic phase (1.61 g, 72%), mp 128–130 °C. ¹H NMR (CDCl₃): δ 7.05 (d, 1H), 6.65 (d, 1H), 3.70 (s, 3H), 3.20 (d, 1H), 3.00–2.80 (m, 2H), 2.65 (d, 1H), 2.30 (s, 3H), 2.20–1.70 (m, 2H), 2.00–1.80 (br d, 2H).

Step b: 2-Amino-2-(aminomethyl)-5-(methylthio)-8-methoxy-1,2,3,4-tetrahydronaphthalene (6t). **6t** was prepared according to the procedure described for **6b**, method A, step b, and obtained as an oil. ¹H NMR (CDCl₃): δ 7.10 (d, 1H), 6.70 (d, 1H), 3.80 (s, 3H), 3.10–2.50 (m, 6H), 2.40 (s, 3H), 2.00–1.50 (m, 2H), 1.80–1.50 (br d, 4H).

Step c: Spiro[(1,3-diazacyclopent-1-ene)-5,2'-(5'-(methylthio)-8'-methoxy-1',2',3',4'-tetrahydronaphthalene)], Fumarate (7t). The compound was obtained as described for **13b** in method A, step c, as a white powder (60% yield), mp 188–190 °C. ¹H NMR (DMSO-*d*₆): δ 8.15 (s, 1H), 7.15 (d, 1H), 6.85 (d, 1H), 6.45 (s, 2H), 3.55 (AB system, 2H), 2.90–2.70 (m, 4H), 2.40 (s, 3H), 2.15–1.75 (m, 2H). Anal. (C₁₄H₁₈N₂OS-C₄H₄O₄) C, H, N, S.

Method C. Step a: 2-(Benzylamino)-3,4-dihydronaphthalene (9a). A solution of 2-tetralone (25 g, 171 mmol), benzylamine (19.40 g, 181 mmol), and *p*-toluenesulfonic acid (100 mg) in toluene (500 mL) was brought to reflux with azeotropic distillation of the water/toluene mixture. After 2 h of reflux, the reaction mixture was cooled and filtered. The enamine **9a** was obtained as an oil (39.50 g, 99%) by evaporation of the solvent; it was used without purification for the next step. ¹H NMR (CDCl₃): δ 7.50–6.80 (m, 9H), 5.30 (s, 1H), 4.30 (br s, 2H), 3.50 (m, 1H), 2.80 (m, 2H), 2.30 (m, 2H).

Step b: 2-(Benzylamino)-2-cyano-1,2,3,4-tetrahydronaphthalene (10a). Trimethylsilyl cyanide (16.67 g, 168 mmol) and ZnI₂ (27.15 g, 85 mmol) were added successively to a solution of **9a** (39.50 g, 168 mmol) in CH₂Cl₂ (600 mL) maintained under an N₂ atmosphere. The mixture was stirred at 20 °C overnight and then washed with water; the organic phase was dried (MgSO₄) and concentrated under vacuum to afford **10a** as an oil (37.85 g, 86%). ¹H NMR (CDCl₃): δ 7.50–6.75 (m, 9H), 4.00 (s, br d, 2H), 3.10 (AB system, 2H), 3.00 (m, 2H), 2.20–2.00 (2m, 2H), 1.60 (m, 1H).

Step c: 2-(Benzylamino)-2-(aminomethyl)-1,2,3,4-tetrahydronaphthalene (11a). A solution of **10a** (37.70 g, 144 mmol) in anhydrous THF (200 mL) was added dropwise to a suspension of LiAlH₄ (10.60 g, 280 mmol) in anhydrous THF

Table 5. Preparation Methods of the Spiro-imidazolines **7a–kk**

compd	starting tetralone ^a	method	last step yield (%)	formula	elemental anal.	mp (°C)
7c	4c ⁴⁴	A	59	C ₁₂ H ₁₃ FN ₂ C ₄ H ₄ O ₄	C,H,N	188–190
7d	4d ⁴⁵	A	63	C ₁₂ H ₁₃ ClN ₂ C ₄ H ₄ O ₄	C,H,N,Cl	138–139
7e	4e ⁴⁶	A	55	C ₁₂ H ₁₃ FN ₂ ·0.6C ₄ H ₄ O ₄	C,H,N	197–200
7f	4f ⁴⁵	A	62	C ₁₂ H ₁₃ ClN ₂ C ₄ H ₄ O ₄	C,H,N,Cl	201–203
7g	4g ⁴⁵	A	56	C ₁₂ H ₁₃ ClN ₂ C ₄ H ₄ O ₄	C,H,N,Cl	204–206
7h	4h ⁴³	A	61	C ₁₂ H ₁₂ Cl ₂ N ₂ C ₄ H ₄ O ₄	C,H,N; Cl: calcd, 19.10; found, 18.54	194–195
7i	4i	C	54	C ₁₃ H ₁₅ ClN ₂ O·C ₄ H ₄ O ₄	C,H,N,Cl	204–207
7j	4j ⁴⁵	A	57	C ₁₂ H ₁₂ Cl ₂ N ₂ C ₄ H ₄ O ₄	C,H,N,Cl	138–140
7k	4k ⁴⁵	A	56	C ₁₂ H ₁₂ Cl ₂ N ₂ C ₄ H ₄ O ₄	C,H,N,Cl	204–205
7l	4l ⁴⁶	A	61	C ₁₂ H ₁₂ F ₂ N ₂ ·0.5C ₄ H ₄ O ₄	C,H,N	118–120
7m	4m ⁴⁷	A	64	C ₁₃ H ₁₆ N ₂ O·C ₄ H ₄ O ₄	C,H,N	186–188
7o	4o	A	70	C ₁₃ H ₁₆ N ₂ O·C ₄ H ₄ O ₄	C,H,N	188–190
7p	4p ⁴⁸	A	66	C ₁₃ H ₁₆ N ₂ O·C ₄ H ₄ O ₄	C,H,N	172–174
7q	4q ⁴⁸	C	63	C ₁₅ H ₂₀ N ₂ O·C ₄ H ₄ O ₄	C,H,N	175–177
7r	4r ⁴⁸	A	58	C ₁₃ H ₁₆ N ₂ O·C ₄ H ₄ O ₄	C,H,N	188–190
7s	4s ⁴³	C	61	C ₁₄ H ₁₈ N ₂ O ₂ ·C ₄ H ₄ O ₄	C,H,N	185–188
7u	4u	C	68	C ₁₅ H ₂₀ N ₂ O ₂ ·C ₄ H ₄ O ₄	C,H,N	174–176
7v	4v	C	66	C ₁₆ H ₂₂ N ₂ O ₂ ·C ₄ H ₄ O ₄	C,H,N	196–198
7w	4w ⁴¹	A	42	C ₁₄ H ₁₈ N ₂ O ₂ ·C ₄ H ₄ O ₄	H,N; C: calcd, 59.66; found, 59.20	138–140
7x	4x ⁴⁹	C	52	C ₁₃ H ₁₆ N ₂ O ₂ ·C ₄ H ₄ O ₄	C,H,N	219–221
7y	4y ⁵⁰	C	66	C ₁₄ H ₁₈ N ₂ O ₂ ·C ₄ H ₄ O ₄	C,H,N	151–154
7z	4z ⁵¹	C	63	C ₁₄ H ₁₈ N ₂ O ₂ ·C ₄ H ₄ O ₄	C,H,N	176–178
7aa	4aa ⁵²	A	71	C ₁₃ H ₁₆ N ₂ ·0.5C ₄ H ₄ O ₄	C,H,N	211–214
7bb	4bb	C	69	C ₁₄ H ₁₈ N ₂ ·C ₄ H ₄ O ₄	C,H,N	148–151
7cc	4cc	C	65	C ₁₅ H ₂₀ N ₂ ·1.5C ₄ H ₄ O ₄	C,H,N	192–194
7dd	4dd	C	67	C ₁₆ H ₂₂ N ₂ ·C ₄ H ₄ O ₄	C,H,N	188–190
7ee	4ee	C	62	C ₁₃ H ₁₃ F ₃ N ₂ ·C ₄ H ₄ O ₄	C,H,N	168–170
7ff	4ff ⁵²	C	59	C ₁₃ H ₁₆ N ₂ ·C ₄ H ₄ O ₄	C,H,N	170–173
7gg	4gg ¹²	C	62	C ₁₄ H ₁₈ N ₂ ·C ₄ H ₄ O ₄	C,H,N	192–194
7hh	4hh	C	64	C ₁₄ H ₁₆ N ₂ ·C ₄ H ₄ O ₄	H,N; C: calcd, 65.44; found, 64.80	189–191
7ii	4ii	C	57	C ₁₅ H ₁₈ N ₂ ·0.5C ₄ H ₄ O ₄	C,H,N	206–209
7jj	4jj	C	62	C ₁₄ H ₁₈ N ₂ ·0.6C ₄ H ₄ O ₄	C,H,N	209–212
7kk	4kk	C	56	C ₁₄ H ₁₈ N ₂ O·C ₄ H ₄ O ₄	C,H,N	178–181

^a Literature reference.

(700 mL) while maintaining a reaction temperature of less than 30 °C. After stirring for 24 h, the mixture was cooled to 0 °C and hydrolyzed by adding successively H₂O (11 mL), 2 N NaOH (11 mL), and H₂O (25 mL). The suspension was filtered and the filtrate concentrated under reduced pressure. The residue was dissolved in EtOAc (400 mL) and washed with water and 1 N HCl. The combined acidic aqueous phases were rendered alkaline with 35% aqueous NaOH and extracted with EtOAc. The organic layer was dried (MgSO₄) and concentrated yielding the amine **11a** as an oil (26 g, 68%). ¹H NMR (CDCl₃): δ 7.50–6.80 (m, 9H), 3.70 (AB system, 2H), 3.00–2.50 (m, 6H), 2.10–1.60 (m, 2H), 1.60–1.30 (2m, 2H).

Step d: 2-Amino-2-(aminomethyl)-1,2,3,4-tetrahydronaphthalene (6a). A vigorously stirred suspension of **11a** (7.98 g, 30 mmol), ammonium formate (7.88 g, 125 mmol), and 10% Pd/C (6 g) in CH₃OH (250 mL) was heated under reflux for 30 min. After cooling, filtration of the catalyst, and evaporation of the solvent, the crude residue was purified by chromatography (SiO₂, CH₂Cl₂/CH₃OH/NH₄OH = 90/10/1) to give the diamine **6a** as an oil (4.65 g, 88%). ¹H NMR (CDCl₃): δ 7.15–6.80 (m, 4H), 3.00–2.50 (m, 6H), 1.70 (m, 2H), 1.60–1.30 (2m, 4H).

Step e: Spiro[(1,3-diazacyclopent-1-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)], Benzenesulfonate (7a). A mixture of diamine **6a** (3.52 g, 20 mmol) and formamidinium acetate (2.08 g, 20 mmol) in EtOH (60 mL) was stirred at 20 °C under an N₂ atmosphere for 12 h. The solvent was evaporated and the residue taken up with 1 N HCl (50 mL); the acidic phase was washed with EtOAc and rendered alkaline with 35% aqueous NaOH. The mixture was extracted with EtOAc, and the combined organic layers were washed with saturated aqueous NaCl and concentrated under vacuum. The solid residue was dissolved in EtOH (50 mL) and treated with 1 equiv of benzenesulfonic acid dissolved in EtOH (20 mL). After evaporation of the solvent and crystallization from an EtOH/Et₂O mixture, the spiranic compound **7a** was isolated as a colorless solid (5.15 g, 74%), mp 124–125 °C. ¹H NMR (DMSO-*d*₆): δ 10.60 (br d, 1H), 10.20 (br d, 1H), 8.50 (s, br d, 1H),

7.60 (m, 2H), 7.30 (m, 3H), 7.10 (m, 4H), 3.65 (AB system, 2H), 3.00 (s, 2H), 2.90 (m, 2H). Anal. (C₁₂H₁₄N₂C₆H₆O₃S) C, H, N, S.

Method D. Spiro[(1,3-diazacyclopent-1-ene)-5,2'-(8'-hydroxy-1',2',3',4'-tetrahydronaphthalene)], Fumarate (7n). A 1 M solution of BBr₃ in CH₂Cl₂ (16.5 mmol) was added dropwise to a cold (–78 °C) solution of **7m** (0.97 g, 4.5 mmol) (method A, step c) in CH₂Cl₂ (30 mL). The temperature of the reaction mixture was brought to 20 °C, and the mixture was then poured into a chilled NaHCO₃ solution. After evaporation of the aqueous phase, the residue was taken up in iPrOH. The resulting suspension was filtered and the filtrate concentrated under vacuum to leave an oily residue which was purified by chromatography (SiO₂, H₂O/dioxane/NH₄OH = 90/10/1 as eluent). The isolated solid was dissolved in EtOH (10 mL) and treated with a solution of fumaric acid (0.46 g, 4 mmol) in EtOH (10 mL). After evaporation and recrystallization of the residue from EtOH, the phenol **7n** was obtained as a white solid (0.56 g, 48%), mp 240–242 °C. ¹H NMR (DMSO-*d*₆): δ 7.70 (s, 1H), 6.90 (t, 1H), 6.65 (d, 1H), 6.55 (d, 1H), 6.40 (s, 1H), 4.10–3.00 (br d, 3H), 3.40 (AB system, 2H), 2.80 (m, 2H), 2.70 (s, 2H), 1.80 (m, 2H). Anal. (C₁₂H₁₄N₂O·C₄H₄O₄) C, H, N.

Following the typical procedures described above (methods A–C), the spiro-imidazolines listed in Table 5 were prepared, starting from the appropriate tetralones.

Spiro[(1-methyl-1,3-diazacyclopent-2-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)], Hydrochloride (25). **Step a: 2-(Methylamino)-2-cyano-1,2,3,4-tetrahydronaphthalene (23).** A solution of 2-tetralone (**4a**) (1.46 g, 10 mmol) in MeOH (20 mL) was stirred for 18 h at room temperature in the presence of KCN (66 mg, 10.2 mmol), methylamine hydrochloride (776.5 mg, 11.5 mmol), and AcOH (1 mL). The solvents were evaporated under reduced pressure, and the residue was taken up in 1 N HCl (10 mL) and ether (100 mL). The aqueous phase was separated, and the organic phase was washed with H₂O (10 mL). The combined aqueous phases were rendered alkaline with 1 N NaOH. Extraction with CH₂

Cl₂ (3 × 20 mL), drying over K₂CO₃, and evaporation under reduced pressure gave the desired compound as an oil (1.49 g, 80% yield). ¹H NMR (CDCl₃): δ 7.20–7.00 (m, 4H), 3.24 (d, 1H), 3.05–2.80 (m, 3H), 2.60 (s, 3H), 2.20 (m, 1H), 2.00 (m, 1H).

Step b: 2-(Methylamino)-2-(aminomethyl)-1,2,3,4-tetrahydronaphthalene (24). A solution of **23** (7.4 g, 39.2 mmol) in anhydrous ether (50 mL) was added dropwise to an ice-cooled suspension of LiAlH₄ (4.5 g, 64.2 mmol) in anhydrous ether (500 mL). After addition, the suspension was heated to reflux for 1 h and then cooled (0 °C) and the excess hydride destroyed by careful addition of H₂O (4.5 mL), 2 N NaOH (4.5 mL), and H₂O (9 mL). The solid was filtered and washed extensively with ether (4 × 100 mL), and the combined ether solutions were evaporated under reduced pressure to provide the black oil **24** (5.5 g, 74% yield). ¹H NMR (CDCl₃): δ 7.10 (m, 4H), 3.00–2.40 (m, 6H), 2.30 (s, 3H), 1.60–2.00 (m, 2H).

Step c: Spiro[(1-methyl-1,3-diazacyclopent-2-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)], Hydrochloride (25). A solution of **24** (2 g, 10.5 mmol) and formamide acetate (1.2 g, 11 mmol) in EtOH (50 mL) was stirred for 20 h at room temperature. The EtOH was evaporated off, and the residue was taken up in a mixture of 1 N HCl (15 mL) and ether (100 mL). The aqueous phase was separated; the organic phase was washed with H₂O (10 mL). Combined aqueous phases were made basic with 1 N NaOH. Extraction with CH₂Cl₂ (3 × 20 mL), drying over K₂CO₃, and evaporation under reduced pressure gave a complex mixture, separated by chromatography on silica gel using a gradient of EtOH/NH₄OH (9/1) in CH₂-Cl₂. The desired compound **25** was eluted as an oil by a mixture of CH₂Cl₂/EtOH/NH₄OH (90/9/1). The oil was dissolved in EtOH, and a solution of HCl in ether was added; by dilution with ether, a solid precipitated which was filtered, crystallized from acetonitrile, and dried under vacuum at 50 °C (300 mg, 12% yield), mp 197 °C. ¹H NMR (CDCl₃): δ 10.55 (m, 1H), 8.65 (br s, 1H), 7.15 (m, 4H), 3.85 (d, 1H), 3.50 (d, 1H), 3.25 (d, 1H), 3.15 (s, 3H), 3.00 (m, 2H), 2.85 (dd, 1H), 2.20 (m, 1H), 1.95 (m, 1H). Anal. (C₁₃H₁₆N₂HCl) C, H, N, Cl.

Spiro[(1,2-dimethyl-1,3-diazacyclopent-2-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)], Fumarate (26). A solution of **24** (2 g, 10.5 mmol) and acetamide hydrochloride (1.04 g, 11 mmol) in EtOH (50 mL) was stirred for 20 h at room temperature. The EtOH was evaporated off, and the residue was taken up in a mixture of 1 N HCl (15 mL) and EtOAc (100 mL). The aqueous phase was separated off and washed with more EtOAc (50 mL) and CH₂Cl₂ (30 mL). The aqueous solution was made basic with 1 N NaOH and extracted with CH₂Cl₂ (3 × 20 mL). The combined organic phase was dried over K₂CO₃ and evaporated under reduced pressure. The residual oil was dissolved in EtOH, and 1 equiv of fumaric acid was added; by dilution with ether, a solid precipitated which was filtered and dried under vacuum (800 mg, 23% yield), mp 168 °C. ¹H NMR (CDCl₃): δ 7.15 (m, 4H), 6.45 (s, 2H), 3.55 (dd, 2H), 3.20 (d, 1H), 3.00 (s, 3H), 3.00–2.70 (2dd, d, 3H), 2.25 (s, 3H), 2.15 (m, 1H), 1.90 (m, 1H). Anal. (C₁₄H₁₈N₂C₄H₄O₄) C, H, N.

Spiro[(2-methyl-1,3-diazacyclopent-1-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)], Fumarate (27). A solution of **6a** (1.65 g, 9.36 mmol) and acetamide hydrochloride (0.898 g, 9.5 mmol) in EtOH (50 mL) was stirred for 20 h at room temperature. The EtOH was evaporated off, the residue was taken up in a mixture of 1 N HCl (15 mL) and ether (100 mL), and the aqueous phase was separated and washed with more ether (50 mL). The aqueous phase was rendered basic with 1 N NaOH and extracted with CH₂Cl₂ (3 × 20 mL) and the organic phase dried over K₂CO₃ and evaporated to dryness. The residual solid was crystallized from acetonitrile and dissolved in EtOH and 1 equiv of fumaric acid added; by dilution with diisopropyl ether, a solid precipitated which was filtered and dried under vacuum (1.5 g, 51% yield), mp 137 °C. ¹H NMR (CDCl₃): δ 7.15 (m, 4H), 6.40 (s, 2H), 3.60 (dd, 2H), 3.00 (s, 2H), 2.40 (m, 2H), 2.15 (s, 3H), 2.00 (m, 2H), 1.90 (m, 1H). Anal. (C₁₃H₁₆N₂C₄H₄O₄) C, H, N.

Spiro[(3-methyl-1,3-diazacyclopent-1-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)], Fumarate (28). A solution of **7a** (700 mg, 3.76 mmol) and triethylamine (522 μ L, 3.76 mmol)

in CH₂Cl₂ (40 mL) was cooled to –78 °C, and dimethyl sulfate (357 μ L, 3.76 mmol) was added dropwise. The reaction mixture was stirred and allowed to warm to room temperature. The solution was washed with 0.1 N NaOH (50 mL), dried over K₂CO₃, and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel by elution with a mixture of CH₂Cl₂/EtOH/NH₄OH (95/4.5/0.5). The first fractions contained the desired compound **28** (150 mg, 20% yield) followed by the 1-methyl isomer **25** (75 mg, 10% yield) and starting material **7a** (350 mg, 50% yield). The compound **28** was dissolved in MeOH, and 1 equiv of fumaric acid was added; by dilution with ether, a solid precipitated which was filtered and dried under vacuum, mp 197 °C. ¹H NMR (CDCl₃): δ 10.55 (m, 1H), 8.05 (s, 1H), 7.1 (m, 4H), 6.45 (s, 2H), 3.55 (d, 1H), 3.38 (d, 1H), 3.05 (s, 3H), 3.10–2.65 (m, 4H), 1.95 (m, 2H). Anal. (C₁₃H₁₆N₂C₄H₄O₄) C, H, N.

Spiro[(1-oxa-3-azacyclopent-2-ene)-4,2'-(1',2',3',4'-tetrahydronaphthalene)] (32). **Step a: 2-Amino-2-(aminocarbonyl)-1,2,3,4-tetrahydronaphthalene (29a).** A solution of amino nitrile **5a** (prepared as described for **7b**, method A, step a) (4.30 g, 25 mmol) in HCO₂H (80 mL) was cooled to 0 °C and saturated with anhydrous HCl. After evolution of gas had ceased, the solvent was evaporated and the residue taken up in acetone (50 mL). Filtration of the white crystalline solid afforded compound **29a** (4.04 g, 85%), mp 185–187 °C. ¹H NMR (DMSO-*d*₆): δ 8.70–7.70 (m, 5H), 7.25–7.05 (m, 4H), 3.20 (AB system, 2H), 2.85 (m, 2H), 2.50–2.20 (m, 1H), 2.20–2.00 (m, 1H).

Step b: 2-Amino-1,2,3,4-tetrahydro-2-naphthalenecarboxylic Acid (30a). A suspension of **29a** (4 g, 21 mmol) in 6 N aqueous HCl (25 mL) was refluxed until homogeneous. The solvent was evaporated off, and the residue was taken up with iPrOH and reconcentrated. The solid residue was dissolved in water and the mixture neutralized by addition of 1 N NaOH. The desired acid **30a** was obtained after filtration and drying (3.85 g, 96%), mp 205–206 °C. ¹H NMR (DMSO-*d*₆): δ 7.20–6.90 (m, 4H), 3.00 (AB system, 2H), 2.80–2.60 (m, 2H), 2.20–2.00 (m, 2H), 2.00–1.70 (m, 1H).

Step c: 2-Amino-2-(hydroxymethyl)-1,2,3,4-tetrahydronaphthalene (31a). A suspension of LiAlH₄ (1.44 g, 38 mmol) in anhydrous THF (100 mL) was added dropwise under an N₂ atmosphere and at room temperature to a solution of acid **30a** (3.63 g, 19 mmol) in anhydrous THF (50 mL). The mixture was brought to reflux for 1 h, and after cooling to 0 °C, H₂O (1.5 mL), 2.5 N NaOH (1.5 mL), and H₂O (3 mL) were added successively. The resulting suspension was filtered and washed with THF. The combined filtrates were concentrated under vacuum to yield the amino alcohol **31a** as an oil (2.53 g, 75%). ¹H NMR (CDCl₃): δ 7.15–6.90 (m, 5H), 3.45 (s, 2H), 3.00–2.50 (br d, 2H), 2.90 (m, 2H), 2.75 (AB system, 2H), 1.75 (m, 2H).

Step d: Spiro[(1-oxa-3-azacyclopent-2-ene)-4,2'-(1',2',3',4'-tetrahydronaphthalene)] (32). A solution of **31a** (0.62 g, 3.5 mmol) in methyl formate (5 mL) is brought to reflux for 1 h. The excess formate is then evaporated and the residue dissolved in CH₂Cl₂ (40 mL). To this new solution, diethylaminosulfur trifluoride (DAST) (1.21 g, 7.5 mmol) in CH₂Cl₂ (3 mL) was added dropwise, at –10 °C, under an N₂ atmosphere. After stirring for 2 h at –10 °C, 25% NH₄OH solution (25 mL) was added. The organic layer was separated, dried (MgSO₄), and concentrated. The solid residue was purified by chromatography (SiO₂, CH₂Cl₂/EtOH = 99/1 as eluent) and, after crystallization from hexane at –30 °C, afforded the spiro derivative **32a** as a colorless solid (0.25 g, 38%), mp 45–46 °C. ¹H NMR (CDCl₃): δ 7.20–7.00 (m, 4H), 6.85 (s, 1H), 4.00 (AB system, 2H), 3.05 (m, 1H), 2.95 (AB system, 2H), 2.10 (m, 1H), 1.85 (m, 1H). Anal. (C₁₂H₁₃NO) C, N; H: calcd, 7.00; found, 7.42.

Spiro[(2-amino-1,3-diazacyclopent-1-ene)-5,2'-(1',2',3',4'-tetrahydronaphthalene)], Hydrobromide (33). A solution of **6a** (1 g, 5.7 mmol) in CH₂Cl₂ (50 mL) was cooled to 0 °C and a solution of cyanogen bromide (690 mg, 5.7 mmol) in CH₂-Cl₂ (5 mL) added in dropwise fashion. The reaction mixture was stirred overnight at room temperature. The white precipitate was collected by filtration and crystallized from 2-propanol (500 mg, 31% yield), mp 209 °C. ¹H NMR

(CDCl₃): δ 7.3–7.0 (m, 4H), 3.45 (dd, 2H), 2.95 (s, 2H), 2.90 (m, 2H), 1.95 (m, 2H). Anal. (C₁₂H₁₅N₃HBr) C, H, N, Br.

Spiro[(1-oxa-2-amino-3-azacyclopent-2-ene)-4,2'-(1',2',3',4'-tetrahydronaphthalene)], Hydrochloride (34a). A solution of BrCN (1.27 g, 12 mmol) in CH₂Cl₂ (5 mL) was added rapidly at 0 °C to a solution of amino alcohol **31a** (1.76, 10 mmol) in CH₂Cl₂ (20 mL). The mixture was stirred overnight at room temperature and the precipitated solid filtered off and washed with CH₂Cl₂. The filtrate was washed with KHCO₃ solution, dried, and evaporated under vacuum; the residue was purified by chromatography (SiO₂, CH₂Cl₂/CH₃OH/NH₄OH = 95/4.5/0.5 as eluent). The oil obtained was dissolved in Et₂O and treated with ethereal HCl. The precipitate was then filtered off and recrystallized from an iPrOH/Et₂O mixture to give **34a** as a white powder (1.55 g, 65%), mp 205–206 °C. ¹H NMR (DMSO-*d*₆): δ 11.00–8.50 (br d, 2H), 7.25–7.00 (m, 4H), 4.55 (AB system, 2H), 3.10 (s, 2H), 3.00–2.80 (m, 2H), 2.10–1.90 (m, 2H). Anal. (C₁₂H₁₄N₂O·HCl) C, H, Cl; N: calcd, 11.73; found, 10.96.

Spiro[(1-oxa-2-amino-3-azacyclopent-2-ene)-4,2'-(8'-chloro-1',2',3',4'-tetrahydronaphthalene)], Hydrochloride (34b). Following the procedure described for **34a** but starting from **5b**, the spiro derivative **34b** was obtained as a white powder, mp 210–214 °C. ¹H NMR (DMSO-*d*₆): δ 10.80–9.10 (br d, 3H), 7.35–7.05 (m, 3H), 4.60 (AB system, 2H), 3.10 (m, 2H), 3.10–2.70 (m, 2H), 2.20–1.90 (m, 2H). Anal. (C₁₂H₁₃ClN₂O·HCl) C, H, N, Cl.

Acknowledgment. The authors thank the expert technical assistance of Michele Hipeaux, Annie Detrift-Aillet, Pascal Vrillaud, Yvette Menant, Elisabeth Adnet, and Veronique Barou, Angela D. Morris for assistance in preparation of the manuscript, and Catherine Dacquet for performing the binding assays.

References

- Weiner, N. Drugs that Inhibit Adrenergic Nerves and Block Adrenergic Receptors. In *The Pharmacological basis of Therapeutics*, 7th ed.; Gilman, A. G., Goodman, L. S., Rall, T. W., Murad, F., Eds.; Macmillan Publishing Co.: New York, 1985; pp 181–214.
- Timmermans, P. B. M. W. M.; van Zwieten, P. A. α_2 Adrenoceptors: Classification, Localization, Mechanism, and Targets for Drugs. *J. Med. Chem.* **1982**, *25*, 1389–1401.
- Timmermans, P. B. M. W. M.; Chiu, A. T.; Thoolen, M. J. M. C. α -Adrenergic Receptors. In *Comprehensive Medicinal Chemistry*; Hansh, C., Ed.; Pergamon: Oxford, U.K., 1990; pp 133–185.
- Ruffolo, R. R.; Stadel, J. M.; Hieble, J. P. α -Adrenoceptors: Recent Developments. *Med. Res. Rev.* **1994**, *14*, 229–270.
- Flavahan, N. A.; Lindblad, L. E.; Verbeuren, T. J.; Sheperd, J. T.; Vanhoutte, P. M. Cooling and α_1 and α_2 -Adrenergic Responses in Cutaneous Veins: Role of Receptor reserve. *Am. J. Physiol.* **1985**, *249*, H950–H955.
- Tsai, T. H.; Beitman, R. E.; Gibson, J. P.; Larson, E. J.; Freime, H.; Fontaine, R. Acute, Subacute and Chronic Toxicity/Carcinogenicity of Lofexidine. *Arzneim.-Forsch.* **1982**, *32*, 955–962.
- De Jonge, A.; Van Meel, J. C. A.; Timmermans, P. B. M. W. M.; van Zwieten, P. A. A Lipophilic, selective α_1 -Adrenoceptor agonist: 2-(2-chloro-5-trifluoromethyl-phenylimino)imidazole (St 587). *Life Sci.* **1981**, *28*, 2009–2016.
- Schmitt, H.; Laubie, M. Adrenoceptors and Central Cardiovascular Regulation. In *Adrenoceptors and Catecholamine Action*; Kunos, G., Ed.; Wiley-Interscience: New York, 1983; Vol. 1, Part B, pp 219–248.
- Pullman, B.; Coubelis, J. L.; Couriere, P. H.; Gervois, J. P. Quantum Mechanical Studies of the Conformational Properties of Phenethylamines and Biochemical and Medicinal Interest. *J. Med. Chem.* **1972**, *15*, 17–23.
- Jackman, L. M.; Jen, T. ¹H and ¹³C Nuclear Magnetic Resonance Studies on the Tautomerism, General Isomerism, and Conformation of some Cyclic Amidines, Guanidines, and related Systems. *J. Am. Chem. Soc.* **1975**, *97*, 2811–2818.
- Timmermans, P. B. M. W. M.; van Zwieten, P. A. Quantitative Structure-Activity Relationships in Centrally Acting Imidazolines Structurally Related to Clonidine. *J. Med. Chem.* **1977**, *20*, 1636–1401.
- DeMarinis, R. M.; Bryan, W. M.; Shah, D. H.; Hieble, J. P.; Pendelton, R. G. α Adrenergic Agents. 1. Direct-Acting α_1 Agonists Related to Methoxamine. *J. Med. Chem.* **1981**, *24*, 1432–1437.
- Wieland, T.; Müller, R.; Niemann, E.; Birkhofer, L.; Schöberl, A.; Wagner, A.; Söll, H. Methoden zur Herstellung und Umwandlung von Aminosäuren und Derivaten. (Amino Acids and Derivatives: Preparation and Transformation Methods.) In *Houben-Weyl Methoden der Organischen Chemie Band XI/2*; Müller, E., Ed.; Georg Thieme Verlag: Stuttgart, Germany, 1958; pp 305–307.
- Mowry, D. T. The Preparation of Nitriles. *Chem. Rev.* **1948**, *42*, 189–280.
- Matier, W. L.; Owens, D. A.; Comer, W. T.; Deitchman, D.; Ferguson, H. C.; Seidehamel, R. J.; Young, J. R. Antihypertensive Agents. Synthesis and Biological Properties of 2-Amino-4-aryl-2-imidazolines. *J. Med. Chem.* **1973**, *16*, 901–908.
- Ojima, I.; Inaba, S.; Nakatsugawa, K. A New Route to Amino-nitriles via Cyanosilylation of Schiff Bases and Oximes. *Chem. Lett.* **1975**, 331–334.
- Cordi, A. A.; Lacoste, J.-M.; Hennig, P. A Reinvestigation of the Intramolecular Buchner Reaction of 1-Diazo-4-phenylbutan-2-ones Leading to 2-Tetralones. *J. Chem. Soc., Perkin Trans. 1* **1993**, 3–4.
- Grimmett, M. R. Advances in Imidazole Chemistry. In *Advances in Heterocyclic Chemistry*; Katritzky, A. R., Boulton, A. J., Eds.; Academic Press Inc.: New York, 1980; Vol. 27, pp 292–296.
- Janssens, W. J.; Verbeuren, T. J.; Vanhoutte P. M. Effect of Moderate Cooling on Adrenergic Neuroeffector Interaction in Canine Cutaneous Veins. *Blood Vessels* **1981**, *18*, 281–295.
- Flavahan, N. A.; Vanhoutte, P. M. Heterogeneity of alpha-Adrenergic Responsiveness in Vascular Smooth Muscle: Role of Receptor Subtypes and Receptor Reserve. In *The alpha-1 Adrenergic Receptors*; Ruffolo, R. R., Ed.; Humana Press: Clifton, NJ, 1987; pp 351–403.
- Mutschler, E. *Arzneimittelwirkungen*, 5th ed.; Wissenschaftliche Verlagsgesellschaft: Stuttgart, Germany, 1986; pp 254–268.
- Cavero, I.; Roach A. G. The Pharmacology of Prazosin, a Novel Antihypertensive Agent. *Life Sci.* **1980**, *27*, 1525–1540.
- Goldberg, M. R.; Robertson, D. Yohimbine: a Pharmacological Probe for the Study of the α_2 -Adrenoceptor. *Pharmacol. Rev.* **1983**, *35*, 143–180.
- Nozulak, J.; Vigouret, J. M.; Jaton, A. L.; Hofmann, A.; Dravid, A. R.; Weber, H. P.; Kalkman, H. O.; Walkinshaw, M. D. Centrally Acting α_1 -Adrenoceptor Agonist Based on Hexahydro-naph[2,3-b]-1,4-oxazines and Octahydrobenzof[quinolines. *J. Med. Chem.* **1992**, *35*, 480–489.
- DeMarinis, R. M.; Shah, D. H.; Hieble, J. P.; Pendelton, R. G. α -Adrenergic Agents. 2. Synthesis and α_1 Agonist Activity of 2-Aminotetralins. *J. Med. Chem.* **1982**, *25*, 136–141.
- Ruffolo, R. R. Drug Receptor Theory. *Drugs News Perspect.* **1993**, *6*, 12–23.
- Ariens, E. J. Receptors: from Fiction to Fact. *Trends Pharmacol. Sci.* **1979**, *1*, 11–15.
- Ariens, E. J.; Rodrigues De Miranda, J. F. The Receptor Concept: Recent Experimental and Theoretical Developments. In *Recent Advances in Receptor Chemistry*; Gualtieri, F., Giannelli, M., Melchiorre, C., Eds.; Elsevier/North-Holland Biomedical Press: Amsterdam, The Netherlands, 1979; pp 1–36.
- Glossmann, H.; Hornung, R.; Presek, P. The Use of Ligand Binding for the Characterisation of Alpha Adrenoceptors. *J. Cardiovasc. Pharmacol.* **1980**, *2* (Suppl. 3), S303–S324.
- Lorenz, W.; Lomasney, J. W.; Collins, S.; Regan, J. W.; Caron, M. G.; Leffkowitz, R. J. Expression of Three alpha-2-Adrenoceptor Subtypes in Rat Tissues: Implications for alpha-2 Receptor Classification. *Mol. Pharmacol.* **1990**, *38*, 599–603.
- Renouard, A.; Widdowson, P. S.; Millan, M. J. Multiple Alpha-2 Receptor Subtypes. I. Comparison of [³H]RX821002-Labeled Rat R_{Alpha-2A} Adrenergic Receptors in Cerebral Cortex to Human H_{Alpha-2A} Adrenergic Receptor and Other Populations of Alpha-2 Adrenergic Subtypes. *J. Pharmacol. Exp. Ther.* **1994**, *270*, 946–957.
- Cheng, Y. C.; Prussoff, W. H. Relationship Between the Inhibition Constant (K_i) and the Concentration of Inhibitor which Causes 50 Percent Inhibition (IC₅₀) of an Enzyme Reaction. *Biochem. Pharmacol.* **1973**, *22*, 3099–3108.
- Imai, S.; Sano, K. Preparation of 2-Alkoxy-5-methoxy-benzaldehydes. Ger. Pat. Appl. DE 2 211 436, Sept. 28, 1972.
- Johnson, F.; Kim, K. S. 3-(2',5'-Dialkoxy-4'-substituted phenyl) Pthalides and Anthracene or Naphthalene Compounds Obtained from Them. Eur. Pat. Appl. EP 7 400, Feb. 6, 1980.
- Screttas, C. G.; Micha-Screttas, M. Hydrolythiation of α -Olefins by a Regiospecific Two-Step Process. Transformation of Alkyl Phenyl Sulfides to Alkylolithium Reagents. *J. Org. Chem.* **1978**, *43*, 1064–1071.
- Cannon, J. G.; Walker, K. A.; Mentanari, A.; Lang, J. P.; Flynn, J. R. Monoethyl Ether Derivatives of 7,8-Dihydroxy- and 8,9-Dihydro-4-n-propyl-1,2,3,4,4a,5,6,10b-octahydrobenzof[quinolines as Possible Products of Metabolism by Catechol-O-Methyltransferase. *J. Med. Chem.* **1990**, *33*, 2000–2006.
- Bartmann, W.; Konz, E.; Rüger, W. Synthesis of Di- and Tetraalkyl-3-piperazineisoquinolines and Related Compounds as Potential Antidepressant Agents. *J. Heterocycl. Chem.* **1987**, *24*, 677–682.

- (38) Mumakata, H.; Keiko, Y.; Iyata, K.; Kashima, T.; Uhara, H. Cyclic Acylation of 3-Phenylpropanoic and 4-Phenylbutanoic Acids in Liquid Hydrogen Fluoride. Kinetics, Mechanism and Substituent effects. *Recl. Trav. Chim. Pays-Bas.* **1974**, *93*, 189-194.
- (39) Hoefnagel, A. J.; Monshouwer, J. C.; Snorn, E. C. G.; Wepster, B. M. Dissociation Constants of β -Arylpropionic Acids, β -Aryl-isovaleric Acids, N-Arylglycines, Aryloxyacetic Acids, N-Aryl- β -Alanines and some Related Systems. *J. Am. Chem. Soc.* **1973**, *95*, 5350-5356.
- (40) Futaki, R. Tracer Studies on the Mechanism of Phenol-Phenol rearrangement. *Tetrahedron Lett.* **1986**, *59*, 5245-5248.
- (41) Munavalli, S.; Anaut Bhat, G.; Viel, C. Synthesis and Properties of New 6,7-Dimethoxy-3,4-dihydro-or-1,2,3,4-Tetrahydroisoquinolines. III. Isomers Chloromethoxybenzylated and Phenethylated at Position 1. *Bull. Soc. Chim. Fr.* **1966**, 3311-3318.
- (42) Soffer, M. D.; Cavagnol, J. C.; Gellerson, H. E. Syntheses in the Direction of Morphine I. 7-Methoxy- and 7,8-Dimethoxy-2-tetralone. *J. Am. Chem. Soc.* **1949**, *71*, 3857.
- (43) Ashwarden, W.; Imhof, R.; Jakob, R.; Kyburz, E. New Benzazecine-diones and Benzoquinolines used in Control of Cerebral Insufficiency and Improvement of Cognitive Functions. U.K. Pat. Appl. GB 2 185 744A, July 29, 1987.
- (44) Nixon, J. A.; Ploch, R. P.; Shaus, J. M.; Titus, R. D. New Ring-Substituted 2-Amino-1,2,3,4-Tetrahydronaphthalenes as Selective Inhibitors of Serotonin Reuptake. Eur. Pat. Appl. EP 343 830, May 16, 1989.
- (45) Rosowsky, A.; Battaglia, I.; Chen, K. K. N.; Modest, E. J. Synthesis of New Chlorine-Substituted Derivatives of 2-Tetralone. *J. Org. Chem.* **1968**, *33*, 4288-4290.
- (46) Adcock, W.; Bettess, P. D.; Rizvi, S. Q. A. Fluorine-19 Substituent Chemical Shifts. *Aust. J. Chem.* **1970**, *23*, 1921-1937.
- (47) Johnson, D. W.; Mander, L. N. Studies on Intramolecular Alkylation VI: Ortho-Alkylation in Phenolic Diazoketones: the Preparation of Intermediates Containing the Cyclohexa-2,4-dienone Moiety Suitable for Gibberellin Synthesis. *Aust. J. Chem.* **1974**, *27*, 1277-1286.
- (48) Cornforth, J. W.; Cornforth, R. H.; Robinson, R. The Preparation of β -Tetralones from β -Naphthol and some Analogous Transformations. *J. Chem. Soc.* **1942**, 689-691.
- (49) Horn, A. S.; Dijkstra, D.; Mulder, T. B. A.; Rollema, H.; Westerink, B. H. C. Synthesis and Activity of a Metabolite of the Dopaminergic Agonist 6,7-ADTN. *Eur. J. Med. Chem.* **1981**, *16*, 469-472.
- (50) Cameron, D. W.; Feutrill, G. I.; Pietersz, G. A. Coloring Matters of the Aphidoidea. XLV. Synthesis of (+,.)-Deoxyquinone A Dimethyl Ether and its Epimer (+,.)-cis-7,9-Dimethoxy-1,3-dimethyl-3,4,5,10-tetrahydro-1H-naphthof[2,3-c]pyran-5,10 dione. *Aust. J. Chem.* **1982**, *35*, 1481-1487.
- (51) Horn, A. S.; Grol, C. J.; Dijkstra, D.; Mulder, A. H. Facile Syntheses of Potent Dopaminergic Agonists and Their Effects on Neurotransmitter Release. *J. Med. Chem.* **1978**, *21*, 825-828.
- (52) Sims, J. S.; Cadogan, M.; Selman, L. H. Improvement of a β -Tetralone Synthesis: 5-,6-,7- and 8-Methyl- β -Tetralones. *Tetrahedron Lett.* **1971**, *14*, 951-954.

JM950205W