



Figure 5. Stereoview of the CoMFA contour map (standard deviation * coefficients) for analysis D. Areas where positively charged substituents are favored are colored blue; regions where negatively charged substituents should bind preferably are colored red. 2-Carboxy-1-naphthol (colored green) is shown as a reference.

JM970015G
S0022-2623(97)00015-0

1996, Volume 39

Cyrus J. Ohnmacht,* Keith Russell, James R. Empfield, Cathy A. Frank, Keith H. Gibson, Daniel R. Mayhugh, Frances M. McLaren, Howard S. Shapiro, Frederick J. Brown, Diane A. Trainor, Christopher Ceccarelli, Margaret M. Lin, Brian B. Masek, Janet M. Forst, Robert J. Harris, James M. Hulsizer, Joseph J. Lewis, Stuart M. Silverman, Reed W. Smith, Paul J. Warwick, Sen T. Kau, Alexa L. Chun, Thomas L. Grant, Burton B. Howe, Jack H. Li, Shephali Trivedi, Tracy J. Halterman, Christopher Yochim, Martin C. Dyroff, M.Kirkland, and Kathleen L. Neilson: *N*-Aryl-3,3,3-trifluoro-2-hydroxy-2-methylpropanamides: K_{ATP} Potassium Channel Openers. Modifications on the Western Region.

Page 4599. In the X-ray Diffraction paragraph, the following italic Roman letters should be converted to Greek characters: line 9, *q* should be θ ; line 14, *y* should be ψ ; lines 31–41, *h* should be η ; lines 35, 40, and 41, *s* should be σ .

JM9700373
S0022-2623(97)00037-X