

Additions and Corrections

1997, Volume 40

Takahiro Tsukida, Yasuyuki Hiramatsu,* Hideki Tsujishita, Takao Kiyoi, Masahiro Yoshida, Kiriko Kurokawa, Hideki Moriyama, Hiroshi Ohmoto, Yukihiisa Wada, Tadayuki Saito, and Hiro-sata Kondo*: Studies on Selection Blockers. 5. Design, Synthesis, and Biological Profile of Sialyl Lewis X Mimetics Based on Modified Serine–Glutamic Acid Dipeptides.

Page 3539. The second sentence of the first paragraph under Inhibition Assay of E-, P-, and L-selectin-sLe^x Binding should read as follows: The wells were washed twice with distilled water, blocked with 5% BSA (bovine serum albumin)–1 mM CaCl₂/50 mM imidazole buffer (pH 7.2) for 1 h at room temperature, and washed three times with 50 mM imidazole buffer (pH 7.2).

Page 3539. The second sentence of the second paragraph under Inhibition Assay of E-, P-, and L-selectin-sLe^x Binding should read as follows: The test compounds were dissolved in distilled water and/or DMSO at 10 mM and finally diluted by 1 mM CaCl₂/50 mM imidazole buffer (pH 7.2) to final concentrations at 20, 10, 5, 2.5, 1.25, 0.63, 0.31, and 0.16 μM, respectively.

JM970797D

Published on Web 01/15/1998