

# Journal of Medicinal Chemistry

© Copyright 2001 by the American Chemical Society

Volume 44, Number 20

September 27, 2001

## Letters

### Oxidation of *N*<sup>ω</sup>-Hydroxyarginine Analogues by NO-Synthase: The Simple, Non Amino Acid *N*-Butyl *N*-Hydroxyguanidine Is Almost as Efficient an NO Precursor as *N*<sup>ω</sup>-Hydroxyarginine

Sylvie Dijols,<sup>†</sup> Céline Perollier,<sup>†</sup>  
David Lefevre-Groboillot,<sup>†</sup> Stéphanie Pethe,<sup>†</sup>  
Roger Attias,<sup>†</sup> Jean-Luc Boucher,<sup>†</sup>  
Dennis J. Stuehr,<sup>‡</sup> and Daniel Mansuy\*<sup>†</sup>


UMR 8601, Université Paris V, 45 Rue des Saints-Peres,  
75270 Paris Cedex 06, France, and Department of  
Immunology, Lerner Research Foundation, Cleveland Clinic,  
9500 Euclid Avenue, Cleveland, Ohio 44195

Received June 19, 2001

**Introduction.** Nitric oxide (NO) is a key inter- and intracellular messenger molecule involved in the maintenance of vascular tone, neuronal signaling, and host response to infection.<sup>1,2</sup> The biosynthesis of NO is catalyzed by constitutively expressed neuronal and endothelial nitric oxide synthases (NOS I and NOS III, respectively) and by inducible NOS (NOS II) that is expressed in macrophages following induction by inflammatory mediators.<sup>2-5</sup> All three NOSs produce NO and L-citrulline from the oxidation of L-arginine (L-arg) by NADPH and O<sub>2</sub> with formation of *N*<sup>ω</sup>-hydroxy-L-arginine (NOHA) as an intermediate.<sup>6,7</sup>

Because of the great importance of either an excess of NO or a deficit of NO in many physiopathological situations,<sup>2-5</sup> many groups have worked during these last years to find selective NOS inhibitors and substrates. Whereas a great number of strong NOS inhibitors have been described,<sup>8</sup> only a few NOS substrates have been reported so far. Most of them are α-amino acids closely related to L-arg or NOHA, such as homo-

Scheme 1



L-arg and homo-NOHA,<sup>9,10</sup> or E-dehydro-L-arg,<sup>11</sup> as small changes of the L-arg or NOHA structure completely abolish the NOS-dependent formation of NO.<sup>9-14</sup> However, it has been recently reported<sup>15</sup> that some *N*-aryl *N*-hydroxyguanidines, such as *N*-*p*-chlorophenyl *N*-hydroxyguanidine, are oxidized by NOS II with concomitant formation of the corresponding *N*-aryl urea and NO in a 1:1 molar ratio.

To better understand the structural factors that are important for a substrate to be oxidized by NOS with formation of NO, and to find new selective and efficient NO donors after in situ oxidation by NOS, we have synthesized a series of NOHA analogues and studied their NOS-catalyzed oxidation. We here report that, although the removal of the α-NH<sub>2</sub> or the α-COOH groups of NOHA leads to a dramatic decrease of NO formation, removal of both groups leads to a simple non α-amino acid compound, *N*-butyl *N*-hydroxyguanidine, which acts as a substrate almost as efficient as NOHA itself for NO production. Our results also show that it is so far the best NOS substrate among a series of *N*-alkyl *N*-hydroxyguanidines.

**Chemistry.** A series of *N*-substituted *N*-hydroxyguanidines (Scheme 1) related to NOHA, **3a**, and its previously described analogues,<sup>10</sup> dinor-NOHA, **1a**, nor-NOHA, **2a**, and homo-NOHA, **4a**, have been synthesized, by using the general procedure previously reported for the preparation of *N*-hydroxyguanidines starting from the corresponding amines<sup>16</sup> (Scheme 2).

\* To whom correspondence should be addressed. Tel: 33 1 42862187. Fax: 33 1 42 86 83 87. E-mail: Daniel.Mansuy@biomedicale.univ-paris5.fr.

<sup>†</sup> Université Paris V.

<sup>‡</sup> Cleveland Clinic.



**Table 2.** Kinetic Constants for NO Formation from Oxidation of Some *N*-Hydroxyguanidines by NOS II<sup>a</sup>


compd	$K_m^b$	$k_{cat}^b$	$k_{cat}/K_m^b$
NOHA, <b>3a</b>	40 ± 10	480 ± 60	12
<b>3d</b>	55 ± 10	320 ± 50	5.8
<b>4a</b>	146 ± 20	410 ± 50	2.8
<b>4d</b> <sup>c</sup>	310 ± 50	280 ± 50	0.9
pClC <sub>6</sub> H <sub>4</sub> NOHG <sup>d</sup>	500 ± 50	100 ± 20	0.2

<sup>a</sup> Activities were measured as indicated in Table 1. <sup>b</sup>  $K_m$  in  $\mu\text{M}$ ,  $k_{cat}$  in  $\text{min}^{-1}$ ,  $k_{cat}/K_m$  in  $\mu\text{M}^{-1} \text{min}^{-1}$ . <sup>c</sup> Data for homo-NOHA from ref 10. <sup>d</sup> Data for *N*-*p*-chlorophenyl *N*-hydroxyguanidine from ref 15.

cavity close to the heme,<sup>21</sup> which is also found in the complex of NOS III with *N*-*p*-chlorophenyl *N*-hydroxyguanidine.<sup>22</sup> Interaction of the hydrophobic chain of substrates with the 338-valine residue of this cavity could play a crucial role in determining substrate affinity.<sup>21</sup> *N*-hydroxyguanidines bearing an  $\alpha$ -amino acid function at an appropriate position, NOHA and homo-NOHA, bind well to the  $\alpha$ -amino acid NOS binding site, whereas nor-NOHA and desamino- or descarboxy-NOHA exhibit structures that neither permit their efficient binding to the  $\alpha$ -amino acid binding site nor permit binding to the hydrophobic cavity. The alkyl group of *N*-alkyl *N*-hydroxyguanidines should prefer to bind to the small NOS hydrophobic cavity, provided that it is not too long (<6 carbons).


More detailed kinetic studies were performed on the NOS II-catalyzed oxidations of the most active compounds **3d** and **4d** by comparison with NOHA and homo-NOHA. Formations of NO (from the hemoglobin assay)<sup>17</sup> and of NO<sub>2</sub><sup>-</sup> (using the Griess assay),<sup>23</sup> the stable end product of NO under aerobic conditions, from oxidation of **3d** or **4d** exhibited characteristics very similar to those of NOS II-catalyzed oxidation of NOHA (data not shown): (1) they absolutely required the presence of NOS II containing BH<sub>4</sub>, NADPH, and O<sub>2</sub>, (2) they were strongly inhibited by classical NOS inhibitors such as *N*<sup>o</sup>-nitro-L-arginine and *S*-ethyl-*iso*-thiourea,<sup>8</sup> and (3) they were not significantly inhibited by superoxide dismutase and catalase, indicating that they were not due to oxidations by O<sub>2</sub><sup>•-</sup> or H<sub>2</sub>O<sub>2</sub> derived from the oxidase function of NOS.<sup>24</sup> NOS II-catalyzed oxidation of compounds **3d** and **4d** exhibited classical saturation kinetics and Lineweaver–Burk plots. Table 2 shows that the  $k_{cat}$  values calculated for **3d** and **4d** were only slightly lower than the one found for NOHA (320 and 280 instead of 480 min<sup>-1</sup>, respectively). However, whereas the  $K_m$  value observed for **3d** was only slightly greater than that found for NOHA, the  $K_m$  value calculated for **4d** was 8 times higher. Thus, the catalytic efficiency of NOS II-dependent oxidation of **3d** was only 2 times lower than that of NOHA ( $k_{cat}/K_m = 5.8$  instead of 12  $\mu\text{M}^{-1} \text{min}^{-1}$ ), whereas those calculated for the oxidations of **4d** and of the previously reported substrate *N*-*p*-chlorophenyl *N*-hydroxyguanidine<sup>15</sup> were very much lower ( $k_{cat}/K_m$  of 0.9 and 0.2  $\mu\text{M}^{-1} \text{min}^{-1}$ ).

Another parameter that is important for the evaluation of the quality of a substrate for NOS is the level of coupling between the electron transfer from NADPH and the transfer of an oxygen atom from O<sub>2</sub> to the substrate. This level is optimal in NOS II-catalyzed oxidation of NOHA, with consumption of 0.5 mol of NADPH per mol of NO produced, as expected for the three-electron oxidation of NOHA to citrulline and NO,



**Figure 1.** NADPH consumption for NO formation during oxidation of compound **3d** by NOS II as a function of **3d** concentration. Rates of NADPH consumption were measured by following the decrease in absorbance at 340 nm, under conditions identical to those described in Table 1, except for 0.2 mM NADPH. Values for the ratio moles of NADPH consumed/moles of NO formed are means ± SD from three experiments.

which requires the consumption of one electron and 1 mol of O<sub>2</sub><sup>6,10</sup> (eq 1). Figure 1 shows that NOS II-



catalyzed oxidation of **3d** to NO consumed 0.5 mol of NADPH per mole of NO produced at saturating **3d** concentrations. Interestingly, this level of 0.5 mol of NADPH consumed per mole of NO is already almost reached at 50  $\mu\text{M}$  **3d**. By comparison, formation of 1 mol of NO upon oxidation of *N*-*p*-chlorophenyl *N*-hydroxyguanidine consumes 2 mol of NADPH when using a substrate concentration of 500  $\mu\text{M}$  ( $K_m$  value, Table 2).<sup>15</sup>

**Conclusion.** Our results identify a new class of structurally simple, NOS-mediated NO donors that is based on non  $\alpha$ -amino acid *N*-alkyl, *N*-hydroxyguanidines. NOS II-catalyzed oxidations of NOHA and **3d** exhibit strikingly similar efficiencies, as shown by their very close  $k_{cat}/K_m$  values (12 and 5.8  $\mu\text{M}^{-1} \text{min}^{-1}$ , Table 2) and the lack of any decoupling between electron transfer and oxygen atom transfer (consumption of only 0.5 mol of NADPH) in both cases. These results open the way toward the research of new NO donors based on selective substrates of each class of NOS.

**Supporting Information Available:** Experimental section. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Kerwin, J. F.; Lancaster, J. R.; Feldman, P. L. Nitric oxide: A new paradigm for second messengers. *J. Med. Chem.* **1995**, *38*, 4343–4362.
- (2) Pfeiffer, S.; Mayer, B.; Hemmens, B. Nitric oxide. Chemical puzzles posed by a biological messenger. *Angew. Chem., Int. Ed.* **1999**, *38*, 1714–1731.
- (3) Knowles, R. G.; Moncada, S. Nitric Oxide Synthases in Mammals. *Biochem. J.* **1994**, *298*, 249–258.



- (4) Forstermann, U.; Closs, E. I.; Pollock, J. S.; Nakane, M.; Schwarz, P.; Gath, I.; Kleinert, H. Nitric oxide synthase isozymes – Characterization, purification, molecular cloning, and functions. *Hypertension* **1994**, *23*, 1121–1131.
- (5) Masters, B. S. S.; McMillan, K.; Sheta, E. A.; Nishimura, J. S.; Roman, L. J.; Martasek, P. Neuronal nitric oxide synthase, a modular enzyme formed by convergent evolution: Structure studies of a cysteine thiolate-ligated heme protein that hydroxylates L-arginine to produce NO as a cellular signal. *FASEB J.* **1996**, *10*, 552–558.
- (6) Stuehr, D. J.; Kwon, N. S.; Nathan, C. F.; Griffith, O. W.; Feldman, P. L.; Wiseman, J. N<sup>ω</sup>-Hydroxy-L-arginine is an intermediate in the biosynthesis of nitric oxide from L-arginine. *J. Biol. Chem.* **1991**, *266*, 6259–6263.
- (7) Klatt, P.; Schmidt, K.; Uray, G.; Mayer, B. Multiple Catalytic Functions of Brain Nitric Oxide Synthase – Biochemical Characterization, Cofactor-Requirement, and the Role of N<sup>G</sup>-Hydroxy-L-Arginine as an Intermediate. *J. Biol. Chem.* **1993**, *268*, 14781–14787.
- (8) Babu, B. R.; Griffith, O. W. Design of isoform-selective inhibitors of nitric oxide synthase. *Curr. Opin. Chem. Biol.* **1998**, *2*, 491–500.
- (9) Stuehr, D. J.; Griffith, O. W. Mammalian nitric oxide synthases. *Adv. Enzymol. Relat. Areas Mol. Biol.* **1992**, *65*, 287–346.
- (10) (a) Moali, C.; Boucher, J. L.; Sari, M. A.; Stuehr, D. J.; Mansuy, D. Substrate specificity of NO synthases: Detailed comparison of L-arginine, homo-L-arginine, their N-omega-hydroxy derivatives, and N-omega-hydroxy-nor-L-arginine. *Biochemistry* **1998**, *37*, 10453–10460. (b) Moali, C.; Brollo, M.; Custot, J.; Sari, M. A.; Boucher, J. L.; Stuehr, D. J.; Mansuy, D. Recognition of α-amino acids bearing various C=NOH functions by nitric oxide synthase and arginase involves very different structural determinants. *Biochemistry* **2000**, *39*, 8208–8218.
- (11) Lee, Y.; Marletta, M. A.; Martasek, P.; Roman, L. J.; Masters, B. S. S.; Silverman, R. B. Conformationally restricted arginine analogues as alternative substrates and inhibitors of nitric oxide synthases. *Bioorg. Med. Chem.* **1999**, *7*, 1097–1104.
- (12) Grant, S. K.; Green, B. G.; Stiffey-Wilusz, J.; Durette, P. L.; Shah, S. K.; Kozarich, J. W. Structural requirements for human inducible nitric oxide synthase substrates and substrate analogue inhibitors. *Biochemistry* **1998**, *37*, 4174–4180.
- (13) Feldman, P. L.; Chi, S.; Sennequier, N.; Stuehr, D. J. Synthesis of the L-arginine congener L-indsopicine and evaluation of its interaction with nitric oxide synthase. *Bioorg. Med. Chem. Lett.* **1996**, *6*, 111–114.
- (14) Vadon, S.; Custot, J.; Boucher, J. L.; Mansuy, D. Synthesis and effects on arginase and nitric oxide synthase of two novel analogues of N-omega-hydroxyarginine, N-omega-hydroxy-indsopicine and p-hydroxy amidinophenylalanine. *J. Chem. Soc., Perkin Trans. 1* **1996**, 645–648.
- (15) Renodon-Corniere, A.; Boucher, J. L.; Dijols, S.; Stuehr, D. J.; Mansuy, D. Efficient formation of nitric oxide from selective oxidation of N-aryl N'-hydroxyguanidines by inducible nitric oxide synthase. *Biochemistry* **1999**, *38*, 4663–4668.
- (16) Schantl, J. G.; Türk, W. 1-(4-Chlorophenyl)-3-hydroxyguanidine and O-acyl-derivatives. *Sci. Pharm.* **1989**, *57*, 375–380.
- (17) (a) Murphy, M. E.; Noack, E. Nitric oxide assay using hemoglobin method. *Methods Enzymol.* **1994**, *233*, 240–250. (b) Hevel, J. M.; Marletta, M. A. Nitric-oxide synthase assays. *Methods Enzymol.* **1994**, *233*, 250–258. (c) NOS I was prepared as described previously.<sup>10</sup> (d) For preparation of NOS II see: Wu, C.; Zhang, J.; Abu-Soud, H.; Ghosh, D. K.; Stuehr, D. J. High-level expression of mouse inducible nitric oxide synthase in *Escherichia coli* requires coexpression with calmodulin. *Biochem. Biophys. Res. Commun.* **1996**, *222*, 439–444. (e) For preparation of NOS III, see: Ghosh, S.; Gachhui, R.; Crooks, C.; Wu, C.; Lisanti, M. P.; Stuehr, D. J. Interaction between caveolin-1 and the reductase domain of endothelial nitric oxide synthase. *J. Biol. Chem.* **1998**, *273*, 22267–22271.
- (18) A synthesis of this compound was previously reported. Ishikawa, T.; Misonou, T.; Ikeno, M.; Sato, K.; Sakamaki, T. N<sup>ω</sup>-Hydroxy agmatine: A novel substance causing endothelium dependent vasorelaxation. *Biochem. Biophys. Res. Commun.* **1995**, *214*, 145–151. It was found to exhibit relaxing properties towards rat aorta rings.
- (19) Crane, B. R.; Arvai, A. S.; Ghosh, S.; Getzoff, E. D.; Stuehr, D. J.; Tainer, J. A. Structures of the N-omega-hydroxy-L-arginine complex of inducible nitric oxide synthase oxygenase dimer with active and inactive pterins. *Biochemistry* **2000**, *39*, 4608–4621.
- (20) Raman, C. S.; Li, H. Y.; Martasek, P.; Kral, V.; Masters, B. S. S.; Poulos, T. L. Crystal structure of constitutive endothelial nitric oxide synthase: A paradigm for pterin function involving a novel metal center. *Cells* **1998**, *95*, 939–950.
- (21) Li, H.; Raman, C. S.; Martasek, P.; Kral, V.; Masters, B. S. S.; Poulos, T. L. Mapping the active site polarity in structures of endothelial nitric oxide synthase heme domain complexed with isothioureas. *J. Inorg. Biochem.* **2000**, *81*, 133–139.
- (22) Raman, C. S.; Li, H.; Martasek, P.; Kral, V.; Masters, B. S. S.; Poulos, T. L. Structure of the catalytic heme domain of neuronal nitric oxide synthase. *Nitric Oxide Biol. Chem.* **2000**, *4*, 202.
- (23) Green, L. C.; Wagner, D. A.; Glogowski, J.; Skipper, P. L.; Wishnok, J. S.; Tannenbaum, S. R. Analysis of nitrate, nitrite, and [<sup>15</sup>N] nitrate in biological fluids. *Anal. Biochem.* **1982**, *126*, 131–138.
- (24) Moali, C.; Boucher, J. L.; Renodon-Corniere, A.; Stuehr, D. J.; Mansuy, D. Oxidations of N<sup>ω</sup>-hydroxyarginine analogues and various N-hydroxyguanidines by NO Synthase II: Key role of tetrahydrobiopterin in the reaction mechanism and substrate selectivity. *Chem. Res. Toxicol.* **2001**, *14*, 202–210.

JM0155446