

Articles

Structure–Activity Relationships at Human and Rat A_{2B} Adenosine Receptors of Xanthine Derivatives Substituted at the 1-, 3-, 7-, and 8-Positions

Soon-Ai Kim,[†] Melissa A. Marshall,[‡] Neli Melman,[†] Hak Sung Kim,[†] Christa E. Müller,[§] Joel Linden,[‡] and Kenneth A. Jacobson^{*†}

Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes & Digestive & Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892, Departments of Internal Medicine and Molecular Physiology & Biological Physics, University of Virginia, Box MR4 6012, Health Science Center, Charlottesville, Virginia 22908, and Pharmazeutisches Institut Pharmazeutische Chemie, Universität Bonn, Bonn, Germany

Received September 11, 2001


In the search for improved selective antagonist ligands of the A_{2B} adenosine receptor, which have the potential as antiasthmatic or antidiabetic drugs, we have synthesized and screened a variety of alkylxanthine derivatives substituted at the 1-, 3-, 7-, and 8-positions. Competition for ¹²⁵I-ABOPX (¹²⁵I-3-(4-amino-3-iodobenzyl)-8-(phenyl-4-oxyacetate)-1-propylxanthine) binding in membranes of stably transfected HEK-293 cells revealed uniformly higher affinity (<10-fold) of these xanthines for human than for rat A_{2B} adenosine receptors. Binding to rat brain membranes expressing A₁ and A_{2A} adenosine receptors revealed greater A_{2B} selectivity over A_{2A} than A₁ receptors. Substitution at the 1-position with 2-phenylethyl (or alkyl/olefinic groups) and at N-3 with hydrogen or methyl favored A_{2B} selectivity. Relative to enprofylline **2b**, pentoxifylline **35** was equipotent and 1-propylxanthine **3** was >13-fold more potent at human A_{2B} receptors. Most N-7 substituents did not enhance affinity over hydrogen, except for 7-(2-chloroethyl), which enhanced the affinity of theophylline by 6.5-fold to 800 nM. The A_{2B} receptor affinity-enhancing effects of 7-(2-chloroethyl) vs 7-methyl were comparable to the known enhancement produced by an 8-aryl substitution. Among 8-phenyl analogues, a larger alkyl group at the 1-position than at the 3-position favored affinity at the human A_{2B} receptor, as indicated by 1-allyl-3-methyl-8-phenylxanthine, with a K_i value of 37 nM. Substitution on the 8-phenyl ring indicated that an electron-rich ring was preferred for A_{2B} receptor binding. In conclusion, new leads for the design of xanthines substituted in the 1-, 3-, 7-, and 8-positions as A_{2B} receptor-selective antagonists have been identified.

Introduction

Four extracellular G protein-coupled receptors for adenosine have been identified as follows: A₁, A_{2A}, A_{2B}, and A₃.¹ A_{2B} receptors, which are coupled to stimulation of adenylyl cyclase^{2,3} and also lead to a rise in intracellular calcium,⁴ are involved in the control of vascular tone,⁵ hepatic glucose balance,⁶ cell growth and gene expression,⁷ mast cell degranulation,⁸ and intestinal water secretion.⁹ Activation of A_{2B} receptors in human retinal endothelial cells may lead to neovascularization by a mechanism involving increased vascular endothelial growth factor expression.⁷ Selective xanthine antagonists of the A_{2B} receptor have recently been reported.^{10,11} Such antagonists are potentially useful in the treatment of asthma,^{12,13} intestinal disorders,⁹ and diabetes (through improved glucose uptake in skeletal muscle and suppression of glucose production in hepatocytes).⁶

On the basis of adenosine receptor binding assays, we

have identified several new xanthines with improved potency and selectivity for human A_{2B} receptors.^{10,14} A *p*-cyanoanilide derivative in this series, *N*-(4-cyanophenyl)-2-[4-(2,3,6,7-tetrahydro-2,6-dioxo-1,3-dipropyl-1H-purin-8-yl)phenoxy]acetamide **1**, was shown to be 204-, 255-, and 289-fold selective for human A_{2B} receptors vs human A₁, A_{2A}, and A₃ receptors, although less selective vs rat A₁ and A_{2A} receptors. Compound **1** (100 nM) was shown to completely inhibit calcium mobilization stimulated by 1 μM NECA in HEK-293 (human embryonic kidney) cells expressing human A_{2B} receptors.¹⁰



Nonselective radioligands have been used to characterize binding to recombinant human A_{2B} receptors overexpressed in cell lines such as HEK-293. These include ¹²⁵I-ABOPX (¹²⁵I-3-(4-amino-3-iodobenzyl)-8-(phenyl-4-oxyacetate)-1-propylxanthine),¹⁵ [³H]8-cyclo-

* To whom correspondence should be addressed. Tel.: (301)496-9024. Fax: (301)480-8422. E-mail: kajacobs@helix.nih.gov.

[†] National Institutes of Health.

[‡] University of Virginia.

[§] Universität Bonn.

Table 1. Affinities of Xanthine Derivatives in a Radioligand Binding Assay at Human A_{2B} Receptors

compd	R ₁	R ₃	R ₇	R ₈	K _i ^a (nM)	compd	R ₁	R ₃	R ₇	R ₈	K _i ^a (nM)
2a (theophylline)	Me	Me	H	H	9070 ± 1490	31	CH ₂ OCH ₃	Me	Me	H	14 000 ± 2400
2b (enprofylline)	H	Pr	H	H	4730 ± 270	32	CH ₂ CN	Me	Me	H	21 600 ± 5200
3	Pr	H	H	H	360 ± 70	33	(CH ₂) ₂ OC-OCH ₃	Me	Me	H	8710 ± 1320
4	propargyl	H	H	H	552 ± 76	34	CH ₂ COCH ₃	Me	Me	H	20 800 ± 4800
5	allyl	H	H	H	461 ± 18 (4)	35	(CH ₂) ₄ CO-CH ₃	Me	Me	H	5180 ± 2010
6	n-Bu	H	H	H	421 ± 38 (4)	36	Me	propargyl	Me	H	9480 ± 1340
7	2-phenylethyl	H	H	H	408 ± 67 (4)	37	Pr	H	H	cyclopentyl	34 ± 10
8	cyclopentyl	H	H	H	965 ± 64 (4)	38	(CPX)	Pr	H	cyclopentyl	63.8 ± 8.3
9	Et	Et	H	H	1770 ± 260	39	(8-PT)	Me	H	C ₆ H ₅	415 ± 219
10	n-Pr	n-Pr	H	H	1110 ± 330 (4)	40	(SPT)	Me	H	C ₆ H ₄ -4-S-O ₃ H	1330 ± 220
11	allyl	allyl	H	H	1330 ± 240	41	Me	Me	H	C ₆ H ₄ -4-OH	60.7 ± 3.1
12	n-hexyl	n-hexyl	H	H	7580 ± 1900	42	Me	Me	H	C ₆ H ₄ -4-OCCH ₃	394 ± 38
13	Et	Me	H	H	1620 ± 440	43	Me	Me	H	C ₆ H ₄ -4-NO ₂	6940 ± 1960
14a	propargyl	Me	H	H	511 ± 52 (4)	44	Me	Me	H	C ₆ H ₄ -4-N-(CH ₃) ₂	289 ± 11
14b	benzyl	Me	H	H	10 200 ± 2000	45	Me	Me	H	C ₆ H ₄ -2-CO ₂ H	11 100 ± 4700
14c	2-phenylethyl	Me	H	H	646 ± 15	46	Me	Me	H	C ₆ H ₄ -3-CO ₂ H	5530 ± 640 (4)
14d	3-phenylpropyl	Me	H	H	2330 ± 250	47	Me	Me	H	C ₆ H ₃ -2,4-(NO ₂) ₂	15 400 ± 6600
15	Me	i-Pr	H	H	4890 ± 870	48	Et	Et	H	C ₆ H ₅	62.0 ± 11.4
16	n-Bu	Et	H	H	1720 ± 160	49	Pr	Pr	H	C ₆ H ₅	18.9 ± 3.2 (4)
17	i-amyl	i-Bu	H	H	6240 ± 1340	50	Pr	Pr	H	pyrazine	79.6 ± 11.3
18 (caffeine)	Me	Me	Me	H	10 400 ± 1800	51	allyl	Me	H	C ₆ H ₅	37 ± 6
19a	Me	Me	2-Cl-ethyl	H	1390 ± 200	52	i-amyl	i-Bu	H	C ₆ H ₅	9890 ± 1440
19b	Me	Me	3-Cl-propyl	H	5620 ± 980	53	Pr	Pr	H	C ₆ H ₃ -2,6-F ₂	310 ± 42
19c	Me	Me	4-Cl-butyl	H	15 000 ± 1100	54	Pr	Pr	H	C ₆ H ₃ -2,3-(OH) ₂	125 ± 10
20	Me	Me	propargyl	H	9460 ± 1880	55	Pr	Pr	H	C ₆ H ₃ -2,4-(OH) ₂	20.7 ± 2.0
21	Me	Me	allyl	H	9490 ± 1220	56	Pr	Pr	H	C ₆ H ₃ -2,5-(OH) ₂	138 ± 6.9
22	Me	Me	benzyl	H	19 400 ± 4140	57	Pr	Pr	H	C ₆ H ₃ -2-OH-4-OCCH ₃	46.5 ± 0.7
23	Me	Me	CH ₂ COOH	H	27 600 ± 11 400	58	Pr	Pr	H	C ₆ H ₃ -3,5-(OH) ₂ -4-OCCH ₃	59.5 ± 7.0
24	Me	Me	CH ₂ CN	H	14 500 ± 300	59	Pr	Pr	H	C ₆ H ₄ -4-NHC-OCH ₃	36.7 ± 9.5 (4)
25	Me	Me	(CH ₂) ₂ NH ₂	H	23 500 ± 6700	60	Pr	Pr	H	C ₆ H ₃ -2-OMe-4-Cl	6530 ± 570 (4)
26	Me	Me	(CH ₂) ₂ OH	H	17 300 ± 2900	61	Pr	Pr	H	C ₆ H ₄ -4-OCCH ₂ -CO ₂ H	40 ± 4
27	Me	Me	(CH ₂) ₂ OCO-CH ₃	H	23 200 ± 8300	62	Pr	Pr	H	C ₆ H ₃ -2-OH-4-OCCH ₂ -CO ₂ H	188 ± 22
28	allyl	allyl	Me	H	3390 ± 760	63	allyl	Me	Me	C ₆ H ₅	11 400 ± 1000
29	propargyl	Me	Me	H	4130 ± 1040	64	allyl	Me	Me	C ₆ H ₅	311 ± 22
30	CH ₂ COOH	Me	Me	H	23 500 ± 7600	65	Pr	Pr	2-Cl-ethyl 2-Cl-ethyl	C ₆ H ₄ C ₆ H ₄	691 ± 72

^a Displacement of specific [¹²⁵I]ABOPX binding at human A_{2B} receptors expressed in HEK-293 cells, in membranes, expressed as K_i ± SEM. n = 3, unless noted. Cases in which n = 4 are noted in parentheses.

Table 2. Synthetic Yields and Characterization of Xanthine Derivatives

compd	formula	yield (%)	analysis
14b	C ₁₃ H ₁₂ N ₄ O ₂	44	C, H, N
14c	C ₁₄ H ₁₄ N ₄ O ₂ ·0.2H ₂ O	32	C, H, N
14d	C ₁₅ H ₁₆ N ₄ O ₂	60	C, H, N
19b	C ₁₀ H ₁₃ ClN ₄ O ₂	88	C, H, N
19c	C ₁₁ H ₁₅ ClN ₄ O ₂	95	C, H, N
64	C ₁₇ H ₁₈ ClN ₄ O ₂	66	HRMS ^a
65	C ₁₉ H ₂₄ ClN ₄ O ₂	32	HRMS ^a

^a High-resolution mass in positive ion FAB mode determined to be within acceptable limits. Compound **64**: calcd, 345.1118; found, 345.1110. Compound **65**: calcd, 375.1588; found, 375.1573. HPLC demonstrated >95% purity, retention times (mobile phase, min): **64**, (A) 22.69; (B) 23.63; **65**, (A) 23.58; (B) 30.18.

of theophylline examined, i.e., **20–27**, including charged and heteroatom substitution, produced K_i values of 10–20 μ M. The notable exception to this pattern was 2-chloroethyltheophylline **19a** (K_i value of 1.4 μ M), which displayed a 7.5-fold greater affinity than caffeine. Homologues of **19a** were also prepared and studied in binding (Table 3). As illustrated in Figure 1, the potency order of 7-substituted compounds was 2-Cl-ethyl **19a** > 3-Cl-propyl **19b** > 4-Cl-butyl **19c**. A 7-methyl group slightly reduced the affinity of 1,3-diallylxanthine (**28** vs **11**). Among caffeine analogues in which the 1-substituent was modified, including charged and heteroatom substitution in **29–35**, some variation in affinity was observed (K_i values of 4–21 μ M), with the 1-propargyl analogue **29** as the most potent. The relatively




Figure 1. Competition by 7-substituted xanthines for ¹²⁵I-ABOPX binding to recombinant human A_{2B} receptors. The indicated antagonists (see Table 1) were incubated to equilibrium (2 h) in 100 μ L with radioligand (0.7 nM ¹²⁵I-ABOPX) and human HEK-A_{2B} cell membranes (25 μ g of membrane protein). Each point is the mean \pm SEM of triplicate determinations. Where not shown, the SEM is smaller than the symbol. K_i values derived from triplicate experiments are summarized in Table 1.

high affinity of pentoxifylline **35** ($K_i = 5.2 \mu$ M) is notable since this compound is used therapeutically to treat intermittent claudication. An isomer **36** was less potent than **29**.

For the 8-cyclopentyl analogues **37** and **38**, the absence of a 3-alkyl group slightly increased affinity. An 8-phenyl substituent, well-explored as a means of enhancing affinity at A₁ and A_{2A} receptors,^{19–21} generally enhanced affinity at the human A_{2B} receptor by

Table 3. Affinities of Xanthine Derivatives in Radioligand Binding Assays at Rat A₁ and A_{2A} Receptors and Selected Compounds at Other Subtypes^{a–d}

compd	K_i (nM)			K_i ratio	
	rA ₁ ^a	rA _{2A} ^b	rA _{2B} ^c	rA ₁ /rA _{2B}	rA _{2A} /rA _{2B}
1	16.8 \pm 3.6 ^f	612 \pm 287 ^f	12.8 \pm 1.1	1.3	48
2a	6920 \pm 160 ^f	6700 \pm 320 ^f	15 100 \pm 2700	0.46	0.44
3	3310 \pm 580	<10% (10 ⁻⁵)	1880 \pm 190	1.8	>10
4	12 700 \pm 5600	<10% (10 ⁻⁵)			
5^e	24 300 \pm 4300	<10% (10 ⁻⁵)			
6	8890 \pm 1090	<10% (10 ⁻⁵)			
7^e	37 600 \pm 8900	<10% (10 ⁻⁵)			
8	5570 \pm 760	<10% (10 ⁻⁴)			
13	24 700 \pm 8100	9780 \pm 220			
14a^e	5830 \pm 250	33 600 \pm 7700	2150 \pm 260	2.7	16
14b	1660 \pm 620	3190 \pm 1970	10 200 \pm 1500	0.16	0.31
14c^e	27 600 \pm 3900	<10% (10 ⁻⁴)	5530 \pm 610	5.0	>20
14d	24 400 \pm 4800	<10% (10 ⁻⁴)	15 300 \pm 2000	1.6	>10
16	2170 \pm 940	<10% (10 ⁻⁴)			
19a	15 200 \pm 1100	4230 \pm 1580	5390 \pm 780	2.8	0.78
19b	23 600 \pm 3900	<10% (10 ⁻⁴)	14 200 \pm 100	1.7	>10
19c	17 700 \pm 3700	33 300 \pm 4300	18 700 \pm 500	0.95	1.8
37	7.68 \pm 1.46				
38	1.0	500	186 \pm 5	0.005	3
51^e	302 \pm 49	1920 \pm 400	174 \pm 35	1.7	11
54	39.0 \pm 5.2	<10% (10 ⁻⁵)			
55	220 \pm 11	107 \pm 10			
56	7.46 \pm 0.58	17 300 \pm 6500			
57	97.7 \pm 2.7	5500 \pm 760			
58	30.3 \pm 3.5	1340 \pm 680			
59	39.2 \pm 10.8	<10% (10 ⁻⁵)			
62	757 \pm 109	2720 \pm 470			
63	<10% (10 ⁻⁵)	23 500 \pm 9800			
64^e	35 \pm 4% (10 ⁻⁵)	3570 \pm 1360	1700 \pm 90	>6	2.1
65^e	1660 \pm 130	16 700 \pm 4300	3360 \pm 530	0.49	5.0

^a Displacement of specific [³H]R-PIA binding in rat brain membranes in HEK-293 cells, expressed as $K_i \pm$ SEM or percent displacement at the indicated concentration ($n = 3–5$). ^b Displacement of specific [³H]CGS 21680 binding in rat striatal membranes, expressed as $K_i \pm$ SEM or percent displacement at the indicated concentration ($n = 3–5$). ^c Rat A_{2B} receptor expressed in HEK cells, using [¹²⁵I]I-ABOPX as radioligand. ^d Human A₃ receptor, $K_i \pm$ SEM (μ M), using [¹²⁵I]I-AB-MECA as radioligand: **3**, 2.37 \pm 0.08; **4**, 3.08 \pm 0.43; **5**, 16.1 \pm 2.9; **6**, 4.61 \pm 1.26; **7**, 7.51 \pm 1.64; **8**, 0.639 \pm 0.158; **14a**, 10.9; **14b**, 4.44 \pm 0.89; **14c**, 6.88 \pm 0.47; **14d**, 8.65 \pm 2.73. ^e **5**, MRS 1973; **7**, MRS 1975; **14a**, MRS 1980; **14c**, MRS 3005; **51**, MRS 1916; **64**, MRS 3000; **65**, MRS 1995. ^f Values from refs 10 and 14.




Figure 2. Correlation in the affinities of various xanthines for binding to human and rat A_{2B} adenosine receptors. The pK_i values of compounds that were evaluated for binding to rat and human receptors (Table 2) are plotted and fit by linear regression.

roughly 20-fold. 8-Phenyltheophylline **39** (K_i value of 415 nM) displayed a 22-fold greater human A_{2B} receptor affinity than theophylline **2a**. Substitution of the 8-phenyl ring indicated that an electron-rich ring was preferred in receptor binding. The rank order of potency of substitution at the 8-aryl ring in theophylline analogues **39–47** was 4-hydroxy > 4-dimethylamino > 4-methoxy > 4-sulfo > 3-carboxy and 4-nitro > 2-carboxy and 2,4-dinitro.

Homologation of 1,3-dialkyl groups in 8-phenylxanthines enhanced affinity, by 22-fold in the case of 1,3-dipropyl analogue **49** in comparison to **39**. As with 8-H xanthines, a 1-allyl substituent greatly increased affinity at the A_{2B} receptor in the 8-aryl series. Thus, 1-allyl-3-methyl-8-phenylxanthine **51** displayed a K_i value of 37 nM. Branched substitution, in **52** as before, greatly decreased affinity. Phenyl ring-substituted analogues of **49** were also tested: The electron-withdrawing 2,6-difluoro and 4-chloro groups on the 8-aromatic ring, in **53** and **60**, reduced affinity. Various di- and trisubstituted phenyl rings, containing electron-donating groups, displayed considerable affinity but not exceeding that of **49**. The functionalized congener XCC (xanthine carboxylic congener, 8-[4-[(carboxymethyl)oxy]phenyl]-1,3-dipropylxanthine, **61**), which served as the basis of A_{2B} receptor-selective amide derivatives such as **1**,¹⁰ was nearly as potent as **49**.

The 7-(2-chloroethyl) group of **65** decreased the affinity of compound **49** by 36-fold to a K_i value of 690 nM. However, in comparison to the 7-methyl modification of 8-phenylxanthines, this group increased affinity. Compound **64**, the 7-(2-chloroethyl) derivative of the 8-phenylxanthine derivative **63**, displayed a 37-fold greater affinity at human A_{2B} receptors.

Selected compounds were tested in binding experiments at rat A_{2B} receptors (Table 3). A close correlation between relative potencies at human and rat A_{2B} receptors was found (Figure 2). Affinities were usually higher by 2–9-fold at the human vs the rat A_{2B} receptor, with more pronounced differences observed for the more potent compounds. Compound **14c**, a 1-phenylethyl derivative, showed slight A_{2B} receptor selectivity in the rat. The reference compound **1** was still highly potent at the rat A_{2B} receptor, with a K_i value of 12.8 nM, although not selective for A_{2B} vs A_1 receptors in the rat. Several other compounds were selective for A_{2B} vs A_{2A} but not A_1 receptors in the rat: **3**, **14a**, **14d**, **19b**, and **51**. Selected compounds were also tested at human A_3 receptors (Table 3, footnote e), and most were found to

be approximately an order of magnitude more selective for human A_{2B} vs A_3 receptors, except compound **8**, which was equipotent at the two subtypes.

Discussion

Theophylline **2a** is widely used as an antiasthmatic drug, although its mechanism of action is uncertain. The related xanthine, enprofylline^{13,14} **2b**, which is also therapeutically efficacious in the treatment of asthma, was earlier thought to act through a nonadenosine receptor-mediated mechanism due to its low affinity at A_1 and A_{2A} receptors. However, the discovery that enprofylline has greater than anticipated affinity and slight selectivity at the A_{2B} subtype¹³ supports the hypothesis that A_{2B} receptor antagonism may contribute to antiasthmatic activity of xanthines.^{12,26,27} This hypothesis was strengthened by functional effects of A_{2B} receptor activation observed in mast cells of dog and human.^{8,12,28} Thus, potent and/or selective A_{2B} receptor antagonists may provide new therapeutic agents.

The SAR of xanthines in binding to A_{2B} adenosine receptors was investigated in the present study. Substitution at the 1-position by a group larger than at the 3-position (H or Me) favored affinity at the A_{2B} receptor. The affinity of xanthines in binding to the human A_{2B} receptor was enhanced by substitution with propyl, butyl, allyl, propargyl, or 2-phenylethyl groups at N-1, while hydrogen or methyl at N-3 favored affinity at this subtype.

A 7-(2-chloroethyl) substituent strikingly favored A_{2B} receptor selectivity. Most N-7 substituents did not enhance affinity over hydrogen, except for 2-chloroethyl, which enhanced the human A_{2B} receptor affinity of theophylline by 6.5-fold. Thus, 7-(2-chloroethyl)theophylline displayed a K_i value of 800 nM. Previously, 7-(2-chloroethyl)theophylline was noted to be slightly selective for adenosine A_{2A} vs A_1 receptors.²⁹ At the A_{2B} receptor, the effects of the 7-(2-chloroethyl) group were compatible with the enhancement produced by 8-aryl substitution. For example, the affinity of an 8-aryl-7-chloroethyl analogue **64** was intermediate between those of the corresponding 7-H and 7-methyl analogues. Compound **64** was roughly 35-fold more potent at the human A_{2B} receptor than the 7-methyl analogue **63** and 8-fold less potent than the 7-H analogue, **51**.

An 8-phenyl substituent, well-explored as a means of enhancing affinity at A_1 and A_{2A} receptors,^{19–24} enhanced affinity at A_{2B} receptors by roughly 20-fold. The detail in the present study concerning 8-aryl ring substitution and compatibility with other xanthine substituents greatly extended previous findings.^{10,14} As for simpler xanthines, even in the 8-phenyl series, a larger alkyl group at the 1-position than at the 3-position favored affinity at the A_{2B} receptor. This was illustrated by 1-allyl-3-methyl-8-phenylxanthine, **51**, which displayed a K_i value of 37 nM at A_{2B} receptors. Substitution of the 8-phenyl ring indicated that an electron-rich ring (e.g., 4-hydroxy substitution, **41**) was preferred in receptor binding. Consistent with this finding, nitro and other electron-withdrawing substituents greatly reduced affinity.

The initial screen of A_{2B} receptor affinity in the present study utilized the human subtype, and selected compounds were also evaluated at rat A_{2B} receptors. We

found uniformly higher affinity of substituted xanthines for human than for rat A_{2B} adenosine receptors, but these differences were consistently less than 10-fold. Compounds having various degrees of selectivity for rat A_{2B} receptors were **3–7** (1-substituted xanthines), **14a–d** (1-substituted theophyllines), **51** (1-allyl-3-methyl-8-phenylxanthine), and **64** (1-allyl-7-chloroethyl-3-methyl-8-phenylxanthine). The 1-phenylethyl group appeared to enhance A_{2B} receptor selectivity in the rat.

Selectivity for the human A_{2B} receptor has been easier to achieve than selectivity for the rat homologue, mainly due to the large species differences at A_1 and A_{2A} receptors.¹⁰ To further define the selectivity of A_{2B} receptor antagonists, additional characterization at receptor homologues of various species will be needed.

In conclusion, we have identified new leads for the design of xanthine derivatives as selective antagonists of the A_{2B} receptor, including asymmetric substitution at the 1- and 3-positions, 7-haloalkyl groups, and 8-aryl ring substitution patterns. Furthermore, the affinities at rat and human A_{2B} receptors have been compared. Optimal combinations of substituents on the 1-, 3-, 7-, and 8-positions of xanthines may lead to more potent and selective antagonists. Such selective compounds will aid in the elucidation of the physiological role of this receptor and possibly lead to therapeutically useful agents for treating asthma, diabetes, and other diseases.^{6,10,30,31}

Experimental Section

Synthetic Methods. ¹H nuclear magnetic resonance (NMR) spectra were obtained with a Varian Gemini-300 spectrometer using CDCl₃ or DMSO-*d*₆ as solvent. The chemical shifts are expressed as parts per million (ppm) downfield from tetramethylsilane or as relative ppm from CDCl₃ (7.27 ppm) and dimethyl sulfoxide (DMSO) (2.5 ppm). FAB (fast atom bombardment) mass spectrometry was performed with a JEOL SX102 spectrometer using 6-kV Xe atoms. CI-NH₃ (chemical ionization) mass spectra were carried out with a Finnigan 4600 mass spectrometer. Elemental analysis ($\pm 0.4\%$ acceptable) was performed by Atlantic Microlab Inc. (Norcross, GA). All melting points were determined with a Unimelt capillary melting point apparatus (Arthur H. Thomas Co., PA) and were uncorrected. All xanthine derivatives were homogeneous as judged using thin-layer chromatography (TLC) (MK6F silica, 0.25 mm, glass backed; Whatman, Clifton, NJ). Where needed, determinations of purity were performed with a Hewlett-Packard 1090 high-performance liquid chromatography (HPLC) system using an SMT OD-5–60 C18 analytical column (250 mm \times 4.6 mm, Separation Methods Technologies, Inc., Newark, DE) in two different linear gradient solvent systems. One solvent system (A) was 0.1 M triethylammonium acetate buffer:CH₃CN in ratios of 80:20 to 20:80 in 30 min with a flow rate of 1 mL/min. The other (B) was H₂O:CH₃OH in ratios of 60:40 to 10:90 in 30 min, 10:90 after 30 min, with a flow rate of 1 mL/min. Peaks were detected by UV absorption (250 nm) using a diode array detector.

6-Amino-1-methyl-3-benzyl-1H-pyrimidine-2,4-dione (66b). To a suspension of 6-amino-1-methyluracil (706 mg, 5 mmol) in methanol (15 mL) was added 10% aqueous NaOH solution (2 mL, 5 mmol) and benzyl bromide (0.65 mL, 5.5 mmol). The reaction mixture was refluxed for 3 h and quenched by the addition of saturated NH₄Cl solution. The solvent was evaporated under reduced pressure, and the residue was diluted with chloroform (50 mL). The organic phase was washed with water and brine and dried over sodium sulfate, filtered, and evaporated under reduced pressure. The residue was purified by flash column chromatography (silica, chloroform:methanol = 10:1) to give 97 mg of **66b** (33%, based on recovered starting material) as a white foaming solid. ¹H

NMR (CDCl₃): δ 3.33 (s, 3H), 5.00 (s, 1H), 5.08 (s, 2H), 7.23–7.42 (m, 5H). MS (positive ion FAB): 232 [M + H]⁺.

6-Amino-1-methyl-5-nitroso-3-benzyl-1H-pyrimidine-2,4-dione (67b). Compound **66b** (53 mg, 0.23 mmol) was dissolved in a mixed solvent of water (1 mL), acetic acid (0.2 mL), and 6 N HCl (0.05 mL) and treated with NaNO₂ (24 mg, 0.35 mmol) in small portions over a period of 10 min. The mixture was filtered, and the filter cake was washed with water to afford **67b** (30 mg, 50%) as a violet solid. ¹H NMR (DMSO-*d*₆): δ 3.25 (s, 3H), 5.10 (s, 2H), 7.23–7.40 (m, 5H). MS (positive ion FAB): 261 [M + H]⁺.

1-Benzyl-3-methylxanthine (14b). Sodium hydrosulfite (120 mg, 0.69 mmol) was added in small portions to a suspension of **67b** (30 mg, 0.12 mmol) in ethyl acetate (4 mL) and water (2 mL), until the violet color disappeared. The organic layer was separated, and the aqueous layer was extracted with ethyl acetate (2 \times 10 mL). The combined organic layer was washed with brine and dried over sodium sulfate, filtered, and evaporated under reduced pressure to dryness. The crude product was suspended in trimethylorthoformate (2 mL) and refluxed for 2 h. Trimethylorthoformate was removed by a nitrogen stream, and the residue was purified by preparative TLC (chloroform:methanol = 10:1). The solid was recrystallized from ethyl acetate/hexane to furnish xanthine **14b** (13 mg, 44%) as a white solid having a mp 224–226 °C. ¹H NMR (CDCl₃): δ 3.65 (s, 3H), 5.28 (s, 2H), 7.27–7.35 (m, 3H) 7.49 (d, 2H, *J* = 6.6 Hz), 7.73 (s, 1H), 11.93 (bs, 1H). MS (positive ion FAB): 257 [M + H]⁺.

6-Amino-1-methyl-3-phenethyl-1H-pyrimidine-2,4-dione (66c). Compound **66c** was prepared by the procedure described for **66b**, except using ethanol as solvent, instead of methanol, and obtained in 21% yield. ¹H NMR (CDCl₃): δ 2.91 (m, 2H), 3.42 (s, 3H), 4.12 (m, 2H), 4.56 (bs, 2H), 5.00 (s, 1H), 7.20–7.31 (m, 5H). MS (positive ion FAB): 246 [M + H]⁺.

6-Amino-1-methyl-5-nitroso-3-phenethyl-1H-pyrimidine-2,4-dione (67c). Compound **67c** was prepared by the procedure described for **67b** in 51% yield. ¹H NMR (DMSO-*d*₆): δ 2.88 (t, 2H, *J* = 7.8 Hz), 3.26 (s, 3H), 4.10 (t, 2H, *J* = 8.0 Hz), 7.20–7.38 (m, 5H). MS (positive ion FAB): 275 [M + H]⁺.

3-Methyl-1-(2-phenylethyl)xanthine (14c). Compound **14c** was prepared by the procedure described for **14b** in 32% yield; mp 240–243 °C. ¹H NMR (CDCl₃): δ 2.97 (m, 2H), 3.66 (s, 3H), 4.29 (m, 2H), 7.22–7.29 (m, 2H) 7.32 (d, 3H, *J* = 4.4 Hz), 7.78 (s, 1H), 11.45 (bs, 1H). MS (positive ion FAB): 271 [M + H]⁺.

6-Amino-1-methyl-3-(3-phenylpropyl)-1H-pyrimidine-2,4-dione (66d). Compound **66d** was prepared by the procedure described for **66c** in 45% yield. ¹H NMR (CDCl₃): δ 1.97 (m, 2H), 2.68 (t, 2H, *J* = 8.0 Hz), 3.39 (s, 3H), 3.97 (t, 2H, *J* = 7.4 Hz), 4.45 (bs, 2H), 4.97 (s, 1H), 7.10–7.28 (m, 5H). MS (positive ion FAB): 260 [M + H]⁺.

6-Amino-1-methyl-5-nitroso-3-(3-phenylpropyl)-1H-pyrimidine-2,4-dione (67d). Compound **67d** was prepared by the procedure described for **67b** in 78% yield. ¹H NMR (DMSO-*d*₆): δ 1.91 (m, 2H), 2.66 (t, 2H, *J* = 7.7 Hz), 3.22 (s, 3H), 3.95 (t, 2H, *J* = 7.3 Hz), 7.11–7.30 (m, 5H). MS (CI/NH₃): 289 [M + H]⁺.

3-Methyl-1-(3-phenylpropyl)xanthine (14d). Compound **14d** was prepared by the procedure described for **14b** in 60% yield; mp 176–180 °C. ¹H NMR (CDCl₃): δ 2.06 (quintet, 2H, *J* = 7.4 Hz), 2.75 (t, 2H, *J* = 7.4 Hz), 3.63 (s, 3H), 4.14 (t, 2H, *J* = 7.4 Hz), 7.12–7.30 (m, 5H), 7.74 (s, 1H), 12.28 (bs, 1H). MS (positive ion FAB): 285 [M + H]⁺.

7-(3-Chloropropyl)theophylline (19b). Sodium hydride (24 mg, 60% in mineral oil, 0.61 mmol) was added to a solution of theophylline (100 mg, 0.56 mmol) in DMF (2 mL), and the reaction mixture was stirred at room temperature for 10 min. 1-Bromo-3-chloropropane (0.07 mL, 0.70 mmol) was then added dropwise. The reaction mixture was stirred at room temperature for 24 h, and the solvent was removed by a nitrogen stream. The residue was purified using preparative TLC (ethyl acetate:hexane:methanol = 10:10:1) to give **19b** (126 mg, 88%) as a white solid. The solid was recrystallized from ethyl acetate/*n*-hexane; mp 122–124 °C. ¹H NMR

(CDCl₃): δ 2.39 (m, 2H), 3.42 (s, 3H), 3.50 (t, 2H, $J = 5.9$ Hz), 3.61 (s, 3H), 4.49 (t, 2H, $J = 6.5$ Hz), 7.62 (s, 1H). MS (positive ion FAB): 257 [M + H]⁺.

7-(4-Chlorobutyl)theophylline (19c). Compound **19c** was prepared by the procedure described for **19b** in 95% yield; mp 115–117 °C. ¹H NMR (CDCl₃): δ 1.81 (m, 2H), 2.07 (m, 2H), 3.42 (s, 3H), 3.57 (t, 2H, $J = 6.3$ Hz), 3.60 (s, 3H), 4.35 (t, 2H, $J = 7.1$ Hz), 7.56 (s, 1H). MS (positive ion FAB): 271 [M + H]⁺.

1-Allyl-7-chloroethyl-3-methyl-8-phenylxanthine (64). Sodium hydride (10 mg, 60% in mineral oil, 0.25 mmol) was added dropwise to a solution of 1-allyl-3-methyl-8-phenylxanthine **51**³⁶ (6 mg, 0.02 mmol) in DMF (1 mL), and the reaction mixture was stirred at room temperature for 10 min. 1,2-Dichloroethane (0.05 mL, 0.6 mmol) was added, and the reaction mixture was stirred at 70 °C for 36 h. The solvent was removed by a nitrogen stream. The residue was purified using preparative TLC (dichloromethane:2-propanol = 100:1) to give **64** (4.8 mg, 70%) as a white solid; mp 130–133 °C. ¹H NMR (CDCl₃): δ 3.66 (s, 3H), 3.94 (t, 2H, $J = 6.0$ Hz) 4.66 (t, 2H, $J = 6.3$ Hz) 4.68 (d, 2H, $J = 5.5$ Hz) 5.22 (dd, 1H, $J = 10.2, 1.1$ Hz) 5.31 (dd, 1H, $J = 17.3, 1.1$ Hz) 5.96 (m, 1H) 7.54–7.56 (m, 3H) 7.64–7.68 (m, 2H). High-resolution MS (positive ion FAB) calcd for C₁₇H₁₈ClN₄O₂ [M + H]⁺, 345.1118; found, 345.1110. HPLC indicated 99% purity (retention times (min): A, 22.7; B, 23.6).

7-Chloroethyl-1,3-dipropyl-8-phenylxanthine (65). To a solution of 1,3-dipropyl-8-phenylxanthine **49**²⁵ (13 mg, 0.042 mmol) in DMF (2 mL) was added NaH (18 mg, 60% in mineral oil, 0.45 mmol), and the reaction mixture was stirred for 10 min at room temperature. 1,2-Dichloroethane (0.10 mL, 1.27 mmol) was added dropwise, and the reaction mixture was stirred at 70 °C for 48 h. The solvent was removed by a nitrogen stream. The residue was purified using preparative TLC (dichloromethane:2-propanol = 100:1) to give **65** (5.0 mg, 32%) as a white solid; mp 128–132 °C. ¹H NMR (CDCl₃): δ 1.09 (t, 3H, $J = 7.4$ Hz), 1.10 (t, 3H, $J = 7.7$ Hz), 1.82 (m, 2H), 1.95 (m, 2H), 4.06 (t, 2H, $J = 6.3$ Hz), 4.11 (dd, 2H, $J = 7.4, 7.7$ Hz), 4.23 (dd, 2H, $J = 7.7, 7.4$ Hz), 4.74 (t, 2H, $J = 6.2$ Hz), 7.64–7.66 (m, 3H), 7.74–7.77 (m, 2H). High-resolution MS (positive ion FAB) calcd for C₁₉H₂₄ClN₄O₂ [M + H]⁺, 375.1588; found, 375.1573. HPLC indicated 96% purity (retention times (min): A, 23.6; B, 30.2).

Pharmacological Methods. Cloning of A_{2B} Receptor. The rat A_{2B} receptor was cloned by Dr. Eric Yuan-Ji Day (University of Virginia) by the following method: Sprague–Dawley rats were euthanized with an overdose of pentobarbital. The urinary bladders were removed and stored in RNALater (Qiagen). mRNA was extracted from rat bladder tissue using a Qiagen mRNA extraction kit. The rat A_{2B} receptor was amplified by RT-polymerase chain reaction (PCR) with the forward primer GGCCATGCAGCTAGACGACGAC and reverse primer TAGGTCACAAGCTCAGACTGA and then cloned into TOPO2.1 and confirmed by sequencing. The confirmed PCR product was then subcloned in a pDouble-Trouble vector using the restriction enzyme sites for HindIII and EcoRV.

Stable Transfection. HEK-293 cells were grown in DMEM/F12 with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin to 50% confluence in six well dishes. Cells were transfected with 2 μ g of plasmid DNA and 5 μ L of lipofectamine in accordance with the manufacturer's instructions (GibcoBRL). Transfected cells were grown for 48 h prior to selection by the addition of 1 mg/mL G418 in the medium. Resistant colonies were isolated and screened using radioligand binding with ¹²⁵I-ABOPX. Clonal lines with high specific binding were expanded and maintained in 0.5 mg/mL G418. Several clones with expression levels of approximately 20 000 fmol/mg were preserved. The average K_D for ¹²⁵I-ABOPX for the rat A_{2B} receptor was determined to be 40.9 nM (SEM = 4.4, $n = 7$).

Binding Assays. Membranes from HEK-293 cells stably expressing the human or rat A_{2B} receptor were used for competition binding assays with ¹²⁵I-ABOPX (2200 Ci/mmol).^{10,32}

Radioligand binding experiments were performed in triplicate with 20–25 μ g of membrane protein in a total volume of 0.1 mL of HE buffer (10 mM HEPES and 1 mM EDTA, pH 7.4) supplemented with 1 U/mL adenosine deaminase and 5 mM MgCl₂. Nonspecific binding was measured in the presence of 100 μ M NECA. Xanthine derivatives for competition assays were diluted in HE buffer with 10% DMSO. The incubation time was 3 h at 21 °C. Competition experiments were carried out using between 0.5 and 1.0 nM ¹²⁵I-ABOPX. Membranes were filtered on Whatman GF/C filters using a Brandel cell harvester (Gaithersburg, MD) and washed three times during 15–20 s with ice-cold buffer (10 mM Tris, 1 mM MgCl₂, pH 7.4). K_i values for different compounds were derived from IC₅₀ values as described, assuming a K_D value for ¹²⁵I-ABOPX of 36 nM at the human A_{2B} receptor. Data from replicate experiments were tabulated as means \pm SEM. Nonspecific binding, measured in the presence of 100 μ M NECA (RBI-Sigma, St. Louis, MO), was 25% of total binding. All nonradioactive compounds were initially dissolved in DMSO and diluted with buffer to the final concentration, with the amount of DMSO in the final assay tubes consistently \leq 0.5%.

For competition experiments, at least six different concentrations of competitor, spanning 3 orders of magnitude adjusted appropriately for the IC₅₀ of each compound, were used. IC₅₀ values, calculated with the nonlinear regression method implemented in the Prism program (GraphPAD, San Diego, CA), were converted to apparent K_i values.³³ Equilibrium binding competition experiments at rat A₁, rat A_{2A}, and human A₃ adenosine receptors were carried out as previously reported.^{34–36}

Acknowledgment. We thank Dr. John Daly (NID-DK) for helpful discussions and for the gift of xanthine derivatives.

References

- (1) Linden, J.; Jacobson, K. A. Molecular biology of recombinant adenosine receptors. In *Cardiovascular Biology of Purines*; Burnstock, G., Dobson, J. G., Liang, B. T., Linden, J., Eds.; Kluwer: Norwell, MA, 1998; pp 1–20.
- (2) Daly, J. W.; Butts-Lamb, P.; Padgett, W. Subclasses of adenosine receptors in the central nervous system: interaction with caffeine and related methylxanthines. *Cell. Mol. Neurobiol.* **1983**, *3*, 69–80.
- (3) Hide, I.; Padgett, W. L.; Jacobson, K. A.; Daly, J. W. A_{2A}-Adenosine receptors from rat striatum and rat pheochromocytoma PC12 cells: Characterization with radioligand binding and by activation of adenylate cyclase. *Mol. Pharmacol.* **1992**, *41*, 352–359.
- (4) Feoktistov, I.; Murray, J. J.; Biaggioni, I. Positive modulation of intracellular Ca²⁺ by adenosine A_{2B}, prostacyclin and prostaglandin E₂ receptors via a cholera toxin-sensitive mechanism in human erythroleukemia cells. *Mol. Pharmacol.* **1994**, *45*, 1160–1167.
- (5) Rubino, A.; Ralevic, V.; Burnstock, G. Contribution of P1-(A_{2B} subtype) and P2-purinoceptors to the control of vascular tone in the rat isolated mesenteric arterial bed. *Br. J. Pharmacol.* **1995**, *115*, 648–652.
- (6) Harada, H.; Asano, O.; Hoshino, Y.; Yoshikawa, S.; Matsukura, M.; Kabasawa, Y.; Nijima, J.; Kotake, Y.; Watanabe, N.; Kawata, T.; Inoue, T.; Horio, T.; Yasuda, N.; Minami, H.; Nagata, K.; Murakami, M.; Nagaoka, J.; Kobayashi, S.; Tanaka, I.; Abe, S. 2-Alkynyl-8-aryl-9-methyladenines as novel adenosine receptor antagonists: Their synthesis and structure–activity relationships toward hepatic glucose production induced via agonism of the A_{2B} receptor. *J. Med. Chem.* **2001**, *44*, 170–179.
- (7) Grant, M. B.; Tarnuzzer, R. W.; Caballero, S.; Ozeck, M. J.; Davis, M. I.; Spoerri, P. E.; Feoktistov, I.; Biaggioni, I.; Shryock, J. C.; Belardinelli, L. Adenosine receptor activation induces vascular endothelial growth factor in human retinal endothelial cells. *Circ. Res.* **1999**, *85*, 699–706.
- (8) Auchampach, J. A.; Jin, J.; Wan, T. C.; Caughey, G. H.; Linden, J. Canine mast cell adenosine receptors: cloning and expression of the A₃ receptors and evidence that degranulation is mediated by the A_{2B} receptor. *Mol. Pharmacol.* **1998**, *52*, 846–860.
- (9) Strohmeier, G. R.; Reppert, S. M.; Lencer, W. I.; Madara, J. L. The A_{2B} adenosine receptor mediates cAMP responses to adenosine receptor agonists in human intestinal epithelia. *J. Biol. Chem.* **1995**, *270*, 2387–2394.

- (10) Kim, Y.-C.; Ji, X.-D.; Melman, N.; Linden, J.; Jacobson, K. A. Anilide derivatives of an 8-phenylxanthine carboxylic congener are highly potent and selective antagonists at human A_{2B} adenosine receptors. *J. Med. Chem.* **2000**, *43*, 1165–1172.
- (11) de Zwart, M.; Vollinga, R. C.; von Frijtag Drabbe Künzel, J. K.; Beukers, M.; Slegers, D.; IJzerman, A. P. Potent antagonists for the Human Adenosine A_{2B} receptor. Derivatives of the triazolotriazine adenosine receptor antagonist ZM241385 with high affinity. *Drug Dev. Res.* **1999**, *48*, 95–103.
- (12) Feoktistov, I.; Biaggioni, I. Adenosine A_{2B} Receptors. *Pharmacol. Rev.* **1997**, *49*, 381–402.
- (13) Robeva, A. S.; Woodard, R.; Jin, X.; Gao, Z.; Bhattacharya, S.; Taylor, H. E.; Rosin, D. L.; Linden, J. Molecular characterization of recombinant human adenosine receptors. *Drug Dev. Res.* **1996**, *39*, 243–252.
- (14) Jacobson, K. A.; IJzerman, A. P.; Linden, J. 1,3-Dialkylxanthine derivatives having high potency as antagonists at human A_{2B} adenosine receptors. *Drug Dev. Res.* **1999**, *47*, 45–53.
- (15) Linden, J. Molecular characterization of A_{2A} and A_{2B} receptors. *Drug Dev. Res.* **1998**, *43*, 2.
- (16) Ji, X.-d.; Jacobson, K. A. Use of the triazolotriazine [3 H]-ZM241385 as a radioligand at recombinant human A_{2B} adenosine receptors. *Drug Des. Discovery* **1999**, *16*, 217–226.
- (17) Ji, X.-d.; Kim, Y.-C.; Ahern, D. G.; Linden, J.; Jacobson, K. A. [3 H]MRS 1754, a selective antagonist radioligand for A_{2B} adenosine receptors. *Biochem. Pharmacol.* **2001**, *61*, 657–663.
- (18) Daly, J. W.; Hide, I.; Müller, C. E.; Shamim, M. Caffeine analogues: structure–activity relationships at adenosine receptors. *Pharmacology* **1991**, *42*, 309–321.
- (19) Shamim, M. T.; Ukena, D.; Padgett, W. L.; Daly, J. W. Effects of 8-phenyl and 8-cycloalkyl substituents on the activity of mono-, di-, and trisubstituted alkylxanthines with substitution at the 1-, 3-, and 7-positions. *J. Med. Chem.* **1989**, *32*, 1231–1237.
- (20) Jacobson, K. A.; de la Cruz, R.; Schulick, R.; Kiriassias, L.; Padgett, W.; Pfeleiderer, W.; Kirk, K. L.; Neumeyer, J. L.; Daly, J. W. 8-Substituted xanthines as antagonists at A_1 - and A_2 -adenosine receptors. *Biochem. Pharmacol.* **1988**, *37*, 3653–3661.
- (21) Shamim, M. T.; Ukena, D.; Padgett, W. L.; Hong, O.; Daly, J. W. 8-Aryl- and 8-cycloalkyl-1,3-dipropylxanthines: further potent and selective antagonists for A_1 -adenosine receptors. *J. Med. Chem.* **1988**, *31*, 613–617.
- (22) Ukena, D.; Shamim, M. T.; Padgett, W. L.; Daly, J. W. Analogues of caffeine: antagonists with selectivity for A_2 adenosine receptors. *Life Sci.* **1986**, *39*, 743–750.
- (23) Daly, J. W.; Padgett, W. L.; Shamim, M. T. Analogues of 1,3-dipropyl-8-phenylxanthine: enhancement of selectivity at A_1 -adenosine receptors by aryl substituents. *J. Med. Chem.* **1986**, *29*, 1520–1524.
- (24) Daly, J. W.; Padgett, W. L.; Shamim, M. T. Analogues of caffeine and theophylline: effect of structural alterations on affinity at adenosine receptors. *J. Med. Chem.* **1986**, *29*, 1305–1308.
- (25) Müller, C. E.; Shi, D.; Manning, M., Jr.; Daly, J. W. Synthesis of paraxanthine analogues (1,7-disubstituted xanthines) and other xanthines unsubstituted at the 3-position: structure–activity relationships at adenosine receptors. *J. Med. Chem.* **1993**, *36*, 3341–3349.
- (26) Feoktistov, I.; Polosa, R.; Holgate, S. T.; Biaggioni, I. Adenosine A_{2B} receptors: a novel therapeutic target in asthma? *Trends Pharmacol. Sci.* **1998**, *19*, 148–153.
- (27) Fozard, J. R.; Hammon, J. P. Adenosine receptor ligands: potential as therapeutic agents in asthma and COPD. *Pulm. Pharmacol. Ther.* **1999**, *12*, 111–114.
- (28) Kirshenbaum, A. S.; Hettlinger, B.; Day, Y.-J.; Gilfillan, A. M.; Metcalfe, D. D.; Kim, Y.-C.; Linden, J.; Jacobson, K. A. *MRS 1754, a Selective A_{2B} Adenosine Receptor Antagonist, Inhibits Degranulation of Cultured Human but Not Murine Mast Cells*. American Academy of Allergy, Asthma, and Immunology 57th Annual Meeting, New Orleans, LA, 2001; abstr. 20000803.
- (29) Coffin, V. L.; Spealman, R. D. Psychomotor-stimulant effects of 3-isobutyl-1-methylxanthine: comparison with caffeine and 7-(2-chloroethyl)theophylline. *Eur. J. Pharmacol.* **1989**, *170*, 35–40.
- (30) Cristalli, G. Synthesis and characterization of potent ligands at human recombinant adenosine receptors. *Drug Dev. Res.* **1998**, *43*, 23.
- (31) Feoktistov, I.; Goldstein, A. E.; Biaggioni, I. Cyclic AMP and protein kinase A stimulate Cdc42: role of A_2 adenosine receptors in human mast cells. *Mol. Pharmacol.* **2000**, *58*, 903–910.
- (32) Linden, J.; Thai, T.; Figler, R.; Jin, X.; Robeva, A. S. Characterization of human A_{2B} adenosine receptors: radioligand binding, western blotting, and coupling to G_q in human embryonic kidney 293 cells and HMC-1 mast cells. *Mol. Pharmacol.* **1999**, *56*, 705–713.
- (33) Linden, J. Calculating the dissociation constant of an unlabeled compound from the concentration required to displace radiolabel binding by 50%. *J. Cycl. Nucl. Res.* **1982**, *8*, 163–172.
- (34) Schwabe, U.; Trost, T. Characterization of adenosine receptors in rat brain by (–) [3 H]N 6 -phenylisopropyladenosine. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1980**, *313*, 179–187.
- (35) Jarvis, M. F.; Schutz, R.; Hutchison, A. J.; Do, E.; Sills, M. A.; Williams, M. [3 H]CGS 21680, an A_2 selective adenosine receptor agonist directly labels A_2 receptors in rat brain tissue. *J. Pharmacol. Exp. Ther.* **1989**, *251*, 888–893.
- (36) Olah, M. E.; Gallo-Rodriguez, C.; Jacobson, K. A.; Stiles, G. L. [125 I]AB-MECA, a high affinity radioligand for the rat A_3 adenosine receptor. *Mol. Pharmacol.* **1994**, *45*, 978–982.
- (37) Daly, J. W.; Padgett, W. L.; Shamim, M. T.; Butts-Lamb, P.; Waters, J. 1,3-Dialkyl-8-(*p*-sulfophenyl)xanthines: Potent water-soluble antagonists for A_1 - and A_2 -adenosine receptors. *J. Med. Chem.* **1985**, *28*, 487–492.

JM0104318