


Rational Design, Synthesis, and Characterization of Novel Inhibitors for Human β 1,4-Galactosyltransferase

Kenji Takaya, Noriko Nagahori, Masaki Kurogochi, Tetsuya Furuike, Nobuaki Miura, Kenji Monde, Yuan Chuan Lee, and Shin-Ichiro Nishimura

J. Med. Chem., **2005**, 48 (19), 6054-6065 • DOI: 10.1021/jm0504297 • Publication Date (Web): 27 August 2005

Downloaded from <http://pubs.acs.org> on March 28, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Rational Design, Synthesis, and Characterization of Novel Inhibitors for Human β 1,4-Galactosyltransferase^{||}

Kenji Takaya,[†] Noriko Nagahori,[†] Masaki Kuroguchi,[†] Tetsuya Furuike,[‡] Nobuaki Miura,[†] Kenji Monde,[†] Yuan Chuan Lee,[§] and Shin-Ichiro Nishimura^{*,†,¶}

Division of Biological Sciences, Frontier Research Center for Post-Genome Science and Technology, Graduate School of Science, Hokkaido University, N-21, W-11, Sapporo 001-0021, Japan, Division of Bioscience, Graduate School of Environmental Earth Science, Hokkaido University, N-10, W-5, Sapporo 060-0810, Japan, Department of Biology, Johns Hopkins University, Baltimore, Maryland 21218, and National Institute of Advanced Industrial Science and Technology (AIST), Sapporo 062-8517, Japan

Received May 6, 2005

An affinity labeling reagent, uridine 5'-(6-amino-{2-[(7-bromomethyl-2-naphthyl)methoxycarbonylmethoxy]ethoxy}acetyl-6-deoxy- α -D-galactopyranosyl) diphosphate (**1a**), was designed on the basis of 3D docking simulation and synthesized to investigate the functional role of Trp310 residue located in the small loop near the active site of human recombinant galactosyltransferase (β GalT-1). Mass spectrometric analysis revealed that the Trp310 residue of β GalT1 can be selectively modified with the naphthylmethyl group of compound **1a** at the C-3 position of the indole ring. This result motivated us to synthesize novel uridine-5'-diphosphogalactose (UDP-Gal) analogues as candidates for mechanism-based inhibitors for β GalT-1. We found that uridine 5'-(6-O-[10-(2-naphthyl)-3,6,9-trioxadecanyl]- α -D-galactopyranosyl) diphosphate (**2**) is the strongest inhibitor ($K_i = 1.86 \mu\text{M}$) against UDP-Gal ($K_m = 4.91 \mu\text{M}$) among compounds reported previously. A cold spray ionization time-of-flight mass spectrometry study demonstrated that the complex of this inhibitor and β GalT-1 cannot interact with an acceptor substrate in the presence of Mn^{2+} .

Introduction

Glycosylation reactions conducted by glycosyltransferases is one of the crucial biological processes for the post-translational modifications of protein(s) and lipid(s) functions that greatly influence various molecular recognition processes including bacterial and viral infections, cell adhesion in inflammation, immune response, cellular differentiation, development, regulation, and many other intercellular communication and signal transductions.¹ Selective inhibitors of glycosyltransferases are therefore of interest because they may lead to the development of novel and potential therapeutic reagents.^{2–9} However, it seems that the difficulty of the development of practical glycosyltransferase inhibitors might be due to the complex and dynamic reaction mechanism that involves many components such as a sugar nucleotide as a donor substrate, an acceptor sugar substrate, and a metal ion.

Galactosyltransferases (Gal-T), the most important and ubiquitous superfamily of sugar elongation enzymes in eukaryotes, transfers galactose from uridine-5'-diphosphogalactose (UDP-Gal) to an acceptor substrate. It was known that β Gal-T1 undergoes critical conformational changes upon glycosyl donor substrate binding

from an open conformation (conformation I) to a closed conformation (conformation II).^{10,11} Upon donor substrate binding, tryptophan 314 (Trp314; GenBank accession number X14558) in the small flexible loop plays an important role for the conformational changes in the long flexible loop (residues 345–365) by moving toward the catalytic pocket to interact with the complex of donor substrate and Mn^{2+} ion.¹² This conformational change seems to be an essential event for the subsequent successful binding with an acceptor sugar substrate because the mutation of Trp to Ala drastically reduced both the binding affinity to the sugar nucleotide and catalytic activity of this enzyme.¹³ In human β Gal-T1 (GenBank accession number X55415), Trp310 located in the consensus sequence (³⁰⁹Gly-Trp-Gly-Gly-Glu-Asp³¹⁴) is thought to act as a key amino acid for inducing this crucial conformational change (see Supporting Information). It is noteworthy that the movement of Trp314 (Trp310 in human β Gal-T1) to the active site of β Gal-T1 might be indispensable both for the binding with the donor substrate and for the conformational change in the long flexible loop required for opening the door of the guest room.

Our interest has been focused on the functional role of this Trp310 residue in the dynamic mechanism of the human β Gal-T1 activity. We considered that compounds prohibiting the access of the Trp310 existing far from the active site would become a new class of the potent inhibitors for β Gal-T1. In the present study, we describe an efficient strategy for the rational design and synthesis of the potent inhibitors of β Gal-T1 by utilizing a designated affinity labeling reagent for the Trp310 residue.

^{||} A part of this work was presented at 84th Annual Meeting of the Chemical Society Japan on March 2004, held in Nishinomiya, Hyogo, Japan.

* To whom correspondence should be addressed. Phone: +81-11-706-9043. Fax: +81-11-706-9042. E-mail address: shin@glyco.sci.hokudai.ac.jp.

[†] Graduate School of Science, Hokkaido University.

[‡] Graduate School of Environmental Earth Science, Hokkaido University.

[§] Johns Hopkins University.

[¶] National Institute of Advanced Industrial Science and Technology.




Figure 1. Dynamic conformational change of β Gal-T1 along with the binding with UDPGal-Mn²⁺. Upon a donor substrate binding, Trp314 in the small flexible loop (residues 313–316, yellow) plays a crucial role in the conformational changes of the long flexible loop (residues 345–365, red) by moving toward the catalytic pocket to interact with β -phosphate oxygen of UDP-Gal: (a) open conformation docked with UDP-galactose (conformation I); (b) closed conformation (conformation II). The flipping of Trp314 to the active site of β 1,4Gal-T1 is essential both for binding with the donor substrate and for the conformational change in the long flexible loop required for opening the door of the guest room (conformation II). This figure was prepared using UCSF Chimera. UCSF Chimera is a highly extended interactive molecular graphics program. For detail information, see <http://www.cgl.ucsf.edu/chimera/>.


Results and Discussion

A. Affinity Labeling of Trp310 Residue. To estimate the distance between UDP-Gal in the active sites of β Gal-T1 and Trp314, UDP-Gal was docked into the binding cavity of conformation I (open conformation, pdb entry: 1FR8). As shown in Figure 1a, it was indicated from X-ray crystallographic data¹⁰ that the distance between Trp314 of the small loop (residues 313–316) and β -phosphate group of the β Gal-T1 was found to be 15.5 Å. In addition, that from the 2- or 6-OH group of UDP-Gal was estimated to be 18.8 and 15.7 Å, respectively. In contrast, it was suggested that Trp314 in conformation II (closed conformation, pdb entry: 1O0R) apparently is located near (2.7 Å) the β -phosphate group of the UDP in the binding pocket (Figure 1b).¹¹ It is likely that the NH (indole ring) group of the Trp314 residue of β Gal-T1 may bind with a complex of Mn²⁺- β -phosphate oxygen atoms of UDP moiety in the donor binding pocket through hydrogen bonding and contribute to hydrophobic interactions with the C5–C6 of galactose of UDP-Gal.¹⁴

This result prompted us to design and synthesize the site-directed affinity labeling reagent based on the chemical modification at the C-6 position of UDP-Gal as the starting material, since tolerance at the C-6 or C-2 position has been demonstrated in synthetic studies on the donor substrates for β Gal-T1.^{15–17} As illustrated in Scheme 1, we synthesized the UDP-Gal derivative **1a** having a bromomethyl substituted naphthalene group¹⁸ as an aromatic affinity tag for the indole side chain of the Trp310 residue. The key intermediate **15** derived from a known 2,3,4-tri-*O*-acetyl-6-azide-6-deoxygalactopyranose¹⁹ was coupled with the active esters **10** and **9** to give the affinity labeling reagent **1a** and **1b** as a chemically stable analogue, respectively. According to the result of the manual docking shown in Figure 1a, the distance between the bromomethyl substituted

naphthalene group and the carbon atom at the C-6 position of UDP-Gal was adjusted to be approximately 15 Å by connecting them with 3,6-dioxaoctandioic acid. When a highly reactive bromomethyl naphthalene group of compound **1a** can access the Trp310 in the short loop of human β Gal-T1, we expect that nucleophilic attack by a spatially adjacent C-3 position of the indole ring on** this reactive group might afford β Gal-T1 covalently modified with the affinity labeling reagent **1a** at Trp310 residue.

As shown in the ESI-TOF mass spectrometry of β Gal-T1 in the presence of compound **1a** (Figure 2), advent of the new ion peaks at m/z 36 785 and 36 076 clearly suggests that compound **1a** acts as a specific tag for β Gal-T. They could be assigned to a product generated by the direct nucleophilic substitution of β Gal-T with **1a** (m/z 36 785, peak γ) and its deacylated form (m/z 36 076, peak β). It was also suggested that the stable analogue **1b** showed a moderate inhibitory effect ($K_i = 22.3 \mu\text{M}$) on β Gal-T activity by a competitive mechanism as indicated in Figure 2d. Further precise analysis of the site labeled by **1a** was carried out by using MALDI-TOF/TOF MS²⁰ of the tryptic digests of the affinity labeled human recombinant β Gal-T1 as shown in Figure 3. An unknown parent ion peak at m/z 3273.149 was subjected to the fragmentation experiment to allow for structural characterization of this peptide fragment. It was finally revealed, by using a series of b- and y-ions, that a new precursor ion peak at m/z 3273.149 was the ion peak due to the sequence ²⁹⁵QQFLTINGFPNNY-WGWWGEDDDIFNR³²⁰ including the small loop region in which the Trp310 residue was selectively modified with the naphthylmethyl group at the C-3 position of the indole ring (two possible isomers were indicated in Figure 3b). Although careful elucidation of the detectable ion peaks in spectrum 2 of Figure 3a has been carried out by TOF/TOF analysis, no significant candi-

Scheme 1^a

^a Reagents and conditions: (i) (1) Cs₂CO₃/MeOH–H₂O, room temp; (2) 2-bromomethyl-7-methylnaphthalene/DMF, 50 °C, 46%; (ii) pentafluorophenyl trifluoroacetate, pyridine/DMF, room temp, 53%; (iii) NBS, AIBN/CCl₄, 80 °C, 27%; (iv) 2-chloro-4*H*-benzodioxaphosphorin 4-one, Et₃N/THF-dioxane, 0 °C; (v) H₃PO₃/CH₃CN, room temp, 47% vol; (vi) Dowex 50W-X8 H⁺/THF, room temp; (vii) *t*BuOOH, I₂/THF, room temp, 65%; (viii) UMP-morpholidate, 1*H*-tetrazole/pyridine, room temp, 62%; (ix) NaOMe/MeOH, –10 °C, 70%; (x) 10% Pd–C, H₂/H₂O, room temp; (xi) **14** (for **1b**) or **15** (for **1a**), Et₃N/H₂O–dioxane, room temp.

date bearing the naphthylmethyl group except the above-mentioned peptide was obtained, indicating that the Trp310 is a specific amino acid reacting with this affinity labeling reagent **1a**. Therefore, we considered that the aromatic stacking between naphthalene moieties of **1a,b** and the indole ring of the Trp310 may be crucial and feasible for designing a new class of potent inhibitors of β Gal-T1.

B. Inhibitor Design Based on Affinity Labeling of Trp310. The results of the affinity labeling of Trp310 by **1a** and the potent inhibitory effect by **1b** motivated us to design much more effective inhibitors than **1b** on the basis of the modification study of UDP-Gal. Considering the structural features of compound **1b**, it seemed that an amide bond and an ester bond may reduce the flexibility of the linker moiety that influences the affinity with Trp310 in the flexible small loop. Therefore, we decided to synthesize compounds **2** and **3** bearing a naphthalene group linked by a flexible trioxadecanyl moiety, in which UDP-Gal and naphthalene were combined by ether bonds. Scheme 2 indicates

the synthetic route of compound **2** (schemes and experimental details of the synthesis of compounds **3–7** are described in the Supporting Information). Thiophenylglycoside **21** was obtained by a key substitution reaction of **18** with an alcohol **20** in 82% yield. Conversion of the thiophenyl group of triol **22** into the phosphate group (five steps) at the anomeric position followed by coupling with UMP-morpholidate afforded compound **2** as an amorphous powder. The K_i values of UDP-Gal derivatives investigated in this paper are listed in Figure 4. It was revealed that compound **2** exhibited the highest inhibitory activity against this enzyme among compounds used in this study. To the best of our knowledge, it was demonstrated that **2** is the strongest competitive inhibitor ($K_i = 1.86 \mu\text{M}$) against UDP-Gal ($K_m = 4.91 \mu\text{M}$) among inhibitors reported (Figure 4a).^{4,7} Analogues **4**, **5**, and **7** lost (or reduced) the inhibitory effect on human recombinant β Gal-T1, indicating that both the flexible trioxadecanyl [(CH₂CH₂O)₃] linker and aromatic (naphthyl) terminal groups as well as two ether bonds are indispensable in achieving the satisfac-




Figure 2. Irreversible binding of **1a** on β Gal-T1: (a) ESI mass spectrum of β Gal-T1 in the negative control experiment; (b) in the presence of $30 \mu\text{M}$ compound **1a**; (c) in the presence of $100 \mu\text{M}$ compound **1a**; (d) inhibitory effect of compound **1b**. Conditions are the following: LC/MS analysis was conducted in using 2% acetic acid; 100 mM ammonium carbonate (pH 6.8) was used as washing buffer; linear gradient mode was 2%; acetic acid and acetonitrile were the eluents.

tory inhibition. Although compound **3** showed a little improved inhibition in comparison with compound **1b**, modification at the C-6 position seemed to be much more effective than that at the C-2 position of Gal residue. Since we have not synthesized a suicide substrate analogue of compound **3**, the mechanism of the inhibition by **3** is not clear.

Our hypothesis explaining the inhibitory effect observed in the present study is that the synthetic analogues of UDP-Gal, having an aromatic (hydrophobic) group through a flexible linker moiety (**1–4**), may disturb the access of the NH (indole ring) group of the Trp310 residue of the human β Gal-T1 to a complex of Mn^{2+} - β -phosphate oxygen atoms of their own UDP moiety in the donor binding pocket. As a result, it seems that an acceptor substrate cannot bind efficiently with the complex of β Gal-T1-UDP-Gal analogues because access of Trp310 is thought to be an essential step to induce the change from conformation I to conformation II required for the successful binding of an acceptor substrate.^{12,21} If this hypothesis is correct, the complex of β Gal-T1-**2** does not show binding capacity with an acceptor substrate. A snapshot of the results from electrospray ionization mass spectrometry under non-destructive conditions (ESI-TOF MS)²² revealed that the complex of human recombinant β Gal-T1 with **2** and Mn^{2+} ($[\text{M} + 12\text{H}]^{12+}$, m/z 3066.6) exhibited no binding

capacity toward an acceptor substrate **8**, while a signal due to the complex of β Gal-T1 with UDP-Gal and Mn^{2+} (m/z 3044.1) shifted to the peak at m/z 3030.1 because of the complex of β Gal-T-UDP- Mn^{2+} and to the peak at m/z 3073.1 corresponding to the molecular mass of the complex for β Gal-T-UDP- Mn^{2+} acceptor substrate **8** under the same conditions (Figure 5). These results are still preliminary but at least justified our hypothesis that these compounds might inhibit the binding of the acceptor substrate with β Gal-T1 by perturbing a key functional role of the Trp310 in the small loop.

Conclusion

Novel inhibitors for β Gal-T1 were designed and synthesized on the basis of the affinity labeling of the Trp310 located in the small flexible loop near the active site. Among UDP-Gal analogues studied herein, compound **2** showed the highest inhibitory effect ($K_i = 1.86 \mu\text{M}$). In general, highly conserved Trp residues (GWGGE and similar sequences) seem to play an essential role in the enzymatic reactions of many glycosyltransferases, namely, β 1,4-galactosyltransferase,²³ β 1,3-galactosyltransferase,²⁴ α 1,4-*N*-acetyl-D-hexosaminyltransferase,²⁵ clostridial cytotoxin glycosyltransferases,²⁶ and polypeptide *N*-acetyl-D-galactosaminyltransferase.²⁷ As for other glycosyltransferases, individual 3D structural information of the flexible loop near the active site will become




Figure 3. Affinity labeling and proteomics analysis of Trp310 by using a suicide substrate **1a**. A precursor ion peak at m/z 3273.149 corresponding to the mass of the peptide fragment Q295-R320 was further employed for precise fragmentation analysis based on a MALDI-TOF/TOF MS method according to the procedure described in our previous report.¹⁸ It was demonstrated that Trp310 can be modified with compound **1a** through a nucleophilic substitution at C-3 position of this indole ring. There is the possibility of a nucleophilic addition reaction by main chain amide nitrogen after the first substitution reaction at C-3 indole, though it is difficult to distinguish these isomers.


key to the rational design of the dynamic inhibitors. We are currently investigating crystallization of the complex of the human recombinant β Gal-T1 with compound **2**, and the results of the structural characterization will be reported as soon as possible.

Experimental Section

Chemicals. Solvents were dried with 4 Å molecular sieves before use. Silica gel column chromatography was performed with Kanto chemical 60N (100–210 μ m diameter) and Iatrobed column from Iatron Laboratories, 6RS-8060. All chemical reagents were purchased from Wako Pure Chemical Co. Ltd. Recombinant human β 1,4-galactosyltransferase-1 was purchased from Toyobo Co. Ltd. UDP and UDP-galactose were obtained from Yamasa Corporation. MALDI matrixes [2,5-

dehydroxybenzoic acid (DHB), α -cyano-4-hydroxycinnamic acid (CHCA), sinapinic acid (SA)] and MALDI peptide calibration standard mixture containing angiotensin II, bombesin, ACTH (18-39), and somatostatin were purchased from Bruker Daltonics GmbsH.

Structural Characterization. ¹H and ¹³C NMR spectra were recorded on Bruker AVANCE 600 and 500 MHz (150 and 125 MHz for ¹³C, respectively). TMS for organic solvents and TMSP for D₂O were used as the internal standards. Cold spray ionization mass spectrometry (CSI-TOF MS) analyses were performed by using a JMS-T100CS spectrometer (AccuTOF CS, JEOL Co. Ltd., Japan) equipped with an orthogonal acceleration time-of-flight (oa-TOF) mass analyzer. MALDI-TOF mass spectra were measured by using an Ultraflex TOF/TOF mass spectrometer equipped with a reflector and controlled by the Flexcontrol 1.2 software package (Bruker

Scheme 2^a

^a Reagents and conditions: (i) 2-bromomethylnaphthalene, NaH/DMF, 0 to room temp, 95%; (ii) SOCl₂, pyridine/DMF, room temp, 85%; (iii) *p*-methoxybenzyl chloride, NaH/DMF, 0 °C to room temp, 95%; (iv) 1 N HCl/MeOH, CH₂Cl₂, room temp, 92% vol, 18, NaH/DMF, 0 °C to room temp, 82%; (v) PhSH, SnCl₂/CH₂Cl₂, -78 °C, 61%; (vi) Ac₂O/pyridine, room temp, 96%; (viii) NBS/acetone-H₂O, room temp, 77%; (ix) *N,N*-diallyl-diisopropyl phosphoramidite, 1*H*-tetrazole/THF, -40 °C, 49%; (x) *t*BuOOH/THF, -10 °C, 98%; (xi) tetrakis(triphenylphosphine)palladium, Et₂NH/THF, room temp, 91%; (xii) Et₂NH/MeOH, room temp, 88%; (xiii) UMP-morpholidate, 1*H*-tetrazole/pyridine, room temp, 41%.

Daltonics GmbsH, Bremen, Germany). Optical rotation was measured by a Perkin-Elmer polarimeter, model 343. High- and low-resolution mass spectrometry and elemental analysis were carried out at the Center for Instrumental Analysis, Hokkaido University. A high-performance liquid chromatography (HPLC) system was used and was equipped with the following: HITACHI L-7100 pump, HITACHI ODS reverse-phase column (46 mm × 250 mm), and HITACHI F-1050 fluorescence detector.

{2-[(7-Methyl-2-naphthyl)methoxycarbonylmethoxy]ethoxy}acetic Acid (9). To a solution of 3,6-dioxaoctandioic acid (500 mg, 2.81 mmol) in a mixed solvent of MeOH (10 mL) and H₂O (10 mL) was added Cs₂CO₃ (914 mg, 2.81 mmol) at room temperature. After being stirred for 20 min at room temperature, the reaction mixture was concentrated in vacuo, the obtained residue was redissolved in a mixed solvent of DMF (20 mL) and H₂O (5 mL), and 2-bromomethyl-7-methylnaphthalene (110 mg, 468 μmol) was added at room temperature. After being stirred for 24 h at 70 °C, the reaction mixture was concentrated and the residue obtained was purified by a Wakogel 50C18 reverse-phase column (water as a eluent) to give compound **9** (82 mg, 46% calculated from 2-bromomethyl-7-methylnaphthalene). ¹H NMR (600 MHz, CD₃OD, 27 °C) δ 7.79 (d, 1H, aromatic), 7.72 (m, 2H, aromatic),

7.59 (s, 1H, aromatic), 7.38 (d, 1H, aromatic), 7.31 (d, 1H, aromatic), 5.32 (s, 2H, OCH₂naph), 4.24 (s, 2H, OCH₂C(=O)Onaphthyl), 3.90 (s, 2H, OCH₂C(=O)OH), 3.63–3.70 (m, 4H, OCH₂CH₂O), 2.47 (s, 3H, CH₃); ¹³C NMR (150 MHz, CD₃OD, 27 °C) δ 176.14, 170.99, 135, 92, 133.51, 133.12, 131.54, 128.25, 127.74, 127.20, 126.55, 126.53, 124.66, 70.49, 70.31, 69, 38, 67, 91, 66.50, 20.35; HRMS calcd for [M + H]⁺ C₁₈H₂₁O₆ 333.1338, found 333.1338.

{2-[(7-Methyl-2-naphthyl)methoxycarbonylmethoxy]ethoxy}acetic Acid Pentafluorophenyl Ester (10). To a solution of **9** (216 mg, 650 μmol) in DMF (10 mL) was added pyridine (155 μL, 1.9 mmol) and pentafluorophenyl trifluoroacetate (218 μL, 1.27 mmol) at room temperature.²⁸ After being stirred overnight at room temperature, the reaction mixture was concentrated and the obtained residue was purified by silica gel column chromatography (hexane/EtOAc = 2:1) to give **10** (170 mg, 53%). ¹H NMR (600 MHz, CDCl₃, 27 °C) δ 7.77 (d, 1H, aromatic), 7.71 (m, 2H, aromatic), 7.57 (s, 1H, aromatic), 7.36 (d, 1H, aromatic), 7.32 (d, 1H, aromatic), 5.34 (s, 2H, OCH₂naph), 4.52 (s, 2H, OCH₂C(=O)Opfp), 4.24 (s, 2H, OCH₂C(=O)O-naphthyl), 3.81–3.86 (m, 4H, OCH₂CH₂O), 2.50 (s, 3H, CH₃); ¹³C NMR (150 MHz, CDCl₃, 27 °C) δ 170.62, 166.64, 136.22, 133.42, 132.60, 131.52, 128.79, 128.22, 127.51,

mL, 46 mmol) and concentrated in vacuo. Purification of the concentrate was conducted by silica gel column chromatography ($\text{CHCl}_3/\text{MeOH} = 25:1$ to $10:1$) to give the pure α anomer compound **12** (1.4 g, 66%) as the mono- Et_3N salt. ^1H NMR (500 MHz, CD_3OD , 27°C) δ 5.86 (dd, 1 H, $J_{1,p} = 8.9$ Hz, $J_{1,2} = 3.4$ Hz, H-1), 5.42–5.45 (m, 2 H, H-3, H-4), 5.21 (m, 1 H, H-2), 4.42 (t, 1 H, H-5), 3.44 (dd, 1 H, $J_{5,6b} = 7.6$ Hz, $J_{6a,6b} = 12.7$ Hz, H-6b), 3.20 (dd, 1 H, $J_{5,6a} = 5.2$ Hz, H-6a), 3.09 (m, 6 H, $(\text{CH}_3\text{CH}_2)_3\text{N}$), 2.16 (s, 3 H, Ac), 2.08 (s, 3 H, Ac), 1.98 (s, 3 H, Ac), 1.35 (m, 9 H, $(\text{CH}_3\text{CH}_2)_3\text{N}$); ^{13}C NMR (125 MHz, CDCl_3 , 27°C) δ 170.27, 170.25, 169.94, 91.81, 91.78, 68.79, 68.74, 67.69, 67.65, 67.53, 50.66, 45.58, 20.80, 20.66, 8.53; HRMS calcd for $[\text{M} + \text{Et}_3\text{N} + \text{H}]^+ \text{C}_{18}\text{H}_{34}\text{N}_4\text{O}_{10}\text{P}$ 497.2013, found 497.2013.

2,3,4-Tri-O-acetyl-6-azido-6-deoxy- α -D-galactopyranosyl Phosphate Mono- Et_3N Salt (13). Compound **12** (1.2 g, 2.4 mmol) dissolved in THF (10 mL) was passed through the cation-exchange resin column (Dowex 50W-X8 H^+ form) with additional THF (30 mL) as eluent. To the solution was added *t*BuOOH (580 μL of 5 M solution in decane) and I_2 (4 mg) at room temperature.³¹ After being stirred for 24 h at room temperature in an argon atmosphere, the reaction mixture was neutralized by Et_3N (338 μL , 2.4 mmol) and then concentrated. The concentrate was purified by a Wakogel 50C18 reverse-phase column (water as a eluent) to give the compound **13** (810 mg, 65%). ^1H NMR (500 MHz, CD_3OD , 27°C) δ 5.78 (dd, 1 H, $J_{1,p} = 7.5$ Hz, $J_{1,2} = 3.4$ Hz, H-1), 5.47 (dd, 1 H, $J_{4,5} = 1.1$ Hz, H-4), 5.42 (dd, 1 H, $J_{3,4} = 3.4$ Hz, H-3), 5.14 (m, 1 H, $J_{2,3} = 10.7$ Hz, H-2), 4.45 (t, 1 H, $J_{5,6} = 6.9$ Hz, H-5), 3.50 (dd, 1 H, $J_{5,6a} = 6.0$ Hz, $J_{6a,6b} = 12.5$ Hz, H-6a), 3.27 (dd, 1 H, $J_{5,6b} = 7.3$ Hz, H-6b), 3.09 (m, 6 H, $(\text{CH}_3\text{CH}_2)_3\text{N}$), 2.16 (s, 3 H, Ac), 2.08 (s, 3 H, Ac), 1.98 (s, 3 H, Ac), 1.33 (m, 9 H, $(\text{CH}_3\text{CH}_2)_3\text{N}$); ^{13}C NMR (125 MHz, CDCl_3 , 27°C) δ 170.71, 170.28, 169.91, 92.00, 91.96, 68.67, 67.87, 67.82, 67.65, 67.50, 50.32, 45.43, 20.86, 20.65, 8.46; HRMS calcd for $[\text{M} + \text{Et}_3\text{N} + \text{H}]^+ \text{C}_{18}\text{H}_{34}\text{N}_4\text{O}_{11}\text{P}$ 513.1962, found 513.1954.

Uridine 5'-(2,3,4-Tri-O-acetyl-6-azido-6-deoxy- α -D-galactopyranosyl) Diphosphate Monoammonium Salt (14). To a solution of **13** (760 mg, 1.48 mmol) in dry pyridine (5 mL) was added UMP-morpholidate (2.0 g, 2.96 mmol) at room temperature. The solution was concentrated in vacuo to complete dryness (removal of water). To the obtained mixture redissolved in dry pyridine (5 mL) was added 1*H*-tetrazole (455 mg, 5.9 mmol) at room temperature, and the mixture was stirred for 24 h in an argon atmosphere.³² To the reaction solution was added additional UMP-morpholidate (700 mg, 960 μmol), and the mixture was stirred for 2 days at room temperature. The reaction mixture was concentrated in vacuo, and the obtained residue was subjected to purification by ion exchange column chromatography (DEAE-Sephacell, 50 mM NH_4HCO_3 as the eluent). Further purification was conducted by a Wakogel 50C18 reverse-phase column (water as the eluent) to give the compound **14** (650 mg, 62%) as the monoammonium salt. ^1H NMR (600 MHz, D_2O , 27°C) δ 7.98 (d, 1 H, uracil), 5.96–5.98 (m, 2 H, uracil, H'-1), 5.80 (dd, 1 H, $J_{1',p} = 7.7$ Hz, $J_{1',2'} = 3.5$ Hz, H''-1), 5.52 (d, 1 H, H''-4), 5.38 (dd, 1 H, $J_{2',3'} = 10.7$ Hz, $J_{3',4'} = 3.3$ Hz, H''-3), 5.22 (m, 1 H, H''-2), 4.51 (t, 1 H, $J_{5',6'} = 6.4$ Hz, H''-5), 4.34–4.38 (m, 2 H, H'-2, H'-3), 4.25–4.28 (m, 2 H, H'-4, H'-5 \times 1), 4.19 (m, 1 H, H'-5 \times 1), 3.56 (dd, 1 H, $J_{5',6'b} = 6.5$ Hz, $J_{6'a,6'b} = 12.9$ Hz, H''-6b), 3.44 (dd, 1 H, $J_{5',6'a} = 6.4$ Hz, H''-6a), 2.22 (s, 3 H, Ac), 2.15 (s, 3 H, Ac), 2.02 (s, 3 H, Ac); ^{13}C NMR (150 MHz, D_2O , 27°C) δ 176.06, 175.78, 175.47, 168.83, 154.49, 144.46, 105.34, 95.31, 95.27, 91.27, 85.88, 85.82, 76.51, 72.29, 71.66, 70.85, 70.80, 70.45, 70.39, 67.61, 67.57, 52.33, 23.00, 22.76, 22.70; HRMS calcd for $[\text{M} + \text{NH}_3 + \text{H}]^+ \text{C}_{21}\text{H}_{33}\text{N}_6\text{O}_{19}\text{P}_2$ 735.1276, found 735.1280.

Uridine 5'-(6-Azido-6-deoxy- α -D-galactopyranosyl) Diphosphate Monoammonium Salt (15). To a solution of **14** (300 mg, 423 μmol) in MeOH (40 mL) was added NaOMe (114 mg, 2.1 mmol) in 6 mL of MeOH at -10°C . After being stirred for 24 h at -10°C , to the reaction mixture was added additional NaOMe (10 mg, 184 μmol in 1 mL of MeOH 1 mL) daily for 4 days. The reaction mixture was neutralized by the

addition of cation-exchange resin (Dowex 50W-X8 H^+) and filtered by glass filter. The filtrate was concentrated and purified by ion exchange column chromatography (DEAE-Sephacell, 50 mM NH_4HCO_3 as the eluent). Further purification was conducted by gel permeation chromatography (Sephadex G-10, water as the eluent) to give the compound **15** (179 mg, 70%) as the monoammonium salt. ^1H NMR (600 MHz, D_2O , 27°C) δ 7.95 (d, 1 H, uracil), 5.97–5.98 (m, 2 H, uracil, H'-1), 5.61 (dd, 1 H, $J_{1',p} = 7.1$ Hz, $J_{1',2'} = 3.5$ Hz, H''-1), 4.35–4.38 (m, 2 H, H'-2, H'-3), 4.18–4.28 (m, 4 H, H'-4, H'-5 \times 2, H''-5), 3.99 (d, 1 H, H''-4), 3.91 (dd, 1 H, $J_{3',4'} = 3.4$ Hz, H''-3), 3.77 (dt, 1 H, $J_{2',3'} = 10.3$ Hz, H''-2), 3.56 (dd, 1 H, $J_{5',6'b} = 7.3$ Hz, $J_{6'a,6'b} = 12.8$ Hz, H''-6b), 3.46 (dd, 1 H, $J_{5',6'a} = 6.1$ Hz, H''-6a); ^{13}C NMR (150 MHz, D_2O , 27°C) δ 168.90, 154.42, 144.37, 105.37, 98.47, 98.43, 91.20, 85.97, 85.91, 76.49, 72.84, 72.34, 71.93, 71.90, 71.03, 70.97, 67.65, 67.61, 62.11, 53.02; HRMS calcd for $[\text{M} + \text{NH}_3 + \text{H}]^+ \text{C}_{15}\text{H}_{27}\text{N}_6\text{O}_{16}\text{P}_2$ 609.0959, found 609.0953.

Uridine 5'-(6-Amino-6-deoxy- α -D-galactopyranosyl) Diphosphate Monoammonium Salt (16). To a solution of **15** (4.5 mg, 7.4 μmol) in H_2O (3 mL) was added 10% Pd-C (10 mg), and the mixture was stirred for 30 min at room temperature in a hydrogen atmosphere. The reaction solution was filtered and the filtrate was concentrated to give compound **16** (4.1 mg, 96%). ^1H NMR (600 MHz, D_2O , 27°C) δ 7.92 (d, 1 H, uracil), 5.95–5.98 (m, 2 H, uracil, H'-1), 5.65 (dd, 1 H, $J_{1',p} = 6.5$ Hz, $J_{1',2'} = 3.7$ Hz, H''-1), 4.35–4.40 (m, 3 H, H'-2, H'-3, H''-5), 4.21–4.29 (m, 3 H, H'-4, H'-5 \times 2), 4.03 (d, 1 H, H''-4), 3.95 (dd, 1 H, $J_{3',4'} = 3.3$ Hz, H''-3), 3.84 (dt, 1 H, $J_{2',3'} = 10.3$ Hz, H''-2), 3.25 (m, 2 H, H''-6a, H''-6b); ^{13}C NMR (150 MHz, D_2O , 27°C) δ 168.87, 154.48, 144.35, 105.32, 98.38, 98.34, 91.27, 85.85, 85.79, 76.41, 72.34, 72.15, 71.63, 70.69, 70.61, 67.80, 67.76, 62.13, 42.87; HRMS calcd for $[\text{M} + \text{H}]^+ \text{C}_{15}\text{H}_{26}\text{N}_3\text{O}_{16}\text{P}_2$ 566.0788, found 566.0793.

Uridine 5'-(6-Amino-2-[(7-bromomethyl-2-naphthyl)methoxycarbonylmethoxy]ethoxy)acetyl-6-deoxy- α -D-galactopyranosyl) Diphosphate Monoammonium Salt (1a). To a solution of **16** (2.5 mg, 4.3 μmol) in H_2O (2 mL) was added Et_3N (18 μL , 12.9 μmol) and compound **11** (3 mg, 5.3 μmol in 2 mL of THF) at room temperature. After the mixture was stirred for 30 min at room temperature, 5 mL of 20 mM ammonium acetate buffer (pH 6.0) was added and then concentrated. The obtained mixture was used directly for the affinity labeling experiment without purification because 30–40% hydrolysis at the bromomethyl group was confirmed after keeping the temperature at 4°C for 24 h as judged by TLC and ESI MS (data not shown). ESI MS (negative ion mode) calcd for $[\text{M} - \text{H}]^- \text{C}_{33}\text{H}_{41}\text{BrN}_3\text{O}_{21}\text{P}_2$ 956.1, found 956.

Uridine 5'-(6-Amino-2-[(7-methyl-2-naphthyl)methoxycarbonylmethoxy]ethoxy)acetyl-6-deoxy- α -D-galactopyranosyl) Diphosphate Monoammonium Salt (1b). To a solution of **16** (10 mg, 17 μmol) in H_2O (2 mL) was added Et_3N (4.8 μL , 34 μmol) and compound **10** (25 mg, 51 μmol in dioxane 4 mL) at room temperature. After being stirred for 30 min at room temperature, the reaction mixture was concentrated and the residue obtained was purified by a Wakogel 50C18 reverse-phase column (60% MeOH as a eluent). The obtained compound was passed through a cation-exchange resin column (Dowex 50W-X8, Na^+ form) to give compound **1b** (18 mg, 84%). ^1H NMR (600 MHz, D_2O , 27°C) δ 7.66–7.87 (m, 5 H, uracil, aromatic \times 4), 7.41–7.45 (m, 2 H, aromatic \times 2), 5.81–5.83 (m, 2 H, H'-1, uracil), 5.60 (dd, 1 H, $J_{1',p} = 7.1$ Hz, $J_{1',2'} = 3.6$ Hz, H''-1), 5.36 (s, 2 H, OCH_2naph), 4.15–4.31 (m, 7 H, H'-2, H'-3, H'-4, H'-5a, H'-5b, $\text{OCH}_2\text{C}=\text{O} \times 2$), 4.13 (t, 1 H, H''-5, $J_{4',5'} = 6.7$ Hz), 4.00 (m, 2H, $\text{OCH}_2\text{C}=\text{O} \times 1$, H''-4), 3.86–3.88 (m, 2H, $\text{OCH}_2\text{C}=\text{O} \times 1$, H''-3), 3.76–3.78 (m, 3H, $\text{OCH}_2\text{CH}_2\text{O}$, H''-2), 3.70–3.72 (m, 2H, $\text{OCH}_2\text{CH}_2\text{O}$), 3.48 (dd, 1 H, $J_{6'a,6'b} = 13.8$ Hz, $J_{5',6'a} = 5.9$ Hz, H''-6a), 3.31 (dd, 1H, $J_{5',6'b} = 7.4$ Hz, H''-6b), 2.48 (s, 3H, CH_3naph); ^{13}C NMR (150 MHz, D_2O , 27°C) δ 175.36, 174.96, 168.59, 144.07, 139.64, 135.79, 135.58, 133.78, 131.68, 130.91, 130.28, 129.41, 129.36, 127.76, 105.13, 98.48, 98.43, 91.31, 85.77, 85.71, 76.49, 73.07, 72.31, 72.18, 72.01, 71.91, 71.06, 71.00,

70.82, 70.12, 67.55, 67.51, 57.05, 41.88, 23.47, 21.77, 17.67; HRMS calcd for $[M + H]^+$ $C_{33}H_{44}N_3O_{21}P_2$ 880.1943, found 880.1924.

10-(2-Naphthyl)-3,6,9-trioxadecanol (17). A solution of triethyleneglycol (399 μ L, 3 mmol) in DMF (8 mL) was cooled to 0 °C, and sodium hydride (54 mg, 1.5 mmol) was added. After the mixture was stirred for 30 min at 0 °C, 2-bromo-methylnaphthalene (276 mg, 1.2 mmol) was added to the solution, and the mixture was stirred for 1 h at 0 °C and was allowed to warm to room temperature for further stirring. After the mixture was stirred for 2 h at room temperature, the reaction was stopped by the addition of MeOH and concentrated. The residue obtained was purified by silica gel column chromatography (toluene/EtOAc = 3:1 to 1:1) to give **17** (202 mg, 70%). 1H NMR (600 MHz, $CDCl_3$, 27 °C) δ 7.79–7.83 (dd, 4 H, naphthalene), 7.46–7.49 (m, 3 H, naphthalene), 4.73 (s, 2 H, OCH_2 -naphthalene), 3.70 (m, 10 H, $[OCH_2CH_2]$, OCH_2CH_2OH), 3.61 (m, 2 H, $[OCH_2CH_2]$); ^{13}C NMR (125 MHz, $CDCl_3$, 27 °C) δ 136.00, 133.62, 133.35, 128.52, 128.21, 128.04, 126.89, 126.43, 126.20, 126.16, 73.73, 72.87, 71.03, 71.00, 70.75, 69.74, 62.13; HRMS calcd for $[M + H]^+$ $C_{17}H_{23}O_4$ 291.1596, found 291.1588.

1-Chloro-10-(2-naphthyl)-3,6,9-trioxadecane (18). To a solution of **17** (200 mg, 688 μ mol) in CH_2Cl_2 (5 mL) was added pyridine (100 μ L) and thionyl chloride (200 μ L, 2.75 mmol) at room temperature. After being stirred for 24 h, the reaction solution was concentrated and the residue obtained was purified by silica gel column chromatography (toluene/EtOAc = 2:1) to give **18** (190 mg, 89%). 1H NMR (600 MHz, $CDCl_3$, 27 °C) δ 7.79–7.83 (m, 4 H, naphthalene), 7.46–7.48 (m, 3 H, naphthalene), 4.73 (s, 2 H, OCH_2 -naphthalene), 3.74 (t, 2 H, OCH_2CH_2-Cl), 3.67–3.72 (m, 8 H, $[OCH_2CH_2]$), 3.61 (t, 2 H, $[OCH_2CH_2]$); ^{13}C NMR (150 MHz, $CDCl_3$, 27 °C) δ 135.74, 133.25, 132.97, 128.14, 127.84, 127.67, 126.46, 126.07, 125.83, 125.78, 73.34, 71.36, 70.74, 70.68, 70.65, 69.43, 42.72; HRMS calcd for $[M]^+$ $C_{17}H_{21}O_3Cl$ 308.1179, found 308.1170.

Phenyl 2,3,4-*p*-Methoxybenzyl-1-thio-6-*O*-triphenyl-methyl- β -*D*-galactopyranoside (19). To a solution of phenyl 1-thio-6-*O*-triphenylmethyl- β -*D*-galactopyranoside³³ (5.66 g, 11 mmol) in dry DMF (50 mL) was added sodium hydride (1.8 g, 49.5 mmol) at 0 °C. After the mixture was stirred for 30 min, *p*-methoxybenzyl chloride (4.9 mL, 36.3 mmol) was added to the solution and further stirred overnight at room temperature. The reaction was stopped by addition of MeOH, and the mixture was concentrated in vacuo. The residue obtained was dissolved in $CHCl_3$, and the organic layer was washed with brine, dried by $MgSO_4$, filtered, and concentrated. The residual syrup was purified by silica gel column chromatography (hexane/EtOAc = 5:1) to give **19** (9.4 g, 98%) as a syrup; $[\alpha]^{20}_D +20.9^\circ$ (*c* 0.85, $CHCl_3$). 1H NMR (500 MHz, $CDCl_3$, 27 °C) δ 7.53 (m, 2 H, aromatic), 7.37–7.40 (m, 6 H, aromatic), 7.21–7.31 (m, 17 H, aromatic), 7.14 (dd, 3 H, aromatic), 7.05 (d, 2 H, aromatic), 6.84–6.87 (dd, 4 H, aromatic), 6.74 (d, 2 H, aromatic), 4.74 (d, 1 H, OCHH-phenyl), 4.63–4.70 (m, 4 H, OCH_2 -phenyl \times 2), 4.57 (d, 1 H, H-4), 4.44 (d, 1 H, OCHH-phenyl), 3.83 (t, 1 H, $J_{3,4} = 9.1$ Hz, H-3), 3.77–3.82 (m, 10 H, H-1, $OCH_3 \times 3$), 3.53–3.56 (dd, 1 H, $J_{5,6b} = 6.3$ Hz, $J_{6a,b} = 9.8$ Hz, H-6b), 3.46 (dd, 1 H, $J_{2,3} = 9.2$ Hz, H-2), 3.28 (t, 1 H, H-5), 3.14 (dd, 1 H, $J_{5,6a} = 6.3$ Hz, H-6a); ^{13}C NMR (150 MHz, $CDCl_3$, 27 °C) δ 159.32, 159.25, 158.98, 143.95, 134.73, 131.51, 130.94, 130.86, 130.64, 130.55, 130.37, 130.03, 129.46, 129.32, 129.23, 128.78, 128.69, 128.53, 127.96, 127.87, 127.29, 127.05, 126.76, 113.84, 113.79, 113.49, 87.79, 86.96, 83.95, 77.77, 75.29, 73.72, 72.58, 63.25, 55.32; HRMS calcd for $[M + Na]^+$ $C_{55}H_{54}O_8NaS$ 897.3437, found 897.3456.

Phenyl 2,3,4-*p*-Methoxybenzyl-1-thio- β -*D*-galactopyranoside (20). To a solution of **19** (9.1 g, 10.4 mmol) in a mixed solvent ($CHCl_3$ /MeOH = 2:1, 75 mL) was added 1 N HCl/MeOH (20.8 mL, 20.8 mmol) at room temperature. After being stirred for 90 min, the mixture was neutralized by addition of Et_3N (3.49 mL, 25.0 mmol) and concentrated. The residue obtained was purified by silica gel column chromatography (hexane/EtOAc = 2:1) to give **20** (6.08 g, 92%) as a crystal (from EtOH and hexane); $[\alpha]^{20}_D -4.9^\circ$ (*c* 1.00, $CHCl_3$). 1H NMR (500

MHz, $CDCl_3$, 27 °C) δ 7.53 (d, 2 H, aromatic), 7.20–7.33 (m, 9 H, aromatic), 6.85–6.90 (m, 6 H, aromatic), 4.89 (d, 1 H, OCHH-phenyl), 4.75 (d, 1 H, OCHH-phenyl), 4.66–4.71 (m, 3 H, OCHH-phenyl \times 3), 4.62 (d, 1 H, H-4), 4.56 (d, 1 H, OCHH-phenyl), 3.90 (t, 1 H, $J_{3,4} = 9.5$ Hz, H-3), 3.76–3.84 (m, 11 H, H-1, H-5, $OCH_3 \times 3$), 3.56 (dd, 1 H, $J_{2,3} = 9.2$ Hz, H-2), 3.45 (dd, 1 H, $J_{6a,b} = 11.3$ Hz, H-6b), 3.39 (dd, 1 H, H-6a); ^{13}C NMR (150 MHz, $CDCl_3$, 27 °C) δ 159.32, 134.14, 131.46, 130.53, 130.43, 130.35, 129.95, 129.89, 129.27, 128.84, 127.12, 113.91, 113.81, 113.78, 87.84, 83.99, 78.77, 77.33, 75.32, 73.70, 72.83, 72.77, 62.32, 55.29; HRMS calcd for $[M + H]^+$ $C_{36}H_{41}O_8S$ 633.2522, found 633.2532.

Phenyl 2,3,4-*p*-Methoxybenzyl-6-*O*-(10-[2-naphthyl]-3,6,9-trioxadecanyl)-1-thio- β -*D*-galactopyranoside (21). To a solution of **20** (2.3 g, 3.6 mmol) in dry DMF (8 mL) was added sodium hydride (262 mg, 7.2 mmol) at 0 °C.³⁴ After the mixture was stirred for 30 min, compound **18** (2.22 g, 7.2 mmol) in dry DMF (8 mL) was added to the reaction mixture at 0 °C. After the mixture was stirred for 12 h, the reaction was stopped by addition of MeOH and the solution was concentrated. The residue obtained was dissolved in $CHCl_3$, washed with brine, dried by $MgSO_4$, filtered, and concentrated. The residual syrup was purified by silica gel column chromatography ($CHCl_3$ /MeOH = 100:1) to give **21** (2.8 g, 82%) as a syrup; $[\alpha]^{20}_D +13.8^\circ$ (*c* 0.9, $CHCl_3$). 1H NMR (500 MHz, $CDCl_3$, 27 °C) δ 7.77–7.82 (m, 4 H, aromatic), 7.54 (m, 2 H, aromatic), 7.45–7.47 (m, 3 H, aromatic), 7.19–7.31 (m, 9 H, aromatic), 6.86 (m, 6 H, aromatic), 4.87 (d, 1 H, OCHH-phenyl), 4.56–4.73 (m, 9 H, H-4, OCH_2 -naphthalene, OCH_2 -phenyl \times 3), 3.88 (m, 2 H, H-3, OCH_2 CHH-Gal), 3.76–3.83 (m, 9 H, $OCH_3 \times 3$), 3.63–3.67 (m, 9 H, H-1, $[OCH_2CH_2]$), 3.54–3.57 (m, 6 H, H-2, H-5, H-6, $[OCH_2CH_2]$, OCH_2 CHH-Gal), 3.48 (m, 1 H, H-6); ^{13}C NMR (150 MHz, $CDCl_3$, 27 °C) δ 159.20, 159.10, 135.77, 134.42, 133.27, 132.98, 131.35, 131.00, 130.63, 130.49, 129.95, 129.57, 129.15, 128.73, 128.13, 127.85, 127.68, 126.91, 126.44, 126.07, 125.82, 125.77, 113.82, 113.74, 113.58, 87.80, 83.91, 77.35, 75.24, 73.96, 73.32, 73.16, 72.37, 70.74, 70.69, 70.66, 70.58, 70.42, 69.72, 69.46, 60.38, 55.27; HRMS calcd for $[M + Na]^+$ $C_{57}H_{60}O_8NaS$ 927.3754, found 927.3741.

Phenyl 6-*O*-(10-[2-Naphthyl]-3,6,9-trioxadecanyl)-1-thio- β -*D*-galactopyranoside (22). To a solution of **21** (2.45 g, 2.7 mmol) in CH_2Cl_2 (25 mL) was added thiophenol (994 μ L, 9.72 mmol), and the mixture was cooled to –78 °C. A solution of $SnCl_4$ (1 M solution of $SnCl_4/CH_2Cl_2$, 8.50 mmol) was added to the reaction mixture and stirred for 20 min at –78 °C.³⁵ The reaction was stopped by addition of saturated $NaHCO_3$ and $CHCl_3$. The organic layer was washed with saturated $NaHCO_3$ and brine and then concentrated. The residue obtained was purified by silica gel column chromatography ($CHCl_3$ /MeOH = 40:1) to give **22** (900 mg, 61%) as a syrup; $[\alpha]^{20}_D +18.7^\circ$ (*c* 1.00, $CHCl_3$). 1H NMR (500 MHz, CD_3OD , 27 °C) δ 7.83–7.86 (m, 4 H, naphthalene), 7.45–7.54 (m, 5 H, naphthalene, phenyl), 7.20–7.30 (m, 3 H, phenyl), 4.73 (s, 2 H, OCH_2 -naphthalene), 4.58 (d, 1 H, $J_{1,2} = 9.7$ Hz, H-1), 3.86 (d, 1 H, H-4), 3.59–3.76 (m, 16 H, H-2, H-5, H-6a, H-6b, $[OCH_2CH_2]$), 3.48 (dd, 1 H, $J_{3,4} = 3.3$ Hz, H-3); ^{13}C NMR (150 MHz, $CDCl_3$, 27 °C) δ 135.55, 133.25, 133.00, 132.07, 128.89, 128.20, 127.88, 127.70, 127.64, 126.61, 126.11, 125.89, 125.86, 88.53, 74.76, 73.31, 70.78, 70.56, 70.49, 70.36, 70.13, 69.75, 69.33, 68.61; HRMS calcd for $[M + H]^+$ $C_{29}H_{37}O_8S$ 545.2209, found 545.2204.

Phenyl 2,3,4-Tri-*O*-acetyl-6-*O*-(10-[2-naphthyl]-3,6,9-trioxadecanyl)-1-thio- β -*D*-galactopyranoside (23). Compound **22** (1.06 g, 1.94 mmol) was dissolved in pyridine (10 mL), and to the solution was added acetic anhydride (6 mL) at room temperature. After being stirred for 12 h, the reaction mixture was concentrated and the residual syrup was purified by silica gel column chromatography (hexane/EtOAc = 4:1 to 2:1) to give **23** (1.3 g, 99%) as a syrup. $[\alpha]^{20}_D -0.9^\circ$ (*c* 1.00, $CHCl_3$). 1H NMR (500 MHz, $CDCl_3$, 27 °C) δ 7.77–7.83 (m, 4 H, naphthalene), 7.45–7.50 (m, 5 H, naphthalene, phenyl), 7.26–7.32 (m, 3 H, phenyl), 5.44 (d, 1 H, H-4), 5.23 (t, 1 H, $J_{2,3} = 9.9$ Hz, H-2), 5.03 (dd, 1 H, $J_{3,4} = 3.3$ Hz, H-3), 4.73 (s, 2 H, OCH_2 -naphthalene), 4.70 (d, 1 H, $J_{1,2} = 10.0$ Hz, H-1),

3.86 (t, 1 H, $J_{5,6} = 6.2$ Hz, H-5), 3.62–3.70 (m, 8 H, [OCH₂-CH₂]), 3.48–3.61 (m, 6 H, H-6a, H-6b, [OCH₂CH₂]), 2.10 (s, 6 H, Ac \times 2), 2.00 (s, 3 H, Ac); ¹³C NMR (150 MHz, CDCl₃, 27 °C) δ 170.12, 169.99, 169.50, 135.82, 133.29, 132.99, 132.08, 128.90, 128.14, 127.88, 127.69, 126.47, 126.07, 125.81, 86.62, 76.02, 73.33, 72.21, 71.13, 70.69, 70.66, 70.52, 69.48, 69.20, 67.81, 67.56, 20.85, 20.67, 20.61; HRMS calcd for [M + H]⁺ C₃₅H₄₃O₁₁S 671.2526, found 671.2548.

2,3,4-Tri-O-acetyl-6-O-(10-[2-naphthyl]-3,6,9-trioxadecanyl)-D-galactopyranose (24). To a solution of **23** (500 mg, 745 μ mol) in mixed solvent (acetone/H₂O = 20:1, 10 mL) was added *N*-bromosuccinimide (398 mg, 2.23 mmol) at room temperature. After being stirred for 90 min, the mixture was concentrated and dissolved in CHCl₃. The solution was washed with saturated NaHCO₃ and brine, and the organic layer was dried by MgSO₄, filtered, and concentrated. The residue obtained was purified by silica gel column chromatography (hexane/EtOAc = 3:1 to 1:2) to give **24** (333 mg, 77%) as a syrup. ¹H NMR (600 MHz, CDCl₃, 27 °C) δ 7.83 (m, 4 H, naphthalene), 7.45–7.48 (m, 3 H, naphthalene), 5.49 (m, 0.8 H, H-3 α), 5.41–5.43 (m, 1 H, H-1 α , H-4 β), 5.16 (dd, 0.8 H, H-2 α), 5.09 (dd, 0.2 H, H-2 β), 5.05 (dd, 0.2 H, H-3 β), 4.74 (s, 2 H, OCH₂-naphthalene), 4.67 (t, 0.2 H, H-1 β), 4.48 (dd, 0.8 H, H-5 α), 4.24 (s, 0.8 H, H-4 α), 3.86 (t, 0.2 H, H-5 β), 3.57–3.72 (m, 12 H, [OCH₂CH₂] \times 3), 3.43–3.55 (m, 2 H, H-6a, H-6b), 1.92–2.13 (m, 9 H, Ac \times 3); ¹³C NMR (150 MHz, CDCl₃, 27 °C) δ 170.75, 170.70, 170.53, 170.45, 170.25, 169.84, 135.95, 135.90, 133.51, 133.23, 128.40, 128.11, 127.93, 126.74, 126.32, 126.10, 126.04, 95.88, 90.73, 73.52, 72.39, 71.26, 71.21, 71.15, 71.05, 70.90, 70.87, 70.82, 70.73, 70.67, 70.62, 70.14, 70.05, 69.62, 69.34, 68.89, 67.94, 67.32, 21.05, 20.92, 20.89, 20.83; HRMS calcd for [M + Na]⁺ C₂₉H₃₈O₁₂Na 601.2261, found 601.2237.

Diallyl 2,3,4-Tri-O-acetyl-6-O-(10-[2-naphthyl]-3,6,9-trioxadecanyl)- α -D-galactopyranosyl Phosphite (25). To a solution of **24** (106 mg, 183 μ mol) in CH₃CN (2 mL) was added 1*H*-tetrazole (23 mg, 329 μ mol), and the mixture was cooled to –40 °C followed by the addition of *N,N*-diallyl-diisopropyl phosphoramidite (73 μ L, 274 μ mol).³⁶ After the mixture was stirred for 7 h at –40 °C, EtOAc was added and the mixture was washed with saturated NaHCO₃ and brine. The organic layer was dried by MgSO₄, filtered, and concentrated. The residue obtained was purified by silica gel column chromatography (hexane/EtOAc = 4:1 to 3:1, containing 0.5 vol % Et₃N) to give **25** (65 mg, 49%). [α]_D²⁰ +17.0° (c 1.00, CHCl₃). ¹H NMR (500 MHz, CDCl₃, 27 °C) δ 7.78–7.83 (m, 4 H, naphthalene), 7.26–7.48 (m, 3 H, naphthalene), 5.88–5.96 (m, 2 H, CH₂=CHCH₂– \times 2), 5.74 (dd, 1 H, $J_{1,P} = 8.4$ Hz, $J_{1,2} = 3.4$ Hz, H-1), 5.50 (d, 1 H, H-4), 5.36 (dd, 1 H, $J_{2,3} = 10.8$ Hz, $J_{3,4} = 3.3$ Hz, H-3), 5.17–5.33 (m, 5 H, CH₂=CHCH₂– \times 2, H-2), 4.73 (s, 2 H, OCH₂-naphthalene), 4.34–4.41 (m, 5 H, CH₂=CHCH₂– \times 2, H-5), 3.54–3.71 (m, 12 H, [OCH₂CH₂]), 3.44–3.54 (m, 2 H, $J_{6a,b} = 11.0$ Hz, H-6a, H-6b), 2.12 (s, 3 H, Ac), 2.04 (s, 3 H, Ac), 1.98 (s, 3 H, Ac); ¹³C NMR (125 MHz, CDCl₃, 27 °C) δ 170.25, 170.12, 170.00, 135.81, 134.37, 134.35, 133.29, 132.99, 128.14, 127.87, 127.69, 126.47, 126.06, 125.82, 116.82, 116.75, 90.93, 90.80, 73.34, 71.01, 70.68, 70.64, 70.46, 69.47, 69.19, 68.48, 68.26, 67.61, 63.47, 63.40, 63.32, 63.26, 20.77, 20.71, 20.66; HRMS calcd for [M]⁺ C₃₅H₄₇O₁₄P 722.2703, found 722.2692.

Diallyl 2,3,4-Tri-O-acetyl-6-O-(10-[2-naphthyl]-3,6,9-trioxadecanyl)- α -D-galactopyranosyl Phosphate (26). To a solution of **25** (59 mg, 82 μ mol) in THF (2 mL) was added Et₃N (114 μ L, 820 μ mol) and cooled to –10 °C followed by the addition of *tert*-butyl hydroperoxide (49 μ L, 246 μ mol). After being stirred for 4 h at –10 °C, the reaction mixture was concentrated and the residue obtained was dissolved in CHCl₃. The solution was washed with saturated NaHCO₃ and brine. The organic layer was dried by MgSO₄, filtered, and concentrated to give **26** (60 mg, 98%). [α]_D²⁰ +47.8° (c 1.00, CHCl₃). ¹H NMR (600 MHz, CDCl₃, 27 °C) δ 7.78–7.82 (m, 4 H, naphthalene), 7.45–7.48 (m, 3 H, naphthalene), 5.91–5.95 (m, 3 H, H-1, CH₂=CHCH₂– \times 2), 5.52 (d, 1 H, H-4), 5.33–5.40 (m, 3 H, CH₂=CHCH₂–, H-3), 5.23–5.28 (m, 3 H, CH₂=

CHCH₂–, H-2), 4.72 (s, 2 H, OCH₂-naphthalene), 4.55–4.60 (m, 4 H, CH₂=CHCH₂– \times 2), 4.35–4.37 (t, 1 H, $J_{5,6a} = 6.3$ Hz, H-5), 3.56–3.69 (m, 13 H, [OCH₂CH₂], H-6b), 3.46–4.9 (dd, 1 H, $J_{6a,b} = 10.2$ Hz, H-6a), 2.12 (s, 3 H, Ac), 2.08 (s, 3 H, Ac), 1.99 (s, 3 H, Ac); ¹³C NMR (150 MHz, CDCl₃, 27 °C) δ 170.03, 169.94, 169.87, 135.82, 133.28, 132.98, 132.24, 132.19, 132.13, 128.11, 127.85, 127.67, 126.43, 126.05, 125.80, 125.79, 118.66, 118.53, 94.56, 73.30, 71.02, 70.66, 70.44, 69.57, 69.47, 68.93, 68.54, 68.51, 68.45, 68.41, 68.02, 67.27, 67.22, 67.17, 20.69, 20.63, 20.58; HRMS calcd for [M + H]⁺ C₃₅H₄₈O₁₅P 739.2731, found 739.2701.

2,3,4-Tri-O-acetyl-6-O-(10-[2-naphthyl]-3,6,9-trioxadecanyl)- α -D-galactopyranosyl Phosphate Mono-Et₃N Salt (27). To a solution of **26** (60 mg, 83 μ mol) in THF (2 mL) was added Et₃NH (172 μ L, 1.66 mmol) and tetrakis(triphenylphosphine)palladium (19 mg, 16.6 μ mol) at room temperature.³⁷ After being stirred for 2 h at room temperature, the solution was concentrated and the residue obtained was purified by Iatrobeds column chromatography (CHCl₃/MeOH = 2:1, containing 0.5 vol % Et₃N) to give **27** (57 mg, 91%). [α]_D²⁰ +40.9° (c 1.00, CHCl₃). ¹H NMR (500 MHz, CDCl₃, 27 °C) δ 7.82–7.86 (m, 4 H, naphthalene), 7.31–7.52 (m, 3 H, naphthalene), 5.79 (dd, 1 H, $J_{1,P} = 7.7$ Hz, $J_{1,2} = 3.3$ Hz, H-1), 5.53 (d, 1 H, H-4), 5.44 (dd, 1 H, $J_{3,4} = 3.3$ Hz, H-3), 5.16 (dt, 1 H, $J_{2,3} = 10.7$ Hz, H-2), 4.76 (s, 2 H, OCH₂-naphthalene), 4.49 (t, 1 H, $J_{5,6a} = 6.6$ Hz, H-5), 3.53–3.73 (m, 13 H, [OCH₂CH₂], H-6a), 3.48 (dd, 1 H, $J_{5,6b} = 7.8$ Hz, $J_{6a,b} = 9.5$ Hz, H-6b), 3.03 (dd, 6 H, CH₃CH₂N \times 3), 2.15 (s, 3 H, Ac), 2.10 (s, 3 H, Ac), 1.99 (s, 3 H, Ac), 1.30 (t, 9 H, CH₃CH₂N \times 3); ¹³C NMR (150 MHz, CDCl₃, 27 °C) δ 170.79, 170.20, 169.96, 135.86, 133.29, 132.99, 128.13, 127.87, 127.68, 126.45, 126.07, 125.83, 92.04, 91.99, 73.30, 70.77, 70.62, 70.59, 70.54, 70.34, 69.50, 68.88, 68.65, 68.20, 68.16, 67.81, 67.50, 45.44, 41.44, 20.95, 20.72, 11.17, 8.95; HRMS calcd for [M + Na]⁺ C₂₉H₃₉O₁₅PNa 681.1924, found 681.1937.

6-O-(10-[2-Naphthyl]-3,6,9-trioxadecanyl)- α -D-galactopyranosyl Phosphate Mono-Et₃N Salt (28). To a solution of **27** (57 mg, 75 μ mol) in MeOH (2 mL) was added Et₃NH (500 μ L), and the mixture was stirred for 3 days at room temperature. The reaction mixture was concentrated, and the residue obtained was purified by ion exchange column chromatography (DEAE Sephasell, 10–50 mM NH₄HCO₃ as a eluent). The obtained residue redissolved in H₂O was passed through a cation-exchange resin column (Dowex 50W-X8, H⁺ form followed by Et₃NH⁺ form) to give **28** (44 mg, 88%). [α]_D²⁰ +41.6° (c 1.00, D₂O). ¹H NMR (500 MHz, D₂O, 27 °C) δ 7.87–7.95 (m, 4 H, naphthalene), 7.52–7.59 (m, 3H, naphthalene), 5.52 (dd, 1H, $J_{1,P} = 7.3$ Hz, $J_{1,2} = 3.6$ Hz, H-1), 4.72 (s, 2 H, OCH₂-naphthalene), 4.20 (t, 1 H, $J_{5,6} = 6.0$ Hz, H-5), 3.93 (d, 1 H, H-4), 3.88 (dd, 1 H, $J_{3,4} = 3.3$ Hz, H-3), 3.80 (dt, 1 H, $J_{2,3} = 10.3$ Hz, H-2), 3.61–3.71 (m, 14 H, H-6a, H-6b, [OCH₂CH₂]), 3.18 (dd, 8 H, CH₃CH₂N \times 4), 1.27 (t, 12 H, CH₃CH₂N \times 4); ¹³C NMR (125 MHz, D₂O, 27 °C) δ 137.70, 135.65, 135.52, 131.05, 130.65, 130.43, 129.97, 129.33, 129.24, 129.14, 97.67, 97.62, 75.52, 72.60, 72.34, 72.27, 72.20, 72.16, 72.00, 71.73, 71.42, 71.06, 71.00, 49.36, 10.93; HRMS calcd for [M + H + Et₃N]⁺ C₂₉H₄₉NO₁₂P 634.2992, found 634.3003.

Uridine 5'-(6-O-[10-{2-Naphthyl}-3,6,9-trioxadecanyl]- α -D-galactopyranosyl) Diphosphate Disodium Salt (2). Compound **28** (26 mg, 39 μ mol) was dissolved in dry pyridine (1 mL) and concentrated in vacuo three times to remove water. To the residual syrup **28** in dry pyridine (1 mL) was added 1*H*-tetrazole (11 mg, 156 μ mol) and UMP-morpholidate (54 mg, 78 μ mol, 1 mL in pyridine solution) at room temperature. After being stirred for 2 days, the mixture was evaporated and the residue obtained was subjected to purification by ion exchange column chromatography (DEAE-Sephasell, 10–150 mM NH₄-HCO₃ as the eluent). Further purification by gel permeation column chromatography (Sephadex G-15, water as an eluent) and cation-exchange resin column (Dowex 50W-X8, Na⁺ form) afforded compound **2** (14 mg, 41%). The compound was lyophilized to give an amorphous powder. ¹H NMR (500 MHz, D₂O, 27 °C) δ 7.87–7.97 (m, 6 H, naphthalene, H-5), 7.55–7.61 (m, 3 H, naphthalene), 5.90 (d, 1 H, $J_{1,2'} = 4.8$ Hz, H'-1),

5.88 (d, 1 H, H-6), 5.64 (dd, 1 H, $J_{1',P} = 7.4$ Hz, $J_{1',2'} = 3.5$ Hz, H''-1), 4.77 (s, 2 H, OCH₂-naphthalene), 4.19–4.34 (6 H, H'-2, H'-3, H'-4, H'-5a, H'-5b, H''-5), 3.94 (d, 1 H, H''-4), 3.89 (dd, 1 H, $J_{2',3'} = 10.2$ Hz, $J_{3',4'} = 3.2$ Hz, H''-3), 3.61–3.80 (m, 15 H, H''-2, H''-6a, H''-6b, [OCH₂CH₂]); ¹³C NMR (150 MHz, D₂O, 27 °C) δ 165.85, 151.38, 141.26, 134.77, 132.69, 132.56, 128.11, 127.71, 127.49, 127.07, 126.39, 126.30, 126.23, 102.33, 95.63, 95.59, 88.27, 82.93, 82.87, 73.58, 72.64, 69.69, 69.51, 69.44, 69.37, 69.31, 69.18, 68.97, 68.87, 68.54, 68.24, 68.19, 64.63, 64.66; HRMS calcd for [M + Na]⁺ C₃₂H₄₄N₂O₂₀NaP₂ 861.1860, found 861.1843.

Construction of the Complex Model of UDP-Galactose and β1,4-Galactosyltransferase (Conformation I). Since the crystal structure of β1,4GalT (PDB entry 1FR8), which is in conformation I, includes UDP in the active site, UDP-galactose extracted from conformation II (PDB entry 100R) was superimposed onto the UDP in 1FR8 so that the root-mean-square deviation (rmsd) between the UDP part of UDP-galactose and UDP in 1FR8 becomes the smallest. Construction of the complex model was carried out by using the SYBYL 6.9 program (Tripos, Inc.).

Affinity Labeling of βGal-T1. Recombinant human β1,4-galactosyltransferase-1 (50 μg, NP_001488 from NCBI database, <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi>) was incubated in the presence or absence of the bromomethyl substrate **1a** (final concentration of 0, 30, 100 μM for the calculation of labeling efficiency and 500 μM for the MS/MS analysis) in 100 μL of 20 mM Tris-maleate buffer (pH 6.5) containing 50 mM NaCl and 2 mM MnCl₂ at 37 °C for 10 h. The reaction mixture was directly reduced and alkylated (10 mM DTT for 1 h at 40 °C and then 50 mM iodoacetamide for 2 h at room temperature), and then the protein was obtained with acetone precipitation. The protein was digested with 0.5 μg of trypsin in 100 mM ammonium bicarbonate (pH 8.3). MALDI-TOF MS and LIFT TOF/TOF spectra were measured after desalting by means of 10 μL of C-18 ZipTips (Millipore) according to the manufacturer's instructions.

Detection of the Affinity Labeled βGal-T1 by ESI-MS. The LC/MS analysis was conducted by JMS-T100CS (AccuTOF CS, JEOL Co. Ltd., Japan) equipped with Inertsil C8 (GL Sciences Inc., Japan). The HPLC system was equilibrated with 2% acetic acid at 50 μL/min flow speed before sample injection. After the sample injection (1 μL, corresponding to about 14 pmol of protein), sample washing was conducted with 2% acetic acid for 5 min and then with 100 mM ammonium carbonate buffer (pH 6.8) for 10 min. The protein was eluted by the linear gradient mode of 2% acetic acid and acetonitrile (acetonitrile from 0% to 100% in 10 min).

Preparation of Matrix Crystals for MALDI. The matrix solutions were prepared as follows: DHB (10 mg) was dissolved in water (1 mL), and CHCA was prepared as a saturated solution in 3:1 (v/v) of acetonitrile/water. An amount of 0.5 μL of matrix solution was applied to the target spot of the Anchorchip plate (Bruker Daltonics), which was equipped with 384 hydrophilic anchors on an otherwise hydrophobic surface, and 1 μL of the sample solution was added and then dried at room temperature. We employed these samples (about 1–10 pmol) with TOF/TOF modes using the above preparation procedure.

MALDI-TOF MS. MALDI-TOF mass spectra were measured by using an Ultraflex TOF/TOF mass spectrometer equipped with a reflector and controlled by the Flexcontrol 1.2 software package (Bruker Daltonics GmbSH, Bremen, Germany). In the MALDI-TOF MS reflector mode, ions generated by a pulsed UV laser beam (nitrogen laser, λ = 337 nm, 5 Hz) were accelerated to a kinetic energy of 23.5 kV. Metastable ions generated by laser-induced decomposition of the selected precursor ions were analyzed without any additional collision gas.

In the MALDI-TOF/TOF mode, precursor ions were accelerated to 8 kV and selected in a timed ion gate. The fragments were further accelerated by 19 kV in the LIFT cell (LIFT means "lifting" the potential energy for the second acceleration of ion source), and their masses were analyzed after the ion

reflector passage. Masses were automatically annotated by using the FlexAnalysis 2.0 software package. External calibration of MALDI mass spectra was carried out using singly charged monoisotopic peaks of a mixture of human angiotensin II (*m/z* 1046.542), bombesin (*m/z* 1619.823), ACTH (18-39) (*m/z* 2465.199), and somatostatin 28 (*m/z* 3147.472). The mixture of these peptides was measured at the central spot of a 3 × 3 square by using external calibration. To achieve mass accuracy better than 60 ppm, internal calibration was carried out by doping the matrix solution with a mixture of the calibration peptides. Calibration of these mass spectra was performed automatically by utilizing a customized macro command of the Flex Control 2.1 software package. The macro command was used for the calibration of the monoisotopic singly charged peaks of the above-mentioned peptides. TOF/TOF spectra were annotated with the BioTools 2.1 software package.

Determination of Inhibitory Constants. The inhibitory activity of the inhibitors **1b–7** for human βGal-T1 was measured by the following procedure. UDP-galactose (40 or 100 μM), inhibitors **1b–7** (0–200 μM), and the dansyl fluorescently labeled *N*-acetylglucosamine **8** (50 μM) were dissolved in the reaction buffer (50 mM Hepes, pH 6.5, 10 mM MnCl₂, 0.25 mg/mL bovine serum albumin, total volume of 100 μL). The enzymatic reaction was started by the addition of βGal-T1 (40–150 μU) and incubated at 37 °C for 10 min. Inactivation was quickly done by heating the reaction solution at 100 °C for 3 min. The reaction mixture was centrifuged for 5 min, and the supernatant (20 μL) was injected into the HPLC system to detect the product (Galβ1,4GlcNAc-dansyl) formation by measuring fluorescence at 525 nm under excitation at 340 nm. The obtained data were plotted and analyzed by employing Dixon and Webb's method.

Cold Spray Ionization Time-of-Flight Mass Spectrometry (CSI-TOF MS). Commercially available recombinant human βGal-T1 was further purified by dialysis with 50 mM ammonium acetate buffer (pH 6.0) to remove salts and detergent before CSI-TOF MS measurements. A solution of purified enzyme was concentrated by ultrafiltration with microcon YM-10 (Millipore). The contents and concentration of sample solutions for CSI-TOFMS analysis are as follows: total volume (20 μL), buffer [25 mM ammonium acetate (pH 6.0)], βGal-T1 (95 pmol), MnCl₂ (2 nmol), UDP (400 pmol), and acceptor substrate **8** (450 pmol). All CSI-TOF mass spectra were measured and recorded at 37 °C according to the conditions described in our previous paper.¹⁹

Acknowledgment. This work was supported partly by grants from the National Project of "Research and Development on Glycocluster Controlling Biomolecules" from NEDO and from the National Project of "Functional Glycoconjugate Research Aimed at Developing for New Industry" from the Ministry of Education, Science, and Culture of Japan. We appreciate Prof. Monica, M. Palcic, and Prof. Ole Hindsgaul of the Carlsberg Laboratory for helpful suggestions and comments for this work.

Supporting Information Available: Synthesis procedures and experimental data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Dwek, R. A. Glycobiology: toward understanding the function of sugars. *Chem. Rev.* **1996**, *96*, 683–720.
- Lee, L. V.; Mitchell, M. L.; Huang, S.-J.; Fokin, V. V.; Sharpless, K. B.; Wong, C.-H. A potent and highly selective inhibitor of human α-1,3-fucosyltransferase via click chemistry. *J. Am. Chem. Soc.* **2003**, *125*, 9588–9589.
- Waldscheck, B.; Streiff, M.; Notz, W.; Kinzy, W.; Schmidt, R. R. α(1,3)-Galactosyltransferase inhibition based on a new type of disubstrate analogue. *Angew. Chem., Int. Ed.* **2001**, *40*, 4007–4011.
- Compain, P.; Martin, O. R. Carbohydrate mimetics-based glycosyltransferase inhibitors. *Bioorg. Med. Chem.* **2001**, *9*, 3077–3092.

- (5) Takayama, S.; Chung, S. J.; Igarashi, Y.; Ichikawa, Y.; Sepp, A.; Lechler, R. I.; Wu, J.; Hayashi, T.; Siuzdak, G.; Wong, C.-H. Selective inhibition of β -1,4- and α -1,3-galactosyltransferases: donor sugar-nucleotide based approach. *Bioorg. Med. Chem.* **1999**, *7*, 401–409.
- (6) Kim, Y. J.; Ichikawa, M.; Ichikawa, Y. A rationally designed inhibitor of α -1,3-galactosyltransferase. *J. Am. Chem. Soc.* **1999**, *121*, 5829–5830.
- (7) Hashimoto, H.; Endo, T.; Kajihara, Y. Synthesis of the first tricomponent bisubstrate analogue that exhibits potent inhibition against GlcNAc: β -1,4-galactosyltransferase. *J. Org. Chem.* **1997**, *62*, 1914–1915.
- (8) Palcic, M. M.; Heerze, L. D.; Srivastava, O. P.; Hindsgaul, O. A bisubstrate analog inhibitor for α (1,2)-fucosyltransferase. *J. Biol. Chem.* **1989**, *264*, 17174–17181.
- (9) Yuasa, H.; Palcic, M. M.; Hindsgaul, O. Synthesis of the carbocyclic analog of uridine 5'-(α -D-galactopyranosyldiphosphate)(UDP-Gal) as an inhibitor of β (1-4)-galactosyltransferase. *Can. J. Chem.* **1995**, *73*, 2190–2195.
- (10) Gastinel, L. N.; Cambillau, C.; Bourne, Y. Crystal structures of the bovine β 4galactosyltransferase catalytic domain and its complex with uridine diphosphogalactose. *EMBO. J.* **1999**, *18*, 3546–3557.
- (11) Ramakrishnan, B.; Qasba, P. K. Crystal structure of lactose synthase reveals a large conformational change in its catalytic component, the β 1,4-galactosyltransferase-I. *J. Mol. Biol.* **2001**, *310*, 205–218.
- (12) Ramakrishnan, B.; Boeggeman, E.; Ramasamy, V.; Qasba, P. K. Structure and catalytic cycle of β -1,4-galactosyltransferase. *Curr. Opin. Struct. Biol.* **2004**, *14*, 593–600.
- (13) Ramasamy, V.; Ramakrishnan, B.; Boeggeman, E.; Qasba, P. K. The role of tryptophan 314 in the conformational changes of β 1,4-galactosyltransferase-I. *J. Mol. Biol.* **2003**, *331*, 1065–1076.
- (14) Ramakrishnan, B.; Balaji, P. V.; Qasba, P. K. Crystal structure of β 1,4-galactosyltransferase complex with UDP-Gal reveals an oligosaccharide acceptor binding site. *J. Mol. Biol.* **2002**, *318*, 491–502.
- (15) Bülter, T.; Schumacher, T.; Namdjou, D.-J.; Gallego, R. G.; Clausen, H.; Elling, L. Chemoenzymatic synthesis of biotinylated nucleotide sugars as substrates for glycosyltransferases. *Chem-BioChem.* **2001**, *2*, 884–894.
- (16) Ernst, B.; Oehrlein, R. Substrate and donor specificity of glycosyltransferases. *Glycoconjugate J.* **1999**, *16*, 161–170.
- (17) Palcic, M. M.; Hindsgaul, O. Glycosyltransferases in the synthesis of oligosaccharide analogs. *Trends Glycosci. Glycotechnol.* **1996**, *8*, 37–49.
- (18) Loudon, G. M.; Koshland, D. E. The chemistry of a reporter group: 2-hydroxy-5-nitrobenzyl bromide. *J. Biol. Chem.* **1970**, *245*, 2247–2254.
- (19) Joosten, J. A. F.; Evers, B.; Van Summeren, R. P.; Kamerling, J. P.; Vliegthart, J. F. G. Synthesis of β -D-Galp-(1-4)- β -D-GlcNAc-(1-2)- α -D-Manp-(1-O)(CH₂)₇CH₃ mimics to explore the substrate specificity of sialyltransferases and *trans*-sialidases. *Eur. J. Org. Chem.* **2003**, *18*, 3569–3586.
- (20) Kuroguchi, M.; Nishimura, S.-I.; Lee, Y. C. Mechanism-based fluorescent labeling of β -galactosidases. *J. Biol. Chem.* **2004**, *279*, 44704–44712.
- (21) Qasba, P. K.; Ramakrishnan, B.; Boeggeman, E. Substrate-induced conformational changes in glycosyltransferases. *Trends Biochem. Sci.* **2005**, *30*, 53–62.
- (22) Nishimura, S.-I.; Nagahori, N.; Takaya, K.; Tachibana, Y.; Miura, N.; Monde, K. Direct observation of sugar–protein, sugar–sugar, sugar–water complexes by using cold-spray ionization time-of-flight mass spectrometry. *Angew. Chem., Int. Ed.* **2005**, *44*, 571–575.
- (23) Aoki, D.; Appert, H. E.; Johnson, D.; Wong, S. S.; Fukuda, M. N. Analysis of the substrate binding sites of human galactosyltransferase by protein engineering. *EMBO J.* **1990**, *9*, 3171–3178.
- (24) Malissard, M.; Dinter, A.; Berger, E. G.; Hennes, T. Functional assignment of motifs conserved in β 1,3-glycosyltransferases. *Eur. J. Biochem.* **2002**, *269*, 233–239.
- (25) Pedersen, L. C.; Dong, J.; Taniguchi, F.; Kitagawa, H.; Krahn, J. M.; Pedersen, L. G.; Sugahara, K.; Negishi, M. Crystal structure of an α 1,4-N-acetylhexosaminyltransferase (EXTL2), a member of the exostosin gene family involved in heparin sulfate biosynthesis. *J. Biol. Chem.* **2003**, *278*, 14420–14428.
- (26) Busch, C.; Hofmann, F.; Gerhard, R.; Aktories, K. Involvement of a conserved tryptophan residue in the UDP-glucose binding of large clostridial cytotoxin glycosyltransferases. *J. Biol. Chem.* **2000**, *275*, 13228–13234.
- (27) Fritz, T. A.; Hurley, J. H.; Trinh, L.-B.; Shiloach, J.; Tabak, L. A. The beginning of mucin biosynthesis: The crystal structure of UDP-GalNAc: polypeptide α -N-acetylgalactosaminyltransferase-T1. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 15307–15312.
- (28) Gayo, L. M.; Suto, M. J. Use of pentafluorophenyl esters for one-pot protection/activation of amino and thiol carboxylic acids. *Tetrahedron Lett.* **1996**, *37*, 4915–4918.
- (29) Westerduin, P.; Veeneman, G. H.; Van der Marel, G. A.; Van Boom, J. H. Synthesis of the fragment GlcNAc- α (1-P-6)-GlcNAc of the cell wall polymer of *Staphylococcus lactis* having repeating N-acetyl-D-glucosamine phosphate units. *Tetrahedron Lett.* **1986**, *27*, 6271–6274.
- (30) Ross, A. J.; Ivanova, I. A.; Ferguson, M. A. J.; Nikolaev, A. V. Parasite glycoconjugates. Part 11. Preparation of phosphodi-saccharide synthetic probes, substrate analogues for the elongating α -D-mannopyranosylphosphate transferase in the *Leishmania*. *J. Chem. Soc., Perkin Trans. 1* **2001**, *1*, 72–81.
- (31) Byun, H.-S.; Erukulla, R. K.; Bittman, R. Synthesis of sphingomyelin and ceramide 1-phosphate from ceramide without protection of the allylic hydroxyl group. *J. Org. Chem.* **1994**, *59*, 6495–6498.
- (32) Wittmann, V.; Wong, C.-H. 1H-Tetrazole as catalyst in phosphomorpholidate coupling reactions: efficient synthesis of GDP-fucose, GDP-mannose, and UDP-galactose. *J. Org. Chem.* **1997**, *62*, 2144–2147.
- (33) Kakarla, R.; Ghosh, M.; Anderson, J. A.; Dulina, R. G.; Sofia, M. J. Expedient route to F unit building block of moenomycin A. *Tetrahedron Lett.* **1999**, *40*, 5–8.
- (34) Dermer, O. C. Metallic salts of alcohols and alcohol analogs. *Chem. Rev.* **1934**, *3*, 385–430.
- (35) Yu, W.; Su, M.; Gao, X.; Yang, Z.; Jin, Z. A facile chemoselective deprotection of the *p*-methoxybenzyl group. *Tetrahedron Lett.* **2000**, *41*, 4015–4017.
- (36) Sim, M. M.; Kondo, H.; Wong, C.-H. Synthesis and use of glycosyl phosphites: An effective route to glycosyl phosphates, sugar nucleotides, and glycosides. *J. Am. Chem. Soc.* **1993**, *115*, 2260–2267.
- (37) Bannwarth, W.; Küng, E. Bis(allyloxy)(diisopropylamino)-phosphine as a new phosphorylation reagent for the phosphorylation of hydroxy functions. *Tetrahedron Lett.* **1989**, *30*, 4219–4222.