


Brief Article

**1-Benzhydryl-3-phenylurea and 1-Benzhydryl-3-phenylthiourea Derivatives:
New Templates among the CB Cannabinoid Receptor Inverse Agonists**

Giulio G. Muccioli, Johan Wouters, Gerhard K. E. Scriba,
Wolfgang Poppitz, Jacques H. Poupaert, and Didier M. Lambert

J. Med. Chem., **2005**, 48 (23), 7486-7490 • DOI: 10.1021/jm0503906 • Publication Date (Web): 21 October 2005

Downloaded from <http://pubs.acs.org> on March 29, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 7 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

1-Benzhydryl-3-phenylurea and 1-Benzhydryl-3-phenylthiourea Derivatives: New Templates among the CB₁ Cannabinoid Receptor Inverse Agonists

Giulio G. Muccioli,[†] Johan Wouters,[#] Gerhard K. E. Scriba,[‡] Wolfgang Poppitz,[§] Jacques H. Poupaert,[†] and Didier M. Lambert^{*,†}

Unité de Chimie pharmaceutique et de Radiopharmacie, Ecole de Pharmacie, Faculté de Médecine, Université catholique de Louvain, Avenue E. Mounier 73, UCL-CMFA 7340, B-1200 Bruxelles, Belgium, Laboratoire de Chimie Biologique Structurale, Faculté des Sciences, University of Namur, Rue de Bruxelles 61, 5000 Namur, Belgium, Department of Pharmaceutical Chemistry, University of Jena, Philosophenweg 14, D-07743 Jena, Germany, and Department of Inorganic and Analytical Chemistry, University of Jena, August-Bebel-Strasse 2, D-07743 Jena, Germany

Received April 25, 2005

New 1-benzhydryl-3-phenylurea derivatives and their 1-benzhydryl-3-phenylthiourea isosteres were synthesized and evaluated for their human CB₁ and CB₂ cannabinoid receptor affinity. These compounds proved to be selective CB₁ cannabinoid receptor ligands, acting as inverse agonists in a [³⁵S]-GTPγS assay. The affinity of 3,5,5'-triphenylimidazolidine-2,4-dione and 3,5,5'-triphenyl-2-thioximidazolidin-4-one derivatives, possessing the 1-benzhydryl-3-phenylurea and 1-benzhydryl-3-phenylthiourea moiety, respectively, was also evaluated. In conclusion, the 1-benzhydryl-3-phenylurea scaffold seems to be a new interesting template of CB₁ cannabinoid receptor inverse agonists.

Introduction


The synthesis of the first potent CB₁ cannabinoid receptor antagonist/inverse agonist SR141716A¹ (rimonabant) and the breeding of knock-out CB₁ mice^{2,3} confirmed the high therapeutic potential for modulating CB₁ cannabinoid receptor activity. Several CB₁ cannabinoid receptor inverse agonists were shown to reduce food intake,^{4,5} ethanol consumption,⁶ and nicotine reward^{7,8} in animals. Other studies suggested a beneficial effect on septic shock in rats⁹ and in gastrointestinal disorders.¹⁰ Subsequent phase II or III clinical trials confirmed the usefulness of such compounds in treating obesity^{11,12} and nicotine¹¹ addictions. Among the cannabinoid receptor antagonists/inverse agonists available, most are constructed around a central heterocyclic ring (pyrazole, triazole, indole, imidazolidinedione, or thioximidazolidinone) bearing further substituted aromatic rings.¹³

In this paper, we report the CB₁ cannabinoid receptor inverse agonist properties of new compounds that are not built around such a central ring. The 1-benzhydryl-3-phenylurea derivatives described herein exhibited interesting affinities and potencies for the CB₁ cannabinoid receptor. The 1-benzhydryl-3-phenylthiourea isosteres also showed affinity and inverse agonist properties at the CB₁ cannabinoid receptor. The compounds were then compared with the corresponding 3-phenyl-5,5'-diphenylimidazolidine-2,4-dione and 3-phenyl-5,5'-diphenyl-2-thioximidazolidin-4-one derivatives.

Results and Discussion

Chemistry. Benzoin condensation starting from substituted benzaldehydes was used for the synthesis of

Scheme 1^a



^a All X₂ substituents are in the para position except that for **8** (X₂ is in ortho). Reagents and conditions: (i₁) H₂O/EtOH, NaCN, reflux 3 h, CHCl₃ extraction of the oily benzoin intermediate; (i₂) HNO₃, reflux, 2 h; (i) CH₃COOH/H₂O, NaCNO, room temp; (i') acetone, NH₄OH, room temp; (ii) DMSO/aqueous KOH, nine microwave pulses.

the substituted benzils (**1**). Phenylurea (**2**, X' = O) and phenylthiourea (**2**, X' = S) derivatives were easily synthesized from the corresponding aniline and phenylisothiocyanate, respectively (Scheme 1). Target 1-benzhydryl-3-phenylurea derivatives (**3–21**) were obtained by reaction of the respective benzil (**1**) and phenylurea (**2**, X' = O), following a microwave-enhanced method previously described¹⁴ (Scheme 1). The corresponding thio derivatives (**22–25**) were obtained similarly by reacting **1** and phenylthiourea (**2**, X' = S). The synthesis of the thio derivatives, reported here for the first time, will allow the oxo and thio isostere comparison at the CB₁ and CB₂ cannabinoid receptors. The 3,5,5'-triphenylimidazolidine-2,4-dione (**26–29**) and 3,5,5'-triphenyl-2-thioximidazolidin-4-one derivatives (**30, 31**) were obtained by reaction of **1** and phenylurea (**2**, X' = O) or phenylthiourea (**2**, X' = S) in ethanol in the presence of ethanolate (Scheme 2).

Pharmacology. An initial screening, performed at 10 μM, using human CB₁ and CB₂ cannabinoid receptors expressed in Chinese hamster ovarian (CHO) cells


* To whom correspondence should be addressed. Phone: +32 2 7647347. Fax: +32 2 7647363. E-mail: lambert@cmfa.ucl.ac.be.

[†] Université Catholique de Louvain.

[#] University of Namur.

[‡] Department of Pharmaceutical Chemistry, University of Jena.

[§] Department of Inorganic and Analytical Chemistry, University of Jena.

Scheme 2^a

^a Reagents and conditions: EtOH, Na, reflux 12 h.

and either [³H]-SR141716 or [³H]-CP-55,940 as radioligands for the CB₁ or CB₂ cannabinoid receptors, respectively, allowed the establishment of some preliminary structure–affinity relationships (Table 1). First, the benzhydryl moiety must be substituted to achieve CB₁ cannabinoid receptor affinity. Indeed compounds **3**, **4**, and **6–9** displaced less than 25% of [³H]-SR141716A specific binding. Furthermore, the preferred benzhydryl substituents are chlorine (**12**) and bromine (**13**). Second, comparing the radioligand displacements obtained with **14–21** revealed that chlorine and bromine are the preferred substituents for the phenyl moiety as well. Compounds **3–21** showed only poor displacement of the CB₂ cannabinoid receptor bound radioligand ([³H]-CP-55,940). *K_i* values of **12–18**, which showed the highest radioligand displacement at 10 μM at the CB₁ cannabinoid receptor, were obtained (Table 1). Substitution of the phenyl moiety (X₂ in Scheme 1) greatly increased the affinity for the CB₁ cannabinoid receptor as illustrated by **12** (*K_i* = 7050 ± 550 nM) and **14** (*K_i* = 1100 ± 100 nM) or by **13** (*K_i* = 2900 ± 250 nM) and **16** (*K_i* = 650 ± 50 nM). With respect to the bromobenzhydryl derivatives (**13**, **16**, **17**, **19–21**), the bromine substituent (X₂ in Scheme 1) proved to be the preferred substituent at the para position of the phenyl ring (**17**, *K_i* = 500 ± 50 nM). Smaller and larger halogen atoms resulted in decreased affinity, as illustrated by **16** (*K_i* = 650 ± 50 nM) and **19** (32 ± 2% of displacement at 10 μM), respectively. Replacing the bromobenzhydryl (X₁ in Scheme 1) by a chlorobenzhydryl (**15**, *K_i* = 1250 ± 100 nM) or an iodobenzhydryl (**18**, *K_i* = 750 ± 100 nM) moiety also decreased the affinity. Four 1-benzhydryl-3-phenylthioureas (**22–25**) were obtained to estimate the effect of the oxygen–sulfur substitution (Table 1). The affinity of the unsubstituted 3-phenyl derivatives was increased by the replacement of the oxygen atom by the sulfur as illustrated by **12** (*K_i* = 7050 ± 550 nM) and **22** (*K_i* = 2900 ± 250 nM) or by **13** (*K_i* = 2900 ± 250 nM) and **23** (*K_i* = 1800 ± 150 nM). However, the affinity of the substituted 3-phenyl derivatives (X₂ in Scheme 1) is decreased for the thio derivatives as shown by **14** (*K_i* = 1100 ± 100 nM) and **24** (*K_i* = 1900 ± 200 nM) or by **16** (*K_i* = 650 ± 50 nM) and **25** (*K_i* = 1500 ± 150 nM). Thus, replacement of the oxygen by a sulfur atom in 1-[bis(4-bromophenyl)methyl]-3-(4-chlorophenyl)urea (**16**) resulted in decreased affinity.

We previously described the structure–affinity relationships at the CB₁ cannabinoid receptor of alkyl and alkylaryl 3-substituted 5,5'-diphenylimidazolidine-2,4-diones¹⁵ and 5,5'-diphenyl-2-thioxoimidazolidin-4-ones¹⁶ as selective CB₁ cannabinoid receptor inverse agonists. Therefore, four 3,5,5'-triphenylimidazolidine-2,4-diones (**26–29**) and two 3,5,5'-triphenyl-2-thioxoimidazolidin-4-one derivatives (**30**, **31**) were synthesized (Scheme 2) and assayed for their CB₁ cannabinoid receptor affinity (Table 1). The four imidazolidine-2,4-diones assayed at

Table 1. Percentage Displacement of [³H]-SR141716A and [³H]-CP-55,940 Specific Binding (10 μM) and Affinities (*K_i*, If Given) for **3–31**^a on hCB₁ and hCB₂ Cannabinoid Receptors^b

| compd | X' | X ₁ | X ₂ | % displacement | | |
|---------------------|----|----------------|--------------------------------|---------------------------|---------------------------|--|
| | | | | hCB ₁ receptor | hCB ₂ receptor | hCB ₁ <i>K_i</i> (nM) |
| 3 | O | H | H | <20 | <10 | ND |
| 4 | O | H | F | 23 ± 2 | <10 | ND |
| 5 | O | H | Cl | 43 ± 2 | <20 | ND |
| 6 | O | H | Br | <20 | <20 | ND |
| 7 | O | H | I | <10 | <10 | ND |
| 8 | O | H | 2-F | <20 | <10 | ND |
| 9 | O | H | OMe | <20 | <20 | ND |
| 10 | O | H | C ₆ H ₁₃ | 55 ± 2 | 24 ± 2 | ND |
| 11 | O | F | H | <20 | <10 | ND |
| 12 | O | Cl | H | 70 ± 3 | 23 ± 2 | 7050 ± 550 |
| 13 | O | Br | H | 70 ± 2 | 38 ± 3 | 2900 ± 250 |
| 14 | O | Cl | Cl | 95 ± 1 | 24 ± 2 | 1100 ± 100 |
| 15 | O | Cl | Br | 80 ± 2 | 29 ± 3 | 1250 ± 100 |
| 16 | O | Br | Cl | 83 ± 1 | 31 ± 2 | 650 ± 50 |
| 17 | O | Br | Br | 75 ± 3 | 22 ± 2 | 500 ± 50 |
| 18 | O | I | Br | 60 ± 2 | 19 ± 2 | 750 ± 100 |
| 19 | O | Br | I | 32 ± 2 | 34 ± 1 | ND |
| 20 | O | Br | CH ₂ OH | 49 ± 2 | <20 | ND |
| 21 | O | Br | C ₆ H ₁₃ | <20 | <20 | ND |
| 22 | S | Cl | H | 77 ± 4 | 23 ± 2 | 2900 ± 250 |
| 23 | S | Br | H | 86 ± 2 | 29 ± 3 | 1800 ± 150 |
| 24 | S | Cl | Cl | 85 ± 1 | 23 ± 2 | 1900 ± 200 |
| 25 | S | Br | Cl | 71 ± 3 | 29 ± 4 | 1500 ± 150 |
| 26 | O | H | H | 28 ± 3 | <20 | ND |
| 27 | O | H | Br | <20 | <20 | ND |
| 28 | O | F | H | 27 ± 3 | <20 | ND |
| 29 | O | Br | H | 29 ± 4 | <20 | ND |
| 30 | S | Br | H | 73 ± 2 | <20 | 3450 ± 300 |
| 31 | S | Br | Cl | 62 ± 2 | <20 | 2750 ± 250 |
| Reference Compounds | | | | | | |
| SR141716A | | | | ND | ND | 5.4 ± 0.2 |
| WIN-552122 | | | | ND | ND | 3800 ± 150 |
| HU-210 | | | | ND | ND | 19 ± 2 |

^a All X₂ substituents are in the para position except that for **8** (X₂ is in ortho). ^b Mean ± SEM of at least four experiments performed in duplicate. The *K_i* values were obtained from non-linear analysis of competition curves using [³H]-SR141716A as radioligand.

10 μM, 3,5,5'-triphenylimidazolidine-2,4-dione (**26**), 3-(4-bromophenyl)-5,5'-diphenylimidazolidine-2,4-dione (**27**), 5,5'-bis(4-fluorophenyl)-3-phenylimidazolidine-2,4-dione (**28**), and 5,5'-bis(4-bromophenyl)-3-phenylimidazolidine-2,4-dione (**29**), displaced less than 30% of the [³H]-SR141716A specifically bound to the CB₁ cannabinoid receptor. The 2-thioxoimidazolidin-4-ones, 5,5'-bis(4-bromophenyl)-3-phenyl-2-thioxoimidazolidin-4-one (**30**) and 5,5'-bis(4-bromophenyl)-3-(4-chlorophenyl)-2-thioxoimidazolidin-4-one (**31**), exhibited *K_i* values of 3450 ± 300 and 2750 ± 250 nM, respectively. When these values are compared with those obtained for the corre-




Figure 1. [^{35}S]-GTP γ S binding stimulation assay of selected compounds and reference cannabinoid ligands (10 μM) on hCB $_1$ cannabinoid receptor. Data are the mean \pm SEM of at least three experiments performed in duplicate. Statistical significance was assessed by one-way ANOVA followed by a Dunnett post-test: (***) $P < 0.01$.

sponding 1-benzhydryl-3-phenylurea and thiourea derivatives, it appears that the orientation of the 1-benzhydryl-3-phenylurea and thiourea phenyl rings imposed by the additional carbonyl led to a decrease in the affinity for the CB $_1$ cannabinoid receptor. This is illustrated by **13** and **29** (70% and 29% of displacement at 10 μM , respectively) or by **25** and **31** (K_i of 1500 \pm 150 and 2750 \pm 250 nM, respectively).

The 1-benzhydryl-3-phenylurea and thiourea derivatives were further characterized in a functional [^{35}S]-GTP γ S assay as previously described.¹⁷ The assay relies on the binding of [^{35}S]-GTP γ S, a radiolabeled nonhydrolyzable GTP analogue, to the G protein upon activation by an agonist of the G-protein-coupled receptor.¹⁸ The assay distinguishes between agonists (increasing nucleotide binding), antagonists (not affecting the binding), and inverse agonists (decreasing nucleotide binding). HU-210, a potent cannabinoid agonist, the inverse agonist SR141716A, and the 1-benzhydryl-3-phenylurea and thiourea derivatives **12–18** and **22–25** were screened at 10 μM . The cannabinoid receptor agonist HU-210 induced an increase in the [^{35}S]-GTP γ S binding (200 \pm 15% compared to basal level), while SR141716A and the tested compounds significantly decreased the nucleotide binding, thus acting as inverse agonists of the human CB $_1$ cannabinoid receptor (Figure 1). For instance, 1-[bis(4-bromophenyl)methyl]-3-(4-chlorophenyl)urea (**16**) and its thio derivative 1-[bis(4-bromophenyl)methyl]-3-(4-chlorophenyl)thiourea (**25**) decreased [^{35}S]-GTP γ S binding by 68 \pm 1% and 61 \pm 1% compared to basal level, respectively. To further explore the inverse agonist properties, the potency of **14–17** and **25** was determined (Table 2). The 1-benzhydryl-3-phenylureas **14–17** and the thiourea derivative **25** dose-dependently decreased the [^{35}S]-GTP γ S binding, showing EC $_{50}$ values of the same magnitude compared to their respective K_i values.

In conclusion, the 1-benzhydryl-3-phenylurea and 1-benzhydryl-3-phenylthiourea derivatives reported here selectively bind to the CB $_1$ cannabinoid receptor acting as inverse agonists. The structure–affinity relationships highlighted the importance of halogen substituents, especially bromine. Thus, 1-[bis(4-bromophenyl)methyl]-3-(4-bromophenyl)urea (**17**) exhibited the highest affinity for the CB $_1$ cannabinoid receptor within this series of compounds. Interestingly, the crucial role of halogen

Table 2. Determination of the Potency (EC $_{50}$) and Percentages of Basal Maximal Stimulation (E_{max}) of [^{35}S]-GTP γ S Binding at hCB $_1$ Cannabinoid Receptors for **14–17**, **25**, HU-210, and SR141716A^a

| compd | EC $_{50}$ (nM) | E_{max} (% above basal) |
|----------------------|-----------------|----------------------------------|
| Urea Derivatives | | |
| 14 | 1050 \pm 100 | -74 \pm 2 |
| 15 | 700 \pm 50 | -66 \pm 1 |
| 16 | 450 \pm 50 | -72 \pm 2 |
| 17 | 500 \pm 50 | -61 \pm 1 |
| Thiourea Derivatives | | |
| 25 | 1250 \pm 100 | -61 \pm 3 |
| Reference Compounds | | |
| SR141716A | 10.1 \pm 0.7 | -84 \pm 2 |
| HU-210 | 0.6 \pm 0.04 | 200 \pm 15 |

^a Mean \pm SEM of at least three experiments performed in duplicate. Statistical significance of [^{35}S]-GTP γ S assay results was assessed using a one-way ANOVA followed by a Dunnett post-test.

substitution of the phenyl rings is also observed in the structure–affinity relationships of SR141716A and other CB $_1$ cannabinoid receptor inverse agonists.¹³ Finally, despite a lower affinity compared to the affinity of the reference inverse agonist SR141716A, the 1-benzhydryl-3-phenylurea structure constitutes a new, interesting template for CB $_1$ cannabinoid receptor inverse agonists. Indeed the noncyclic core scaffold CB $_1$ cannabinoid receptor inverse agonist described herein differs from most of the reported CB $_1$ cannabinoid receptor antagonists/inverse agonists, which are built around a cyclic central unit.

Experimental Section

All reagents, used without further purification, were purchased from Sigma-Aldrich or Acros. Solvents were of analytical grade. A commercial household microwave oven (2.45 GHz) was used. Melting points (uncorrected) were determined in open capillaries using an Electrothermal 9100 apparatus. ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker AM-300 spectrometer at room temperature and analyzed using WIN-NMR software. Chemical shifts (δ) are reported relative to the tetramethylsilane peak set at 0.00 ppm. Signals were abbreviated as follows: s, singlet; d, doublet; t, triplet; m, multiplet. For multiplets, signals are reported as intervals. Coupling constants are expressed in hertz. Mass spectra were recorded on a Finnigan MAT 44S, with an ionization voltage of 70 eV. Elemental analyses were performed on a Carlo Erba EA 1108 analyzer (Carlo Erba, Milano, Italy) and are within $\pm 0.4\%$ of the theoretical values. Infrared (IR) spectra (ν in cm^{-1}) were recorded on a Perkin-Elmer FT-IR 286 spectrometer (Supporting Information).

Synthesis: General Procedure for the Synthesis of 1-Benzhydryl-3-phenylurea Derivatives. Compounds **3**, **5**, **7**, **11–13**, **15**, **17**, and **19** were previously described.¹⁴ The other 1-benzhydryl-3-phenylurea derivatives were similarly obtained. To a solution of benzil (7.15 mmol) and phenylurea (14.3 mmol) in 25 mL of DMSO was added with stirring 25 mL of 1.2 M aqueous KOH. Following 90 s of 750 W microwave irradiation, the mixture was stirred for an additional 5 min. Additional 30 s pulses were applied at 6, 9, 12, 15, 18, 21, 24, and 30 min. Between pulses, the mixture was stirred. After the completion of the sequence, the mixture was poured onto cold water and the resulting precipitate was filtered, dried, and crystallized. All microwave irradiations were carried out in an open system.

1-Benzhydryl-3-(4-fluorophenyl)urea (4). Yield: 52%. Mp 229.1–230.2 $^\circ\text{C}$. MS DEI (Desorption Electron Impact): 320 [M] $^+$. ^{13}C NMR (DMSO- d_6) δ 56.71 (CH), 114.88, 115.20, 119.02, 126.78, 127.43, 128.34, 136.43, 143.02 (C and CH arom), 154.21 (C=O). Anal. (C $_{20}$ H $_{17}$ FN $_2$ O) C, H, N.

1-Benzhydryl-3-(4-bromophenyl)urea (6). Yield: 50%. Mp 233.4–234.1 °C. MS DEI: 381 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 56.97 (CH), 112.68, 119.67, 127.04, 128.59, 131.57, 143.16 (C and CH arom), 154.28 (C=O). Anal. (C₂₀H₁₇BrN₂O) C, H, N.

1-Benzhydryl-3-(2-fluorophenyl)urea (8). Yield: 55%. Mp 229.5–230.2 °C. MS DEI: 320 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 56.87 (CH), 114.58, 114.84, 119.69, 121.44, 121.57, 124.35, 126.81, 126.94, 128.43, 142.99 (C and CH arom), 153.01 (C=O), 153.92.

1-Benzhydryl-3-(4-methoxyphenyl)urea (9). Yield: 51%. Mp 215.0–215.9 °C. MS DEI: 332 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 55.06 (CH₃), 56.74 (CH), 113.87, 119.11, 126.81, 128.36, 133.28, 143.25 (C and CH arom), 153.92 (C=O), 154.44. Anal. (C₂₁H₂₀N₂O₂) C, H, N.

1-Benzhydryl-3-(4-hexylphenyl)urea (10). Yield: 48%. Mp 172.1–173.2 °C. MS DEI: 386 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 13.84, 21.99, 28.14, 31.06, 34.35, and 40.12 (CH₂), 56.74 (CH), 117.56, 126.81, 128.36, 135.05, 137.74, 143.18 (C and CH arom), 154.31 (C=O). Anal. (C₂₆H₃₀N₂O) C, H, N.

1-[Bis(4-chlorophenyl)methyl]-3-(4-chlorophenyl)urea (14). Yield: 46%. Mp >300 °C. MS DEI: 406 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 56.20 (CH), 119.60, 120.70, 125.29, 128.99, 129.70, 131.89, 139.47, 142.38 (C and CH arom), 154.54 (C=O). Anal. (C₂₀H₁₅Cl₃N₂O) C, H, N.

1-[Bis(4-bromophenyl)methyl]-3-(4-chlorophenyl)urea (16). Yield: 49%. Mp >300 °C. MS DEI: 494 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 55.81 (CH), 119.34, 124.97, 128.21, 128.66, 131.90, 139.20, 141.73 (C and CH arom), 154.28 (C=O). Anal. (C₂₀H₁₅Br₂ClN₂O) C, H, N.

1-[Bis(4-iodophenyl)methyl]-3-(4-bromophenyl)urea (18). Yield: 43%. Mp 246.1–247.0 °C. MS DEI: 633 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 56.13 (CH), 93.20, 112.74, 119.73, 129.43, 131.38, 131.51, 137.39, 137.65, 139.59, 142.51 (C and CH arom), 154.21 (C=O). Anal. (C₂₀H₁₅I₂BrN₂O) C, H, N.

1-[Bis(4-bromophenyl)methyl]-3-(4-hydroxymethylphenyl)urea (20). Yield: 52%. Mp 213.4–214.3 °C. MS DEI: 490 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 56.19 (CH), 60.98 (CH₂), 120.31, 121.86, 127.49, 128.07, 129.44, 131.57, 137.91, 142.38 (C and CH arom), 154.67 (C=O). Anal. (C₂₁H₁₈Br₂N₂O₂*1/2H₂O) C, H, N.

1-[Bis(4-bromophenyl)methyl]-3-(4-hexylphenyl)urea (21). Yield: 49%. Mp 202.8–203.3 °C. MS DEI: 544 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 14.08, 18.67, 22.17, 28.37, 31.16, and 34.59 (CH₂), 55.87 (CH), 117.92, 120.31, 128.53, 129.31, 131.57, 135.39, 137.78, 142.31 (C and CH arom), 154.47 (C=O). Anal. (C₂₆H₂₈Br₂N₂O) C, H, N.

Synthesis of 1-Benzhydryl-3-phenylthiourea Derivatives. The procedure described for the synthesis of 1-benzhydryl-3-phenylurea derivatives was used except that phenylthiourea was substituted for phenylurea.

1-[Bis(4-chlorophenyl)methyl]-3-phenylthiourea (22). Yield: 43%. Mp 185.1–185.8 °C. MS DEI: 387 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 59.23 (CH), 122.64, 123.94, 128.27, 128.40, 129.11, 131.76, 139.34, 140.50 (C and CH arom), 180.23 (C=S). Anal. (C₂₀H₁₆Cl₂N₂S) C, H, N.

1-[Bis(4-bromophenyl)methyl]-3-phenylthiourea (23). Yield: 44%. Mp 206.8–207.5 °C. MS DEI: 476 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 59.37 (CH), 120.25, 122.64, 123.94, 128.27, 129.44, 131.31, 139.34, 140.89 (C and CH arom), 180.23 (C=S). Anal. (C₂₀H₁₆Br₂N₂S) C, H, N.

1-[Bis(4-chlorophenyl)methyl]-3-(4-chlorophenyl)thiourea (24). Yield: 41%. Mp 200.8–201.4 °C. MS DEI: 422 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 59.30 (CH), 119.03, 124.20, 127.69, 128.40, 129.12, 131.77, 138.36, 140.37 (C and CH arom), 180.22 (C=S). Anal. (C₂₀H₁₅Cl₃N₂S) C, H, N.

1-[Bis(4-bromophenyl)methyl]-3-(4-chlorophenyl)thiourea (25). Yield: 43%. Mp 230.0–230.3 °C. MS DEI: 511 [M]⁺. ¹³C NMR (DMSO-*d*₆) δ 59.43 (CH), 120.31, 124.20, 127.69, 128.07, 129.05, 131.31, 138.30, 140.67 (C and CH arom), 180.22 (C=S). Anal. (C₂₀H₁₅Br₂ClN₂S) C, H, N.

General Procedure for the Synthesis of 3-Substituted 5,5'-Diphenylimidazolidine-2,4-dione Derivatives. Sodium (1 g) was stirred in ethanol (50 mL) for 15 min before benzil (9.5 mmol) and phenylurea (19 mmol) were added. The

resulting solution was refluxed for 12 h and after cooling was poured onto ice. The precipitate was filtered, dried, and recrystallized from ethanol.

3,5,5'-Triphenylimidazolidine-2,4-dione (26). Yield: 62%. Mp 204.2–204.9 °C. MS DEI: 329 [M + H]⁺. ¹³C NMR (DMSO-*d*₆): δ 69.21 (C), 117.66, 121.24, 126.97, 128.33, 128.59, 128.79, 129.04, 131.82, 139.66 (C and CH arom), 154.28 (C=O), 172.46 (C=O).

3-(4-Bromophenyl)-5,5'-diphenylimidazolidine-2,4-dione (27). Yield: 58%. Mp 214.4–215.1 °C. MS DEI: 408 [M + H]⁺. ¹³C NMR (DMSO-*d*₆): δ 69.65 (C), 121.54, 127.23, 128.79, 129.11, 129.24, 131.44, 132.35, 139.86 (C and CH arom), 154.22 (C=O), 172.52 (C=O). Anal. (C₂₁H₁₅Br₂N₂O₂) C, H, N.

5,5'-Bis(4-fluorophenyl)-3-phenylimidazolidine-2,4-dione (28). Yield: 57%. Mp 242.9–244.0 °C. MS DEI: 364 [M]⁺. ¹³C NMR (DMSO-*d*₆): δ 68.03 (C), 115.27, 115.6, 126.72, 128.07, 128.72, 128.85, 131.44, 135.45, 160.11, 163.34 (C and CH arom), 153.83 (C=O), 172.01 (C=O). Anal. (C₂₁H₁₄F₂N₂O₂) C, H, N.

5,5'-Bis(4-bromophenyl)-3-phenylimidazolidine-2,4-dione (29). Yield: 58%. Mp 162.5–163.7 °C. MS DEI: 486 [M]⁺. ¹³C NMR (DMSO-*d*₆): δ 68.45 (C), 121.94, 126.56, 128.46, 128.96, 129.54, 129.87, 131.78, 138.45 (C and CH arom), 154.05 (C=O), 171.76 (C=O). Anal. (C₂₁H₁₄Br₂N₂O₂) C, H, N.

Synthesis of 3-Substituted 5,5'-Diphenyl-2-thioxoimidazolidin-4-one Derivatives. These derivatives were obtained similarly to the 3-substituted 5,5'-diphenylimidazolidine-2,4-dione derivatives starting from the corresponding phenylthiourea.

5,5'-Bis(4-bromophenyl)-3-phenyl-2-thioxoimidazolidin-4-one (30). Yield: 40%. Mp 293.0–293.9 °C. MS DEI: 502 [M]⁺. ¹³C NMR (DMSO-*d*₆): δ 70.89 (C), 122.26, 128.79, 128.98, 131.83, 132.67, 136.94 (C and CH arom), 172.33 (C=O), 181.00 (C=S). Anal. (C₂₁H₁₄Br₂N₂OS) C, H, N.

5,5'-Bis(4-bromophenyl)-3-(4-chlorophenyl)-2-thioxoimidazolidin-4-one (31). Yield: 41%. Mp >300 °C. MS DEI: 536 [M]⁺. ¹³C NMR (DMSO-*d*₆): δ 71.53 (C), 122.83, 129.37, 131.18, 132.02, 132.41, 134.22, 137.40 (C and CH arom), 172.72 (C=O), 181.19 (C=S). Anal. (C₂₁H₁₃Br₂ClN₂OS) C, H, N.

Competition Binding Assay. CHO cells stably expressing the human CB₁ or the human CB₂ cannabinoid receptors were donated by Drs. M. Detheux and P. Nokin, respectively (Euroscreen s.a., Gosselies, Belgium). Cells membranes were obtained as previously described.¹⁶ Stock solutions of the compounds were prepared in DMSO and further diluted (100×) with the binding buffer to the desired concentration. Under these conditions *B*_{max} was 57 pmol/mg protein and *K*_d was 1.13 ± 0.13 nM for the hCB₁ cannabinoid receptor ([³H]-SR141716A). The competitive binding experiments were performed using [³H]-SR141716A (1 nM, 52 Ci/mol, Amersham, Roosendaal, The Netherlands) or [³H]-CP-55940 (1 nM, 101 Ci/mol, from NEN Life Science, Zaventem, Belgium) as radioligands for the hCB₁ and the hCB₂ cannabinoid receptor, respectively, at 30 °C in plastic tubes, and 40 μg of membranes per tube was resuspended in 0.5 mL (final volume) of binding buffer (50 mM Tris-HCl, 3 mM MgCl₂, 1 mM EDTA, 0.5% bovine serum albumine, pH 7.4). Test compounds were present at varying concentrations, and nonspecific binding was determined in the presence of 10 μM HU-210 (Tocris, Bristol, U.K.). After 1 h of incubation, solutions were rapidly filtered through 0.5% PEI pretreated GF/B glass fiber filters (Whatman, Maidstone, U.K.) on a M-48T Brandell cell harvester and washed twice with 5 mL of ice-cold binding buffer without serum albumin. Radioactivity was measured in a Pharmacia Wallac 1410 β-counter in 10 mL of Aqualuma (PerkinElmer, Schaesberg, The Netherlands) after 10 s of shaking and 3 h of resting. Assays were performed at least in triplicate.

[³⁵S]-GTPγS Assay. Binding experiments were performed at 30 °C in plastic tubes containing 40 μg of protein in 0.5 mL (final volume) of binding buffer (50 mM Tris-HCl, 3 mM MgCl₂, 1 mM EDTA, 100 mM NaCl, 0.1% bovine serum albumin, pH 7.4) supplemented with 20 μM GDP and the appropriate

concentration of test compounds. The assay was initiated by the addition of [³⁵S]-GTP γ S (0.05 nM, final concentration, 1173 Ci/mmol, Amersham, Roosendaal, The Netherlands). Following 1 h of incubation, a total of 5 mL of ice-cold washing buffer (50 mM Tris-HCl, 3 mM MgCl₂, 1 mM EDTA, 100 mM NaCl) was added. The suspension was immediately filtered through GF/B filters using a 48-well Brandell cell harvester and washed twice with the same ice-cold buffer. Radioactivity was counted as mentioned above. Nonspecific binding was measured in the presence of 100 μ M Gpp(NH)p. Assays were performed in triplicate.

Data Analysis. IC₅₀ and EC₅₀ values were determined by nonlinear regression analysis performed using the GraphPad Prism 4.0 software (GraphPad Software, San Diego). The K_i values were calculated on the basis of the Cheng-Prusoff equation: $K_i = IC_{50}/(1 + L/K_d)$.

Acknowledgment. G.G.M. is very indebted to "Fonds pour la formation à la Recherche dans l'Industrie et l'Agriculture (FRIA)" for the award of a research fellowship. This work was partially funded by a FRSM grant (Belgian National Fund for Scientific Research) and a FSR grant (Université catholique de Louvain).

Supporting Information Available: General procedures for the synthesis of benzils, phenylureas, and phenylthioureas, ¹H NMR and IR data, and results from elemental analysis of the synthesized compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Rinaldi-Carmona, M.; Barth, F.; Heaulme, M.; Shire, D.; Candland, B.; Congy, C.; Martinez, S.; Maruani, J.; Neliat, G.; Caput, D. SR141716A, a Potent and Selective Antagonist of the Brain Cannabinoid Receptor. *FEBS Lett.* **1994**, *350*, 240–244.
- Zimmer, A.; Zimmer, A. M.; Hohmann, A. G.; Herkenham, M.; Bonner, T. I. Increased Mortality, Hypoactivity, and Hypoalgesia in Cannabinoid CB₁ Receptor Knockout Mice. *Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96*, 5780–5785.
- Ledent, C.; Valverde, O.; Cossu, G.; Petitot, F.; Aubert, J. F.; Beslot, F.; Bohme, G. A.; Imperato, A.; Pedrazzini, T.; Roques, B. P.; Vassart, G.; Fratta, W.; Parmentier, M. Unresponsiveness to Cannabinoids and Reduced Addictive Effects of Opiates in CB₁ Receptor Knockout Mice. *Science* **1999**, *283*, 401–404.
- Ravinet Trillou, C.; Arnone, M.; Delgorge, C.; Gonalons, N.; Keane, P.; Maffrand, J. P.; Soubrie, P. Anti-Obesity Effect of SR141716, a CB₁ Receptor Antagonist, in Diet-Induced Obese Mice. *Am. J. Physiol.* **2003**, *284*, R345–R353.
- Plummer, C. W.; Finke, P. E.; Mills, S. G.; Wang, J.; Tong, X.; Doss, G. A.; Fong, T. M.; Lao, J. Z.; Schaeffer, M. T.; Chen, J.; Shen, C. P.; Stribling, D. S.; Shearman, L. P.; Strack, A. M.; Van der Ploeg, L. H. Synthesis and Activity of 4,5-Diarylimidazoles as Human CB₁ Receptor Inverse Agonists. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 1441–1446.
- Colombo, G.; Vacca, G.; Serra, S.; Carai, M. A.; Gessa, G. L. Suppressing Effect of the Cannabinoid CB₁ Receptor Antagonist, SR141716, on Alcohol's Motivational Properties in Alcohol-Preferring Rats. *Eur. J. Pharmacol.* **2004**, *498*, 119–123.
- Cohen, C.; Perrault, G.; Voltz, C.; Steinberg, R.; Soubrie, P. SR141716, a Central Cannabinoid (CB₁) Receptor Antagonist, Blocks the Motivational and Dopamine-Releasing Effects of Nicotine in Rats. *Behav. Pharmacol.* **2002**, *13*, 451–463.
- Le Foll, B.; Goldberg, S. R. Cannabinoid CB₁ Receptor Antagonists as Promising New Medications for Drug Dependence. *J. Pharmacol. Exp. Ther.* **2005**, *312*, 875–883.
- Kadoi, Y.; Hinohara, H.; Kunimoto, F.; Kuwano, H.; Saito, S.; Goto, F. Effects of AM281, a Cannabinoid Antagonist, on Systemic Haemodynamics, Internal Carotid Artery Blood Flow and Mortality in Septic Shock in Rats. *Br. J. Anaesth.* **2005**, *94*, 563–568.
- Mascolo, N.; Izzo, A. A.; Ligresti, A.; Costagliola, A.; Pinto, L.; Cascio, M. G.; Maffia, P.; Cecio, A.; Capasso, F.; Di Marzo, V. The Endocannabinoid System and the Molecular Basis of Paralytic Ileus in Mice. *FASEB J.* **2002**, *16*, 1973–1975.
- Cleland, J. G.; Ghosh, J.; Freemantle, N.; Kaye, G. C.; Nasir, M.; Clark, A. L.; Coletta, A. P. Clinical Trials Update and Cumulative Meta-Analyses from the American College of Cardiology: WATCH, SCD-HeFT, DINAMIT, CASINO, INSPIRE, STRATUS-US, RIO-Lipids and Cardiac Resynchronisation Therapy in Heart Failure. *Eur. J. Heart Failure* **2004**, *6*, 501–508.
- Van Gaal, L. F.; Rissanen, A. M.; Sheen, A. J.; Ziegler, O.; Rössner, S. Effects of the Cannabinoid-1 Receptor Blocker Rimonabant on Weight Reduction and Cardiovascular Risk Factors in Overweight Patients: 1-Year Experience from the RIO-Europe Study. *Lancet* **2005**, *365*, 1389–1397.
- Muccioli, G. G.; Lambert, D. M. Current Knowledge on the Antagonists and Inverse Agonists of Cannabinoid Receptors. *Curr. Med. Chem.* **2005**, *12*, 1361–1394.
- Muccioli, G. G.; Wouters, J.; Poupaert, J. H.; Norberg, B.; Poppitz, W.; Scriba, G. K. E.; Lambert, D. M. Versatile Access to Benzhydryl-Phenylureas through an Unexpected Rearrangement during Microwave-Enhanced Synthesis of Hydantoins. *Org. Lett.* **2003**, *5*, 3599–3602.
- Ooms, F.; Wouters, J.; Oscari, O.; Happaerts, T.; Bouchard, G.; Carrupt, P.-A.; Testa, B.; Lambert, D. M. Explorations of the Pharmacophore of 3-Alkyl-5-arylimidazolidinediones as New CB₁ Cannabinoid Receptor Ligands and Potential Antagonists: Synthesis, Lipophilicity, Affinity, and Molecular Modeling. *J. Med. Chem.* **2002**, *45*, 1748–1756.
- Muccioli, G. G.; Martin, D.; Scriba, G. K. E.; Poppitz, W.; Poupaert, J. H.; Wouters, J.; Lambert, D. M. Substituted 5,5'-Diphenyl-2-thioxo-imidazolidin-4-one as CB₁ Cannabinoid Receptor Ligands: Synthesis and Pharmacological Evaluation. *J. Med. Chem.* **2005**, *48*, 2509–2517.
- Govaerts, S. J.; Muccioli, G. G.; Hermans, E.; Lambert, D. M. Characterization of the Pharmacology of Imidazolidinediones Derivatives at Cannabinoid CB₁ and CB₂ Receptors. *Eur. J. Pharmacol.* **2004**, *495*, 43–53.
- Harrison, C.; Traynor, J. R. The [³⁵S]-GTP γ S Binding Assay: Approaches and Applications in Pharmacology. *Life Sci.* **2003**, *74*, 489–508.

JM0503906