

Deoxycholic Acid-Derived Tetraoxane Antimalarials and Antiproliferatives¹

Nataša Terzić,[†] Dejan Opsenica,[†] Dragana Milić,[‡] Bernard Tinant,[§] Kirsten S. Smith,^{||} Wilbur K. Milhous,^{||} and Bogdan A. Šolaja^{*,‡}

Institute of Chemistry, Technology and Metallurgy, Belgrade, Serbia, Laboratoire de Chimie Physique et de Cristallographie, Université Catholique de Louvain, Louvain-la-Neuve, Belgium, Division of Experimental Therapeutics, Walter Reed Army Institute of Research, Washington, D.C. 20307-5100, and Faculty of Chemistry, University of Belgrade, Belgrade, Serbia

Received June 12, 2007

The synthesis of deoxycholic acid (DCA)- and cholic acid (CA)-derived mixed tetraoxanes revealed that *N*-(2-dimethylamino)ethyl derivatives are potent antimalarials in vitro and in vivo. The tetraoxanes presented in this paper are dual inhibitors: besides curing mice in vivo without observed toxic effects, they kill cancer cell lines at very low concentrations. For example, DCA and CA derivatives **16** and **25** cured 3/5 (160 mg/kg/day) and 2/5 (40 mg/kg/day, MTD >960 mg/kg), respectively, and they were extremely active against melanoma LOX IMVI cancer, LC₅₀ = 22 nM and 69 nM, respectively.


Introduction

Malaria was successfully reduced after World War II as a consequence of easy access to the cheap insecticide, dichlorodiphenyltrichloroethane (DDT), as well as inexpensive and readily available drugs such as chloroquine (CQ)^a, mefloquine (MFQ), and quinine. CQ and MFQ were both preceded by quinine, which was the first purified natural product used as a drug (in 1821). Malaria is caused by multiplication of the protozoan parasite *Plasmodium* in erythrocytes and is a major health problem in many tropical and subtropical countries. Of the four species of malaria that cause human disease, *P. falciparum*, *P. vivax*, *P. malariae*, and *P. ovale*, *P. falciparum* is the most lethal strain. The present resurgence of malaria and the lack of proper treatment affect 300–500 million people annually causing over 1.5 million deaths.² The development of resistance to chloroquine has severe health implications for countries in malaria-endemic regions. In a recent genetic study³ of *P. falciparum*, it was found that this species is unexpectedly diverse; another study⁴ points to the multiple independent origins of mutations in one parasite gene that confer resistance to the widely used drug CQ. The results show that, in principle, *P. falciparum* could rapidly develop resistance to multiple drugs (CQ: estimated ~6–30 years), additionally justifying a further search for new drugs.

For some time, our research has focused on the generation of analogs containing a 1,2,4,5-tetraoxacyclohexane antimalarial pharmacophore with a cholic acid (CA)-derived carrier.⁵ On the basis of our accumulated evidence regarding the influence of substitution at the spirocycloalkane moiety, it appears that a methyl group at C(4''), Chart 1 affords the best in vitro and in vivo antimalarial activity.^{5b,d}

With concern of the synthesis of mixed tetraoxanes, four distinct procedures based on coupling of a ketone/protected ketone species to a protected/nonprotected *gem*-dihydroperoxide have been used: (a) coupling of TMS-protected *gem*-dihydroperoxide to an aldehyde (catalyst TMSOTf);⁶ (b) coupling of

Chart 1



nonprotected *gem*-dihydroperoxide to a ketone (catalyst H₂SO₄/CH₃CN);^{6b} (c) one-pot *gem*-dihydroperoxide MTO/TFE mediated preparation and coupling to a ketone (catalyst HBF₄);⁷ and (d) coupling of nonprotected *gem*-dihydroperoxides to acetals derived from a suitable ketone (catalyst BF₃/Et₂O).^{8,9} The best reported yields of desired products were obtained using procedures described in refs 7 and 8 (up to 90%); however, the tetraoxanes with meaningful antimalarial activity were synthesized according to the procedure presented in refs 5b–d and 9, with yields of 18–50%. Our approach to the synthesis of cholic acid-derived tetraoxanes utilizes the coupling of a ketone to steroidal *gem*-dihydroperoxide obtained from a steroidal ketone in high yield and with high purity.

We found that the toxicity of steroidal tetraoxanes against healthy cells (PBMC,^{5a,d} VERO^{5d}) is low as compared with their antimalarial activity (SI = 826–33 000) and to their activity against certain types of cancers.^{5d} In addition, preliminary results on nonhemolytic behavior^{5b} as well as their high maximum tolerated doses (MTD) in an antiproliferative screen (400 mg/kg) and an antimalarial screen (>1960 mg/kg)^{5d} provide a strong rationale for further research in this area.

* Corresponding author. E-mail: bsolaja@chem.bg.ac.yu. Phone: +381-11-263-86-06. Fax: +381-11-263-60-61.

[†] Institute of Chemistry, Technology and Metallurgy.


[‡] University of Belgrade.

[§] Université Catholique de Louvain.

^{||} Walter Reed Army Institute of Research.

^a Abbreviations: CQ, chloroquine; MFQ, mefloquine; ART, artemisinin; CA, cholic acid; DCA, deoxycholic acid; MG_MID, mean graph midpoint.

Scheme 1



Accordingly, in this work, we present the results of our study on the preparation, antimalarial and antiproliferative activity, and in vitro metabolism of deoxycholic acid (DCA)- and CA-derived tetraoxanes. We explore the differences in in vitro antimalarial activity and cytotoxicity between the two series and analyze the obtained in vivo results.

Chemistry

Tetraoxanes **3a–c** were prepared according to our procedure described earlier^{5b–d,10} using HCl and an H₂SO₄/CH₃CN system for bishydroperoxyacetalization and peroxyacetalization reactions, respectively (Scheme 1). *Gem*-dihydroperoxide **2** was obtained in high yield from ketone **1** and was further coupled to cyclohexanones to give **3a** and **3b** + **3c** in the yield of 34% and 52%, respectively. Epimeric tetraoxanes, obtained from 4-methylcyclohexanone, were separated as methyl esters **3b** and **3c**. Esters **3** were further transformed into corresponding amides via acids **4** according to a well-established procedure^{5b–d,10} utilizing mixed anhydride intermediates: **3** → **4** → **5–19**. The overall yield of amides in each series starting from *gem*-dihydroperoxide **2** was 19–39%. The obtained compounds have been fully characterized using standard spectroscopic methods.

The configuration at C(4'') in coupled products could not be determined on the basis of NMR spectral data; however, it was assigned by X-ray crystallographic structural analysis of the corresponding amide **14**, and it appears to be *R* (Figure 1). Consequently, tetraoxanes **3b**, **4b**, **12**, **14**, **16**, **16a**, and **18** were




Figure 1. ORTEP plot of tetraoxane **14**.¹¹ Carbons are represented as partially filled ellipsoids; hydrogens are unfilled circles; oxygens are in red; nitrogens are in blue.

assigned to the same 4''*R* series, while the diastereomeric 4''*S* series consists of compounds **3c**, **4c**, **13**, **15**, **17**, **17a**, and **19**.

The asymmetric part of the unit cell of **14** consists of two independent tetraoxane molecules; there are no cocrystallized solvent molecules. The two tetraoxane molecules have a very similar geometry; only differences in the orientation of the extremities of the C(17) side chain are observed (N(25)–C(26)–C(27)–C(28) = –168° in molecule 1 and –62° in molecule 2). One observed intermolecular hydrogen bonds involving the amide bond. In molecule 1: N(25)–H(25)···O(29) (1 – *x*, 0.5 + *y*, 2 – *z*), H···O 2.11(2) Å, N···O = 2.978(4) Å, N–H···O = 168°. In molecule 2: N(25)–H(25)···O(29) (–*x*, 0.5 + *y*, 2 – *z*), H···O 2.13(2) Å, N···O = 2.984(4) Å, N–H···O = 165°.

The cholic acid-based tetraoxanes (Chart 1) were obtained by an analogous procedure starting from known acids^{5b} and *N,N*-dimethylethan-1,2,-diamine (see Experimental Section).

Antimalarial Activity

The synthesized tetraoxanes **3–27** were screened in vitro against three *P. falciparum* strains: D6 (chloroquine-susceptible), W2 (chloroquine-resistant, susceptible to mefloquine), and TM91C235 (Thailand), a multidrug-resistant strain, following the protocol given in ref 5a. In general, spirocyclohexylidene tetraoxanes of the DCA series (Table 1, R = H, compounds **3a**, **4a**, **5–11**) appear to be less active than their C(4'') methyl analogues (Table 1, R = CH₃). Of the two epimeric C(4'') methyl series, 4''*R* is significantly more active. The above findings both hold true for the CA derivatives as well (Tables 1 and 2, and ref 5b).

Members of the DCA series (Tables 1 and 2, R = H, and R = CH₃ (4''*S*)) are moderately active against D6, W2, and TM91C235 *P. falciparum* strains; however, they are noticeably less active than the corresponding 4''*R* derivatives. In the 4''*R* series, tetraoxane **16** was identified as the most potent in vitro of all of the DCA derivatives. Compound **16** possesses an *N*-(2-dimethylamino)ethyl group that should be protonated under the acidic conditions in parasite's food vacuole and was introduced to increase the concentration of the active compound at the location of action. A few features of its in vitro activity should be noted. First, the compound exerts approximately the same

Table 1. In Vitro Antimalarial Activities of Tetraoxanes **3a–27** against *P. falciparum* D6^a and W2^b Strains

| compd | R = H | | | | compd | R = CH ₃ (4''R) | | | | compd | R = CH ₃ (4''S) | | | |
|------------------------|-----------------------|--------|-----------------------|--------|------------|----------------------------|-------|-----------------------|-------|-----------|----------------------------|--------|-----------------------|--------|
| | IC ₅₀ (nM) | | IC ₉₀ (nM) | | | IC ₅₀ (nM) | | IC ₉₀ (nM) | | | IC ₅₀ (nM) | | IC ₉₀ (nM) | |
| | D6 | W2 | D6 | W2 | | D6 | W2 | D6 | W2 | | D6 | W2 | D6 | W2 |
| 3a | 201.43 | 111.69 | 258.37 | 170.27 | 3b | 39.20 | 34.21 | 26.04 | 49.29 | 3c | 37.95 | 38.30 | 62.88 | 80.30 |
| 4a | 106.73 | 93.86 | 310.14 | 178.18 | 4b | 11.55 | 12.05 | 17.08 | 84.75 | 4c | 19.21 | 42.24 | 31.69 | 89.66 |
| 5 | 32.72 | 19.08 | 50.41 | 35.58 | 12 | 13.62 | 18.76 | 18.98 | 29.72 | 13 | 22.89 | 19.43 | 31.35 | 118.00 |
| 6 | 74.28 | 40.76 | 115.17 | 110.94 | 14 | 9.81 | 10.10 | 20.21 | 30.49 | 15 | 29.29 | 33.83 | 39.26 | 53.70 |
| 7 | 221.21 | 104.87 | ND ^c | 276.13 | 16 | 12.83 | 16.83 | 18.66 | 19.77 | 17 | 104.14 | 135.93 | 148.09 | 155.03 |
| 8 | 133.24 | 81.62 | 195.48 | 126.25 | 18 | 23.32 | 23.10 | 33.69 | 38.90 | 19 | 49.99 | 50.82 | 77.45 | 76.86 |
| 9 | 178.86 | 150.14 | 197.37 | 168.75 | | | | | | | | | | |
| 10 | 105.56 | 55.49 | 131.40 | 78.62 | | | | | | | | | | |
| 11 | 346.75 | 346.75 | ND | ND | | | | | | | | | | |
| 24 | 6.70 | 11.24 | 11.70 | 25.93 | 20 | 5.48 | 7.86 | 8.07 | 10.11 | 21 | 19.78 | 23.10 | 29.20 | 54.61 |
| 25 | 28.18 | 14.49 | 45.14 | 51.13 | 22 | 18.63 | 16.67 | 27.78 | 24.80 | 23 | 43.82 | 43.54 | 71.16 | 94.02 |
| 26^d | 11.83 | 4.74 | ND | ND | | | | | | | | | | |
| 27 | 3547 | 1413 | | | | | | | | | | | | |
| ART^e | 9.0 | 6.7 | 12.8 | 11.5 | MFQ | 10.52 | 3.09 | 21.65 | 8.41 | CQ | 13.72 | 349.35 | 17.63 | 491.53 |

^a *P. falciparum* African D6 clone. ^b *P. falciparum* Indochina W2 clone. ^c ND, not determined. ^d Taken from ref 1b. ^e Average of greater than eight replicates.

Table 2. In Vitro Antimalarial Activities of Tetraoxanes **3a–27** against *P. falciparum* TM91C235^a Strain

| compound | R = H | | compound | R = CH ₃ (4R) | | compound | R = CH ₃ (4''S) | |
|------------------------|-----------------------|-----------------------|------------|--------------------------|-----------------------|-----------|----------------------------|-----------------------|
| | IC ₅₀ (nM) | IC ₉₀ (nM) | | IC ₅₀ (nM) | IC ₉₀ (nM) | | IC ₅₀ (nM) | IC ₉₀ (nM) |
| 3a | ND ^b | ND | 3b | 42.40 | 74.87 | 3c | 50.78 | 77.27 |
| 4a | ND | ND | 4b | 19.90 | 47.02 | 4c | 31.76 | 93.05 |
| 5 | 40.94 | 76.72 | 12 | 15.74 | 30.79 | 13 | 27.96 | 53.46 |
| 6 | ND | ND | 14 | 12.35 | 23.37 | 15 | 34.12 | 50.74 |
| 7 | 221.21 | ND | 16 | 14.72 | 24.25 | 17 | 135.85 | 153.01 |
| 8 | ND | ND | 18 | 32.87 | 42.71 | 19 | 66.75 | 108.96 |
| 9 | 168.70 | 202.58 | | | | | | |
| 10 | 84.74 | 124.68 | | | | | | |
| 11 | 346.75 | | | | | | | |
| 24 | 10.26 | 27.78 | 20 | 6.24 | 11.02 | 21 | 27.17 | 69.71 |
| 25 | 19.65 | 37.85 | 22 | 12.24 | 18.47 | 23 | 39.17 | 102.04 |
| 27 | 3547 | | | | | | | |
| ART^c | 13.04 | 17.40 | MFQ | 16.44 | 65.71 | CQ | 144.90 | 268.76 |

^a *P. falciparum* multidrug resistant TM91C235 strain (Thailand). ^b Not determined. ^c Average of greater than eight replicates.

activity against both D6 and W2 (Table 1, IC₉₀ ~ 19 nM), and at the same time, it shows a small IC₉₀/IC₅₀ (W2) ratio of 1.23 (as compared with 1.72 (W2) of artemisinin and compared with 2.15 (W2) of mefloquine). Second, compound **16** was 10 times more active than CQ against the multidrug-resistant strain TM91C235, with an IC₉₀ = 24.25 nM, and its activity was comparable to that of artemisinin (IC₉₀ = 17.40 nM, Table 2). In addition, of the three *N*-(2-dimethylamino)ethyl CA derivatives **22**, **23**, and **25** that have been prepared, the 4''R derivative **22** was again the most potent in the series (Tables 1 and 2, Chart 1). The anilides of CA series, **20**, **21**, and **24**, exert even higher activity when compared with the related *N*-(2-dimethylamino)ethyl derivatives, as well as to the corresponding DCA-derived tetraoxanes **8**, **18**, and **19**. Poor activity of tetraoxanes **9** and **10** clearly supports our earlier finding¹² that tertiary amides do not confer any appreciable antimalarial activity, while tetraoxane **27** is totally inactive in the in vitro antimalarial screen. The observed inactivity of fullerene derivative **27** indicates that the fullerene part acts as a radical sponge.¹³ It is feasible to envisage that radicals generated upon reaction of tetraoxane moiety with Fe(II) were trapped by the fullerene part of the molecule within the cell.

It became obvious that the additional C(7) acetyloxy group of cholic acid confers significant in vitro antimalarial activity to the analogues and also plays an important role in our pharmacophore model.¹⁴ In order to better comprehend the activity difference between the CA and the DCA series, we submitted amides **16**, **22**, **23**, and **25** to mice infected with *P. berghei* using a modified Thompson test. In addition, in vitro

metabolism studies were performed for the same compounds to assess the bioavailability of drug candidates upon oral administration. Metabolic stability assays and metabolite identification were done using human and mouse liver microsomes to help gauge the first-pass metabolism of the drug candidates in relevant species.^{5d}

The mice were infected on day 0, and the tested compounds were administered orally on days 3, 4, and 5 postinfection (Table 3). The data showed that cyclohexylidene CA-derived tetraoxane **25** (Chart 1, R = H, X = NHCH₂CH₂N(CH₃)₂) cured all mice at 320 mg/kg/day without any parasitemia on day 31. Compound **25** also cured 3/5 and 2/5 mice at the lower doses of 80 and 40 mg/kg/day, respectively. Having the quite active *N*-(2-dimethylamino)ethyl derivative **25** and being limited by the quantity of substances, we tested **16**, **22**, and **23**, at doses ≤ 320 mg/kg/day. DCA derivative **16** cured 3/5 mice at 160 mg/kg/day; however, the activity sharply declined when the lower dose of 40 mg/kg/day was administered (0/5 cured and no delay in patency). Of the two epimeric CA-based tetraoxanes **22** and **23**, the former 4''R is more active with a minimal curative dose of 40 mg/kg/day.

Antiproliferative Activity

Fifteen tetraoxanes were chosen by NIH–NCI for in vitro screening using a diverse panel of ~60 human cancer cell lines starting at a concentration of 10⁻⁴ M.¹⁵ The results, summarized in Table 4, reveal that most compounds tested in antiproliferative screen demonstrated considerable growth inhibition (GI₅₀ < 100 μM; 50% growth inhibitory activity) against nearly all of the

Table 3. Activity of Tetraoxanes **16**, **22**, **23**, and **25** against *P. Berghei* in Vivo (po)^a

| compound | mg/kg/day | mice dead/day died | mice alive day 31/ total | survival time (days) ^b | met. stab., <i>t</i> _{1/2} (min) | metabolite identity |
|--------------------------------|-----------|-----------------------|-----------------------------|-----------------------------------|--|---------------------|
| 16 (4''R) | 160 | 2/27 | 3/5 | 29 | human, 110 | hydroxylation (1) |
| | 40 | 2/8, 2/9, 1/10 | 0/5 | 9 | mouse, >120 | dihydroxylation (1) |
| 22 (4''R) | 40 | 1/9, 1/13, 1/19, 1/24 | 1/5 | 19 | human, 42 | hydroxylation (1) |
| | 20 | 3/8, 1/9, 1/10 | 0/5 | 9 | mouse, 77 | dehydration (2) |
| 23 (4''S) | 320 | 1/12, 1/14, 2/15 | 1/5 | 17 | human, >120 | hydroxylation (1) |
| | 80 | 1/8, 1/9, 2/10, 1/16 | 0/5 | 11 | mouse, >120 | dihydroxylation (1) |
| 25 | 320 | | 5/5 | 31 | human, 40 | dehydration |
| | 80 | 1/13, 1/20 | 3/5 | 25 | | none detected |
| | 40 | 1/11, 1/13, 1/17 | 2/5 | 21 | mouse, 40 | hydroxylation (2) |
| | 10 | 1/7, 4/8 | 0/5 | 8 | | dihydroxylation (3) |
| infected controls ^c | 0 | 7–9 | 0/5 | | | hydroxylation (1) |

^a Groups of five *P. berghei* (KBG 173 strain) infected CD-1 mice were treated on days 3, 4, and 5 postinfection with tetraoxanes suspended in 0.5% hydroxyethylcellulose–0.1% Tween 80. Mice alive on day 31 with no parasites in a blood film are considered cured. ^b Including cured mice. ^c All non-infected age controls survived (5/5).

Table 4. Summary of the NCI–DTP 60-Cell-Line Screening for Tetraoxanes **6**, **7**, **12**, **15–17a**, **19–23**, **25**, and **26**

| compd | no. of cell lines ^a | no. of cell lines with GI ₅₀ < 100 μM ^b | range of GI ₅₀ (μM) |
|------------|--------------------------------|---|--------------------------------|
| 6 | 55 | 52 | 66.10 to 0.11 |
| 7 | 53 | 53 | 23.40 to <0.01 |
| 12 | 54 | 53 | 2.34 to 0.11 |
| 15 | 51 | 51 | 35.48 to 0.062 |
| 16 | 51 | 51 | 2.63 to <0.0050 |
| 16a | 59 | 59 | 13.18 to 0.015 |
| 17 | 51 | 51 | 9.33 to <0.0050 |
| 17a | 58 | 58 | 2.69 to 0.13 |
| 19 | 53 | 14 | 39.81 to <0.0050 |
| 20 | 44 | 29 | 54.95 to 0.16 |
| 21 | 49 | 35 | 67.60 to 0.18 |
| 22 | 49 | 49 | 3.02 to 0.18 |
| 23 | 48 | 45 | 12.88 to 0.057 |
| 25 | 53 | 53 | 19.95 to <0.01 |
| 26 | 54 | 54 | 12.30 to <0.01 |

^a Cell lines for which results were reported by NIH–NCI. ^b GI₅₀ refers to 50% growth inhibitory activity.

cancer cell lines reported by NIH–NCI. It is of particular importance that each of the 15 tested tetraoxanes inhibited at least 1 of the cancer cell lines on a submicromolar scale, occasionally at 10 nM, and 3 times even at <5 nM. As expected, the screened compounds exhibited variable growth inhibition and cytotoxicity against different cancer cell lines. A more stringent analysis is shown in Table 5, in which only the results satisfying the following criteria are presented: GI₅₀ < 100 nM or/and TGI (total growth inhibition) < 1 μM or/and LC₅₀ < 6 μM. The assessed antiproliferative activity, expressed as GI₅₀, TGI, LC₅₀ (concentration of the compound at which 50% of the cells are killed) were obtained applying the 48 h continuous drug exposure protocol using the SRB (sulforhodamine B) protein assay.¹⁵ The most susceptible cell lines were melanoma (LOX IMVI), renal (UO-31), and non-small cell lung cancer (HOP-92/62). Out of 80 entries, eight possess GI₅₀ values below 10 nM, and in 23 instances, GI₅₀ values <100 nM were observed (Table 5). TGI at concentrations <800 nM was observed on 44 instances, and on 7 instances ≤20 nM. The LC₅₀ values confirm the potency of our compounds with 26 instances of activity <1 μM. The obtained results indicate that our compounds exhibit very high antiproliferative activity in a dose-dependent manner.

The two most active compounds, **16** and **25**, possessed an *N*-(2-dimethylamino)ethyl terminus (Table 5). Deoxycholic derivative **16** totally arrests cancer cell growth (TGI) between

0.4 and 0.007 μM, and it is most active against colon cancer and melanoma cell lines with LC₅₀ = 47 nM and 22 nM, respectively (Figure 2). Its high general toxicity against cancer cells is reflected by an LC₅₀ MG-MID value of –5.39 (mean graph midpoint, see Supporting Information for details).

Cholic acid-derived tetraoxane **25**, which also possesses an *N*-(2-dimethylamino)ethyl terminus, most efficiently inhibits melanoma, NSCL, colon, CNS, and renal cancer cells in the submicromolar range (Table 5). The compound was most active against NSCL HOP-62 and melanoma M14 cancer cells with LC₅₀ values of 83 nM and 69 nM, respectively (Figure 3a). Highly specific compounds are the *N*-(2-dimethylamino)ethyl-containing compound **17** and the primary amide **26**. Tetraoxane **17** totally arrests melanoma LOX IMVI cell growth at ~5.2 nM (TGI) and kills them with LC₅₀ ~56 nM (log LC₅₀ = –7.25; MG-MID = –4.65); **26** is highly specific against renal UO-31 cancer cells (Figure 3b,c) with LC₅₀ = 40 nM (log LC₅₀ = –7.40; MG-MID = –4.88).

Discussion

In this report, we present the results of an extensive antimalarial and antiproliferative study of the cholic and deoxycholic acid-derived tetraoxanes seen in Scheme 1 and Chart 1. When tested in vitro against the three *P. falciparum* strains, D6, W2, and TM91C235, the compounds showed moderate to pronounced IC₉₀ activity, with several of them being equally or more active than artemisinin. In general, CA derivatives were more active than related DCA compounds in both the in vitro (**7**, **16**, **17** vs **25**, **22**, **23**, Tables 1 and 2) and the in vivo (**16** vs **22**, Table 3) antimalarial screens. Tetraoxane **22** cured one mouse of five at a dose of 40 mg/kg/day, as compared with none cured using the same dose of **16**. In addition, although all compounds were relatively stable in the presence of human and mouse liver microsomes, with in vitro *t*_{1/2} > 30 min, the more active CA derivative **22** appears considerably less stable metabolically than derivative **16**. Half-lives for **22** were 42 min (human) and 77 min (mouse), in comparison with 110 and >120 min for **16**, respectively (Table 3). In addition, the observed greater in vivo activity of the (4''R)-methyl compound over the (4''S)-CA analogue (**22** vs **23**) is in accordance with the respective in vitro and in vivo results (Tables 1, 2, and 3), and as above, the less active compound **23** is considerably more stable metabolically than its epimer




Table 5. In Vitro Antiproliferative Activities of Fifteen Tetraoxanes against Selected Cell Lines

| compound (MG_MID _{TGI}) | cell line | | activity after 48 h (μM) | | |
|--------------------------------------|----------------------------|-----------------|---------------------------------------|------------------|-------------------------------|
| | | | GI ₅₀ ^a | TGI ^b | LC ₅₀ ^c |
| 6 (-4.64) | leukemia | RMPI-8226 | 0.11 | 0.65 | 6.82 |
| 7 (-5.01) | leukemia | MOLT-4 | <0.010 | >100 | |
| | non-small cell lung cancer | NCI-H460 | 0.536 | 1.90 | 4.70 |
| | renal cancer | 786-0 | 0.159 | 0.298 | 0.559 |
| 12 (-5.43) | leukemia | HL-60(TB) | 0.769 | 1.84 | 4.41 |
| | non-small cell lung cancer | NCI-H23 | 0.660 | 1.71 | 3.39 |
| | CNS cancer | SF-295 | 0.524 | 1.69 | 3.46 |
| | | SF-539 | 0.933 | 1.84 | 3.61 |
| | melanoma | LOX IMVI | 0.111 | 0.259 | - |
| | renal cancer | A498 | 0.923 | 1.92 | 4.00 |
| | | SN12C | 0.833 | 1.58 | 3.00 |
| | prostate cancer | DU-145 | 0.699 | 1.73 | 3.34 |
| | breast cancer | MDA-MB-231/ATCC | 0.821 | 1.76 | 3.77 |
| | | MDA-MB-435 | 0.951 | 1.94 | 3.96 |
| 15 (-5.06) | non-small cell lung cancer | HOP-92 | 0.134 | 0.873 | 4.53 |
| | melanoma | LOX IMVI | 0.116 | 0.391 | 1.68 |
| | renal cancer | CAKI-1 | 0.0622 | 0.163 | 0.429 |
| | | UO-31 | 0.122 | 0.329 | - |
| | prostate cancer | DU-145 | 0.663 | 1.61 | 3.91 |
| 16 (-6.0) | leukemia | CCRF-CEM | 0.056 | 0.17 | 0.50 |
| | non-small cell lung cancer | HOP-62 | 0.054 | 0.23 | 1.16 |
| | colon cancer | COLO 205 | 0.083 | 0.02 | 0.49 |
| | | HCC-2998 | 0.077 | 0.21 | 0.98 |
| | | HCT-116 | 0.0090 | 0.021 | 0.047 |
| | | HCT-15 | 0.094 | | |
| | CNS cancer | U251 | 0.026 | 0.087 | |
| | melanoma | LOX IMVI | <0.005 | 0.0074 | 0.022 |
| | | M14 | 0.078 | 0.019 | 0.46 |
| | renal cancer | 786-0 | 0.060 | 0.15 | 0.39 |
| | breast cancer | T-47D | 0.13 | 0.41 | |
| 16a (-5.14) | leukemia | CCRF-CEM | 0.257 | 0.539 | >10 |
| | | HL-60(TB) | 0.377 | 1.01 | 6.09 |
| | non-small cell lung cancer | HOP-92 | 0.228 | 0.471 | 0.973 |
| | melanoma | LOX IMVI | 0.184 | 0.378 | 0.774 |
| | renal cancer | 786-0 | 0.626 | 2.04 | 5.43 |
| | | UO-31 | 0.206 | 0.421 | 0.864 |
| 17 (-5.62) | melanoma | LOX IMVI | <0.0050 | 0.00518 | 0.0564 |
| | renal cancer | 786-0 | 0.0813 | 0.149 | |
| | | ACHN | 0.439 | 1.05 | 2.30 |
| | | CAKI-1 | 0.602 | 1.40 | 3.25 |
| 17a (-5.9) | leukemia | CCRF-CEM | 0.186 | 0.384 | 0.792 |
| | | HL-60(TB) | 0.264 | 0.477 | 0.863 |
| | | RPMI-8226 | 0.292 | 0.826 | 4.80 |
| | non-small cell lung cancer | HOP-92 | 0.171 | 0.368 | 0.794 |
| | melanoma | LOX IMVI | 0.180 | 0.352 | 0.689 |
| | ovarian cancer | IGROV1 | 0.160 | 0.338 | 0.712 |
| | renal cancer | 786-0 | 0.335 | 1.05 | 3.65 |
| | | UO-31 | 0.165 | 0.318 | 0.613 |
| | prostate cancer | PC-3 | 0.367 | 1.62 | 4.45 |
| | | DU-145 | 0.641 | 1.91 | 4.49 |
| 19 (-4.37) | non-small cell lung cancer | HOP-92 | <0.005 | 0.13 | 15.20 |
| 20 (-4.29) | melanoma | LOX IMVI | 0.161 | 0.417 | 1.89 |
| 22 (-5.45) | leukemia | CCRF-CEM | 0.258 | 0.725 | |
| | non-small cell lung cancer | NCI-H322M | 1.46 | 2.79 | 5.34 |
| | melanoma | LOX IMVI | 0.183 | | |
| | ovarian cancer | OVCAR-3 | 1.19 | 2.47 | 5.11 |
| | renal cancer | CAKI-1 | 1.37 | 2.69 | 5.29 |
| | breast cancer | MDA-MB-435 | 1.53 | 2.91 | 5.52 |
| 21 (-4.46) | melanoma | LOX IMVI | 0.181 | 0.370 | 0.758 |
| | non-small cell lung cancer | A549/ATCC | 0.255 | 0.664 | |
| | | HOP-92 | 0.266 | 0.766 | 8.84 |
| | renal cancer | UO-31 | 0.222 | 0.487 | >10 |
| 23 (-5.55) | leukemia | CCRF-CEM | 0.067 | 0.151 | 0.340 |
| | | HOP-92 | 0.070 | 0.695 | 2.86 |
| | non-small cell lung cancer | NCI-H23 | 0.318 | 1.69 | 3.11 |
| | melanoma | LOX IMVI | 0.057 | 0.130 | 0.295 |
| | renal cancer | ACHN | 0.592 | 1.25 | 2.64 |
| | | SN12C | 0.606 | 1.28 | 2.72 |
| 23 (-5.55) | breast cancer | MDA-MB-435 | 0.650 | 1.34 | 2.78 |
| 25 (-5.31) | non-small cell lung cancer | HOP-62 | <0.01 | 0.0166 | 0.083 |
| | colon cancer | HCT-116 | 0.0831 | 1.69 | 4.62 |
| | CNS cancer | SF-268 | 0.785 | 2.25 | 5.35 |
| | melanoma | LOX IMVI | <0.01 | 0.0224 | 0.0688 |
| | | M14 | 0.0440 | 0.613 | 2.92 |

Table 5. Continued

| compound (MG_MID (TGI)) | cell line | | activity after 48 h (μM) | | |
|----------------------------|----------------------------|----------|---------------------------------------|------------------|-------------------------------|
| | | | GI ₅₀ ^a | TGI ^b | LC ₅₀ ^c |
| 26 (-5.39) | renal cancer | 786-0 | 0.783 | 2.09 | 4.57 |
| | non-small cell lung cancer | NCI-H23 | 0.55 | 2.07 | 5.28 |
| | | NCI-H522 | 0.23 | 0.77 | 3.02 |
| | | IGROV1 | 0.032 | 0.12 | 0.43 |
| | ovarian cancer | CAKI-1 | 0.20 | 1.12 | 3.59 |
| | renal cancer | UO-31 | <0.01 | 0.014 | 0.040 |

^a 50% growth inhibitory activity. ^b Total growth inhibition. ^c Concentration of the compound at which 50% of the cells are killed.



c)

Figure 2. Inhibition of cancer cell growth 48 h after being exposed to tetraoxane deoxycholic derivative **16**. (a) All cancer cell lines, (b) colon cancer cells, and (c) melanoma cells.

22. These results collectively suggest that CA is favored over DCA as a potential tetraoxane antimalarial carrier.

The most active compound *in vivo* of all tested *N*-(2-dimethylamino)ethyl derivatives was cyclohexylidene derivative **25** (Chart 1, R = H, X = NHCH₂CH₂N(CH₃)₂), which has an






Figure 3. Inhibition of cancer cell growth 48 h after being exposed to tetraoxane cholic acid derivatives **25** and **26**. (a) All cancer cell lines as inhibited by compound **25**, (b) all cancer cell lines as inhibited by compound **26**, and (c) renal cancer cell lines inhibited by tetraoxane **26**.

MCD \leq 40 mg/kg/day. Although **25** is the most metabolically labile of the tested analogues ($t_{1/2}$ = 40 min, human, mouse) and more than three metabolites were generated in human and mouse microsomal preparations, it is still a relatively stable compound, particularly in comparison with artemisinins. Furthermore, no toxicity was observed during *in vivo* screening,

and the tetraoxane moiety itself is metabolically stable (no peroxide bond scission was observed in in vitro ADME studies).¹⁶ This indicates that the given tetraoxanes, their metabolic products, and their reductive scission products (upon exerting lethal effect on the parasite) were nontoxic to experimental animals (mice), unlike artesunate and some artemisinin-derived trioxane dimers.¹⁷ This is of particular importance since the investigated compounds were shown to have dual activity, both antimalarial and antiproliferative.¹⁸ Details are given in Table 3 and in the text above; however, it is important to add that the essentially nontoxic tetraoxane **25** (MTD > 960 mg/kg; MCD ≤ 40 mg/kg/day) kills two cancer cell lines (NSCL cancer HOP-62 and melanoma LOX IMVI) at LC₅₀ ~70–80 nM and totally inhibits the cancer growth of the same cells at concentrations as low as 17–22 nM. Along the same lines is DCA derivative **16** (MTD > 480 mg/kg, 3/5 mice cured): it totally inhibits the cancer growth (TGI) of melanoma LOX IMVI cells at 7.4 nM, kills them at LC₅₀ = 22 nM, and at the same time, it is toxic to colon cancer HCT-116 cells (LC₅₀ = 47 nM). The current portfolio of potent antiproliferatives is complemented with primary amide **26**, which is specifically very potent against renal UO-31 cancer cells (LC₅₀ = 40 nM) as well as tetraoxane **17**, which, like its C(4'') epimer **16**, is specifically active against LOX IMVI cells. It is difficult to and it would be speculative to compare the anticancer potential of our tetraoxanes to compounds tested on different cell lines;¹⁹ however, one may assume on the basis of the same test that most of our compounds shown here and elsewhere¹⁸ are considerably more potent than artemisinin (NSC 369397),¹⁵ artemether (NSC 665970),¹⁵ and certain deoxoartemisinin derivatives.²⁰ For example, all 15 compounds given in Table 5 have average MG_MID_{TGI} = -5.22 while corresponding values for artemisinin and artemether are -4.00087 and -4.0044, respectively. In terms of the mode of action, we speculate that tetraoxanes may exert their antitumor activity by acting as an Fe(II)-sensitized radical source that induces apoptosis,¹² in a similar fashion to the artemisinins.^{19,21,22}

To conclude, we presented the results of a study that demonstrated that DCA and CA tetraoxane derivatives are potent nontoxic antimalarials and antiproliferatives. We showed that the antimalarial potency of a mixed (deoxy)cholic acid-based tetraoxane both in vitro and in vivo depends on the stereochemistry of the substitution as well as the pattern at C(4''), which is in line with previous docking calculations.¹⁴ In addition, on the basis of in vitro and in vivo results, as well as on our pharmacophore model,¹⁴ the conclusion may be drawn that a CA-derived carrier is more effective than a DCA one.

Experimental Section^{5b,5d}

Methyl 3,3-Dihydroperoxy-12 α -acetoxy-5 β -cholan-24-oate (2). Ketone **1** (5 g, 9.91 mmol) was dissolved at room temperature (r.t.) in a CH₃CN/CH₂Cl₂ mixture (200 mL, 3:1) followed by 30% H₂O₂ (10.3 mL, 0.1 mol) and a few drops of concd HCl. The reaction mixture was stirred for 2 h at r.t., quenched with water (20 mL), and was worked-up in the usual manner. The obtained crude product (5.15 g, 94%; colorless foam) was used in the following step. An analytical sample was obtained after column chromatography (Lobar B, LichroPrep Si 60, eluent heptane/EtOAc (7:3)). Colorless foam, softness at 68–70 °C. [α]_D²⁰ = +73.8 (*c* = 0.168, CHCl₃). Anal. (C₂₇H₄₄O₈) C, H.

General Procedure for Preparation of Mixed Tetraoxanes 3. To a cold (0 °C) solution of dihydroperoxide **2** (443.9 mg, 0.9 mmol) and ketone (1.80 mmol) in toluene (12.5 mL), 650 μ L of an ice-bath cooled H₂SO₄:CH₃CN mixture (1:10, v/v) was added dropwise. The reaction mixture was stirred at 0 °C for 15 min, and

after the usual workup, the crude product was purified by column chromatography to afford tetraoxane **3**.

Methyl 12 α -Acetoxy-5 β -cholan-24-oate-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (3a). Chromatography: dry-flash SiO₂, eluent heptane/EtOAc (8.5:2.5) and Lobar LichroPrep RP-18, eluent MeOH/H₂O (97.5/2.5). Yield 178 mg (34%). Colorless foam softens at 74–76 °C. [α]_D²⁰ = +65.7 (*c* = 0.07, CHCl₃). Anal. (C₃₃H₅₂O₈ × H₂O) C, H.

Methyl 12 α -acetoxy-5 β -cholan-24-oate-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4''R)- and (4''S)-methylcyclohexane (3b and 3c). According to the general procedure for mixed tetraoxane **3**, *gem*-dihydroperoxide **2** (8.46 g, 17.0 mmol) was transformed into **3b** and **3c**. The crude reaction mixture was purified by column chromatography (dry-flash SiO₂, eluent heptane: EtOAc = 8.5/1.5 and Lobar B, LichroPrep RP-18, eluent MeOH/H₂O = 97/3). Isomers were separated by column chromatography on Lobar B, LichroPrep Si 60, eluent heptane: EtOAc = 95/5. **3b** (4''R): Yield 2.96 g (29%). Colorless foam softens at 78–80 °C. [α]_D²⁰ = +79.2 (*c* = 0.048, CHCl₃). Anal. (C₃₄H₅₄O₈) C, H. **3c** (4''S): Yield 2.36 g (23%). Colorless foam softens at 80–83 °C. [α]_D²⁰ = +73.8 (*c* = 0.084, CHCl₃). Anal. (C₃₄H₅₄O₈ × 0.5 H₂O) C, H.

12 α -Acetoxy-5 β -cholan-24-oic Acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (4a). Methyl ester **3a** (3.65 g, 6.33 mmol) was hydrolyzed at 80 °C with NaOH (380 mg, 9.5 mmol) in *i*-PrOH/H₂O mixture (160 mL, 3:1 v/v). After 15 min, the reaction was cooled and diluted with 50 mL H₂O and 100 mL CH₂Cl₂. The water layer was acidified to pH 2 with diluted HCl, and layers were separated. The water layer was further extracted with CH₂Cl₂ (3 × 30 mL), the combined organic layers were washed with water and brine, dried over anhyd Na₂SO₄, and evaporated to dryness. Yield 3.28 g (92%). Colorless foam softens at 99–102 °C. [α]_D²⁰ = +71.25 (*c* = 0.08, CHCl₃). Anal. (C₃₂H₅₀O₈ × 0.5 H₂O) C, H.

12 α -Acetoxy-5 β -cholan-24-oic Acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4''R)-methylcyclohexane (4b). Using the same procedure as described for **4a**, we hydrolyzed methyl-ester **3b** (1.5 g, 2.54 mmol) to 1.45 g (99%) **8b**. Colorless foam softens at 115–117 °C. [α]_D²⁰ = +80.8 (*c* = 0.078, CHCl₃). Anal. (C₃₂H₅₂O₈ × 0.5 H₂O) C, H.

12 α -Acetoxy-5 β -cholan-24-oic Acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4''S)-methylcyclohexane (4c). Using the same procedure as described for **4a**, we hydrolyzed methyl-ester **3c** (1.5 g, 2.54 mmol) to **4c** 1.45 g (98%) **8c**. Colorless foam softens at 119–122 °C. [α]_D²⁰ = +67.3 (*c* = 0.11, CHCl₃). Anal. (C₃₂H₅₂O₈ × 0.5 H₂O) C, H.

General Procedure for Preparation of Amides 5–19. A solution of **4a** (341.4 mg, 0.6 mmol) in dry CH₂Cl₂ (20 mL), with added Et₃N (84 μ L, 0.6 mmol) and ClCO₂Et (58 μ L, 0.6 mmol), was stirred for 60 min at 0 °C. The given amount of amine was added, and after 30 min of stirring, the reaction mixture was warmed to r.t. After 90 min, it was diluted with H₂O; the layers were separated, and the reaction mixture was worked-up in a usual manner. Crude product was purified by column chromatography.

12 α -Acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (5). Using a suspension of 10 equiv of NH₄Cl and 10 equiv of Et₃N in dry CH₂Cl₂ (20 mL), we obtained 287 mg (84%) of **5**. Column chromatography: Lobar B, LichroPrep Si 60, eluent EtOAc. Colorless foam softens at 106–108 °C. [α]_D²⁰ = +77.5 (*c* = 0.08, CHCl₃). Anal. (C₃₂H₅₁NO₇) C, H.

***N*-(*n*-Propyl)-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (6).** Using a 6 equiv of *n*-PrNH₂ (0.27 mL, 3.30 mmol) in dry CH₂Cl₂ (20 mL), we obtained 270.3 mg (82%) of **6**. Column chromatography: Lobar B, LichroPrep, eluent EtOAc/heptane = 7/3. Colorless foam softens at 87–89 °C. [α]_D²⁰ = +76.4 (*c* = 0.072, CHCl₃). Anal. (C₃₅H₅₇NO₇ × H₂O) C, H.

***N*-(2-Dimethylamino)ethyl-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (7).** Using 6 equiv of *N,N*-dimethyl-ethylenediamine (0.34 mL,

3,16 mmol) in dry CH_2Cl_2 (20 mL), we obtained 262.9 mg (78%) of **7**. Preparative TLC on SiO_2 : eluent $\text{CHCl}_3/\text{MeOH}/\text{NH}_3 = 9/1/1$. Colorless foam softens at 64–66 °C. $[\alpha]_D^{20} = +68.7$ ($c = 0.064$, CHCl_3). HRMS: m/z 633.4440 corresponding to a molecular formula $\text{C}_{36}\text{H}_{61}\text{O}_7\text{N}_2$ (error in ppm: 6.1).

N-Phenyl-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (8). Using 6 equiv of PhNH_2 (0.31 mL, 3.37 mmol) in dry CH_2Cl_2 (20 mL), we obtained 307 mg (86%) of **8**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 2.5/7.5. Colorless foam softens at 122–124 °C. $[\alpha]_D^{20} = +67.1$ ($c = 0.076$, CHCl_3). Anal. ($\text{C}_{38}\text{H}_{55}\text{NO}_7$) C, H.

N,N-Di-(n-propyl)-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (9). Using 6 equiv of (*n*-Pr)₂NH (0.45 mL, 3.27 mmol) in dry CH_2Cl_2 (20 mL), we obtained 298 mg (85%) of **9**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 4/6. Colorless foam softens at 139–140 °C. $[\alpha]_D^{20} = +74.3$ ($c = 0.07$, CHCl_3). Anal. ($\text{C}_{38}\text{H}_{63}\text{NO}_7 \times 0.5\text{H}_2\text{O}$) C, H.

12 α -Acetoxy-5 β -cholan-24-piperidine-24-on-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (10). Using 6 equiv of piperidine (0.32 mL, 3.22 mmol) in dry CH_2Cl_2 (20 mL), we obtained 295 mg (91%) of **10**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 4/6. Colorless foam softens at 72–74 °C. $[\alpha]_D^{20} = +69.4$ ($c = 0.062$, CHCl_3). Anal. ($\text{C}_{37}\text{H}_{59}\text{NO}_7$) C, H.

12 α -Acetoxy-5 β -cholan-24-hydrazide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (11). Using a suspension of 6 equiv of $\text{NH}_2\text{NH}_2 \times 2\text{HCl}$ and 12 equiv of Et_3N in dry CH_2Cl_2 (20 mL), we obtained 209.4 mg (63%) of **11**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 3/7. Colorless foam softens at 178–180 °C. $[\alpha]_D^{20} = +70.3$ ($c = 0.064$, CHCl_3).

12 α -Acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'R)-methylcyclohexane (12). Using a suspension of 10 equiv of NH_4Cl and 10 equiv of Et_3N in dry CH_2Cl_2 (20 mL), we obtained 274 mg (99%) of **12**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 7/3. Mp = 150–153 °C (colorless prisms, Et_2O). $[\alpha]_D^{20} = +75.0$ ($c = 0.096$, CHCl_3). Anal. ($\text{C}_{33}\text{H}_{53}\text{NO}_7 \times 0.5\text{H}_2\text{O}$) C, H, N.

12 α -Acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'S)-methylcyclohexane (13). Using a suspension of 10 equiv of NH_4Cl and 10 equiv of Et_3N in dry CH_2Cl_2 (20 mL), we obtained 294.2 mg (99%) of **13**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 7/3. Mp = 104–106 °C (colorless powder $\text{Et}_2\text{O}/\text{hexane}$). $[\alpha]_D^{20} = +73.3$ ($c = 0.086$, CHCl_3). Anal. ($\text{C}_{33}\text{H}_{53}\text{NO}_7 \times 0.5\text{H}_2\text{O}$) C, H, N.

N-(n-Propyl)-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'R)-methylcyclohexane (14). Using 3 equiv of *n*-PrNH₂ (0.12 mL, 1.46 mmol) in dry CH_2Cl_2 (20 mL), we obtained 253.1 mg (84%) of **14**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 4/6. Mp = 113–115 °C (colorless prisms, $\text{Et}_2\text{O}/\text{hexane}$). $[\alpha]_D^{20} = +70.3$ ($c = 0.074$, CHCl_3). Anal. ($\text{C}_{36}\text{H}_{59}\text{NO}_7 \times 0.5\text{H}_2\text{O}$) C, H, N.

X-ray Analysis of 14. The X-ray intensity data were collected at 100 K with a MAR345 image plate using Mo K α ($\lambda = 0.71069$ Å) radiation. A crystal of approximate dimensions 0.30 × 0.18 × 0.12 mm was chosen, mounted in inert oil, and transferred to the cold gas stream for flash cooling. The unit cell parameters were refined using all of the collected spots after the integration process. Molecular formula = $\text{C}_{36}\text{H}_{59}\text{N O}_7$, Mr = 617.84, monoclinic, $P2_1$, $a = 13.039(4)$, $b = 9.963(3)$, $c = 25.874(8)$ Å, $\beta = 93.76(2)^\circ$, $V = 3354(2)$ Å³, $Z = 4$, $D_x = 1.22$ g cm⁻³, $\mu = 0.083$ mm⁻¹, $F(000) = 1352$, $T = 100$ K.

A total of 13 698 reflections were collected from 131 images taken at a crystal-to-detector distance of 160 mm. There are 7190 independent reflections ($R_{\text{int}} = 0.075$). The structure was solved by direct methods with SHELXS97²³ and refined by full-matrix block least-squares on F^2 using SHELXL97. All of the nonhydrogen

atoms were refined anisotropically. The hydrogen atoms were calculated with AFIX and included in the refinement with a common isotropic temperature factor. Final R values are $R = 0.061$ for 5951 observed reflections, R (all data) = 0.075, $wR = 0.163$, $S = 1.03$. The data have been deposited with the Cambridge Crystallographic Data Centre (Nr CCDC 649757).

N-(n-Propyl)-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'S)-methylcyclohexane (15). Using 3 equiv of *n*-PrNH₂ (0.12 mL, 1.46 mmol) in dry CH_2Cl_2 (20 mL), we obtained 245.7 mg (84%) of **15**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 4/6. Mp = 153–155 °C (colorless powder, $\text{Et}_2\text{O}/\text{hexane}$). $[\alpha]_D^{20} = +69.2$ ($c = 0.078$, CHCl_3). Anal. ($\text{C}_{36}\text{H}_{59}\text{NO}_7 \times 0.5\text{H}_2\text{O}$) C, H, N.

N-(2-Dimethylamino)ethyl-12 α -acetoxy-5 β -cholan-24-amid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'R)-methylcyclohexane (16). Using 2 equiv of *N,N*-dimethyl-ethylenediamine (97 μL , 0.92 mmol) in dry CH_2Cl_2 (20 mL), we obtained 246.3 mg (82%) of **16**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/MeOH/ $\text{NH}_3 = 70/2/1$. Colorless foam softens at 96–97 °C. $[\alpha]_D^{20} = +70.4$ ($c = 0.054$, CHCl_3). HRMS-ESI: m/z 647.4654 corresponding to a molecular formula $\text{C}_{37}\text{H}_{63}\text{O}_7\text{N}_2$ (error in ppm: 2.9).

N-(2-Dimethylamino)ethyl-12 α -acetoxy-5 β -cholan-24-amid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'S)-methylcyclohexane (17). Using 2 equiv of *N,N*-dimethyl-ethylenediamine (0.102 mL, 0.96 mmol) in dry CH_2Cl_2 (20 mL), we obtained 251.6 mg (81%) of **17**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/MeOH/ $\text{NH}_3 = 70/2/1$. Colorless foam softens at 72–74 °C. $[\alpha]_D^{20} = +57.5$ ($c = 0.04$, CHCl_3). HRMS-ESI: m/z 647.4604 corresponding to a molecular formula $\text{C}_{37}\text{H}_{63}\text{O}_7\text{N}_2$ (error in ppm: 4.8).

N-Phenyl-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'R)-methylcyclohexane (18). Using 3 equiv of PhNH_2 (0.13 mL, 1.40 mmol) in dry CH_2Cl_2 (20 mL), we obtained 283 mg (93%) of **18**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 15/85. Colorless foam softens at 109–112 °C. $[\alpha]_D^{20} = +59.8$ ($c = 0.082$, CHCl_3). Anal. ($\text{C}_{39}\text{H}_{57}\text{NO}_7 \times 0.5\text{H}_2\text{O}$) C, H, N.

N-Phenyl-12 α -acetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'S)-methylcyclohexane (19). Using 3 equiv of PhNH_2 (0.13 mL, 1.40 mmol) in dry CH_2Cl_2 (20 mL), we obtained 264.8 mg (87%) of **19**. Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 15/85. Colorless foam softens at 110–113 °C. $[\alpha]_D^{20} = +56.8$ ($c = 0.074$, CHCl_3). Anal. ($\text{C}_{39}\text{H}_{57}\text{NO}_7 \times 0.5\text{H}_2\text{O}$) C, H, N.

N-Phenyl-7 α ,12 α -diacetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'R)-methylcyclohexane (20). Using the above general procedure for preparation of amides, we reacted 7 α ,12 α -diacetoxy-5 β -cholan-24-oic acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'R)-methylcyclohexane^{5b} (372.5 mg, 0.52 mmol) and PhNH_2 (188 μL , 2.06 mmol) to give amide **20** (366.8 mg, 83%). Column chromatography SiO_2 : Lobar B, LichroPrep, eluent EtOAc/heptane = 2.5/7.5. Colorless foam softens at 134–136 °C. $[\alpha]_D^{20} = +45.3$ ($c = 0.064$, CHCl_3). Anal. ($\text{C}_{41}\text{H}_{59}\text{NO}_9 \times 2\text{H}_2\text{O}$) C, H, N.

N-Phenyl-7 α ,12 α -diacetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'S)-methylcyclohexane (21). Using the above general procedure for preparation of amides, we reacted 7 α ,12 α -diacetoxy-5 β -cholan-24-oic acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'S)-methylcyclohexane^{5b} (316.2 mg, 0.50 mmol) and PhNH_2 (182 μL , 1.99 mmol) to give amide **21** (316.2 mg, 89%). Column chromatography: Lobar B, LichroPrep, eluent EtOAc: heptane = 2.5/7.5. Colorless foam softens at 133–136 °C. $[\alpha]_D^{20} = +47.6$ ($c = 0.082$, CHCl_3). Anal. ($\text{C}_{41}\text{H}_{59}\text{NO}_9 \times 2\text{H}_2\text{O}$) C, H, N.

N-(2-Dimethylamino)ethyl-7 α ,12 α -diacetoxy-5 β -cholan-24-amid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4'R)-methylcyclohexane (22). Using the above general procedure for preparation of amides, we reacted 7 α ,12 α -diacetoxy-5 β -cholan-24-oic acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-

((4''R)-methyl)cyclohexane^{5b} (276.0 mg, 0.44 mmol) and *N,N*-dimethylethan-1,2,-diamine (185 μ L, 1.74 mmol) to give amide **22** (259 mg, 84%). Dry-flash chromatography SiO₂: eluent EtOAc/MeOH/NH₃ = 25/1/2. Colorless foam softens at 92–94 °C. [α]_D²⁰ = +41.9 (*c* = 0.074, CHCl₃). Anal. (C₃₉H₆₄N₂O₉ × H₂O × NH₃) C, H, N.

***N*-(2-Dimethylamino)ethyl-7 α ,12 α -diacetoxy-5 β -cholan-24-amid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4''S)-methyl)cyclohexane (23).** Using the above general procedure for preparation of amides, we reacted 7 α ,12 α -diacetoxy-5 β -cholan-24-oic acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spiro-1''-(4''S)-methyl)cyclohexane^{5b} (237.7 mg, 0.37 mmol) and *N,N*-dimethylethan-1,2,-diamine (159 μ L, 1.50 mmol) to give amide **23** (237.1 mg, 81%). Dry-flash chromatography SiO₂: eluent EtOAc/MeOH/NH₃ = 25/1/2. Colorless foam softens at 102–104 °C. [α]_D²⁰ = +50.0 (*c* = 0.076, CHCl₃). Anal. (C₃₉H₆₄N₂O₉ × 1.5 H₂O) C, H, N.

***N*-Phenyl-7 α ,12 α -diacetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (24).** Using the above general procedure for preparation of amides, we reacted 7 α ,12 α -diacetoxy-5 β -cholan-24-oic acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane^{5b} (266.6 mg, 0.43 mmol) and PhNH₂ (157 μ L, 1.72 mmol) to give amide **24** (226.2 mg, 76%). Column chromatography SiO₂: Lobar B, LichroPrep, eluent EtOAc/heptane = 2.5/7.5. Colorless foam softens at 123–125 °C. [α]_D²⁰ = +42.9 (*c* = 0.084, CHCl₃). Anal. (C₄₁H₅₇NO₉ × H₂O) C, H, N.

***N*-(2-dimethylamino)ethyl-7 α ,12 α -diacetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (25).** 7 α ,12 α -Diacetoxy-5 β -cholan-24-oic acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane^{5b} (300 mg, 0.483 mmol) was transformed into amide **25** (139.1 mg, 42%) using Me₂NCH₂CH₂NH₂ (318.4 μ L, 2.9 mmol) in dry CH₂Cl₂ (25 mL). Crude product was purified using column chromatography Lobar B, LichroPrep RP-18; eluent MeOH/H₂O (95/5). Colorless foam softens at 79–85 °C. Anal. (C₃₈H₆₂N₂O₉ × 2 H₂O) C, H, N.

***N*-(2-Pyrrolidino-[3'',4'':1,9](C₆₀-I_h)^{5,6}fullerene-1-yl-ethyl)-7 α ,12 α -diacetoxy-5 β -cholan-24-amide-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane (27).** To a suspension of *N*-(2-pyrrolidino-[3'',4'':1,9](C₆₀-I_h)^{5,6}fullerene)-1-yl-ethanaminium trifluoroacetate²⁴ (50.0 mg, 54.3 μ mol) in dichloromethane (DCM, 10 mL), triethylamine (5.5 mg, 7.5 μ L, 54.3 μ mol) was added, and the reaction mixture was stirred at room temperature for 15 min. In another flask, an ice bath cooled solution of 7 α ,12 α -diacetoxy-5 β -cholan-24-oic acid-3-spiro-6'-(1',2',4',5'-tetraoxacyclohexane)-3'-spirocyclohexane^{5b} (33.4 mg, 54.3 μ mol) in DCM (4 mL) was treated with ethylchloroformate (5.8 mg, 5.2 μ L, 54.3 μ mol) and triethylamine (5.5 mg, 7.5 μ L, 54.3 μ mol), and the reaction mixture was stirred at 0 °C for 5 min. Then, the suspension of fulleramine was slowly added to the solution of the formed mixed anhydride, and the obtained mixture was stirred for 24 h at ambient temperature. After evaporation to dryness, the residue was subjected to dry flash column chromatography. Elution with PhMe/EtOAc 7/3 and subsequent precipitation from a DCM highly concentrated solution with Et₂O gave amide **27** (24.3 mg, 32%) as a brown powder. MALDI-TOF MS Anal. Calcd for C₉₈H₆₀N₂O₉: 1408. Found: 1408 [M]⁺.

In Vitro Antimalarial Activity. The in vitro antimalarial drug susceptibility screen is a modification of the procedures first published by Desjardins et al.,²⁵ with modifications developed by Milhous et al.,²⁶ and the details are given in ref 5a.

In vivo Antimalarial Activity. The *P. berghei* mouse efficacy tests were conducted using a modified version of the Thompson test. Basically, groups of five mice were inoculated intraperitoneally with erythrocytes infected with a drug-sensitive strain of *P. berghei* on day 0. Drugs were suspended in 0.5% hydroxyethylcellulose–0.1% Tween 80 and administered orally once a day beginning on day 3 postinfection. Dosings are given in Table 3. Cure was defined as survival until day 31 posttreatment. Untreated control mice die on day 6–8 postinfection.

Acknowledgment. This work has been supported by the Ministry of Science of Serbia (Grant 142022) and the Serbian Academy of Sciences and Arts. We thank the NIH–NCI's Developmental and Therapeutics program for evaluation of our tetraoxanes. Material has been reviewed by the Walter Reed Army Institute of Research. There is no objection to its presentation or publication. The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting true views of the Department of the Army or the Department of Defense.

Supporting Information Available: Analytical data of synthesized/isolated compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Preliminary results were presented at *4th International Conference of the Chemical Societies of the South-East European Countries*, Belgrade, July 18–21, 2004.
- (2) Malaria Foundation International, <http://www.malaria.org/> (accessed June 2007), and the sites given therein.
- (3) Duan, J. M. J.; Makova, K. D.; Joy, D. A.; Huynh, C. G.; Branch, H.; Li, H.; Su, X.-z. Chromosome-wide SNPs reveal an ancient origin for *Plasmodium falciparum*. *Nature* **2002**, *418*, 323–326.
- (4) Waton, J. C.; Feng, X. G.; Ferdig, M. T.; Cooper, R. A.; Mu, J.; Baruch, D. I.; Magill, A. J.; Su, X.-z. Genetic diversity and chloroquine selective sweeps in *Plasmodium falciparum*. *Nature*, **2002**, *418*, 320–323.
- (5) Selected references: (a) Opsenica, D.; Pocsfalvi, G.; Juranić, Z.; Tinant, B.; Declercq, J.-P.; Kyle, D. E.; Milhous, W. K.; Šolaja, B. A. Cholic Acid Derivatives as 1,2,4,5-Tetraoxane Carriers: Structure and Antimalarial and Antiproliferative Activity. *J. Med. Chem.* **2000**, *43*, 3274–3282. (b) Šolaja, B. A.; Terzić, N.; Pocsfalvi, G.; Gerena, L.; Tinant, B.; Opsenica, D.; Milhous, W. K. Mixed Steroidal 1,2,4,5-Tetraoxanes: Antimalarial and Antimycobacterial Activity. *J. Med. Chem.* **2002**, *45*, 3331–3336. (c) Opsenica, D.; Kyle, E.; Milhous, W. K.; Šolaja, B. A. Antimalarial, antimycobacterial and antiproliferative activity of phenyl substituted mixed tetraoxanes. *J. Serb. Chem. Soc.* **2003**, *68*, 291–302. (d) Opsenica, I.; Terzić, N.; Opsenica, D.; Angelovski, D.; Lehnig, M.; Eilbracht, P.; Tinant, B.; Juranić, Z.; Smith, K. S.; Yang, Y. S.; Diaz, D. S.; Smith, P. L.; Milhous, W. K.; Doković, D.; Šolaja, B. A. Tetraoxane Antimalarials and Their Reaction with Fe(II). *J. Med. Chem.* **2006**, *49*, 3790–3799.
- (6) Kim, H.-S.; Tsuchiya, K.; Shibata, Y.; Wataya, Y.; Ushigoe, Y.; Masuyama, A.; Nojima, M.; McCullough, K. J. Synthetic methods for unsymmetrically-substituted 1,2,4,5-tetraoxanes and of 1,2,4,5,7-pentoxocanes. *J. Chem. Soc., Perkin Trans. 1* **1999**, 1867–1870.
- (7) Iskra, J.; Bonnet-Delpont, D.; Bégue, J.-P. One-pot synthesis of non-symmetric tetraoxanes with the H₂O₂/MTO/fluorous alcohol system. *Tetrahedron Lett.* **2003**, 6309–6312.
- (8) Terent'ev, A. O.; Kutkin, A. V.; Starikova, Z. A.; Antipin, M. Yu.; Ogibin, Yu. N.; Nikishin, G. I. New preparation of 1,2,4,5-tetraoxanes. *Synthesis* **2004**, (14) 2356–2366.
- (9) Recently, the approaches of Iskra and Terent'ev were successfully combined for the synthesis of quite active adamantylidene compounds: Amewu, R.; Stachulski, A. V.; Ward, S. A.; Berry, N. G.; Bray, P. G.; Davies, J.; Labat, G.; Vivas, L.; O'Neill, P. M. Design and synthesis of orally active dispiro 1,2,4,5-tetraoxanes; synthetic antimalarials with superior activity to artemisinin. *Org. Biomol. Chem.* **2006**, *4*, 4431–4436.
- (10) Opsenica, I.; Terzić, N.; Opsenica, D.; Milhous, W. K.; Šolaja, B. 7,8,15,16-Tetraoxa dispiro[5.2.5.2]hexadecane-3-carboxylic acid derivatives and their antimalarial activity. *J. Serb. Chem. Soc.* **2004**, *69*, 919–922.
- (11) Spek, A. L. *PLATON, Molecular Geometry Program*; University of Utrecht, The Netherlands, 1998.
- (12) Opsenica, D.; Angelovski, G.; Pocsfalvi, G.; Juranić, Z.; Žižak, Z.; Kyle, D.; Milhous, W. K.; Šolaja, B. A. Antimalarial and antiproliferative evaluation of bis-steroidal tetraoxanes. *Bioorg. Med. Chem.* **2003**, *11*, 2761–2768.
- (13) (a) Krusic, P. J.; Wasserman, E.; Keizer, P. N.; Morton, J. R.; Preston, K. F. Radical Reactions of C₆₀. *Science*, **1991**, *254*, 1183–1185. (b) Cremonini, M. A.; Lunazzi, L.; Placucci, G.; Krusic, P. J. *J. Org. Chem.* **1993**, *58*, 4735–4738. (c) Liangbing Gan, L.; Huang, S.; Zhang, X.; Zhang, A.; Cheng, B.; Cheng, H.; Li, X.; Shang, G.

- Fullerenes as a tert-butylperoxy radical trap, metal catalyzed reaction of tert-butyl hydroperoxide with fullerenes, and formation of the first fullerene mixed peroxides C₆₀(O)(OOtBu)₄ and C₇₀(OOtBu)₁₀. *J. Am. Chem. Soc.* **2002**, *124*, 13384–13385.
- (14) Bhattacharjee, A. K.; Carvalho, K. A.; Opsenica, D.; Šolaja, B. A. Structure-activity relationship study of steroidal 1,2,4,5-tetraoxane antimalarials using computational procedures. *J. Serb. Chem. Soc.* **2005**, *70*, 329–345.
- (15) Drug discovery and development program, National Cancer Institute, Bethesda, MD (NCI), <http://dtp.nci.nih.gov> (accessed June 2007).
- (16) This is in line with our previous observations,^{5d} where high SI values (in vitro) were confirmed with no toxicity of tested steroidal tetraoxanes in experimental animals (in vivo).
- (17) Very interesting artemisinin dimers were recently prepared in an attempt to develop dual-activity entity; c.f. Posner, G. H.; McRiner, A. J.; Paik, I.-H.; Sur, S.; Borstnik, K.; Xie, S.; Shapiro, A. S.; Alagbala, A.; Foster, B. Anticancer and antimalarial efficacy and safety of artemisinin-derived trioxane dimers in rodents. *J. Med. Chem.* **2004**, *47*, 1299–1301. For further improvement towards antimalarial drug based on artemisinin-derived dimers, see Posner, G. H.; Paik, I.-H.; Chang, W.; Borstnik, K.; Sinishtaj, S.; Rosenthal, A. S.; Shapiro, T. A. Malaria-infected mice are cured by a single dose of novel artemisinin derivatives. *J. Med. Chem.* **2007**, *50*, 2516–2519.
- (18) For our previous reports on selective cytotoxicity of tetraoxane antimalarials against cancer cells as compared with healthy ones, see ref 5d and relevant references cited therein.
- (19) (a) Chen, H.-H.; Zhou, H.-J.; Fang, X. Inhibition of human cancer cell line growth and human umbilical vein endothelial cell angiogenesis by artemisinin derivatives in vitro. *Pharm. Research* **2003**, *48*, 231–236. (b) Liu, Y.; Wong, V. K.-W.; Ko, B. C.-B.; Wong, M.-K.; Che, C.-M. Synthesis and cytotoxicity studies of artemisinin derivatives containing lipophilic alkyl carbon chains. *Org. Lett.* **2005**, *7*, 1561–1564. (c) Alagbala, A. A.; McRiner, A. J.; Borstnik, K.; Labonte, T.; Chang, W.; D'Angelo, J. G.; Posner, G. H.; Foster, B. A. Biological mechanisms of action of novel C-10 non-acetal trioxane dimers in prostate cancer cell lines. *J. Med. Chem.* **2006**, *49*, 7836–7842. (d) Paik, I.-H.; Xie, S.; Shapiro, T.; Labonte, T.; Narducci Sarjeant, A. A.; Baege, A. C.; Posner, G. H. Second generation, orally active, antimalarial, artemisinin-derived trioxane dimers with high stability, efficacy, and anticancer activity. *J. Med. Chem.* **2006**, *49*, 2731–2734.
- (20) Jeyadevan, J. P.; Bray, P. G.; Chadwick, J.; Mercer, A. E.; Byrne, A.; Ward, S. A.; Park, B. K.; Williams, D. P.; Cosstick, R.; Davies, J.; Higson, A. P.; Irving, E.; Posner, G. H.; O'Neill, P. M. Antimalarial and antitumor evaluation of novel C-10 non-acetal dimers of 10β-(2-hydroxyethyl)deoxoartemisinin. *J. Med. Chem.* **2004**, *47*, 1290–1298.
- (21) Sadava, D.; Phillips, T.; Lin, C.; Kane, S. E. Transferin overcomes drug resistance to artemisinin in human small-cell lung carcinoma cells. *Cancer Lett.* **2002**, *179*, 151–156.
- (22) Moore, J. C.; Lai, H.; Li, J.-R.; Ren, R.-L.; McDougall, J. A.; Singh, N. P.; Choua, C.-K. Oral administration of dihydroartemisinin and ferrous sulfate retarded implanted fibrosarcoma growth in the rat. *Cancer Lett.* **1995**, *98*, 83–87.
- (23) Sheldrick, G. M. *SHELXS-97* and *SHELXL-97*, Program for the Solution and Refinement of Crystal Structures; University of Göttingen: Germany, 1997.
- (24) Kordatos, K.; Da Ros, T.; Bosi, S.; Vázquez, E.; Bergamin, M.; Cusan, C.; Pellarini, F.; Tomberli, V.; Baiti, B.; Pantarotto, D.; Georgakilas, V.; Spalluto, G.; Prato, M. Novel Versatile Fullerene Synthons. *J. Org. Chem.* **2001**, *66*, 4915–4920.
- (25) Desjardins, R. E.; Canfield, C. J.; Haynes, D. E.; Chulay, J. D. Quantitative assessment of antimalarial activity in vitro by a semiautomated microdilution technique. *Antimicrob. Agents Chemother.* **1979**, *16*, 710–718.
- (26) Milhous, W. K.; Weatherly, N. F.; Bowdre, J. H.; Desjardins, R. E. In vitro activities of and mechanisms of resistance to antifol antimalarial drugs. *Antimicrob. Agents Chemother.* **1985**, *27*, 525–530.

JM070684M