

Structure-Based Design, Synthesis, Evaluation, and Crystallographic Studies of Conformationally Constrained Smac Mimetics as Inhibitors of the X-linked Inhibitor of Apoptosis Protein (XIAP)[†]

Haiying Sun,^{‡,∇} Jeanne A. Stuckey,^{§,○,▲} Zaneta Nikolovska-Coleska,^{‡,∇} Dongguang Qin,^{‡,∇} Jennifer L. Meagher,[○] Su Qiu,^{‡,∇} Jianfeng Lu,^{‡,∇} Chao-Yie Yang,^{‡,∇} Naoyuki G. Saito,[#] and Shaomeng Wang^{*,‡,§,||,L,∇}

Departments of Internal Medicine, Biological Chemistry, Pharmacology, Medicinal Chemistry, Radiation Oncology, Comprehensive Cancer Center, Life Sciences Institute, Biophysics Research Division, University of Michigan, Ann Arbor, Michigan 48109

Received June 5, 2008

Small molecules designed to mimic the binding of Smac protein to X-linked inhibitor of apoptosis protein (XIAP) are being pursued as a promising new class of anticancer drugs. Herein, we report the design, synthesis, and comprehensive structure–activity relationship studies of a series of conformationally constrained bicyclic Smac mimetics. Our studies led to the discovery of a number of highly potent and cell-permeable Smac mimetics and yielded important new insights into their structure–activity relationship for their binding to XIAP and for their activity in inhibition of cancer cell growth. Determination of the crystal structure of one potent Smac mimetic, compound **21**, in complex with XIAP BIR3 provides the structural basis for its high-affinity binding to XIAP and for the design of highly potent Smac mimetics.

Introduction

Apoptosis is a critical cell process in normal development and homeostasis of multicellular organisms to eliminate unwanted or damaged cells. Inappropriate regulation of apoptosis plays a major role in many human diseases, including cancer.^{1–4} Defects in the apoptosis machinery confers apoptosis resistance on cancer cells to therapeutic agents, makes current anticancer therapies less effective, and leads ultimately to their failure in the clinic.^{2–4} Accordingly, targeting critical apoptosis regulators aimed at overcoming apoptosis resistance of cancer cells is a promising cancer therapeutic strategy.

The X-linked inhibitor of apoptosis protein (XIAP^a) has been identified as a key apoptosis inhibitor, although its role in cells may not be limited to the regulation of apoptosis.^{5–10} XIAP inhibits apoptosis through direct binding to and inhibition of three cysteine proteases, an initiator caspase-9 and the two effectors caspase-3 and -7.^{5–10} XIAP contains three baculoviral IAP repeats (BIR) domains. While the third BIR domain (BIR3) of XIAP selectively targets caspase-9, the BIR2 domain, together with the immediate preceding linker, inhibits both caspase-3 and caspase-7. Because these caspases play a critical role in the execution of apoptosis, XIAP functions as an efficient inhibitor of apoptosis. Consistent with its potent apoptosis-

suppressing function, XIAP is found to be highly expressed in many human tumor cell lines and tumor samples from patients¹¹ and plays an important role in conferring resistance on cancer cells to a variety of anticancer drugs.^{8,9} Because XIAP blocks apoptosis at the downstream effector phase, a point where multiple signaling pathways converge, it represents a particularly attractive molecular target for the design of new classes of anticancer drugs aimed at overcoming the apoptosis resistance of cancer cells.^{8,9,12}

The antiapoptotic function of XIAP is antagonized by second mitochondria-derived activator of caspases or direct IAP binding protein with low pI (Smac/DIABLO), a protein released from mitochondria into the cytosol in response to apoptotic stimuli.^{13,14} Crystal and NMR structures^{15,16} show that Smac, through its N-terminal AVPI (Ala1-Val2-Pro3-Ile4) motif, interacts with the XIAP BIR3 domain at the same site where caspase-9 binds and removes the inhibition of XIAP to caspase-9 by direct competition.^{17,18} The mechanism by which Smac removes the inhibition of XIAP to caspase-3/-7 is not entirely clear. It has been proposed that Smac protein removes the inhibition of XIAP to caspase-3/-7 by binding to the XIAP BIR2 domain through its AVPI motif.¹⁹ Hence, the AVPI binding motif in Smac plays a critical role for the interaction of Smac protein with XIAP and its functional antagonism against XIAP.

Earlier studies using Smac-based peptides containing the AVPI binding motif, tethered to a carrier peptide for intracellular delivery, have demonstrated that such compounds can enhance the antitumor activity of chemotherapeutic agents and of TNF-related apoptosis inducing ligand in vitro and in vivo.^{20–22} Although Smac-based peptides have served as useful tools for important proof-of-concept studies, they are not suitable drug candidates due to their very limited cellular activity and expected poor in vivo stability. To overcome the limitations associated with peptide-based Smac mimetics, a number of laboratories, including ours, have pursued the design of peptidic and nonpeptidic small-molecule Smac mimetics with a goal to obtain more druglike compounds, which may be developed as a new class of anticancer drugs.^{23–30}

[†] Coordinates of compound **21** in complex with XIAP BIR3 and structural factors have been deposited in the RCSB Protein Data Bank (access code 2JK7).

* To whom correspondence should be addressed. Phone: 734-615-0362. Fax: 734-647-9647. E-mail: shaomeng@umich.edu.

[‡] Department of Internal Medicine, University of Michigan.

[§] Department of Biological Chemistry, University of Michigan.

^{||} Department of Pharmacology, University of Michigan.

^L Department of Medicinal Chemistry, University of Michigan.

[#] Department of Radiation Oncology, University of Michigan.

[∇] Comprehensive Cancer Center, University of Michigan.

[○] Life Sciences Institute, University of Michigan.

[▲] Biophysics Research Division, University of Michigan.

^a Abbreviations: IAP, inhibitor of apoptosis protein; XIAP, X-linked IAP; Smac, second mitochondria-derived activator of caspases; DIABLO, direct IAP binding protein with low pI; BIR, baculoviral IAP repeats (BIR) domain; BIR2, the second BIR domain; BIR3, the third BIR domain; FP, fluorescence polarization.




Figure 1. Design and examples of conformationally constrained bicyclic Smac peptide mimetics.

Using a structure-based approach, our laboratory has reported the design of a number of conformationally constrained, bicyclic Smac mimetics.^{23,24,26,30} Our previous studies showed that these designed Smac mimetics can achieve high binding affinities to XIAP and are effective in inhibition of cell growth and induction of apoptosis in cancer cells. For example, SM-131, which contains a [7,5] bicyclic core structure, binds to XIAP BIR3 protein with a K_i of 61 nM in a competitive binding assay and directly antagonizes the XIAP inhibition of caspase-9 activity in a cell-free functional assay.²⁶ This compound also potently inhibits cancer cell growth and induces apoptosis in cancer cells as a single agent.²⁶

Although our previous studies^{23,24,26,30} have led to the discovery of potent and cell-permeable Smac mimetics, our understanding on their structure–activity relationship is still limited. Furthermore, although molecular modeling was employed to predict the binding models of our designed Smac mimetics to XIAP BIR3 protein in our previous studies, the predicted binding models have not been experimentally confirmed.

To gain a more in-depth understanding of the structure–activity relationship for our designed conformationally constrained Smac mimetics for their binding to XIAP and for their cellular activity, we have designed, synthesized, and evaluated a series of new Smac mimetics. To obtain a solid structural basis for the interaction of our designed Smac mimetics with XIAP BIR3, we have determined a high-resolution crystal structure of a potent Smac mimetic (compound **21**) in complex with XIAP BIR3. We report herein structure-based design, synthesis, biochemical and biological evaluation, and crystallographic studies of conformationally constrained Smac mimetics as antagonists of XIAP.

Results and Discussion

Structure-Based Design of Conformationally Constrained Smac Mimetics and Their Structure–Activity Relationship. We have employed a structure-based strategy for the design of conformationally constrained, bicyclic Smac mimetics (Figure 1).^{23,24}

In the crystal structure of Smac protein in complex with XIAP BIR3, the hydrophobic side chain of Ile4 in the Smac AVPI

peptide inserts into a hydrophobic pocket in XIAP BIR3 and the carbonyl group of Ile4 does not have specific interactions with the protein.¹⁵ We thus replaced the Ile4 residue with a simple benzylamine group, which led to compound **1**.^{23,24} Compound **1** binds to XIAP BIR3 protein with a K_i of 0.29 μ M in our fluorescence-polarization (FP) binding assay³² and is 2 times more potent than the natural Smac AVPI peptide ($K_i = 0.58 \mu$ M) under the same assay conditions. Compound **1** was thus used as a new lead in our subsequent design.

The isopropyl side chain in Val2 has no specific interactions with the protein in the crystal structure, while the five-membered ring of Pro3 has van der Waals contacts with the side chain of Trp323 in the protein. Modeling suggested that the isopropyl side chain in Val2 may be fused with the five-membered ring of Pro3 to form a bicyclic lactam structure without causing steric clashes with the protein.^{23,24} Cyclization of the side chain of Val2 and the five-membered ring of Pro3 generates a new chiral center. To probe the stereospecificity of this new chiral center for binding to XIAP, we designed and synthesized stereoisomers **2** and **3**, which contain a [6,5] bicyclic lactam core structure.²⁴ Compound **2** binds to XIAP BIR3 with a K_i value of 4.47 μ M, but compound **3** shows no appreciable binding at concentrations up to 100 μ M. Hence, the binding data for compounds **2** and **3** established that the *S*-configuration (up-configuration) of the chiral center in the bicyclic Smac mimetics is highly preferred for their binding to XIAP BIR3. Accordingly, in our subsequent design and synthesis, we have focused on compounds with the *S*-configuration (up-configuration) for the chiral center.

Although compound **2** binds to XIAP BIR3 with a good affinity, it is 8 times less potent than the Smac AVPI peptide and 15 times less potent than compound **1**. Modeling showed that, although compound **2** largely mimics the bound conformation of the AVPI peptide to XIAP BIR3, its rigid [6,5] bicyclic structure causes some deviations (Figure 2A) that may account for its lower binding affinity to XIAP BIR3. Therefore, we designed and synthesized compounds **4**²³ and **5**, in which the six-membered ring in **2** was expanded to a seven- or eight-membered ring, respectively. Modeling predicted that these two compounds can nicely mimic the bound conformation of the Smac AVPI peptide to XIAP BIR3 protein (Figure 2B,C) and




Figure 2. Predicted binding models of compounds **2**, **4**, **5**, **20**, and **21** to XIAP BIR3 protein in comparison with that for Smac AVPI peptide in the crystal structure. Carbon atoms in compounds **2**, **4**, **5**, **20**, and **21** and AVPI are shown in stick. The carbon atoms are colored in yellow in **2**, **4**, **5**, **20**, and **21** and in green in the AVPI peptide. The carbon atoms in several key residues in XIAP are colored in gray. Nitrogen and oxygen atoms are colored in blue and red, respectively.


suggested that both compounds should bind to XIAP BIR3 with a higher affinity than compound **2**. Indeed, compounds **4** and **5** achieved K_i values of 0.15 and 0.10 μM , respectively, and are 30 or 45 times more potent than compound **2**, confirming the modeling prediction.






















We carried out further modifications in both *N*-terminal (region 1) and *C*-terminal (region 2) regions (Figure 1) in compounds **2**, **4**, or **5** to further explore the structure–activity relationships in these two regions. The results are summarized in Table 1.

In the crystal structure of Smac protein in complex with XIAP BIR3, the methyl group of Ala1 inserts into a small hydrophobic pocket. Compounds **6**, **7**, and **8** were designed and synthesized in which the methyl group in **2** was replaced with an ethyl, *n*-propyl, or isopropyl group, respectively. While compound **6** with an ethyl group is 3 times more potent than **2**, compound **8** with an isopropyl group is 10 times less potent than **2** and compound **7** with an *n*-propyl group shows no appreciable binding up to 100 μM to XIAP BIR3. These data demonstrated that the small hydrophobic pocket in XIAP BIR3 occupied by the methyl group in Ala1 in the Smac AVPI peptide can not

accommodate a group larger than an ethyl group for effective binding. Therefore, only the methyl or ethyl group was employed in this position in our subsequent design and synthesis.

The positively charged primary amino group in Ala1 in the Smac AVPI has strong charge–charge interactions with the negatively charged carboxylic group in the side chain of Glu314 of XIAP BIR3 and is critical for binding of Smac-based peptides to XIAP BIR3. However, we found that Smac mimetics with high binding affinities to XIAP BIR3 but containing this primary amino group have very weak cellular activity.^{23,24,26} Therefore, we replaced this primary amine with a secondary or tertiary amine or a hydroxyl group, which can also form one or more hydrogen bonds with the protein under physiological conditions.²⁶ Their binding data indicated that replacement of the amino group with an *N*-methyl amino group in Smac mimetics containing either a [7,5] or [8,5] bicyclic lactam leads to reduction of the binding affinity by 2–3 times (**13** vs **12**, **20** vs **16**, and **21** vs **17**). However, replacement of the primary amino group with an *N,N*-dimethyl amino (**14**) or a cyclic tertiary amino (**22** and **23**) or a hydroxyl group (**15** and **24**) decreases the binding affinities by 20–800 times. These data show that

Table 1. Chemical Structures of Conformationally Constrained Bicyclic Smac Mimetics and Their Binding Affinities to XIAP BIR3


| Compounds | X | R ₁ | R ₂ | R ₃ | XIAP BIR3 K _i (μM) |
|-----------|---------------------------------|----------------|---|---|----------------------------------|
| AVPI | | | | | 0.58 ²³ |
| 1 | | | | | 0.29 ²³ |
| 2 | CH ₂ | Me | NH ₂ | Bn | 4.47 ²⁴ |
| 3 | | | | | >100 ²⁴ |
| 4 | (CH ₂) ₂ | Me | NH ₂ | Bn | 0.15 ²³ |
| 5 | (CH ₂) ₃ | Et | NH ₂ | Bn | 0.10 |
| 6 | CH ₂ | Et | NH ₂ | Bn | 1.41 ± 0.16 |
| 7 | CH ₂ | <i>n</i> -Pr | NH ₂ | Bn | >100 |
| 8 | CH ₂ | <i>i</i> -Pr | NH ₂ | Bn | 43.11 ± 1.51 |
| 9 | CH ₂ | Me | NH ₂ |  | 2.33 ± 0.68 |
| 10 | CH ₂ | Et | NH ₂ |  | 0.35 ± 0.01 |
| 11 | (CH ₂) ₂ | Me | NH ₂ |  | 0.06 ± 0.02 |
| 12 | (CH ₂) ₂ | Et | NH ₂ |  | 0.025 ± 0.004 |
| 13 | (CH ₂) ₂ | Et | MeNH |  | 0.061 ± 0.006 |
| 14 | (CH ₂) ₂ | Et | Me ₂ N |  | 14.4 ± 0.6 |
| 15 | (CH ₂) ₂ | Et | OH |  | 29.0 ± 1.4 |
| 16 | (CH ₂) ₃ | Me | NH ₂ |  | 0.014 ± 0.003 |
| 17 | (CH ₂) ₃ | Et | NH ₂ |  | 0.039 ± 0.004 |
| 18 | (CH ₂) ₃ | Et | NH ₂ |  | 0.759 ± 0.16 |
| 19 | (CH ₂) ₃ | Et | NH ₂ |  | 0.047 ± 0.009 |
| 20 | (CH ₂) ₃ | Me | MeNH |  | 0.026 ± 0.005 |
| 21 | (CH ₂) ₃ | Et | MeNH |  | 0.067 ± 0.018 |
| 22 | (CH ₂) ₃ | Et |  |  | 0.856 ± 0.075 |
| 23 | (CH ₂) ₃ | Et |  |  | 32.87 ± 4.18 |
| 24 | (CH ₂) ₃ | Me | OH |  | 12.97 ± 2.02 |
| 25 | (CH ₂) ₃ | Me | MeNH |  | 0.014 ± 0.005 |
| 26 | (CH ₂) ₃ | Me | MeNH |  | 0.274 ± 0.037 |
| 27 | (CH ₂) ₃ | Me | MeNH |  | 0.015 ± 0.008 |

the primary and secondary amino groups are highly preferred at this site for Smac mimetics to achieve high binding affinities to XIAP BIR3.

The crystal structure of Smac in complex with XIAP BIR3 showed that the carbonyl group of Ile4 is exposed to solvent and has no specific interaction with the protein.¹⁵ However, we hypothesized that this carbonyl group may contribute to the binding affinity of Smac to XIAP by controlling the orientation of the hydrophobic side chain of Ile4, thus reducing the entropic loss upon binding. To test this hypothesis, we replaced the benzyl group in **2**, **4**, and **5** with a diphenyl methyl group, which led to compounds **9**, **11**, and **16**, respectively. These three compounds (**9**, **11**, and **16**) were found to be three times more potent than their corresponding analogues **2**, **4**, and **5**, supporting our hypothesis.

Our modeling predicted that the pro *R* phenyl group enters the hydrophobic pocket and the pro *S* phenyl group is exposed to solvent. To test this modeling prediction, we have designed compounds **18** and **19** by replacing either the pro *R* or pro *S* phenyl group in **17** with a naphthyl group. It was found that while the *S* form isomer **18** is as potent as **17**, the *R* form isomer **19** is 35 times less potent than **17**, thus confirming our modeling prediction.

We next tested if other structures that can control and restrict the orientation of the phenyl group can also improve the binding affinity. To this end, we have designed and synthesized compounds **25**–**27**, in which the conformation of the phenyl group was further constrained by a fused second ring and was previously employed for the design of potent, Smac peptidomimetics.²⁵ Compound **25**, in which the chiral amine has the *R* configuration, has a K_i value of 14 nM, whereas its stereoisomer **26** has a K_i value of 207 nM. Thus, compound **25** is 15 times more potent than its isomer **26**, indicating that the *R* form is the desirable configuration for the chiral center. Compound **27**, in which the (*R*)-1,2,3,4-tetrahydro-naphthalenyl group in **25** was replaced by a (*R*)-2,3-dihydro-1H-indene group, also binds to XIAP BIR3 with a high affinity and has a K_i value of 15 nM.

Functional Antagonism of Smac Mimetics against XIAP BIR3 Protein in a Cell-Free Functional Assay. On the basis of their expected mode of action, Smac mimetics should antagonize the inhibition of XIAP BIR3 to caspase-9 by binding to the same site where caspase-9 binds. We thus established a cell-free functional assay using recombinant XIAP BIR3 protein and evaluated the ability of our designed Smac mimetics to antagonize the XIAP BIR3 protein.²⁶ In this assay, addition of dATP and cytochrome c into cell lysates activates caspase-9, and recombinant XIAP BIR3 protein inhibits the activity of caspase-9 in a dose-dependent manner. XIAP BIR3 protein at 0.5 μM is capable of completely inhibiting the activity of caspase-9 under the assay conditions (Figure 3).

We evaluated two potent Smac mimetics, compounds **20** and **25**, and the Smac AVPI peptide in this cell-free functional assay for their ability to directly overcome the inhibitory effect of XIAP BIR3 protein to caspase-9 activity (Figure 3). Compound **24**, which is >500-times less potent than compounds **20** and **25** in its binding affinity to XIAP BIR3, was included as a negative control. It was found that compounds **20**, **25**, and the AVPI peptide all effectively and dose-dependently restore the activity of caspase-9, whereas compound **24** has no significant effect at concentrations as high as 100 μM. On the basis of the kinetic data, the calculated EC₅₀ values for compounds **20**, **25**, and the AVPI peptide in restoration of the caspase-9 activity at 60 min are 0.78, 0.63, and 4.8 μM, respectively, consistent with




Figure 3. Functional antagonism of compounds **20**, **24**, **25**, and the AVPI peptide against XIAP BIR3 to promote the activity of caspase-9 in a cell-free functional assay.

their binding data to XIAP BIR3. Hence, our functional assay established that compounds **20** and **25** are potent antagonists of XIAP BIR3 and both of them are also more potent than the Smac AVPI peptide. Consistent with its weak binding affinity, **24** is inactive at concentrations as high as 100 μM in the functional assay.

Evaluation of the Activity of Smac Mimetics in Cell Growth Inhibition and Induction of Cell Death and Apoptosis in Human Cancer Cell Lines. One major motivation for the design and synthesis of nonpeptidic Smac mimetics is to dramatically improve the cellular activity over Smac-based peptides. Although Smac mimetics were initially thought to be primarily useful for sensitizing cancer cells to other anticancer drugs,^{20,21,23,24,29} a number of studies have shown that properly designed Smac mimetics are capable of inhibiting cancer cell growth and inducing apoptosis in a subset of cancer cell lines^{25,30} such as the MDA-MB-231 human breast cancer cell line and SK-OV-3 human ovarian carcinoma cell line.

We have therefore tested compounds **20**, **21**, and **25** for their activity in inhibition of cell growth and induction of cell death and apoptosis as single agents in the MDA-MB-231 and SK-OV-3 cancer cell lines. In our previous study,²⁶ we showed that methylation of the primary amine group in compound **12**, which resulted in compound **13**, did not significantly alter the binding affinity to XIAP BIR3 but led to dramatic improvement in the cellular activity. To further confirm this observation in our Smac mimetics containing the [8,5] bicyclic ring, we have included compound **17**, which contains a primary amine group and differs from compound **21** by the methylation of the amine group. Compound **24** was used as another control in these cellular assays to demonstrate the specificity of compounds **20**, **21**, and **25**. The results are shown in Figure 4.




Figure 4. Inhibition of cell growth by Smac mimetics in the MDA-MB-231 breast cancer and SK-OV-3 ovarian cancer cell lines. Cells were treated for 4 days, and cell growth was determined using WST-based assay.

Compounds **20**, **21**, and **25** potently inhibit cell growth in the MDA-MB-231 breast cancer cell line in a dose-dependent manner with IC_{50} values of 0.41, 0.41, and 0.1 μM , respectively. Similarly, compounds **20**, **21**, and **25** also potently inhibit cell growth in the SK-OV-3 ovarian cancer cell line with IC_{50} values of 0.9, 1.0, and 0.18 μM , respectively. Consistent with its weak binding affinity to XIAP BIR3, compound **24** has IC_{50} values of 7.0 μM and >10 μM , respectively, being >30-times less potent than compound **25** in both cancer cell lines. Consistent with our previous observation, compound **17** has a weak cellular activity in both cell lines, indicating that the methylation on the primary amine group in **17** results in dramatic improvement of the cellular activity.

We next evaluated if our designed Smac mimetics can effectively induce apoptosis in the MDA-MB-231 cell line, and the results are shown in Figure 5. As expected, compounds **20**, **21**, and **25** effectively induce apoptosis in a dose-dependent manner. While all three compounds induce robust apoptosis at 100 nM, compound **25** is the most potent among them, consistent with their potency in inhibition of cell growth. Similar data were obtained using the SK-OV-3 ovarian cancer cell line (data not shown). Our data thus indicate that potent Smac mimetics not only potently inhibit cell growth but also effectively induce apoptosis in cancer cells.

Because XIAP potently inhibits caspase activity, we next determined if compounds **20** and **25** induces apoptosis and cell death in a caspase dependent manner. Using a trypan blue exclusion assay, it was shown that both compounds effectively




Figure 5. Induction of apoptosis by Smac mimetics in the human breast cancer MDA-MB-231 cell line. Cells were treated for 48 h, and apoptosis was determined using Annexin V-propidium iodide (P.I.) staining by flow cytometry. Annexin V (+) and P.I. (-) cells are labeled as early apoptotic cells, whereas Annexin V (+) and P.I. (+) cells are labeled as late apoptotic cells. Percentage of early and late apoptotic cells are shown.

kills the MDA-MB-231 cancer cells in a dose-dependent manner but **25** is more potent (Figure 6). Treatment with **25** at 10, 100, and 1000 nM for 24 h kills 9%, 23%, and 65% of the MDA-MB-231 cells, respectively. The cell killing by both **20** and **25** can be effectively attenuated by Z-DEVD-FMK, a cell-permeable caspase-3/7 inhibitor, indicating that cell death induction by these Smac mimetics is caspase-3/7 dependent (Figure 6).

Determination of the Crystal Structure of Compound 21 in Complex with XIAP BIR3 Domain. To firmly establish the structural basis of the binding of our designed Smac mimetics to XIAP BIR3, we have determined the crystal structure of compound **21** in complex with XIAP BIR3 to 2.8 Å resolution (Figure 7).

The experimentally determined structure of compound **21** in complex with one XIAP BIR3 molecule is in excellent agreement with that predicted by computational modeling. The root-mean-square-deviation (rmsd) between the non-hydrogen atoms of compounds **21** in the crystal structure and the modeled structure is 0.81 Å.

The interactions of compound **21** with XIAP BIR3 closely mimic those between the Smac AVPI peptide and the protein, although some differences are observed. The methyl group in Ala1 side chain in the Smac AVPI peptide is replaced with an ethyl group in compound **21**. This ethyl group makes more extensive van der Waals interactions with Trp310 and Leu307 than the methyl group in Ala1 in Smac, which, in turn, produces a minor displacement of the positively charged *N*-terminal amino group. In Smac, this positively charged amino group in Ala1 has three hydrogen bonds to the side chains of Gln319 and Glu314. In compound **21**, the *N*-methylated amino group loses contact with Gln319 but gains an additional hydrogen bonding interaction with the backbone carbonyl group of Asp309.

Similar to the Smac AVPI peptide, the first carbonyl group in compound **21** forms a less than optimal hydrogen bond to the indole NH group in Trp323. The second amino group in **21** has an optimal hydrogen bond with the backbone carbonyl group of Thr308 and the carbonyl group in the [8,5] lactam ring forms an optimal hydrogen bond with the backbone amino group of Thr308. Another strong hydrogen bond is observed between the third amino group and the backbone carbonyl group of Gly306. For hydrophobic contacts, the ethyl group in **21** inserts into a small hydrophobic pocket formed by the hydrophobic side chains of Leu307 and Trp310, as well as the hydrophobic portion of the side chain in Gln319, mimicking the methyl group in alanine residues in the Smac AVPI peptide. The eight-membered ring in **21** has extensive contacts with the side chain of Trp323. While the five-membered ring in **21** still has contacts with the side chain of Trp323, similar to that in Smac, but the interaction appears to be weaker than that in Smac. However, this five-membered ring in **21** gains additional new contacts with Tyr324. As predicted by modeling, the pro *R* phenyl group inserts into a hydrophobic pocket formed by the hydrophobic side chains of Leu292 and the hydrophobic portion of the side chains in Lys297 and Lys299.

Interestingly, the crystal structure also revealed a remarkable dimer, which consists of two molecules of the protein and compound **21** (Figure 7A). The dimeric interface created between the two compound **21**:XIAP BIR3 complexes in this crystal structure is a unique dimer configuration when compared to other BIR3 three-dimensional structures.^{33,34} This dimeric interface is mainly stabilized by the presence of compound **21**. The pro *S* phenyl group in both molecules of compound **21** makes multiple hydrophobic interactions with Trp323 and Tyr324 in the other protein molecule in the symmetric dimer. In addition, the [8,5] bicyclic lactam ring from each molecule of **21** stacks on the top of each other and has nice hydrophobic contacts. Other protein-protein interactions that stabilize this interface come from the *N*-terminal Phe250 from each molecule. The hydrophobic side chain of each Phe250 residue in the dimeric structure has extensive contacts with the diphenyl groups in compound **21** from the opposite side. This dimeric structure formed between 2 molecules of compound **21** and two XIAP BIR3 molecules provides a structural basis for the design of bivalent Smac mimetics to either induce dimerization for two BIR domains or concurrent targeting of two different BIR domains in the same IAP molecule.

Synthesis of Designed Smac Mimetics. The synthesis for Smac mimetics containing the [6,5] and [7,5] bicyclic lactam systems have been reported previously by our group.^{23,24} The general synthetic method for Smac mimetics containing the [8,5] bicyclic lactam system is shown in Scheme 1.

The key Intermediate **28** was prepared according to a previously reported method.^{30,31} Condensation of **28** with corresponding amines gave a series of amides. After removal of the Boc protecting group in these amides, the resulted ammonium salts were condensed with corresponding *N*-Boc amino acids, followed by deprotection of the Boc protecting group to yield compounds **16–21** and **25–27**. Refluxing of **17** with 1,4-dibromobutane or 2-bromoethyl ether under the existence of *N,N*-diisopropylethylamine in methanol furnished compounds **22** and **23**. Condensation of **28** and aminodiphenylmethane furnished an amide. Removal of the Boc protecting




Figure 6. Induction of cell death by compounds **20** and **25** in the human breast cancer MDA-MB-231 cell line and dependence on caspase-3/7. Cells were treated with different concentrations of Smac mimetic for 48 h and cell viability was determined using trypan blue exclusion assay. To determine the dependence of cell death induction by Smac mimetics on caspase-3/7, a cell-permeable caspase-3/7 inhibitor, Z-DEVD-FMK, was employed.




Figure 7. X-ray crystal structure of compound **21** in complex with human XIAP BIR3, determined to 2.8 Å resolution. (A) Overall representation of compound **21** in complex with XIAP BIR3. Compound **21** is shown in green ball-and-stick. (B) Detailed interactions between compound **21** and XIAP BIR3 residues shown as a 90° rotation of the yellow domain in A. Compound **21** is shown in yellow ball-and-stick. (C) Superposition of modeled and crystal structures of compound **21** in complex with XIAP BIR3.

group in this amide followed by condensation of the resulted ammonium salt with L-lactic acid yielded compound **24**.

Summary

On the basis of the crystal structure of Smac in complex with XIAP BIR3 domain, we have designed and synthesized a series of conformationally constrained, bicyclic Smac mimetics. Our studies have generated important structure–activity relationships for their binding affinities to XIAP BIR3 and for their activity in inhibition of cell growth in human cancer cells. Our efforts have led to the design of a number of potent and cell-permeable Smac mimetics such as **20** (SM-122), **21** (SM-130), and **25** (SM-230). These compounds bind to XIAP BIR3 with high affinities and antagonize XIAP BIR3 in cell-free functional assays. Importantly, they are potent and effective in inhibition of cell growth and in induction of apoptosis in cancer cells. Our determination of the

crystal structure of compound **21** in complex with XIAP BIR3 provides the structural basis for its high-affinity interaction with XIAP BIR3 and for further design and optimization. These potent and cell-permeable Smac mimetics serve as powerful pharmacological tools to further elucidate the role XIAP and other IAP proteins in apoptosis regulation and promising lead compounds for the development of a new class of anticancer drugs.

Experimental Section

1. Chemistry. General Methods. NMR spectra were acquired at a proton frequency of 300 MHz. ¹H chemical shifts are reported with TMS (0.00 ppm) or DHO (4.70 ppm) as internal standards. ¹³C chemical shifts are reported with CDCl₃ (77.00 ppm) or 1,4-dioxane (67.16 ppm) as internal standards. The final products were purified by HPLC on a Waters Sunfire C18 reverse phase semipreparative HPLC

column (19 mm × 150 mm) using solvent A (0.1% of TFA in water) and solvent B (0.1% of TFA in CH₃CN) as eluents with a flow rate of 10 mL/min.

General Synthetic Method for [8,5] Ring Contained Smac Mimetics. To a solution of compound **28** (1 mmol) in 10 mL of CH₂Cl₂ were added corresponding amine (1.2 mmol), EDC (1.2 mmol), HOBt (1.2 mmol), and *N,N*-diisopropylethylamine (3 mmol) at 0 °C. The mixture was stirred at room temperature overnight. After condensation, the residue was purified by chromatography to give an amide. To a solution of this amide in 5 mL of methanol was added 2 mL of HCl solution (4 N in 1,4-dioxane). The solution was stirred at room temperature overnight and then concentrated to give an ammonium salt. To a mixture of this salt in 10 mL of CH₂Cl₂ were added a desired *N*-Boc-L-amino acid (1.2 mmol), EDC (1.2 mmol), HOBt (1.2 mmol), and *N,N*-diisopropylethylamine (3 mmol) at 0 °C. The mixture was stirred at room temperature overnight and then concentrated. The residue was purified by chromatography to give an amide. To a solution of this amide in 5 mL of methanol was added 1 mL of HCl solution (4 N in 1,4-dioxane). The solution was stirred at room temperature overnight and then concentrated. The residue was purified by C18 reverse phase semipreparative HPLC to give the expected Smac mimetic.


(3S,6S,11S)-6-((S)-2-Amino-propionylamino)-5-oxo-decahydro-pyrrolo[1,2- α]jazocine-3-carboxylic Acid Benzhydryl-amide (16). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 M Hz, D₂O) δ 7.30–7.18 (m, 10H), 5.95 (s, 1H), 4.73 (m, 1H), 4.35 (m, 1H), 4.22 (m, 1H), 3.96 (m, 1H), 2.16 (m, 1H), 2.05 (m, 1H), 1.93–1.32 (m, 10 H), 1.39 (d, *J* = 7.5 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.67, 172.65, 169.93, 141.64, 141.45, 129.57, 129.45, 128.34, 128.09, 127.77, 62.50, 61.40, 58.16, 57.55, 51.42, 36.16, 33.30, 32.64, 28.07, 25.36, 22.14, 15.95. ESI MS: *m/z* 463.3 (M + H)⁺. HR ESI MS for C₂₇H₃₅N₄O₃ required, 463.2709; found, 463.2699; Anal. (C₂₇H₃₄N₄O₃ · 1.0HCl · 1.0CF₃COOH) C, H, N.

(3S,6S,11S)-6-((S)-2-Amino-butrylamino)-5-oxo-decahydro-pyrrolo[1,2- α]jazocine-3-carboxylic Acid Benzhydryl-amide (17). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 M Hz, D₂O) δ 7.30–7.18 (m, 10H), 5.95 (s, 1H), 4.71 (m, 1H), 4.34 (m, 1H), 4.22 (m, 1H), 3.83 (m, 1H), 2.14 (m, 1H), 2.03 (m, 1H), 1.93–1.20 (m, 12 H), 0.88 (t, *J* = 7.5 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.67, 172.60, 169.96, 141.64, 141.34, 129.54, 129.45, 128.37, 128.05, 127.78, 62.41, 61.39, 58.18, 54.65, 51.41, 36.22, 33.43, 32.66, 28.09, 25.34, 25.04, 22.16, 9.07. ESI MS: *m/z* 477.3 (M + H)⁺. HR ESI MS for C₂₈H₃₇N₄O₃ required, 477.2866; found, 477.2858. Anal. (C₂₈H₃₆N₄O₃ · 1.0HCl · 2.0H₂O) C, H, N.

(3S,6S,11S)-6-((S)-2-Amino-butrylamino)-5-oxo-decahydro-pyrrolo[1,2- α]jazocine-3-carboxylic Acid ((R)-naphthalen-1-yl-phenyl-methyl)-amide (18). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 MHz, D₂O) δ 7.75 (m, 1H), 7.65 (m, 1H), 7.58 (m, 1H), 7.32–7.10 (m, 4H), 7.10–6.92 (m, 5H), 6.58 (s, 1H), 4.63 (m, 1H), 4.27 (m, 1H), 3.80 (m, 1H), 2.12–1.65 (m, 6H), 1.65–1.20 (m, 8H), 0.86 (t, *J* = 7.5 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.19, 172.59, 169.83, 141.15, 136.53, 134.23, 131.35, 131.11, 129.40, 128.32, 128.04, 127.37, 126.66, 126.08, 124.02, 62.20, 61.18, 54.96, 54.60, 51.33, 36.32, 33.32, 32.61, 27.92, 25.33, 25.01, 22.19, 9.08. ESI MS: *m/z* 527.3 (M + H)⁺. HR ESI MS for C₃₂H₃₉N₄O₃ required, 527.3022; found, 527.3021. Anal. (C₃₂H₃₈N₄O₃ · 1.0HCl · 1.0CF₃COOH · 0.6H₂O) C, H, N.

(3S,6S,11S)-6-((S)-2-Amino-butrylamino)-5-oxo-decahydro-pyrrolo[1,2- α]jazocine-3-carboxylic Acid ((S)-naphthalen-1-yl-phenyl-methyl)-amide (19). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 MHz, D₂O) δ 7.58 (m, 1H), 7.20–6.50 (12H), 4.56 (m, 1H), 4.33 (m, 1H), 3.87 (m, 1H), 3.74 (m, 1H), 1.82–0.90 (m, 12H), 0.80 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 172.22,

Scheme 1. General Synthesis of Conformationally Constrained [8,5] Bicyclic Smac Mimetics^a



^a Reagents and conditions: (a) (i) Amine, EDC, HOBt, *N,N*-diisopropylethylamine, CH₂Cl₂, overnight; (ii) 4 N HCl in 1,4-dioxane, MeOH; (iii) *N*-Boc-L-amino acid, EDC, HOBt, *N,N*-diisopropylethylamine, CH₂Cl₂; (iv) 4 N HCl in 1,4-dioxane, MeOH. (b) 1,4-Dibromobutane or 2-bromoethyl ether, *N,N*-diisopropylethylamine, MeOH, reflux, 67% for **22** and 54% for **23**. (c) (i) Aminodiphenylmethane, EDC, HOBt, *N,N*-diisopropylethylamine, CH₂Cl₂, overnight; (ii) 4 N HCl in 1,4-dioxane, MeOH; (iii) L-lactic acid, EDC, HOBt, *N,N*-diisopropylethylamine, CH₂Cl₂, 71% over three steps.

172.16, 169.64, 141.38, 141.19, 139.58, 137.49, 134.18, 131.30, 129.21, 128.69, 128.18, 127.00, 125.97, 125.34, 124.57, 123.90, 62.01, 60.87, 54.62, 54.18, 51.13, 36.32, 33.58, 32.51, 27.74, 25.37, 25.04, 22.30, 9.10. ESI MS: *m/z* 527.3 (M + H)⁺. HR ESI MS for C₃₂H₃₉N₄O₃ required, 527.3022; found, 527.3013; Anal. (C₃₂H₃₈N₄O₃ · 1.0HCl · 1.0CF₃COOH · 0.8H₂O) C, H, N.

(3S,6S,11S)-6-((S)-2-Methylamino-propionylamino)-5-oxo-decahydro-pyrrolo[1,2- α]jazocine-3-carboxylic Acid Benzhydryl-amide (20). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 M Hz, D₂O) δ 7.30–7.18 (m, 10H), 5.95 (s, 1H), 4.74 (m, 1H), 4.34 (m, 1H), 4.24 (m, 1H), 3.83 (m, 1H), 2.57 (s, 3H), 2.21–1.50 (m, 11H), 1.40 (d, *J* = 7.0 Hz, 3H), 1.38 (m, 1H). ¹³C NMR (75 MHz, D₂O) δ 173.77, 172.62, 169.95, 141.56, 141.28, 129.54, 129.44, 128.40, 128.35, 128.01, 127.82, 62.50, 61.40, 58.16, 57.49, 51.42, 36.16, 33.30, 32.64, 31.61, 28.07, 25.36, 22.14, 15.95. Anal. (C₂₈H₃₆N₄O₃ · 1.0HCl · 1.0CF₃COOH) C, H, N.

(3S,6S,11S)-6-((S)-2-Methylamino-butrylamino)-5-oxo-decahydro-pyrrolo[1,2- α]jazocine-3-carboxylic Acid Benzhydryl-amide (21). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The putridity was checked by analytical HPLC to be over 98%. ¹H NMR (300 M Hz, D₂O) δ 7.32–7.19 (m, 10H), 5.95 (s, 1H), 4.75 (m, 1H), 4.36 (m, 1H), 4.25 (m, 1H), 3.71 (m, 1H), 2.56 (s, 3H), 2.18–1.40 (m, 14H), 0.87 (t, *J* = 7.6 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.76, 172.39, 168.73, 141.60, 141.27, 129.54, 129.44, 128.37, 128.05, 127.76, 62.96, 62.41, 61.44, 58.21, 51.51, 36.18, 33.40, 32.66, 32.04, 28.10, 25.38, 24.08, 22.11, 8.81. ESI MS: *m/z* 491.3 (M + H)⁺. HR ESI MS for C₂₉H₃₉N₄O₃ required, 491.3022; found, 491.3026; Anal. (C₂₉H₃₈N₄O₃ · 1.0HCl · 1.0CF₃COOH) C, H, N.

(3S,6S,11S)-6-((S)-2-pyrrolidin-1-yl-butrylamino)-5-oxo-decahydro-pyrrolo[1,2- α]jazocine-3-carboxylic Acid Benzhydryl-amide (22). To a solution of **17** (salt with HCl, 260 mg, 0.5 mmol) in 10 mL of methanol were added 1,4-dibromobutane (430 mg, 2 mmol) and *N,N*-diisopropylethylamine (0.5 mL). The solution was refluxed overnight and then condensed. The residue was dissolved in 5 mL of methanol. To this solution was added 0.5 mL of TFA. After condensation, the residue was purified by C18 reverse phase semipreparative HPLC to give **22** (215 mg, 67%). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 40 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 M Hz, D₂O) δ 7.30–7.19 (m, 10H), 5.95 (s, 1H), 4.77 (m, 1H), 4.36 (m, 1H), 4.25 (m, 1H), 3.68 (m, 1H), 3.50–2.70 (brs, 4H), 2.23–1.39 (m, 18H), 0.85 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.78, 172.23, 168.82, 141.60, 141.26, 129.54, 129.45, 128.38, 128.06, 127.74, 69.12, 62.40, 61.45, 58.22, 51.52, 36.17, 33.45, 32.66, 28.12, 26.40, 25.38, 23.64, 23.23, 22.10, 8.98. ESI MS: *m/z* 531.3

(M + H)⁺. HR ESI MS for C₃₂H₄₃N₄O₃ required, 531.3335; found, 531.3334. Anal. (C₃₂H₄₂N₄O₃·1.0HCl·1.3H₂O) C, H, N.

(3S,6S,11S)-6-((S)-2-Morpholin-4-yl-butylamino)-5-oxo-decahydro-pyrrolo[1,2- α]azocine-3-carboxylic Acid Benzhydryl-amide (23). To a solution of **17** (salt with HCl, 256 mg, 0.5 mmol) in 10 mL of methanol were added 2-bromoethyl ether (580 mg, 2.5 mmol) and *N,N*-diisopropylethylamine (0.5 mL). The solution was refluxed overnight and then condensed. The residue was dissolved in 5 mL of methanol. To this solution was added 0.5 mL of TFA. After condensation, the residue was purified by C18 reverse phase semipreparative HPLC to give **23** (178 mg, 54%). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 40 min. ¹H NMR (300 MHz, D₂O) δ 7.30–7.19 (m, 10H), 5.95 (s, 1H), 4.79 (m, 1H), 4.37 (m, 1H), 4.25 (m, 1H), 3.86 (brs, 4H), 3.67 (m, 1H), 3.31–3.20 (m, 4H), 2.18–1.39 (m, 14H), 0.85 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.77, 172.16, 167.95, 141.61, 141.26, 129.54, 129.45, 128.38, 128.06, 127.74, 70.21, 64.28, 62.40, 61.45, 58.22, 51.58, 51.02, 36.17, 33.44, 32.66, 28.12, 25.37, 22.09, 21.22, 9.14. ESI MS: *m/z* 547.3 (M + H)⁺. HR ESI MS for C₃₂H₄₃N₄O₄ required, 547.3284; found, 547.3298. Anal. (C₃₂H₄₂N₄O₄·1.0HCl·2.5H₂O) C, H, N.

(3S,6S,11S)-6-((S)-2-Hydroxy-propionylamino)-5-oxo-decahydro-pyrrolo[1,2- α]azocine-3-carboxylic Acid Benzhydryl-amide (24). To a solution of **28** (320 mg, 1 mmol) in 10 mL of CH₂Cl₂ were added aminodiphenylmethane (220 mg, 1.2 mmol), EDC (230 mg, 1.2 mmol), HOBT (160 mg, 1.2 mmol), and *N,N*-diisopropylethylamine (1.2 mL) at 0 °C. The mixture was stirred at room temperature overnight. After condensation, the residue was purified by chromatography to give an amide. To a solution of this amide in 5 mL of methanol was added 2 mL of HCl solution (4 N in 1,4-dioxane). The solution was stirred at room temperature overnight and then condensed to give an ammonium salt. To a mixture of this salt in 10 mL of CH₂Cl₂ were added L-lactic acid (110 mg, 1.2 mmol), EDC (232 mg, 1.2 mmol), HOBT (164 mg, 1.2 mmol), and *N,N*-diisopropylethylamine (1.2 mL) at 0 °C. The mixture was stirred at room temperature overnight and then condensed. The residue was purified by chromatography to give compound **24** (328 mg, 71% over three steps). ¹H NMR (300 MHz, CDCl₃, TMS) δ 7.78 (brd, *J* = 8.4 Hz, 1H), 7.37–7.20 (m, 11H), 6.22 (brd, *J* = 8.5 Hz, 1H), 4.88 (m, 1H), 4.64 (m, 1H), 4.23 (m, 2H), 3.80 (brs, 1H), 2.50 (m, 1H), 2.12 (m, 1H), 2.08–1.75 (m, 4H), 1.74–1.39 (m, 5H), 1.41 (d, *J* = 7.2 Hz, 3H), 1.27 (m, 1H). ¹³C NMR (75 MHz, CDCl₃) δ 174.17, 172.08, 169.77, 141.63, 141.47, 128.60, 128.52, 127.52, 127.36, 127.22, 68.27, 60.32, 59.48, 57.10, 49.55, 36.59, 35.36, 32.16, 25.03, 24.70, 22.94, 20.88. ESI MS: *m/z* 464.3 (M + H)⁺. HR ESI MS for C₂₇H₃₄N₃O₄ required, 464.2549; found, 464.2544. Anal. (C₂₇H₃₃N₃O₄·1.6H₂O) C, H, N.

(3S,6S,11S)-6-((S)-2-Methylamino-propionylamino)-5-oxo-decahydro-pyrrolo[1,2- α]azocine-3-carboxylic Acid ((R)-(1,2,3,4-Tetrahydro-naphthalen-1-yl)-amide (25). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 MHz, D₂O) δ 7.22–7.04 (m, 4H), 4.87 (m, 1H), 4.70 (m, 1H), 4.32–4.16 (m, 2H), 3.90–3.78 (m, 1H), 2.68–2.60 (m, 2H), 2.56 (s, 3H), 2.16–1.42 (m, 16), 1.40 (d, *J* = 7.2 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.62, 172.70, 169.86, 138.61, 136.40, 129.76, 128.93, 128.06, 126.85, 62.76, 61.44, 57.49, 51.48, 48.62, 36.26, 33.30, 32.65, 31.61, 30.08, 29.19, 28.18, 25.44, 22.18, 20.47, 15.94. ESI MS: *m/z* 441.3 (M + H)⁺. HR ESI MS for C₂₅H₃₇N₄O₃ required, 441.2866; found, 441.2862. Anal. (C₂₅H₃₆N₄O₃·1.0HCl·1.0CF₃COOH) C, H, N.

(3S,6S,11S)-6-((S)-2-Methylamino-propionylamino)-5-oxo-decahydro-pyrrolo[1,2- α]azocine-3-carboxylic Acid ((S)-(1,2,3,4-Tetrahydro-naphthalen-1-yl)-amide (26). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 MHz, D₂O) δ 7.20–7.02 (m, 4H), 4.88 (m, 1H), 4.22 (m, 1H), 4.16 (m, 1H), 3.83 (m, 1H), 2.69–2.60 (m, 2H), 2.57 (s, 3H), 2.20–1.52 (m, 15 H), 1.44 (m, 1H), 1.40 (d, *J* = 7.2 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 173.65, 172.61,

169.88, 138.67, 136.46, 129.89, 128.74, 128.13, 126.84, 63.01, 61.39, 57.48, 51.45, 48.44, 36.30, 33.30, 32.71, 31.64, 30.16, 29.10, 28.15, 25.43, 22.21, 20.22, 15.97. ESI MS: *m/z* 442.3 (M + H)⁺. HR ESI MS for C₂₅H₃₇N₄O₃ required, 441.2866; found, 441.2864. Anal. (C₂₅H₃₆N₄O₃·1.0HCl·1.0CF₃COOH) C, H, N.

(3S,6S,11S)-6-((S)-2-Methylamino-propionylamino)-5-oxo-decahydro-pyrrolo[1,2- α]azocine-3-carboxylic Acid ((R)-Indan-1-ylamide) (27). The gradient ran from 70% of solvent A and 30% of solvent B to 50% of solvent A and 50% of solvent B in 30 min. The purity was checked by analytical HPLC to be over 98%. ¹H NMR (300 MHz, D₂O) δ 7.23–7.06 (m, 4H), 5.19 (t, *J* = 7.8 Hz, 1H), 4.75 (m, 1H), 4.32–4.18 (m, 2H), 3.83 (m, 1H), 2.98–2.69 (m, 2H), 2.56 (s, 3H), 2.36 (m, 1H), 2.22–1.50 (m, 12 H), 1.43 (m, 1H), 1.40 (d, *J* = 7.0 Hz, 3H). ¹³C NMR (75 MHz, D₂O) δ 174.18, 172.66, 169.87, 144.29, 143.14, 128.66, 127.35, 125.45, 124.66, 62.78, 61.45, 57.48, 55.33, 51.48, 36.25, 33.39, 33.30, 32.65, 31.60, 30.27, 28.20, 25.44, 22.17, 15.94. ESI MS: *m/z* 427.3 (M + H)⁺. HR ESI MS for C₂₄H₃₅N₄O₃ required, 427.2709; found, 427.2707. Anal. (C₂₄H₃₄N₄O₃·1.0HCl·1.0CF₃COOH) C, H, N.

2. Molecular Modeling. The crystal structure of XIAP BIR3 in a complex with the Smac protein¹⁵ (PDB reference: 1G73) was used to predict the binding models of XIAP BIR3 bound to designed compounds. We used the Amber program suite⁴⁰ (version 7) to perform all molecular dynamics (MD) simulations. A recent version of the Amber force field (ff96)⁴¹ was used for the natural amino acids in the complex and the TIP3P model⁴² was used for water molecules. There is one Zn²⁺ ion covalently bound to C300, C303, H320, and C327 in the XIAP BIR3 domain. This Zn²⁺ ion, while important for structural integrity, has no direct interaction with the ligands. We used parameters developed by Ryde⁴³ for the Zn²⁺ ion and its coordination with the neighboring four residues to model this chelating structure in our simulation. All the MD simulations were carried out at NTP. The SHAKE algorithm⁴⁴ was used to fix the bonds involving hydrogen. The PME method⁴⁵ was used to account for long-range electrostatic interactions, and the nonbonded cutoff distance was set at 10 Å. The time step was 2 fs, and neighboring pairs list was updated after every 20 steps. For the refinement of the structure between compound **20** and XIAP BIR3, the protocol is as follows: A 500-step minimization of the solvated system was performed followed by 6 ps of MD simulation to gradually heat the system from 0 to 298 K. The system was then equilibrated by another 34 ps simulation at 298 K. Finally, the 1 ns production simulation was run and the snapshots of conformations (typically 2000), evenly spaced in time, were collected for structural analysis.

All the initial binding poses of designed ligands with XIAP BIR3 were predicted using the GOLD program⁴⁶ (version 2.2). The center of the binding site for XIAP BIR3 was set at Thr308 and the radius of the binding site was defined as 13 Å, large enough to cover the binding pocket. For each genetic algorithm run, a maximum number of 200000 operations were performed on a population of 5 islands of 100 individuals. Operator weights for crossover, mutation, and migration were set to 95, 95, and 10, respectively. The docking simulations were terminated after 20 runs for each ligand. GoldScore implemented in Gold 2.2 was used as the fitness function to evaluate the docked conformations. The 20 conformations ranked highest by each fitness function were saved for analysis of the predicted docking modes. For the docking poses reported, these were the highest ranked conformations from the docking simulations.

3. Protein Expression and Purification, Binding Assays to XIAP BIR3, and Cell-Free Functional Assays for the Activity of Caspase-9. Protein Expression and Purification. An N-terminal 6xHis-tagged BIR3 domain from human XIAP expressed in abundance of 50 mg of protein per liter of culture in *Escherichia coli* and was readily purified from soluble extracts. The sequence encoding the BIR3 domain (residues 241–356) of the human XIAP gene was cloned into the pET28a expression vector. High protein expression levels were achieved by growing transformed BL21(DE3) cells at 37 °C in 2xYT media containing 30 μ g/mL kanamycin to an optical density (600 nm) of 0.6 and inducing with 0.4 mM

isopropyl-D-thiogalactoside for 4 h at 25 °C. Cells pelleted at 5000g were frozen at -80 °C and resuspended in lysis buffer containing 50 mM Tris pH 7.5, 200 mM NaCl, 50 mM Zn acetate, 3 mM β -mercaptoethanol, and protease inhibitors. The cells were subjected to sonication and the cellular debris pelleted by centrifugation at 30000g for 20 min. The soluble fraction was applied to Ni-NTA agarose (Qiagen) pre-equilibrated with wash buffer containing 50 mM Tris pH 7.5, 200 mM NaCl, 50 mM Zn acetate, and 25 mM imidazole. The BIR3 domain protein was then eluted from the column using wash buffer containing 250 mM imidazole. To reduce the number of protein aggregates, a final concentration of 1 mM DTT was added to the elute before applying it to a Superdex-75 column (Pharmacia) equilibrated with 20 mM Tris pH 7.5, 200 mM NaCl, 50 mM Zn acetate, and 1 mM DTT. The resulting protein was judged to be greater than 98% pure by the presence of a single band on an SDS-PAGE gel.

Fluorescence-Polarization-Based Binding for XIAP BIR3 Protein. A sensitive in vitro binding assay using the fluorescence polarization (FP)-based method³² was used to determine the binding affinity of Smac mimetics to XIAP BIR3 protein. In this assay, 5-carboxyfluorescein was coupled to the lysine side chain of a mutated Smac peptide with the sequence (AbuRPFK-Fam). This fluorescently tagged peptide, named SM5F, was used as the fluorescent tracer in the FP-based binding assay of different compounds to the XIAP BIR3 protein (residues 241–356). The K_d value of SM5F peptide to XIAP BIR3 protein was determined to be 17.9 nM.³²

Dose-dependent binding experiments were carried out with serial dilutions of the tested compounds. An aliquot of the samples and preincubated XIAP BIR3 protein (0.030 μ M) and SM5F peptide (5 nM) in the assay buffer (100 mM potassium phosphate, pH 7.5; 100 μ g/mL bovine gamma globulin; 0.02% sodium azide, purchased from Invitrogen Life Technology), were added to 96-well plates. For each assay, the controls included XIAP BIR3 protein and SM5F (equivalent to 0% inhibition), and SM5F only (equivalent to 100% inhibition). The polarization values were measured after 3 h of incubation using an ULTRA READER (Tecan U.S. Inc., Research Triangle Park, NC). IC₅₀ values, the inhibitor concentration at which 50% of the bound tracer is displaced, were determined from a plot using nonlinear least-squares analysis. Curve fitting was performed using GRAPHPAD PRISM software (GraphPad Software, Inc., San Diego, CA).

Cell-Free Caspase-9 Functional Assay. MDA-MB-231 cell lysates were prepared by solubilizing cells in ice cold buffer (pH 7.5), which contained 50 mM KCl, 5 mM EGTA, 2 mM MgCl₂, 1 mM DTT, 0.2% CHAPS, 50 mM HEPES, and cocktail protease inhibitors and incubated on ice for 10 min. Cytochrome c and dATP were added into the cell lysates and incubated at 30 °C in a water bath for 60 min to activate caspase-9. To inhibit caspase-9 activity, recombinant human XIAP BIR3 protein (0.5 μ M) was added in the cell extract at the same time as Cytochrome c and dATP were added and was found to dose-dependently inhibit the activity of caspase-9. To determine caspase-9 activity, 25 μ M of caspase-9 substrate (Z-LEHD-AFC) purchased from BioVision was added. Fluorescence detection of substrate cleavage by caspase-9 was carried out with an ULTRA READER using an excitation wavelength of 400 nm and an emission wavelength of 505 nm, and the reaction was monitored for 60–120 min. To determine the antagonism of a Smac mimetic, different concentrations of a tested Smac mimetic (from 100 μ M to 1 nM) were added into the reaction mixture together with XIAP BIR3 protein.

IV. Assays for Cell Growth, Apoptosis, and Cell Viability.

Cell Lines. MDA-M-231 and SK-OV-3 cancer cell lines were purchased from the American Type Culture Collection (Manassas, VA), maintained in RPMI 1640 (Invitrogen), supplemented with 10% fetal bovine serum (Invitrogen) and 1% penicillin-streptomycin, at 37 °C in 5% CO₂. Both cell lines were cultured in RPMI 1640 supplemented with 10% fetal bovine serum (Invitrogen) and 1% penicillin-streptomycin in the presence of puromycin from EMD Biosciences (2 μ g/ml).

Table 2. Data Collection, Phasing, and Refinement Statistics of the Crystal Structure of XIAP BIR3:Compound **21**

| data collection statistics | |
|---|---|
| data set | XIAP BIR3: 21 |
| space group | <i>P</i> 6 ₅ 22 |
| unit cell (Å) | <i>a</i> = <i>b</i> = 115.76; <i>c</i> = 61.79 |
| wavelength (Å) | 1.0000 |
| resolution (Å) ^a | 2.8 (2.9–2.8) |
| <i>R</i> _{sym} (%) ^b | 7.2 (30.8) |
| $\langle I/\sigma I \rangle$ ^c | 20 (3) |
| completeness (%) ^d | 99.3 (100) |
| redundancy | >9 |
| refinement statistics | |
| resolution (Å) ^e | 2.8 |
| <i>R</i> -factor (%) ^f | 22.8 (37.5) |
| <i>R</i> _{free} (%) ^g | 26.9 (46.3) |
| protein atoms ^h | 784 |
| water molecules | 38 |
| unique reflections | 6181 |
| rmsd ⁱ | |
| bonds (Å) | 0.007 |
| angles (deg) | 1.203 |

^a Statistics for highest resolution bin of reflections in parentheses. ^b $R_{\text{sym}} = \sum_h \sum_j |I_{hj} - \langle I_h \rangle| / \sum_h \sum_j I_{hj}$, where I_{hj} is the intensity of observation j of reflection h and $\langle I_h \rangle$ is the mean intensity for multiply recorded reflections. ^c Intensity signal-to-noise ratio. ^d Completeness of the unique diffraction data. ^e Resolution cut-off used during heavy-atom refinement and phase calculations. ^f $R\text{-factor} = \sum_h \|F_o\| - \|F_c\| / \sum_h \|F_o\|$, where F_o and F_c are the observed and calculated structure factor amplitudes for reflection h . ^g R_{free} is calculated against a 10% random sampling of the reflections that were removed before structure refinement. ^h Total number of protein atoms refined in the asymmetric unit. ⁱ Root mean square deviation of bond lengths and bond angles.

Cell Growth Assay. The effect of Smac mimetics on cell growth was evaluated by a WST-8 [2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2*H*-tetrazolium monosodium salt] assay (Dojindo Molecular Technologies, Inc.). Cells (3000–4000 cells in each well) were cultured in 96-well tissue culture plates in medium (200 μ L) containing various concentrations of Smac mimetics for the indicated time. At the end of incubation, WST-8 dye (20 μ L) was added to each well and incubated for an additional 1–3 h, and then the absorbance was measured in a microplate reader (Molecular Devices) at 450 nm. Cell growth inhibition was evaluated as the ratio of the absorbance of the sample to that of the control.

Apoptosis Assay. The apoptosis assay was performed with an annexin-V/propidium iodide (PI) apoptosis detection kit (Roche) according to the manufacturer's instructions. Briefly, cells were harvested, washed with ice-cold PBS, and then stained with annexin-V-FITC and PI for 15 min at room temperature in the dark. Stained cells were analyzed in a FACS calibur flow cytometer. Annexin-V (-)/PI (-) cells were classified as live cells; Annexin-V (+)/PI (-) cells were classified as early apoptotic cells; Annexin-V (+)/PI (+) cells were classified as late apoptotic cells; Annexin V (-) and P.I. (+) cells were classified as dead cells.

Cell Viability Assay. Cell viability was determined using a trypan blue exclusion assay. Approximately 10000 MDA-MB-231 cells were seeded in 24-well plates and incubated at 37 °C in an atmosphere of 95% air and 5% CO₂ for 24 h before drug treatments. Cells were treated with different concentrations of compounds **20** and **25** for 48 h. To examine the role of caspase-3/-7 in cell death induction by Smac mimetics, a cell-permeable caspase-3/-7 inhibitor, Z-DEVD-FMK (Calbiochem, San Diego, CA) was employed and cells were treated with a combination of Z-DEVD-FMK with compound **20** or **25** concurrently. To determine cell viability, 1:1 dilution of the suspension of 0.4% Trypan blue reagent (Invitrogen Corporation) was used. Blue cells were counted as nonviable cells. At least 100 cells were counted in the control well, and three independent countings were performed.

4. Crystallographic Studies. Crystallization of XIAP BIR3: Compound 21 Complex. Bipyramidal crystals of the BIR3 domain of human XIAP (residues 241–356) in complex with compound **21** were grown at 20 °C from hanging drop vapor diffusion experiments by mixing 2 μ L of protein (10 mg/ml preincubated with 10 mM compound **21**) in 20 mM Tris pH 7.5, 200 mM NaCl, 50 mM Zn acetate, and 1 mM DTT with 2 μ L of precipitant (20% polyethylene glycol 8000, 0.2 M MgCl₂, and 0.1 M Tris-HCl pH 8.5). Crystals were cryoprotected with paratone-N and data collected at a wavelength of 1.0 Å at the Advanced Photon Source on the COMCAT 32-ID beam line equipped with a MAR165 CCD and processed with DENZO/SCALEPACK.³⁵ The complex crystallized in space group *P*6₅22 with a solvent content of 66%,³⁶ resulting in one molecule per asymmetric unit. The data are 99.3% complete to 2.8 Å resolution with a *R*_{sym} of 7.2%.

Structure Determination and Refinement. The BIR3:compound **21** complex was solved to 2.8 Å resolution by molecular replacement methods in CNS³⁷ using the crystal structure of XIAP BIR3 (residues 256–346) with another Smac mimetic solved in our laboratory as the search model. Difference Fourier maps (*F*_o – *F*_c) calculated after CNS rigid-body refinement of the BIR3 domain against BIR3:compound **21** diffraction data showed the presence of compound **21** and six additional residues at the *N*-terminus (250–255). Multiple rounds of CNS refinement and model building using O³⁸ resulted in a BIR3:compound **21** structure refined to 2.8 Å resolution (Table 1). Simulated annealing omit maps calculated in CNS were used to check for the correct placement of atoms in the structure. The final structure was analyzed in PROCHECK.³⁹ All residues are in the allowed regions of the Ramachandran plot. The *N*-terminal residues (241–249) and the *C*-terminal residues (346–356) were disordered in the structure.

Acknowledgment. The financial support from the Breast Cancer Research Foundation, the National Cancer Institute, NIH (R01CA109025), the Prostate Cancer Foundation, the Department of Defense Prostate Cancer Program (W81XWH-04-1-0213), and Ascenta Therapeutics are greatly appreciated. We thank Dr. G. W. A. Milne and Karen Kreutzer for critical reading of the manuscript.

Supporting Information Available: Elemental analysis data for compounds **16–27**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Kerr, J. F.; Wyllie, A. H.; Currie, A. R. Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br. J. Cancer* **1972**, *26*, 239–257.
- Thompson, C. B. Apoptosis in the pathogenesis and treatment of disease. *Science* **1995**, *267*, 1456–1462.
- Nicholson, D. W. From bench to clinic with apoptosis-based therapeutic agents. *Nature* **2000**, *407*, 810–816.
- Reed, J. C. Apoptosis-based therapies. *Nat. Rev. Drug Discovery* **2002**, *1*, 111–121.
- Deveraux, Q. L.; Reed, J. C. IAP family proteins-suppressors of apoptosis. *Genes Dev.* **1999**, *1*, 239–252.
- Salvesen, G. S.; Duckett, C. S. IAP proteins: blocking the road to death's door. *Nat. Rev. Mol. Cell Biol.* **2002**, *3*, 401–410.
- Riedl, S. J.; Shi, Y. Molecular mechanisms of caspase regulation during apoptosis. *Nat. Rev. Mol. Cell Biol.* **2004**, *5*, 897–907.
- LaCasse, E. C.; Baird, S.; Korneluk, R. G.; MacKenzie, A. E. The inhibitors of apoptosis (IAPs) and their emerging role in cancer. *Oncogene* **1998**, *17*, 3247–3259.
- Holcik, M.; Gibson, H.; Korneluk, R. G. XIAP: Apoptotic brake and promising therapeutic target. *Apoptosis* **2001**, *6*, 253–261.
- Huang, Y.; Park, Y. C.; Rich, R. L.; Segal, D.; Myszka, D. G.; Wu, H. Structural basis of caspase inhibition by XIAP: differential roles of the linker versus the BIR domain. *Cell* **2001**, *104*, 781–790.
- Tamm, I.; Kornblau, S. M.; Segall, H.; Krajewski, S.; Welsh, K.; Kitada, S.; Scudiero, D. A.; Tudor, G.; Qui, Y. H.; Monks, A.; Andreeff, M. Expression and prognostic significance of IAP-family genes in human cancers and myeloid leukemias. *Clin. Cancer Res.* **2000**, *6*, 1796–1803.
- Fulda, S. Inhibitor of apoptosis proteins as targets for anticancer therapy. *Expert Rev. Anticancer Ther.* **2007**, *7*, 1255–64.
- Du, C.; Fang, M.; Li, Y.; Li, L.; Wang, X. Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. *Cell* **2000**, *102*, 33–42.
- Verhagen, A. M.; Ekert, P. G.; Pakusch, M.; Silke, J.; Connolly, L. M.; Reid, G. E.; Moritz, R. L.; Simpson, R. J.; Vaux, D. L. Identification of DIABLO, a mammalian protein that promotes apoptosis by binding to and antagonizing IAP proteins. *Cell* **2000**, *102*, 43–53.
- Wu, G.; Chai, J.; Suber, T. L.; Wu, J. W.; Du, C.; Wang, X.; Shi, Y. Structural basis of IAP recognition by Smac/DIABLO. *Nature* **2000**, *408*, 1008–1012.
- Liu, Z.; Sun, C.; Olejniczak, E. T.; Meadows, R.; Betz, S. F.; Oost, T.; Herrmann, J.; Wu, J. C.; Fesik, S. W. Structural basis for binding of Smac/DIABLO to the XIAP BIR3 domain. *Nature* **2000**, *408*, 1004–1008.
- Srinivasula, S. M.; Hegde, R.; Saleh, A.; Datta, P.; Shiozaki, E.; Chai, J.; Lee, R. A.; Robbins, P. D.; Fernandes-Alnemri, T.; Shi, Y.; Alnemri, E. S. A conserved XIAP-interaction motif in caspase-9 and Smac/DIABLO regulates caspase activity and apoptosis. *Nature* **2001**, *410*, 112–116.
- Shiozaki, E. N.; Chai, J.; Rigotti, D. J.; Riedl, S. J.; Li, P.; Srinivasula, S. M.; Alnemri, E. S.; Fairman, R.; Shi, Y. Mechanism of XIAP-mediated inhibition of caspase-9. *Mol. Cell* **2003**, *11*, 519–527.
- Huang, Y.; Rich, R. L.; Myszka, D. G.; Wu, H. Requirement of both the second and third BIR domains for the relief of X-linked inhibitor of apoptosis protein (XIAP)-mediated caspase inhibition by Smac. *J. Biol. Chem.* **2003**, *278*, 49517–49522.
- Fulda, S.; Wick, W.; Weller, M.; Debatin, K. M. Smac agonists sensitize for Apo2L/TRAIL- or anticancer drug-induced apoptosis and induce regression of malignant glioma in vivo. *Nat. Med.* **2002**, *8*, 808–815.
- Arnt, C. R.; Chiorean, M. V.; Heldebrandt, M. P.; Gores, G. J.; Kaufmann, S. H. Synthetic Smac/DIABLO Peptides Enhance the Effects of Chemotherapeutic Agents by Binding XIAP and cIAP1 in Situ. *J. Biol. Chem.* **2002**, *277*, 44236–44243.
- Yang, L.; Mashima, T.; Sato, S.; Mochizuki, M.; Sakamoto, H.; Yamori, T.; Oh-Hara, T.; Tsuruo, T. Predominant suppression of apoptosis by inhibitor of apoptosis protein in non-small cell lung cancer H460 cells: therapeutic effect of a novel polyarginine-conjugated Smac peptide. *Cancer Res.* **2003**, *63*, 831–837.
- Sun, H.; Nikolovska-Coleska, Z.; Yang, C.-Y.; Xu, L.; Liu, M.; Tomita, Y.; Pan, H.; Yoshioka, Y.; Krajewski, K.; Roller, P. P.; Wang, S. Structure-Based Design of Potent, Conformationally Constrained Smac Mimetics. *J. Am. Chem. Soc.* **2004**, *126*, 16686–16687.
- Sun, H.; Nikolovska-Coleska, Z.; Yang, C.-Y.; Xu, L.; Tomita, Y.; Krajewski, K.; Roller, P. P.; Wang, S. Structure-based design, synthesis, and evaluation of conformationally constrained mimetics of the second mitochondria-derived activator of caspase that target the X-linked inhibitor of apoptosis protein/caspase-9 interaction site. *J. Med. Chem.* **2004**, *47*, 4147–50.
- Oost, T. K.; Sun, C.; Armstrong, R. C.; Al-Assaad, A. S.; Betz, S. F.; Deckwerth, T. L.; Ding, H.; Elmore, S. W.; Meadows, R. P.; Olejniczak, E. T.; Oleksijew, A.; Oltersdorf, T.; Rosenberg, S. H.; Shoemaker, A. R.; Tomaselli, K. J.; Zou, H.; Fesik, S. W. Discovery of potent antagonists of the antiapoptotic protein XIAP for the treatment of cancer. *J. Med. Chem.* **2004**, *47*, 4417–26.
- Sun, H.; Nikolovska-Coleska, Z.; Lu, J.; Qiu, S.; Yang, C.-Y.; Gao, W.; Meagher, J.; Stuckey, J.; Wang, S. Design, Synthesis, and Evaluation of a Potent, Cell-Permeable, Conformationally Constrained Second Mitochondria Derived Activator of Caspase (Smac) Mimetic. *J. Med. Chem.* **2006**, *49*, 7916–7920.
- Zobel, K.; Wang, L.; Varfolomeev, E.; Franklin, M. C.; Elliott, L. O.; Wallweber, H. J.; Okawa, D. C.; Flygare, J. A.; Vucic, D.; Fairbrother, W. J.; Deshayes, K. Design, Synthesis, and Biological Activity of a Potent Smac Mimetic That Sensitizes Cancer Cells to Apoptosis by Antagonizing IAPs. *ACS Chem. Biol.* **2006**, *1*, 525–533.
- Chauhan, D.; Neri, P.; Velankar, M.; Podar, K.; Hideshima, T.; Fulciniti, M.; Tassone, P.; Raje, N.; Mitsiades, C.; Mitsiades, N.; Richardson, P.; Zavel, L.; Tran, M.; Munshi, N.; Anderson, K. C. Targeting mitochondrial factor Smac/DIABLO as therapy for multiple myeloma (MM). *Blood* **2007**, *109*, 1220–1227.
- Li, L.; Thomas, R. M.; Suzuki, H.; De Brabander, J. K.; Wang, X.; Harran, P. G. A Small Molecule Smac Mimetic Potentiates TRAIL- and TNF-Mediated Cell Death. *Science* **2004**, *305*, 1471–1474.
- Sun, H.; Nikolovska-Coleska, Z.; Lu, J.; Meagher, J. L.; Yang, C.-Y.; Qiu, S.; Tomita, Y.; Ueda, Y.; Jiang, S.; Krajewski, K.; Roller,

- P. P.; Stuckey, J. A.; Wang, S. Design, Synthesis, and Characterization of a Potent, Nonpeptide, Cell-Permeable, Bivalent Smac Mimetic That Concurrently Targets Both the BIR2 and BIR3 Domains in XIAP. *J. Am. Chem. Soc.* **2007**, *129*, 15279–15294.
- (31) Duggan, H. M. E.; Hitchcock, P. B.; Young, D. W. Synthesis of 5/7-, 5/8- and 5/9-bicyclic lactam templates as constraints for external beta-turns. *Org. Biomol. Chem.* **2005**, *3*, 2287–2295.
- (32) Nikolovska-Coleska, Z.; Wang, R.; Fang, X.; Pan, H.; Tomita, Y.; Li, P.; Roller, P. P.; Krajewski, K.; Saito, N. G.; Stuckey, J. A.; Wang, S. Development and optimization of a binding assay for the XIAP BIR3 domain using fluorescence polarization. *Anal. Biochem.* **2004**, *332*, 261–273.
- (33) Verdecia, M. A.; Huang, H.; Dutil, E.; Kaiser, D. A.; Hunter, T.; Noel, J. P. Structure of the human anti-apoptotic protein survivin reveals a dimeric arrangement. *Nat. Struct. Biol.* **2000**, *7*, 602–608.
- (34) Franklin, M. C.; Kadkhodayan, S.; Ackerly, H.; Alexandru, D.; Distefano, M. D.; Elliot, L. O.; Flygare, J. A.; Mausisa, G.; Okawa, D. C.; Ong, D.; Vucic, D.; Deshayes, K.; Fairbrother, W. J. Structure and Function Analysis of Peptide Antagonists of Melanoma Inhibitor of Apoptosis (ML-IAP). *Biochemistry* **2003**, *42*, 8223–8231.
- (35) Otwinowski, Z.; Minor, W., *Processing of X-ray Diffraction Data Collected in Oscillation Mode in Methods in Enzymology, Volume 276: Macromolecular Crystallography, Part A*; Carter, C. W., Jr., Sweet, R. M., Eds.; Academic Press: New York, 1997; pp 307–326.
- (36) Matthews, B. W. Solvent content of protein crystals. *J. Mol. Biol.* **1968**, *33*, 491–497.
- (37) Brunger, A. T.; Adams, P. D.; Clore, G. M.; Delano, W. L.; Gros, P.; Grosse-Kunstleve, R. W.; Jiang, J.-S.; Kuszewski, J.; Nigles, N.; Pannu, N. S.; Read, R. J.; Rice, L. M.; Simonson, T.; Warren, G. L. Crystallography and NMR system (CNS): a new software system for macromolecular structure determination. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **1998**, *54*, 905–921.
- (38) Jones, T. A.; Zou, J. Y.; Cowan, S. W.; Kjeldgaard, M. Improved methods for building protein models in electron density maps and the location of errors in these models. *Acta Crystallogr., Sect. A: Found. Crystallogr.* **1991**, *47*, 110–119.
- (39) Laskowski, R. A.; MacArthur, M. W.; Moss, D. S.; Thornton, J. M. PROCHECK—a program to check the stereochemical quality of protein structures. *J. Appl. Crystallogr.* **1993**, *26*, 283–291.
- (40) Case, D. A.; Pearlman, D. A.; Caldwell, J. W.; Cheatham, T. E., III.; Wang, J.; Ross, W. S.; Simmerling, C. L.; Darden, T. A.; Merz, K. M.; Stanton, R. V.; Cheng, A. L.; Vincent, J. J.; Crowley, M.; Tsui, V.; Gohlke, H.; Radmer, R. J.; Duan, Y.; Pitera, J.; Massova, I.; Seibel, G. L.; Singh, U. C.; Weiner, P. K.; Kollman, P. A. *AMBER7*; University of California: San Francisco, 2002.
- (41) Kollman, P. A.; Dixon, R.; Cornell, W.; Fox, T.; Chipot, C.; Pohorille, A. The development/application of a “minimalist” organic/biochemical molecular mechanics force field using a combination of ab initio calculations and experimental data. In *Computer Simulation of Biomolecular Systems, Vol. 3*; Wilkinson, A., Weiner, P., van Gunsteren, W. F., Eds.; Elsevier: New York, 1997 pp 83–96.
- (42) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **1983**, *79*, 926–935.
- (43) Ryde, U. Molecular Dynamics Simulations of Alcohol Dehydrogenase with a four- or five-coordinate catalytic zinc ion. *Proteins* **1995**, *21*, 40–56.
- (44) Rychaert, J. P.; Ciccotti, G.; Berendsen, H. J. C. Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of *n*-alkanes. *J. Comput. Phys.* **1977**, *23*, 327–341.
- (45) Darden, T. A.; York, D. M.; Pedersen, L. Particle mesh Ewald: An $N \log(N)$ method for Ewald sums in large systems. *J. Chem. Phys.* **1993**, *98*, 10089–10092.
- (46) Jones, G.; Willett, P.; Glen, R. C.; Leach, A. R.; Taylor, R. Development and validation of a genetic algorithm for flexible docking. *J. Mol. Biol.* **1997**, *267*, 727–748.

JM8006849