

Structure Mechanism Insights and the Role of Nitric Oxide Donation Guide the Development of Oxadiazole-2-Oxides as Therapeutic Agents against Schistosomiasis

Ganesha Rai,[†] Ahmed A. Sayed,^{‡,§} Wendy A. Lea,[†] Hans F. Luecke,^{||} Harinath Chakrapani,[⊥] Stefanie Prast-Nielsen,[#] Ajit Jadhav,[†] William Leister,[†] Min Shen,[†] James Inglese,[†] Christopher P. Austin,[†] Larry Keefer,[⊥] Elias S. J. Arnér,[#] Anton Simeonov,[†] David J. Maloney,[†] David L. Williams,^{*,‡,∇} and Craig J. Thomas^{*,†}

[†]NIH Chemical Genomics Center, National Human Genome Research Institute, NIH, 9800 Medical Center Drive, MSC 3370, Bethesda, Maryland 20892-3370, [‡]Department of Biological Sciences, Illinois State University, Normal, Illinois 61790, [§]Department of Biochemistry, Ain Shams University, Cairo, Egypt, ^{||}Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, NIH, Bethesda, Maryland 20892-3370, [⊥]Laboratory of Comparative Carcinogenesis, National Cancer Institute, NIH, Frederick, Maryland 21702-1201, [#]Department of Medical Biochemistry and Biophysics, Division of Biochemistry, Karolinska Institutet, Stockholm SE-17177, Sweden, and [∇]Department of Immunology/Microbiology, Rush University Medical Center, 1735 West Harrison Street, Chicago, Illinois 60612

Received July 10, 2009

Schistosomiasis is a chronic parasitic disease affecting hundreds of millions of individuals worldwide. Current treatment depends on a single agent, praziquantel, raising concerns of emergence of resistant parasites. Here, we continue our explorations of an oxadiazole-2-oxide class of compounds we recently identified as inhibitors of thioredoxin glutathione reductase (TGR), a selenocysteine-containing flavoenzyme required by the parasite to maintain proper cellular redox balance. Through systematic evaluation of the core molecular structure of this chemotype, we define the essential pharmacophore, establish a link between the nitric oxide donation and TGR inhibition, determine the selectivity for this chemotype versus related reductase enzymes, and present evidence that these agents can be modified to possess appropriate drug metabolism and pharmacokinetic properties. The mechanistic link between exogenous NO donation and parasite injury is expanded and better defined. The results of these studies verify the utility of oxadiazole-2-oxides as novel inhibitors of TGR and as efficacious antischistosomal agents.

Introduction

The chronic infectious disease schistosomiasis is a major neglected disease caused by trematode flatworms of the genus *Schistosoma*.¹ Estimates suggest that a staggering 200+ million individuals suffer with this disease worldwide, and the sub-Saharan Africa region experiences an estimated 280000 deaths annually from schistosomiasis.^{2,3} Further, schistosomiasis causes altered immune responses to infections and vaccinations, increased perinatal transmission of HIV, and increased susceptibility to and transmission of malaria, tuberculosis, hepatitis C, and HIV.^{1,4} Schistosomiasis ranks among the most prominent neglected diseases due to the lack of interest from commercial pharmaceutical enterprises. It is rampant in impoverished populations due to inadequate sanitation and tainted water supplies, and the burden of infection perpetuates these conditions. There is currently no vaccine for schistosomiasis, and treatment of the disease is accomplished solely through the use of praziquantel (PZQ^a) (I) (Figure 1).^{5,6} The low cost of this agent (<\$0.25 USD/dose) and its efficacy against the adult form of all schistosome

species has led to its widespread use (tens of millions of annual treatments).⁷ Because individuals do not acquire immunity after treatment and are rapidly reinfected, treatment must be administered annually or biannually. Although PZQ's mechanism of action is not fully understood, research suggests that PZQ alters homeostasis of calcium, resulting in a state of schistosomal paralysis that leads to exposure and increased susceptibility to the host immune system.⁸ A recent study has demonstrated that RNAi knockdown of the voltage-operated calcium channel β subunit ablated praziquantel efficacy in *Dugesia japonica*.⁹ The dependence and extensive use of PZQ has triggered concerns about the emergence of drug resistant parasites, and evidence of drug resistance has been reported.¹⁰ Given the solitary and precarious reliance on PZQ for the control of such a widespread disease, there is an urgent need to define novel targets and drugs for the treatment of schistosomiasis.

The schistosome life cycle requires a snail intermediate host and a mammalian definitive host. Schistosome development in humans is complex and begins with cercarial penetration of human skin. Following entry into the circulatory system, parasites travel through the lungs and ultimately situate in the liver where they grow, differentiate, mate, and then migrate to the urogenital system or mesenteric veins of the hepatoportal system to commence egg production. Given their wide tissue distribution and long lifespan (up to 30 years), schistosomes have evolved robust defense mechanisms to mitigate oxidative damage and the host immune response.

*To whom correspondence should be addressed. For C.J.T.: phone, 301-217-4079; fax, 301-217-5736; E-mail, craigt@nhgri.nih.gov. For D.L.W.: phone, 312-942-1375; fax, 312-942-2808; E-mail, david_l_williams@rush.edu.

^a Abbreviations: TGR, thioredoxin glutathione reductase; NO, nitric oxide; PZQ, praziquantel; GR, glutathione reductase; TrxR, thioredoxin reductase; Prx2, peroxiredoxin2; GSH, glutathione.

Most eukaryotes have multiple enzymatic processes to detoxify reactive oxygen species. In humans, these detoxification pathways depend upon two NADPH-dependent flavoenzymes that maintain glutathione and thioredoxin in their reduced state, aptly named glutathione reductase (GR) and thioredoxin reductase (TrxR).^{11,12} Biochemical, reverse genetic, and genomic analysis of *Schistosoma mansoni* provide evidence that reduction of both glutathione and thioredoxin in schistosomes is dependent upon a single multifunctional selenocysteine-containing flavoenzyme designated thioredoxin glutathione reductase (TGR).¹³ A crystal structure of TGR (which contains a total of 13 cysteine residues and one selenocysteine) has been presented and offers a structural rationale for this dual activity.¹⁴ The dependence of schistosomes on a single, biochemically unique redox system is theorized to present an opportunity for drug development. Williams and co-workers revealed TGR as the putative molecular target of potassium antimonyl tartrate (a former treatment option for schistosomiasis) and that RNA interference targeting TGR resulted in rapid larval death (>90% in 4 days).¹⁵ Additionally, the disease-modifying antirheumatic drug auranofin (a gold containing sugar conjugate) was found to be a potent inhibitor of TGR and killing of ex vivo parasites by this agent paralleled inhibition of TGR.^{13,15}

These results present convincing evidence that TGR is an appropriate molecular target for pharmacological intervention and potential chemotherapy for schistosomiasis. We recently reported the development and validation of assays based upon the redox cascade of TGR in concert with peroxiredoxin2 (Prx2) using fluorescence intensity change associated with the consumption of NADPH¹⁶ and via a colorimetric approach utilizing reduction of Ellman's reagent.¹⁷ A quantitative high-throughput screen¹⁸ of the NIH Molecular Libraries¹⁹ Small Molecule Repository was performed, and numerous small molecule actives were identified.¹⁶ The chemotypes identified from this screen showed a range of potencies (including several in the low nanomolar range) and were validated as agents capable of inhibiting TGR in both the screening (Prx2-coupled NADPH consumption) and the follow-up (DTNB reduction) assay formats. Selected compounds were then advanced to investigations involving ex vivo worm cultures from each *Schistosoma* species that infects humans (*S. mansoni*, *S. japonicum*, and *S. hematobium*). One agent, the well-studied oxadiazole-2-oxide designated furoxan (**2**) (Figure 1), was capable of potent ex vivo worm killing at modest concentrations (10 μ M) and, importantly, against all developmental stages of worm (transformed skin larva, lung-stage schistosomula, juvenile liver worms, and adult worms).²⁰ On the basis of this encouraging result, furoxan (**2**) was entered into an in vivo evaluation in *S. mansoni* infected mice. Five consecutive daily intraperitoneal treatments of 10 mg/kg of **2** were performed in mice during the skin-stage infection (day 1–5), the juvenile/liver-stage infection (day 23–27), and the adult/egg-laying stage infection (day 37–41), and efficacy was quantified by adult worm burdens at day 49.²⁰ Significant reduction of worm burdens and hepatomegaly and splenomegaly were noted for treatment groups at each parasite life stage. Both male and female worms were affected, and a lower number of egg-induced granulomas were noted in comparison to control groups. Taken as a whole, the activity of furoxan (**2**) far exceeds the activity criteria set forth by the World Health Organization for candidate therapeutic development for schistosomiasis.²¹




Figure 1. Structures of praziquantel (**1**), furoxan (**2**), and bis(thiophen-2-ylmethanone)-furoxan (**3**).

The oxadiazole-2-oxide core is a well-studied structural motif explored by Ponzio in the early half of the 20th century and more recently expanded upon by Gasco and co-workers as a preeminent class of nitric oxide (NO) donating compounds.^{22,23} NO [the nitrogen–oxygen radical (NO[•]) not to be confused with the nitroxyl anion (NO⁻)]^{24,25} is a well-studied physiological signaling agent best known for its ability to relax smooth muscle tissue, resulting in vasodilatation and increased blood flow.²⁶ NO has been implicated in numerous biochemical processes and, importantly, elevated and sustained levels of NO can be toxic. Gasco first reported the thiol mediated release of NO from furoxan (**2**) and the resulting vasodilatory properties in 1994.²⁷ Since that time, the core oxadiazole-2-oxide moiety has been conjugated to numerous bioactive compounds in attempts to make chimeric small molecules with dual activities including NO-donating β -adrenoceptor agonists,²⁸ NO-donating aspirin conjugates,²⁹ and NO-donating proton pump inhibitors.³⁰

There is ample direct and indirect evidence that NO can act as an antischistosomal and, more broadly, antiparasitic molecule.^{31–33} NO produced by human white cells has been shown to kill larval schistosome parasites.³⁴ NO produced through inducible NO synthase activation is a core component of the immune response, and direct evidence of NO being a primary aspect of host defense against *Echinococcus granulosus* and *Trypanosoma congolense* has been presented.^{35,36} Winberg et al. recently demonstrated that NO production diminished *Leishmania donovani* promastigotes ability to inhibit periphagosomal F-actin breakdown, allowing normal phagosomal maturation and parasite killing.³⁷ Beyond innate activation of the NO synthase, exogenous NO from small molecule NO donors has shown promise in several arenas. A topical formulation of S-nitroso-N-acetylpenicillamine (SNAP) was reported as an effective treatment option for the control of cutaneous leishmaniasis.³⁸ A class of bicyclic nitroimidazoles was recently reported as a deazaflavin-dependent nitroreductase prodrug that releases oxidized nitrogen molecules that are converted to NO and are capable of anaerobic killing of *Mycobacterium tuberculosis*.³⁹ Further, the mainstays of Chagas disease chemotherapy are benznidazole and nifurtimox (agents with heteroaromatic nitro moieties).⁴⁰ Recently, Wilkinson et al. demonstrated that these prodrugs are activated in *Trypanosoma cruzi* by a NADPH-dependent type I nitroreductase.⁴¹ It is entirely plausible that this process results in the generation of NO.

Given the relationship between NO and antiparasitic activity and the prior research on furoxan (**2**) as a NO donor compound, it was sensible to assess the NO generating capacity of these agents. In the presence of a free thiol (cysteine), it was determined that furoxan (**2**) was an effective NO donor compound (as judged by ABTS⁴² oxidation).²⁰ Analysis of NO release by **2** in the presence of TGR was found to be dependent on the presence of NADPH. Interestingly,

several oxadiazole-2-oxide analogues with potent TGR inhibition, for instance bis(thiophen-2-ylmethanone)-furoxan (**3**) (TGR IC_{50} of ~ 60 nM), were capable of NO generation in the presence of cysteine but not in the presence of TGR/NADPH. This ultimately proved significant as only oxadiazole-2-oxides that generated NO in the presence of TGR/NADPH were capable of *ex vivo* worm killing.²⁰ The association between NO release and worm killing was further evidenced by the partial mitigation of furoxan (**2**) induced parasite death in the presence of a NO scavenger molecule C-PTIO.^{20,43} Clearly, there is a relationship between NO release and the activity of this chemotype versus schistosomiasis. Here, we define the molecular pharmacophore of these agents, optimize compounds in terms of NO generation and potency versus TGR and worm killing efficacy, provide mechanistic evidence for TGR inhibition and parasite death, and discuss the selectivity and bioavailability of chosen analogues.

Results and Discussion

Prior to any optimization effort to improve this chemotype's ability to inhibit TGR and effectively act as an anti-schistosomal agent, we needed to establish a mechanistic link between NO donation and TGR inhibition. As such, our first considerations involved the synthetic deconstruction of the molecular structure of **2**. Various synthetic pathways for oxadiazole-2-oxides have been reviewed by Gasco and co-workers.^{22,23} Additionally, Maloney and co-workers have recently described the synthetic elaboration of the small molecule oxadiazole-2-oxides described in this report.⁴⁴ From the outset, an examination of the *N*-oxide moiety was of paramount importance. To better understand this key structural feature of the compound, we examined analogues that lack (derivative **4**) and regiochemically transpose (derivative **5**) the *N*-oxide (Figure 2). As opposed to furoxan (**2**), both **4** and **5** gave no appreciable NO release with TGR nor was either analogue active versus TGR or an effective *ex vivo* worm killing agent. Our next consideration was to determine the importance of the nitrile functionality. To explore the structure–activity relationships (SAR) of this moiety, we considered analogues




Figure 2. Structure–activity relationships of the *N*-oxide moiety.

Table 1. SAR of Nitrile Substitutions





Analogue	R'	TGR ^a		Ex Vivo Parasite Killing ^b						
		IC_{50} (μ M)	Max Resp. (% inh)	2 hr	4 hr	8 hr	16 hr	24 hr	48 hr	72 hr
	CN	6.3	-102	n.e.	curling	25%D	40%D	100%D	100%D	100%D
	CH ₃	> 50	-15	n.e.	n.e.	n.e.	n.e.	n.e.	sluggish	5%D
	CH ₂ OH	11.2	-100	n.e.	n.e.	n.e.	n.e.	sluggish	sluggish	10%D
	CHO	0.11	-105	n.e.	n.e.	n.e.	n.e.	sluggish	3%D	5%D
	COOH	0.63	-95	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.
	CONH ₂	17.8	-75	n.e.	n.e.	12%D	27%D	60%D	100%D	100%D

^a Data represent the results from three separate experiments. ^b Data collected by visual examination of worm movement and shape. n.e. (no effect): all worms are scored as active in culture with typical appearance; curling: worms are seen to curl rather than stretch and contract; sluggish: worm movement is significantly reduced; % D: % of worms dead as judged by lack of movement.

along the scale of oxidation states (i.e., alkyl to acyl). Several analogues were accessible intermediates of synthesis of **2** (including derivatives **7** and **8**), while others required a separate synthetic effort (derivatives **6**, **9**, and **10**).⁴⁴ The SAR of these analogues provided the first indication of the complexity surrounding the relationship between NO release, TGR inhibition, and worm killing (described through visual examination of cultured worms treated with 10μ M of compound over a 72 h time-course). The methyl analogue **6** was inactive as a NO donor and a TGR inhibitor and was essentially inactive in the worm killing evaluation (Table 1). The hydroxyl analogue **7** regained the ability to inhibit TGR (IC_{50} of 11.3μ M) but was only a weak NO donor and possessed minimal ability to kill *ex vivo* worms. The aldehyde **8** and carboxylic acid **9** were noted to be potent inhibitors of TGR (IC_{50} of 0.11 and 0.63μ M, respectively). Surprisingly, these analogues possessed only modest NO donating abilities and were ineffective worm killing agents. Interestingly, the aldehyde moiety in **8** would be capable of entering into electrophilic/nucleophilic reactive mechanisms with cysteine (and selenocysteine) residues in a similar fashion to the nitrile of **2**. The only analogue in this SAR series that retained both TGR inhibition and worm killing ability was the amide **10**, albeit at slightly lower potency and efficacy. With these results in place, it seems reasonable to consider the 3-cyano-1,2,5-oxadiazole-2-oxide core as the pharmacophore for these agents in terms of TGR inhibition, NO donating potential, and worm killing efficacy.

With a verified and consistent definition of the pharmacophore, we next turned to a systematic evaluation of the phenyl ring through the appraisal of standard substitutions (including methyl, methoxy, halo, trifluoromethyl, nitro, hydroxyl, and phenyl) and replacements (including heterocycles furan and thiophene). The results are detailed in Table 2 and show clear evidence that the majority of analogues preserving the core 3-cyano-1,2,5-oxadiazole-2-oxide moiety (**11–18**, **20–29**) retain TGR inhibition and worm killing capacity (**11–18**, **20–28** were additionally noted to be capable NO donating molecules). The SAR trends revealed that electron-withdrawing substitutions (halo, trifluoromethyl, and nitro) provided slight potency enhancements and the meta-position seemed favored (in general, substitutions at the ortho position were not synthetically viable (compound **21** was the lone exception)). Importantly, there was general agreement between TGR inhibition and worm killing efficacy. For instance, analogues **12–17** possessed TGR IC_{50} between 2 and 4μ M and were at least as efficacious as furoxan (**2**) in terms of worm killing. Conversely, several analogues with TGR IC_{50} values between 9 and 18μ M and were generally less active against worms in culture. From this phenyl ring scan, there were several intriguing lessons. The unique difference between

Table 2. SAR of Phenyl Ring Substitutions and Substitutions on the 3 and 4 Carbons of the Oxadiazole Ring

Analogue	R	TGR ^a IC ₅₀ (μM)	TGR ^a Max Resp. (% inh)	Ex Vivo Parasite Killing ^b							
				2 hr	4 hr	8 hr	16 hr	24 hr	48 hr	72 hr	
	2	NA	6.3	-102	n.e.	curling	25%D	40%D	100%D	100%D	100%D
	3	NA	0.04	-101	n.e.	n.e.	n.e.	n.c.	n.e.	n.e.	n.c.
	11	3-NO ₂	2.2	-107	n.e.	n.e.	n.e.	sluggish	10%D	15%D	21%D
	12	3-CF ₃	2.5	-102	n.e.	n.e.	7%D	25%D	54%D	100%D	100%D
	13	3-Br, 4-F	2.8	-102	n.e.	n.e.	curling	5%D	50%D	75%D	100%D
	14	3-Br	2.8	-100	n.e.	n.e.	20%D	39%D	100%D	100%D	100%D
	15	3-Cl	3.5	-101	n.e.	sluggish	15%D	20%D	64%D	80%D	100%D
	16	4-Br	3.5	-99	n.e.	sluggish	39%D	60%D	100%D	100%D	100%D
	17	4-Cl	4.0	-99	n.e.	n.e.	n.e.	sluggish	42%D	60%D	100%D
	18	4-CF ₃	7.1	-103	n.e.	n.e.	12%D	20%D	64%D	80%D	100%D
	19	3-OH	7.1	-96	n.e.	n.e.	ND	n.e.	n.e.	n.e.	n.e.
	20	4-F	7.9	-103	n.e.	n.e.	5%D	15%D	72%D	100%D	100%D
	21	2-OMe	7.9	-96	n.e.	n.e.	4%D	11%D	40%D	50%D	60%D
	22	3,4,5-OMe	8.9	-102	n.e.	n.e.	7%D	sluggish	40%D	70%D	75%D
	23	3-OMe	8.9	-102	n.e.	n.e.	n.e.	n.e.	7%D	12%D	20%D
	24	4-OMe	10.0	-97	n.e.	n.e.	n.e.	10%D	29%D	60%D	100%D
	25	4-Me	11.2	-98	n.e.	n.e.	n.e.	15%D	50%D	55%D	70%D
	26	4-Ph	15.8	-71	n.e.	n.e.	sluggish	sluggish	20%D	40%D	45%D
	27	4-OH	17.9	-64	n.e.	n.e.	sluggish	15%D	40%D	60%D	100%D
	28	O	2.8	-95	n.e.	n.e.	10%D	40%D	70%D	100%D	100%D
	29	S	3.5	-94	n.e.	n.e.	ND	25%D	25%D	100%D	100%D
	30	NA	0.063	-102	n.c.	n.c.	n.c.	n.c.	n.c.	n.c.	n.c.

^aData represent the results from three separate experiments. ^bData collected by visual examination of worm movement and shape. n.e. (no effect): all worms are scored as active in culture with typical appearance; curling: worms are seen to curl rather than stretch and contract; sluggish: worm movement is significantly reduced; % D: % of worms dead as judged by lack of movement. ND = not determined.

substitutions at the meta and para positions was further highlighted by the hydroxyl analogues **19** and **27**. Analogue **19** possessed good TGR inhibition but was inactive as a worm killing agent, while **27** was among the least potent analogues for TGR inhibition yet retained respectable activity versus worms. Replacement of the phenyl ring with heterocycles furan and thiophene (analogues **28** and **29**, respectively) resulted in derivatives with slightly improved potencies for TGR inhibition, good NO release profiles, and good worm killing efficacy. Another important exploration involved the substitution of the 4 position of the oxadiazole ring with a substituted carbonyl. Data from our initial studies on compounds with acyl substitutions on the oxadiazole ring system (such as in **3**) and analogues presented here (such as aldehyde **8**) demonstrate that such functionality confers significant potency enhancements in terms of TGR inhibition. These analogues, however, are devoid of the key 3-cyano-1,2,5-oxadiazole-2-oxide pharmacophore and were markedly less active versus cultured worms. We were, therefore, eager to examine an analogue that possessed both the nitrile moiety and a ketone moiety. As such, we synthesized a novel oxadiazole-2-oxide (analogue **30**) that satisfies these requirements (it should be noted that **30** required an entirely novel synthetic route spanning a total of nine transformations⁴⁴). The inhibition of TGR by **30** was in alignment with our previous results for **3** and **8** (potency of 63 nM). Unfortunately, like the aforementioned analogues, **30** was devoid of activity versus worms in culture.

The results of these studies provide solid evidence that inhibition of TGR by oxadiazole-2-oxides must be coupled with the ability of these small molecules to act as a NO donating compound in order to effect parasite death. A major question, however, concerns the role that molecular NO plays

in TGR inhibition and ultimately worm killing. That NO can modify protein structure and function by the act of S-nitrosylation was first shown in the early 1990s.^{45,46} Becker et al. subsequently showed that human GR can be inhibited by NO donating compounds and an active site cysteine residue is oxidized as a result.⁴⁷ Additionally, S-nitrosylation of cysteine residues on coxsackieviral protease 2A, 3C, and rhinovirus 3C is recognized to affect coxsackievirus and rhinovirus, respectively.^{48–50} Given the aforementioned precedence and the intriguing presence of 13 cysteine residues and one selenocysteine in TGR, we set out to examine the potential role of S-nitrosylation (or Se-nitrosylation) during TGR inhibition by **2**. To explore this possibility, we took advantage of the well-known biotin-switch assay introduced by Snyder and co-workers.^{51,52} A cartoon representation of the principal experiment is outlined in Figure 3A. Here, recombinant TGR was incubated with and without NADPH, furoxan (**2**), and glutathione (GSH). After incubation, the reactants were removed from TGR on a spin column and by acetone precipitation of TGR. All cysteine (or selenocysteine) residues that remain in their reduced sulfhydryl form are then capped by methyl methanethiosulfonate (MMTS). TGR was then treated with reduced ascorbate to return all S-nitrosylated cysteines (or Se-nitrosylated selenocysteines) to their reduced form, followed by reaction with the thiol-modifying reagent biotin-HPDP to allow visualization of labeled protein. After purification of the protein on a spin column and acetone precipitation, it was analyzed by native-Western blotting using a HRP-streptavidin conjugate to identify biotinylated proteins (Figure 3B). The results shown clearly indicate that recombinant TGR is S-nitrosylated by **2** and that the nitrosylation depends on the protein activity (i.e., requires NADPH) (Figure 3B). The experiment was performed with




Figure 3. (A) Cartoon representation of the Biotin-switch assay (see refs 49 and 50). (B) Detection of S-nitrosylation in recombinant *S. mansoni* TGR after reaction with furoxan (**2**) 20 μ M (lane 1) and 10 μ M (lane 2) \pm NADPH. The molecular weight of unmodified recombinant TGR is 64880 Da.

Table 3. SAR of Bis Oxadiazole-2-Oxides

Analogue	X	TGR ^a		Ex Vivo Parasite Killing ^b					
		IC ₅₀ (μ M)	Max Resp. (% inh)	2 hr	4 hr	8 hr	16 hr	24 hr	48 hr
2	NA	6.3	- 102	n.e.	curling	25%D	40%D	100%D	100%D
31	1,3-Phenyl	3.5	- 95	n.e.	curling	45%D	60%D	100%D	100%D
32	1,4-Phenyl	1.0	- 98	n.e.	n.e.	ND	55%D	82%D	100%D
33	5-Fluoro-1,3 Phenyl	0.48	- 95	n.e.	n.e.	ND	55%D	78%D	100%D
34	2,5-Thiophene	0.40	- 102	n.e.	8%D	ND	62%D	82%D	100%D
35	2,4-Thiophene	0.35	- 92	n.e.	15%D	ND	55%D	91%D	100%D

^aData represent the results from three separate experiments. ^bData collected by visual examination of worm movement and shape. n.e. (no effect): all worms are scored as active in culture with typical appearance; curling: worms are seen to curl rather than stretch and contract; sluggish: worm movement is significantly reduced; % D: % of worms dead as judged by lack of movement. ND = not determined.

pure, recombinant TGR and analyzed on a native gel; additional bands are likely the result of both monomeric and dimeric proteins and multiple S-nitrosylation events. No TGR labeling was observed when incubation was with GSH or in the absence of NADPH.

The results from this experiment strongly suggest that NO plays a significant role in TGR inhibition and likely in parasite killing. On the basis of this knowledge, our thoughts regarding the optimization of the oxadiazole-2-oxide chemotype focused on ways to optimize the NO donating qualities of individual compounds. Our previous attempts to optimize the compound through a standard phenyl ring scan (Table 2) provided evidence that the electron-withdrawing nature of the substituent at the 4-position of the oxadiazole ring was beneficial. While this makes mechanistic sense, even the most electron deficient phenyl rings provided only modest potency enhancements in terms of TGR inhibition and no discernible benefit in worm killing. Our next consideration was to explore analogues capable of delivering two equivalents of NO. To this end, we synthesized and evaluated several bis-oxadiazole-2-oxide derivatives linked through substituted phenyl rings and heterocycles. The results for these agents are shown in Table 3 and demonstrate the validity of the theory. All bis-oxadiazole-2-oxide analogues were capable of TGR inhibition at potencies exceeding furoxan (**2**) and worm killing

profiles also demonstrated similarity to furoxan (**2**) and in some cases modest improvements (analogues **33**, **34**, and **35** were examined in a separate experiment). Worm death will not occur until the GSH/GSSG ratio reaches a critical point and the overall worm redox balance is unrecoverable, making it difficult to derive correlations between worm killing and TGR inhibition below a certain level.

With a growing understanding of this chemotype's potential and the advancements of several analogues with improved properties, it was important to consider the mechanism of NO donation from oxadiazole-2-oxides. A mechanistic rationale for NO donation that is initiated through nucleophilic attack (presumably by the sulfhydryl moiety of a cysteine residue or by selenocysteine) at either the 3 or 4 position of the oxadiazole ring and subsequent rearrangement of the heterocycle in a manner that allows release of the nitroxyl anion (NO^-) has been presented.^{22,23} An enzymatic oxidation is posited to transform this agent to NO (NO^*). It is important to note that the nitroxyl anion (NO^-) exists as a triplet that reacts very rapidly with triplet dioxygen to form peroxynitrite ion. Thus, it is unlikely that the transformation from NO^- to NO^* occurs via oxidation by molecular oxygen. Additionally, the previously reported mechanism²⁷ suggests a permanent ligation between the attacking nucleophile and the rearranged carbon scaffold of furoxan (**2**). While this mechanism is plausible,

there are several aspects that are not consistent with the data from our study. Foremost, this mechanism does not account for the release of NO (rather than NO^-) during a reaction with cysteine (i.e., in an environment lacking an enzyme capable of oxidizing NO^- to NO^*). A second issue is the stoichiometry of the processes. Our previous data suggest that TGR acts catalytically on furoxan (**2**) in terms of NO release. A 10 μM concentration of **2** in the presence of 15 nM of TGR and NADPH resulted in the generation of ~ 15 mol % of NO relative to **2**; i.e., ~ 1500 nM NO from only 15 nM of TGR or 100 equiv of NO per enzyme equivalent. Obviously, TGR has the ability to attack furoxan (**2**), effect molecular rearrangement and release of NO, and subsequently release from the heterocycle scaffold in such a manner that it retains the ability to re-enter the cycle with another molecule of furoxan (**2**). Presumably, this cycle continues until a critical concentration of NO is present and the enzyme is sufficiently S-nitrosylated as to lose activity. Given these inconsistencies, we sought evidence of alternate mechanisms.

Recently, Chakrapani et al. presented experimental evidence of NO donation from C-nitroso compounds through a first-order homolytic C–N bond scission event.⁵³ Interestingly, this mechanism required a geminal nitrile moiety to allow homolysis of the C–N bond. Moreover, the reactivity between nitrile groups and free thiols is well established and explored.^{54,55} To further explore the mechanistic release of NO, we examined selected analogues ability to donate NO as judged by the aforementioned ABTS method (see the Supporting Information section for these data). The results




Figure 4. Evaluation of furoxan (**2**) and selected analogues for reactivity with free cysteine as judged by consumption of reagent and the molar ratio of NO release per stoichiometric equivalent of reagent.

confirmed that analogues must be capable of generating NO in the presence of TGR and NADPH to be effective worm killing agents. The ABTS method provides a measure of the oxidation products of NO rather than NO itself. To more exactly quantify the release of NO, we examined the release of NO quantitatively through a well-known chemiluminescent assay that relies upon the reaction between NO and ozone to produce an activated NO_2 species that produces an intense chemiluminescence emission.⁵⁴ Additionally, selected analogues were characterized for their ability to react with free cysteine (as judged by consumption of starting material detected across an HPLC gradient). For these experiments, we selected a number of compounds with specific SAR in terms of TGR inhibition and worm killing ability (Figure 4). Furoxan (**2**) was noted to react fully with cysteine and produce approximately 0.6 equivalents of NO per equivalent of **2**. Gratifyingly, bis-oxadiazole-2-oxide analogues **33** and **34** were both cysteine reactive and produced more NO equivalents relative to furoxan (**2**) (1.55 mol equiv for **33** and 1.19 mol equiv for **34**). Analogue **6**, with the replacement of the nitrile with a methyl group, was not capable of reacting with cysteine nor was **6** capable of NO donation. Most interestingly, derivatives **3** (with the bis-thiophene ketone moiety) and **5** (with a transposed N–O moiety, but retaining the nitrile) did react with cysteine but did not generate NO. An additionally interesting result was that compound **30** (with both thiophene ketone and nitrile moieties) reacted with cysteine and was capable of releasing NO. From this result, the failure of **30** to produce worm death remains unexplained.

Mechanistically, it is not possible to identify the location of the initial nucleophilic attack, but the ultimate result is addition into the oxadiazole ring system alpha to the nitrile (Figure 5). The ring system would likely open resulting in a quaternary alkyl-thio-nitroso-nitrile species that is highly similar to the C-nitroso compounds shown by Chakrapani et al. to be capable of homolytic C–N bond cleavage and release of NO (path 1). Alternatively, this intermediate could undergo transnitrosylation and release of R-SNO and a byproduct of the original oxadiazole-2-oxide heterocycle (path 2). It is further possible that this intermediate could rearrange to form an isoxazol-5(4*H*)-imine type intermediate prior to homolytic C–N bond cleavage and release of NO (path 3). A second mechanistic issue is the ultimate release of the attacking nucleophile from the core oxadiazole scaffold.




Figure 5. Schematic description of potential mechanisms for NO release by oxadiazole-2-oxides via reaction with reactive thiols (or with selenol).

Table 4. Comparative Inhibition of Selected Analogues vs TGR, Human GR, and Rat TrxR

analogue	TGR ^a , IC ₅₀ (μM)	hGR ^b , IC ₅₀ (μM)	rTrxR ^b , IC ₅₀ (μM)
2	6.3	> 50	> 400
6	> 50	> 50	> 400
7	11.2	> 50	> 400
8	0.11	35.2	< 12.5
9	0.63	> 50	> 400
10	17.8	> 50	> 400
11	2.2	> 50	ca. 150
12	2.5	> 50	ca. 250
13	2.8	> 50	ca. 250
14	2.8	> 50	ca. 150
15	3.5	35.7	< 12.5
16	3.5	> 50	ca. 250
17	4.0	> 50	> 400
18	7.1	45.3	< 12.5
20	7.9	> 50	ca. 300
22	8.9	> 50	ca. 150
23	8.9	> 50	> 400
24	10.0	> 50	> 400
25	11.2	> 50	> 400
27	17.9	> 50	> 400
28	2.8	> 50	ca. 350
31	3.5	> 50	ca. 150

^aData represent the results from three separate experiments. ^bMethods for determining inhibitory potential versus hGR and rTrxR are presented in the Supporting Information.

While plausible mechanisms can be envisioned for this deligation event, there is currently no experimental evidence for how it would proceed. Interestingly, we tested this point through the synthesis of a biotinylated version of furoxan (**2**) (conjugated through an ether linkage at the 4-position of the phenyl ring) and subsequent treatment of this analogue with recombinant TGR and NADPH. Gel analysis of the resulting mixture provided evidence that TGR was not cross-linked to biotin, further suggesting that TGR reacts with oxadiazole-2-oxides in a catalytic rather than terminally reactive mechanism (data not shown).

These insights into the interaction of oxadiazole-2-oxides with TGR suggest that the preeminent mechanistic event is the nucleophilic attack by either cysteine or selenocysteine residues on TGR. Selenium has a unique role in biology and the biosynthesis of selenocysteine residues and, ultimately, selenoproteins have been the subject of numerous reviews.⁵⁶ To date, 25 human proteins have been characterized that contain selenocysteine residues and each retain a specified cellular function (over 300 selenoprotein families have been found in nature).⁵⁷ A minimalist view of atomic sulfur and selenium highlight several divergent properties that translate into important biochemical differences for selenocysteine and cysteine residues. Primary among these variations are the relative pK_a values for both amino acids (8.5 for cysteine versus 5.2 for selenocysteine) and the stronger nucleophilicity of selenocysteine.^{58–60} Evolutionary analyses clearly suggest that selenocysteine supports unique roles in selenoproteins that can not easily be maintained by cysteine.⁶¹ The strikingly low pK_a of selenocysteine results in full anionic character for this residue at physiological pH. In several instances, the divergent activities of selenocysteine and cysteine residues have been exploited. For instance, Filipovska and co-workers recently reported the rational design of gold(I) containing *N*-heterocyclic carbene complexes that target the selenocysteine-containing TrxR and are selectively toxic to cancer cells.⁶² Arnér

and co-workers have found that forms of selenium compromised thioredoxin reductase-derived apoptotic proteins (SecTRAPs) may have prooxidant toxic properties in addition to a loss of thioredoxin reducing capacity.⁶³

While there is not yet any direct evidence, it seems reasonable to assume that the enhanced nucleophilicity associated with the selenocysteine residues (including the one found in TGR) makes this moiety a likely candidate for the initial attack on furoxan (**2**) and other oxadiazole-2-oxides. As such, it was important to explore the selectivity of this chemotype with regards to its ability to act as a general electrophile for other selenocysteine containing enzymes. Additionally, it was essential to consider if these compounds inhibit related host enzymes, most significantly GR and TrxR. To resolve both questions, we assayed several analogues versus human GR and rat TrxR (rat TrxR is a selenoprotein⁶⁴ while human GR is not). We found that furoxan (**2**) does not inhibit recombinant human GR (IC₅₀ > 50 μM) or recombinant rat TrxR (IC₅₀ > 400 μM) (Table 4). Of interest was the discovery that three of the agents tested (the aldehyde **8** and phenyl ring analogues **15** and **18**) were found to inhibit both hGR and rat TrxR, albeit with reduced affinity. The aldehyde derivative **8** may yield insight into the mechanism of these agents, and the discovery of inhibitors of both hGR and rat TrxR is an important finding on its own accord. More important for this study, however, is the fact that several of the antischistosomal analogues were found not to be inhibitors of the TGR-related mammalian host enzymes. Taken as a whole, the data presented here provide strong evidence supporting the selective action by oxadiazole-2-oxides against schistosomes and suggests that the interaction of this chemotype with TGR is dependent upon both binding affinity and appropriately aligned reactivities.

We were further interested in determining if the oxadiazole-2-oxide scaffold has suitable properties for oral bioavailability. Because therapies for schistosomiasis are intended to be single dose agents (as opposed to chronic treatments), we were initially and primarily interested in determining selected analogues' potential for passive membrane permeation and stability in liver microsomes. Profiles of these agents binding to the human ether-à-go-go related gene (hERG) potassium channel and inhibition of selected members of the cytochrome P450 class of enzymes were also of interest. Our own investigations of these agents suggested highly variable solubilities in water. We were interested in going beyond conjectural evaluations of this property (i.e., HPLC retention times) and establish more reliable data for selected analogues. Consequently, solubility measurements (in both μM and μg/mL) and LogD determinations were ascertained. The results of these profiles are shown in Table 5. Several compounds had only modest water solubility including furoxan (**2**), **16**, **31**, **32**, and **34**. Analogues containing the thiophene ring were noted as having better solubility profiles, including **29** and **35**. Interestingly, the addition of a single fluorine moiety to the bis-oxadiazole-2-oxide analogue **31** resulted in an impressive gain in solubility (see derivative **33**).

Beyond solubility profiling, several analogues were found to have modest ability to permeate artificial Caco-2 cell monolayers (apical to basolateral (A → B) permeability was typically greater than basolateral to apical (B → A)). Furoxan (**2**) was not found to pass through Caco-2 cells, however, several analogues (including **16**, **29**, **33**, and **35**) were noted to possess adequate Caco-2 profiles. While these data do not guarantee successful realization of these analogues as orally

Table 5. Solubility and Selected AD ME Properties for Chosen Analogues

analogue	solubility ^a (μ M)	solubility ^a (μ g/mL)	LogD ^b	Caco-2 permeability ^c mean A \rightarrow B	Caco-2 permeability ^c mean B \rightarrow A	microsomal stability ^d	hERG FastPatch ^e
2	30.1	5.6	2.42	1.0	0	8/63	13.2%
16	18.6	5.0	3.1	7.9	4.8	10/67	5.5%
29	240.4	46.4	2.54	2.54	9.0	1/64	3.4%
31	5.0	1.5	ND*	ND*	ND*	ND*	7.8%
32	3.7	1.1	ND*	ND	ND	ND	ND
33	89.1	28.0	2.91	7.9	2.7	20/51	29.4%
34	11.5	3.5	2.92	4.0	1.8	3/56	83.9%
35	122.2	36.9	2.92	9.4	3.4	12/52	57.2%

^a Kinetic solubility analysis was performed by Analiza Inc. and are based upon quantitative nitrogen detection as described (www.analiza.com). The data represent results from three separate experiments with an average intraassay %CV of 4.5%. ^b LogD analysis was performed by Analiza Inc. and are based upon octanol/buffer partitioning and quantitative nitrogen detection of sample content as described (www.analiza.com). The data represent results from three separate experiments and is corrected to calibration standards and adjusted to account for fixed amounts of DMSO. ^c Caco-2 permeability analysis was performed by Apredica Inc. and are based upon monolayer Caco-2 cells applied to collagen-coated BioCoat Cell Environment at 24500 cells per well as described (www.apredica.com). Data are expressed in P_{app} (apparent permeability) in 10^{-6} cm s⁻¹. Atenolol was used as a low permeability control (mean A \rightarrow B = 1.2 and mean B \rightarrow A = 1.2) and propranolol was used as a high permeability control (mean A \rightarrow B = 20.3 and mean B \rightarrow A = 17.4). ^d Microsomal stability analysis was performed by Apredica Inc. and are based upon duplicate incubations of test reagent with liver microsomes (human and rat; only human data is shown) over 60 min at 37 °C as described (www.apredica.com). LC/MS/MS is utilized to quantitate remaining test reagent and the data is reported as % remaining. Analysis in the absence and presence of NADPH was performed to assess NADPH free degradation. Data is listed as NADPH(+) degradation/NADPH(-) degradation. ^e hERG FastPatch analysis was performed by Apredica Inc. using an automated patch clamp electrophysiological procedure using HEK293 cells stably transfected with hERG cDNA as described (www.apredica.com). Test reagents were applied at standard concentrations (10 μ M) and data is recorded as mean % inhibition from three separate experiments.

available agents, it does dispel concerns that the oxadiazole-2-oxide scaffolds are incapable of passive membrane permeation. Analysis of these analogues in liver microsomes (over 1 h in the presence/absence of NADPH) suggests that these compounds are only moderately stable and may possess short half-lives in intact animal models. However, the variable data (for instance compound **29** versus **33**) suggests that more robust analogues may be found through additional modification. Further, the hERG and Cyp profiles for these agents were generally reasonable. Taken as a whole, these data suggests that properly substituted oxadiazole-2-oxides may possess efficacy and ADMET properties adequate for clinical application.

Conclusion

Given the widespread morbidity and mortality associated with schistosomiasis and the precarious reliance on PZQ as the exclusive antischistosomal agent, development of new treatments for the disease is critical. Here, we present data from our continued evaluation of the oxadiazole-2-oxide class of reagents as inhibitors of the multifunctional selenocysteine-containing flavoenzyme TGR. Through chemical modification of the lead agent (furoxan (**2**)), we have defined the pharmacophore as the 3-cyano-1,2,5-oxadiazole-2-oxide moiety and established SARs surrounding the phenyl ring of the core structure. Through use of the biotin-switch assay, we have shown that incubation of TGR with furoxan (**2**) leads to S-nitrosylation (or Se-nitrosylation) of TGR. These data allowed us to focus our attention on analogues capable of dual release of NO, resulting in agents with superior activity versus ex vivo parasites. We have further defined the selectivity of these agents as compared to other reductases including human GR and rat TrxR (also a selenoprotein). Finally, a preliminary evaluation of selected in vitro pharmacokinetic properties, including water solubility, Caco-2 cell permeability, and microsomal stability, suggest that the oxadiazole-2-oxide chemotype can possess the properties required for an orally bioavailable clinical reagent. We are currently developing models to examine the oral efficacy of selected agents.

Acknowledgment. We thank Allison Peck for critical reading of this manuscript. This research was supported by the Molecular Libraries Initiative of the National Institutes of Health Roadmap for Medical Research, the Intramural Research Program of the National Human Genome Research Institute, National Cancer Institute, and Center for Cancer Research. Schistosome life stages used in this research were supplied in part by the National Institute of Allergy and Infectious Diseases Schistosomiasis Resource Center at the Biomedical Research Institute (Rockville, Maryland) through NIAID contract N01-AI-30026. This work was partially supported by grant AI065622.

Supporting Information Available: An extended table of TGR inhibition and NO release profiles for selected analogues and experimental methods for TGR biochemical assay protocol, ex vivo worm assay protocol, GR assay protocol, biotin-switch assay protocol, TrxR1 assay protocol (DNTB), NO release protocol, reactivity of oxadiazole-2-oxides with Cys (HPLC Assay), and characterization of compounds are detailed. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Hotez, P. J.; Brindley, P. J.; Bethony, J. M.; King, C. H.; Pearce, E. J.; Jacobson, J. Helminth infections: the great neglected tropical diseases. *J. Clin. Invest.* **2008**, *118*, 1311–1321.
- (2) Steinmann, P.; Keiser, J.; Bos, R.; Tanner, M.; Utzinger, J. Schistosomiasis and water resources development: systematic review, meta-analysis, and estimates of people at risk. *Lancet Infect. Dis.* **2006**, *6*, 411–425.
- (3) van der Werf, M. J.; de Vlas, S. J.; Brooker, S.; Looman, C. W.; Nagelkerke, N. J.; Habbema, J. D.; Engels, D. Quantification of clinical morbidity associated with schistosome infection in sub-Saharan Africa. *Acta Trop.* **2003**, *86*, 125–139.
- (4) King, C. H.; Dangerfield-Cha, M. The unacknowledged impact of chronic schistosomiasis. *Chronic Illness* **2008**, *4*, 65–79.
- (5) Gonnert, R.; Andrews, P. Praziquantel, a new broad spectrum antischistosomal agent. *Z. Parasitenk.* **1977**, *52*, 129–150.
- (6) Hagan, P.; Appleton, C. C.; Coles, G. C.; Kusel, J. R.; Tchuente, L. A. Schistosomiasis control: keep taking the tablets. *Trends Parasitol.* **2004**, *20*, 92–97.
- (7) Fenwick, A.; Webster, J. P. Schistosomiasis: challenges for control, treatment and drug resistance. *Curr. Opin. Infect. Dis.* **2006**, *19*, 577–582.
- (8) Greenberg, R. M. Are Ca²⁺ channels targets of Praziquantel action? *Int. J. Parasitol.* **2004**, *35*, 1–9.

- (9) Nogi, T.; Zhang, D.; Chan, J. D.; Marchant, J. S. A Novel Biological Activity of Praziquantel Requiring Voltage-Operated Ca^{2+} Channel β Subunits: Subversion of Flatworm Regenerative Polarity. *PLoS Neglected Trop. Dis.* **2009**, *3*, e464.
- (10) Doenhoff, M. J.; Cioli, D.; Utzinger, J. Praziquantel: mechanisms of action, resistance and new derivatives for schistosomiasis. *Curr. Opin. Infect. Dis.* **2008**, *21*, 659–667.
- (11) Townsend, D. M.; Tew, K. D.; Tapiero, H. The importance of glutathione in human disease. *Biomed. Pharmacother.* **2003**, *57*, 145–155.
- (12) Lillig, C. H.; Holmgren, A. Thioredoxin and related molecules – from biology to health and disease. *Antioxid. Redox Signaling* **2007**, *9*, 25–47.
- (13) Alger, H. M.; Sayed, A. A.; Stadecker, M. J.; Williams, D. L. Molecular and enzymatic characterisation of *Schistosoma mansoni* thioredoxin. *Int. J. Parasitol.* **2002**, *32*, 1285–1292.
- (14) Angelucci, F.; Miele, A. E.; Boumis, G.; Dimastrogiovanni, D.; Brunori, M.; Bellelli, A. Glutathione reductase and thioredoxin reductase at the crossroad: the structure of *Schistosoma mansoni* thioredoxin glutathione reductase. *Proteins* **2008**, *72*, 936–945.
- (15) Kuntz, A. N.; Davioud-Charvet, E.; Sayed, A. A.; Califf, L. L.; Dessolin, J.; Arnér, E. S.; Williams, D. L. Thioredoxin glutathione reductase from *Schistosoma mansoni*: an essential parasite enzyme and a key drug target. *PLoS Med.* **2007**, *4*, e206.
- (16) Simeonov, A.; Jadhav, A.; Sayed, A. A.; Wang, Y.; Nelson, M. E.; Thomas, C. J.; Inglese, J.; Williams, D. L.; Austin, C. P. Quantitative high-throughput screen identifies inhibitors of the *Schistosoma mansoni* redox cascade. *PLoS Neglected Trop. Dis.* **2008**, *2*, e127.
- (17) Lea, W. A.; Jadhav, A.; Rai, G.; Sayed, A. A.; Cass, C. L.; Inglese, J.; Williams, D. L.; Austin, C. P.; Simeonov, A. A 1536-well-based kinetic HTS assay for inhibitors of *Schistosoma mansoni* thioredoxin glutathione reductase. *Assay Drug Dev. Technol.* **2008**, *6*, 551–555.
- (18) Austin, C. P.; Brady, L. S.; Insel, T. R.; Collins, F. S. NIH Molecular Libraries Initiative. *Science* **2004**, *306*, 1138.
- (19) Inglese, J.; Auld, D. S.; Jadhav, A.; Johnson, R. L.; Simeonov, A.; Yasgar, A.; Zheng, W.; Austin, C. P. Quantitative high-throughput screening: a titration-based approach that efficiently identifies biological activities in large chemical libraries. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 11473–11478.
- (20) Sayed, A. A.; Simeonov, A.; Thomas, C. J.; Inglese, J.; Austin, C. P.; Williams, D. L. Identification of oxadiazoles as new drug leads for the control of schistosomiasis. *Nature Med.* **2008**, *4*, 407–412.
- (21) Nwaka, S.; Hudson, A. Innovative lead discovery strategies for tropical diseases. *Nature Rev. Drug Discovery* **2006**, *5*, 941–955.
- (22) Gasco, A.; Fruttero, R.; Sorba, G.; Di Stilo, A.; Calvino, R. NO donors: focus on furoxan derivatives. *Pure Appl. Chem.* **2004**, *76*, 973–981.
- (23) Gasco, A.; Schoenafinger, K. The NO-releasing heterocycles. In *Nitric Oxide Nonors*; Wang, G., Cai, T. B., Taniguchi, N., Eds.; Wiley-VCH: Weinheim, Germany, 2005.
- (24) Lundberg, J. O.; Weitzberg, E.; Gladwin, M. T. The nitrate–nitrite–nitric oxide pathway in physiology and therapeutics. *Nature Rev. Drug Discovery* **2008**, *7*, 156–167.
- (25) Paolucci, N.; Jackson, M. I.; Lopez, B. E.; Miranda, K.; Tocchetti, C. G.; Wink, D. A.; Hobbs, A. J.; Fukuto, J. M. The pharmacology of nitroxyl (HNO) and its therapeutic potential: not just the janus face of NO. *Pharm. Ther.* **2007**, *113*, 442–458.
- (26) Ignarro, L. J. *Nitric Oxide Biology and Pathobiology*; Academic Press: San Diego, 2000.
- (27) Medana, C.; Ermondi, G.; Fruttero, R.; Di Stilo, A.; Ferretti, C.; Gasco, A. Furoxans as nitric oxide donors. 4-Phenyl-3-furoxan-carbonitrile: thiol-mediated nitric oxide release and biological evaluation. *J. Med. Chem.* **1994**, *37*, 4412–4416.
- (28) Buonsanti, M. F.; Bertinaria, M.; Di Stilo, A.; Cena, C.; Fruttero, R.; Gasco, A. Nitric oxide donor β_2 -agonists: furoxan derivatives containing the fenoterol moiety and related furazans. *J. Med. Chem.* **2007**, *50*, 5003–5011.
- (29) Turnbull, C. M.; Cena, C.; Fruttero, R.; Gasco, A.; Rossi, A. G.; Megson, I. L. Mechanism of action of novel NO-releasing furoxan derivatives of aspirin in human platelets. *Br. J. Pharmacol.* **2006**, *148*, 517–526.
- (30) Sorba, G.; Galli, U.; Cena, C.; Fruttero, R.; Gasco, A.; Morini, G.; Adami, M.; Coruzzi, G.; Brenciaglia, M. I.; Dubini, F. A new furoxan NO-donor rabeprazole derivative and related compounds. *ChemBioChem* **2003**, *4*, 899–903.
- (31) Brunet, L. R. Nitric oxide in parasitic infections. *Int. Immunopharmacol.* **2001**, *1*, 1457–1467.
- (32) Colasanti, M.; Gradoni, L.; Mattu, M.; Persichini, T.; Salvati, L.; Venturini, G.; Ascenzi, P. Molecular basis for the anti-parasitic effect on NO. *Int. J. Mol. Med.* **2002**, *9*, 131–134.
- (33) Rivero, A. Nitric oxide: an antiparasitic molecule of invertebrates. *Trends Parasitol.* **2006**, *22*, 219–225.
- (34) James, S. L.; Glaven, J. Macrophage cytotoxicity against schistosomula of *Schistosoma mansoni* involves arginine-dependent production of reactive nitrogen intermediates. *J. Immunol.* **1989**, *143*, 4208–4212.
- (35) Amri, M.; Aissa, S. A.; Belguendouz, H.; Mezioug, D.; Touil-Boukoffa, C. In vitro antihydatic action of IFN- γ is dependent on the nitric oxide pathway. *J. Interferon Cytokine Res.* **2007**, *27*, 781–787.
- (36) Magez, S.; Radwanska, M.; Drennan, M.; Fick, L.; Baral, T. N.; Allie, N.; Jacobs, M.; Nedospasov, S.; Brombacher, F.; Ryffel, B.; De Baetselier, P. Tumor Necrosis Factor (TNF) Receptor-1 (TNFRp55) Signal Transduction and Macrophage-Derived Soluble TNF are Crucial for Nitric Oxide-Mediated *Trypanosoma congolense* Parasite Killing. *J. Infect. Dis.* **2007**, *196*, 954–962.
- (37) Winberg, M. E.; Rasmusson, B.; Sundqvist, T. *Leishmania donovani*: inhibition of phagosomal maturation is rescued by nitric oxide in macrophages. *Exp. Parasitol.* **2007**, *117*, 165–170.
- (38) López-Jaramillo, P.; Ruano, C.; Rivera, J.; Terán, E.; Salazar-Irigoyen, R.; Esplugues, J. V.; Moncada, S. Treatment of cutaneous leishmaniasis with nitric oxide donor. *Lancet* **1998**, *351*, 1176–1177.
- (39) Singh, R.; Manjunatha, U.; Boshoff, H. I. M.; Ha, Y. W.; Niyomrattanakit, P.; Ledwidge, R.; Dowd, C. S.; Lee, I. Y.; Kim, P.; Zhang, L.; Kang, S.; Keller, T. H.; Jiricek, J.; Barry, C. E. PA-824 Kills Nonreplicating Mycobacterium tuberculosis by Intracellular NO Release. *Science* **2008**, *322*, 1392–1395.
- (40) Mady, C.; Ianni, B. M.; de Souza, J. L., Jr. Benzimidazole and Chagas disease: can an old drug be the answer to an old problem? *Expert Opin. Investig. Drugs* **2008**, *17*, 1427–1433.
- (41) Wilkinson, S. R.; Taylor, M. C.; Horn, D.; Kelly, J. M.; Cheeseman, I. A mechanism for cross-resistance to nifurtimox and benzimidazole in trypanosomes. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 5022–5027.
- (42) Nims, R. W.; Darbyshire, J. F.; Saavedra, J. E.; Christodoulou, D.; Hanbauer, I.; Cox, G. W.; Grisham, M. B.; Laval, F.; Cook, J. A.; Krishna, M. C.; Wink, D. A. Colorimetric methods for the determination of nitric oxide concentration in neutral aqueous solutions. *Methods* **1995**, *7*, 48–55.
- (43) Akaike, T.; Yoshida, M.; Miyamoto, Y.; Sato, K.; Kohno, M.; Sasamoto, K.; Miyazaki, K.; Ueda, S.; Maeda, H. Antagonistic action of imidazolineoxyl N-Oxides against endothelium-derived relaxing factor/NO through a radical reaction. *Biochemistry* **1993**, *32*, 827–832.
- (44) Rai, G.; Thomas, C. J.; Leister, W.; Maloney, D. J. Synthesis of oxadiazole-2-oxide analogues as potential antischistosomal agents. *Tetrahedron Lett.* **2009**, *50*, 1710–1713.
- (45) Lei, S. Z.; Pan, Z. H.; Aggarwal, S. K.; Chen, H. S.; Hartman, J.; Sucher, N. J.; Lipton, S. A. Effect of nitric oxide production on the redox modulatory site of the NMDA receptor-channel complex. *Neuron* **1992**, *8*, 1087–1099.
- (46) Stamler, J. S.; Simon, D. I.; Osborne, J. A.; Mullins, M. E.; Jaraki, O.; Michel, T.; Singel, D. J.; Loscalzo, J. S-Nitrosylation of proteins with nitric oxide: synthesis and characterization of biologically active compounds. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, *89*, 444–448.
- (47) Becker, K.; Savvides, S. N.; Keese, M.; Schirmer, R. H.; Karplus, P. A. Enzyme inactivation through sulfhydryl oxidation by physiologic NO-carriers. *Nat. Struct. Biol.* **1998**, *5*, 267–271.
- (48) Badorf, C.; Fichtlscherer, B.; Rhoads, R. E.; Zeiher, A. M.; Muelsch, A.; Dimmeler, S.; Knowlton, K. U. Nitric oxide inhibits dystrophin proteolysis by coxsackieviral protease 2A through S-nitrosylation. *Circulation* **2000**, *102*, 2276–2281.
- (49) Saura, M.; Zaragoza, C.; McMillian, A.; Quick, R. A.; Hohenadl, C.; Lowenstein, J. M.; Lowenstein, C. J. An antiviral mechanism of nitric oxide: Inhibition of a viral protease. *Immunity* **1999**, *10*, 21–28.
- (50) Xian, M.; Wang, Q. M.; Chen, X.; Wang, K.; Wang, P. G. S-Nitrosothiols as novel, reversible inhibitors of human rhinovirus 3C protease. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2097–2100.
- (51) Jaffrey, S. R.; Snyder, S. H. The biotin switch method for the detection of S-nitrosylated proteins. *Sci. STKE* **2001**, *86*, p11.
- (52) Jaffrey, S. R.; Fang, M.; Snyder, S. H. Nitrosopeptide Mapping: A Novel Methodology Reverses S-nitrosylation of Dexas1 on a Single Cysteine Residue. *Chem. Biol.* **2002**, *9*, 1329–1335.
- (53) Chakrapani, H.; Bartberger, M. D.; Toone, E. J. C-Nitroso Donors of Nitric Oxide. *J. Org. Chem.* **2009**, *74*, 1450–1453.
- (54) Oballa, R. M.; Truchon, J.-F.; Bayly, C. I.; Chauret, N.; Day, S.; Crane, S.; Berthelette, C. A generally applicable method for assessing the electrophilicity and reactivity of diverse nitrile-containing compounds. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 998–1002.

- (55) Fontijn, A.; Sabadell, A. J.; Ronco, R. J. Homogeneous Chemiluminescent Measurement of Nitric Oxide with Ozone. *Anal. Chem.* **1970**, *42*, 575–579.
- (56) Hatfield, D. L.; Carlson, B. A.; Xu, X. M.; Mix, H.; Gladyshev, V. N. Selenocysteine incorporation machinery and the role of selenoproteins in development and health. *Prog. Nucleic Acid Res. Mol. Biol.* **2006**, *81*, 97–142.
- (57) Kryukov, G. V.; Gladyshev, V. N. The prokaryotic selenoproteome. *EMBO Rep.* **2004**, *5*, 538–543.
- (58) Benesch, R. E.; Benesch, R. The acid strength of the SH group in cysteine and related compounds. *J. Am. Chem. Soc.* **1955**, *77*, 5877–5881.
- (59) Huber, R. E.; Criddle, R. S. Comparison of the chemical properties of selenocysteine and selenocystine with their sulfur analogs. *Arch. Biochem. Biophys.* **1967**, *122*, 164–173.
- (60) Johansson, L.; Gafvelin, G.; Arnér, E. S. J. Selenocysteine in proteins—properties and biotechnological use. *Biochem. Biophys. Acta.* **2005**, *1726*, 1–13.
- (61) Castellano, S.; Andrés, A. M.; Bosch, E.; Bayes, M.; Guigó, R.; Clark, A. G. Low Exchangeability of Selenocysteine, the 21st Amino Acid, in Vertebrate Proteins. *Mol. Biol. Evol.* **2009**, *26*, 2031–2040.
- (62) Hickey, J. L.; Ruhayel, R. A.; Barnard, P. J.; Baker, M. V.; Berners-Price, S. J.; Filipovska, A. Mitochondria-Targeted Chemotherapeutics: The Rational Design of Gold(I) *N*-Heterocyclic Carbene Complexes that are Selectively Toxic to Cancer Cells and Target Protein Selenols in Preference to Thiols. *J. Am. Chem. Soc.* **2008**, *130*, 12570–12571.
- (63) Anestál, K.; Prast-Nielsen, S.; Cenas, N.; Arnér, E. S. J. Cell Death by SecTRAPs: Thioredoxin Reductase as a Prooxidant Killer of Cells. *PLoS One* **2008**, *3*, e1846.
- (64) Zhong, L.; Arnér, E. S. J.; Ljung, J.; Aslund, F.; Holmgren, A. Rat and calf thioredoxin reductase are homologous to glutathione reductase with a carboxyl-terminal elongation containing a conserved catalytically active penultimate selenocysteine residue. *J. Biol. Chem.* **1998**, *273*, 8581–8591.