

## Discovery of Trp-His and His-Arg Analogues as New Structural Classes of Short Antimicrobial Peptides

Rohit K. Sharma,<sup>†</sup> Ravi P. Reddy,<sup>†</sup> Werner Tegge,<sup>‡</sup> and Rahul Jain<sup>\*†</sup>

<sup>†</sup>Department of Medicinal Chemistry, National Institute of Pharmaceutical Education and Research, Sector 67, SAS Nagar 160 062, Punjab, India, and <sup>‡</sup>Department of Chemical Biology, Helmholtz Centre for Infection Research, Helmoltzstrasse 7, D-38124 Braunschweig, Germany

Received May 4, 2009

Naturally occurring antimicrobial peptides contain a large number of amino acid residues, which limits their clinical applicability. In search of short antimicrobial peptides, which represent a possible alternative for lead structures to fight antibiotic resistant microbial infections, a series of synthetic peptide analogues based on Trp-His and His-Arg structural frameworks have been prepared and found to be active against several Gram-negative and Gram-positive bacterial strains as well as against a fungal strain with MIC values of the most potent structures in the range of 5–20  $\mu\text{g}/\text{mL}$  ( $\text{IC}_{50}$  in the range of 1–5  $\mu\text{g}/\text{mL}$ ). The synthesized peptides showed no cytotoxic effect in an MTT assay up to the highest test concentration of 200  $\mu\text{g}/\text{mL}$ . A combination of small size, presence of unnatural amino acids, high antimicrobial activity, and absence of cytotoxicity reveals the synthesized Trp-His and His-Arg analogues as promising candidates for novel antimicrobial therapeutics.

### Introduction

Microbial infections continue to be a major cause of morbidity and mortality worldwide. Moreover, the problems have been continuously exacerbated by widespread antibiotic resistance, emergence of new pathogens in addition to the resurgence of old ones, and the lack of effective new therapeutics. The scientific efforts of the last 50 years centered around about a dozen antimicrobial core chemotype scaffolds.<sup>1,2</sup> The major ones among these, which have become available on the market, are the oxazolidinone synthetic core (e.g., linezolid), the lipopeptides (e.g., daptomycin), and the ketolides (e.g., telithromycin), which are modified macrolides.<sup>3,4</sup> In the recent years, many reports in the literature have suggested that naturally occurring antimicrobial peptides (AMPs<sup>a</sup>), which constitute a major component of the innate self-defense system, have the potential to represent such a class of antibiotics.<sup>5–7</sup> These AMPs are not only lethal to a broad spectrum of pathogens but also have a unique low tendency for resistance development. Although the exact mode of action of AMPs is still not completely understood, it has been well established that AMPs interact with the cell membrane of susceptible microorganisms, where either their accumulation in the membrane causes increased permeability and loss of barrier function or they cross the membrane to access

cytoplasmic targets.<sup>8,9</sup> The selective action of generally positively charged AMPs is demonstrated by preferential interaction with the anionic phospholipids of the bacterial cell membrane rather than with the neutral mammalian cell membrane, which is made of zwitterionic phospholipids and cholesterol.<sup>10–12</sup>


However, in spite of the rapid action of naturally occurring AMPs against a broad spectrum of microorganisms and the fact that the development of resistance by the microorganisms against them is slow because it requires substantial changes in the lipid composition of cell membranes of the microorganisms, there are some serious drawbacks that limit their practical use. One main disadvantage with naturally occurring AMPs is their large size, which poses several challenges regarding synthesis, metabolic stability, immunogenicity, bioavailability, route of administration, and production costs. One approach that could be adopted to resolve these problems is to design and develop smaller synthetic peptidomimetics having unnatural residues without compromising the minimum requirement for being antimicrobial in nature. Thus, the focus of the present work has been to discover small synthetic antimicrobial peptidomimetics that can ultimately lead to promising candidates for novel antimicrobial therapeutics.

In our earlier research endeavor that has been recently reported,<sup>13</sup> we tried to ascertain the minimum pharmacophore by employing the HipHop module<sup>14</sup> of the software package CATALYST (Accelrys, San Diego, CA). For this purpose, the experimental data reported by Svendsen and co-workers was used wherein the synthesis and evaluation of smaller cationic AMPs composed of two to six natural and synthetic amino acids has been reported.<sup>15,16</sup> In absence of any substantial report regarding the presence of a particular receptor or protein target for cationic AMPs, the study was restricted to include only positively ionizable (PI) and hydrophobic (HYD) features for pharmacophore generation, as


\*To whom correspondence should be addressed. Phone: +91-172-2214682. Fax: +91-172-2214692. E-mail: rahuljain@niper.ac.in.

<sup>a</sup> Abbreviations: AMP, antimicrobial peptide; CDI, 1,1'-carbonyldiimidazole; DCC, *N,N'*-dicyclohexylcarbodiimide; DIC, *N,N'*-diisopropylcarbodiimide; HPLC, high-pressure liquid chromatography; HONB, *endo-N*-hydroxy-5-norbornene-2,3-dicarboximide; HYD, hydrophobic;  $\text{IC}_{50}$ , inhibitory concentration that affords 50% inhibition of microbial growth; MIC, minimum inhibitory concentration; MRSA, methicillin-resistant *Staphylococcus aureus*; MRSE, methicillin resistant *S. epidermidis*; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2*H*-tetrazolium bromide; PI, positively ionizable; YPD, yeast extract/peptone/dextrose.




Scheme 1<sup>a</sup>

<sup>a</sup> Reagents and conditions: (i) C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>NH<sub>2</sub>, DCC, DMF, 4 °C, 12 h; (ii) 3 N HCl in 1,4-dioxane, rt, 15 min; (iii) 7 N NH<sub>3</sub> in MeOH, 0 °C, 10 min; (iv) Boc-Trp-OH, HONB, DIC, DMF, 4 °C, 48 h; (v) 3 N HCl in 1,4-dioxane, rt, 15 min.

Scheme 2<sup>a</sup>


<sup>a</sup> Reagents and conditions: (i) MeOH, HCl gas, 4 °C, 2 h; reflux, 8 h; (ii) 7 N NH<sub>3</sub> in MeOH, 0 °C, 10 min; (iii) Boc-Trp-OH, HONB, DIC, DMF, 4 °C, 48 h; (iv) 3 N HCl in 1,4-dioxane, rt, 15 min.

Scheme 3<sup>a</sup>

<sup>a</sup> Reagents and conditions: (i) Boc-L-His(2-alkyl)-OH (**1a-f**); HONB, DIC, DMF, 4 °C, 48 h; (ii) 3 N HCl in 1,4-dioxane, rt, 15 min.


protective groups to obtain the desired amide in several steps. However, we required an amide in which the  $\alpha$ -amino group of Arg remained unprotected and thus initiated a study to achieve the amidation reaction on fully unprotected

Arg. A study that has been recently reported by us proved successful with identification of 1,1'-carbonyldiimidazole (CDI) as a useful reagent for the desired purpose.<sup>24</sup> The unprotected arginine upon reaction with benzylamine in

Scheme 4<sup>a</sup>

<sup>a</sup> Reagents and conditions: (i) H-Arg-OMe (**13**), HONB, DIC, DMF, 4 °C, 48 h; (ii) 3 N HCl in 1,4-dioxane, rt, 15 min.

**Table 1.** Antimicrobial Activities of Synthesized Dipeptides (Series 1)




no.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	MRSA		MRSE		<i>E. coli</i>		<i>K. pneumoniae</i>		<i>P. aeruginosa</i>		<i>C. albicans</i>	
				MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>
4a	CH(CH <sub>3</sub> ) <sub>2</sub>	NHBzl	<i>t</i> -Boc	200	10.26	> 200		> 200		> 200		> 200		> 200	
4b	C(CH <sub>3</sub> ) <sub>3</sub>	NHBzl	<i>t</i> -Boc	> 200		> 200		> 200		> 200		> 200		> 200	
4c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	NHBzl	<i>t</i> -Boc	200	22.36	> 200		> 200		> 200		> 200		200 33.17	
4d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	NHBzl	<i>t</i> -Boc	200	31.69	200	23.81	> 200		> 200		> 200		> 200	
4e	adamantan-1-yl	NHBzl	<i>t</i> -Boc	100	7.59	200	29.07	> 200		200		> 200		200 28.34	
4f	H	NHBzl	<i>t</i> -Boc	> 200		200	15.62	> 200		200	21.24	200	40.70	200 9.70	
5a	CH(CH <sub>3</sub> ) <sub>2</sub>	NHBzl	H	> 200		100	12.80	> 200		> 200		> 200		> 200	
5b	C(CH <sub>3</sub> ) <sub>3</sub>	NHBzl	H	> 200		100	9.24	> 200		> 200		> 200		> 200	
5c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	NHBzl	H	100	27.10	50	8.51	200	22.49	> 200		> 200		100 10.30	
5d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	NHBzl	H	10	4.80	50	7.54	> 200		50	17.37	> 200		100 8.41	
5e	adamantan-1-yl	NHBzl	H	100	15.96	50	4.70	200	21.34	200	31.99	> 200		50 8.25	
5f	H	NHBzl	H	200	12.26	200	28.98	> 200		> 200		> 200		200 15.17	
8a	CH(CH <sub>3</sub> ) <sub>2</sub>	OMe	<i>t</i> -Boc	> 200		> 200		> 200		> 200		> 200		> 200	
8b	C(CH <sub>3</sub> ) <sub>3</sub>	OMe	<i>t</i> -Boc	50	4.35	100	8.12	200	16.27	> 200		> 200		> 200	
8c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	OMe	<i>t</i> -Boc	100	2.95	100	5.44	200	24.61	> 200		> 200		> 200	
8d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	OMe	<i>t</i> -Boc	200	13.96	50	6.29	200	20.87	> 200		> 200		200 17.70	
8e	adamantan-1-yl	OMe	<i>t</i> -Boc	20	1.69	50	4.03	100	6.59	> 200		> 200		200 26.61	
8f	H	OMe	<i>t</i> -Boc	200	16.37	> 200		> 200		> 200		> 200		200 31.39	
9a	CH(CH <sub>3</sub> ) <sub>2</sub>	OMe	H	200	13.13	> 200		200	12.19	> 200		> 200		200 15.62	
9b	C(CH <sub>3</sub> ) <sub>3</sub>	OMe	H	10	4.21	100	6.73	200	19.03	> 200		> 200		> 200	
9c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	OMe	H	200	9.54	200	14.52	200	9.48	> 200		> 200		200 18.25	
9d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	OMe	H	100	10.12	200	12.46	> 200		200	29.38	> 200		> 200	
9e	adamantan-1-yl	OMe	H	20	2.30	20	4.59	> 200		200	24.41	> 200		> 200	
9f	H	OMe	H	> 200		> 200		200	18.52	> 200		> 200		200 20.62	

<sup>a</sup> In  $\mu\text{g/mL}$ . Standard used: *C. albicans* (amphotericin B, IC<sub>50</sub> = 0.34  $\mu\text{g/mL}$ ), MRSA (vancomycin, IC<sub>50</sub> = 0.14  $\mu\text{g/mL}$ ), MRSE (vancomycin, IC<sub>50</sub> = 0.14  $\mu\text{g/mL}$ ), *E. coli* (streptomycin, IC<sub>50</sub> = 0.73  $\mu\text{g/mL}$ ), *K. pneumoniae* (neomycin, IC<sub>50</sub> = 0.6  $\mu\text{g/mL}$ ), *P. aeruginosa* (ciprofloxacin, IC<sub>50</sub> = 1.18  $\mu\text{g/mL}$ ).

water in the presence of CDI conveniently produced the desired amide product. The reaction was found to be stereo-conservative in nature, and the side chain functionality was found to be relatively unaffected by the reagents and reaction conditions. Further, L-Arg-NHBzl (**10**) upon coupling with *N*- $\alpha$ -Boc-2-alkyl-L-histidines (**1a-f**) in the presence of DIC and HONB in DMF produced the Boc protected dipeptides **11a-f** (Scheme 3). As discussed earlier, removal of the Boc group afforded the desired dipeptides **12a-f**. On the other hand, the synthesis of similar peptides having a methyl ester group at the C-terminus was carried out by coupling

compounds **1a-f** with commercially available L-Arg-OMe (**13**) in the same way as earlier to produce protected dipeptides **14a-f**, which finally afforded the desired dipeptides **15a-f** upon cleavage of the Boc group (Scheme 4).

**Antimicrobial Activity.** The results of antibacterial and antifungal activities of the Trp-His peptides (series 1) are provided in Table 1. Analogues **8e**, **9e**, **8c**, **8b**, and **5d** were most potent against methicillin-resistant *Staphylococcus aureus* (MRSA) with IC<sub>50</sub> values of 1.69, 2.30, 2.95, 4.35, and 4.80  $\mu\text{g/mL}$ , respectively. Thus, it can be interpreted that the presence of hydrophobicity (bulk) at the C-2 position of

**Table 2.** Antimicrobial Activities of Synthesized Dipeptides (Series 2)

no.	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	MRSA		MRSE		<i>E. coli</i>		<i>K. pneumoniae</i>		<i>P. aeruginosa</i>		<i>C. albicans</i>	
				MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>	MIC <sup>a</sup>	IC <sub>50</sub> <sup>a</sup>
11a	CH(CH <sub>3</sub> ) <sub>2</sub>	NHBzl	<i>t</i> -Boc	100	17.95	100	12.66	200	23.69	> 200		200	45.15	> 200	
11b	C(CH <sub>3</sub> ) <sub>3</sub>	NHBzl	<i>t</i> -Boc	200	17.52	> 200		100	12.03	> 200		> 200		200	33.15
11c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	NHBzl	<i>t</i> -Boc	> 200		200	19.30	> 200		200	20.81	> 200		> 200	
11d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	NHBzl	<i>t</i> -Boc	100	12.07	100	18.42	200		> 200		> 200		> 200	
11e	adamantan-1-yl	NHBzl	<i>t</i> -Boc	20	7.47	100	11.50	> 200		100	15.08	100	27.21	> 200	
11f	H	NHBzl	<i>t</i> -Boc	> 200		100	8.79	> 200		> 200		> 200		> 200	
12a	CH(CH <sub>3</sub> ) <sub>2</sub>	NHBzl	H	100	4.85	200	12.76	> 200		> 200		200	27.76	> 200	
12b	C(CH <sub>3</sub> ) <sub>3</sub>	NHBzl	H	20	4.18	10	3.13	> 200		> 200		> 200		> 200	
12c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	NHBzl	H	50	6.99	50	3.94	> 200		> 200		> 200		100	9.95
12d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	NHBzl	H	20	2.06	5	1.00	200	17.51	200	23.52	> 200		200	18.00
12e	adamantan-1-yl	NHBzl	H	5	2.15	10	1.19	200	9.29	100	16.15	> 200		50	7.46
12f	H	NHBzl	H	100	7.04	100	8.07	> 200		> 200		> 200		200	16.64
14a	CH(CH <sub>3</sub> ) <sub>2</sub>	OMe	<i>t</i> -Boc	200	12.45	200	13.10	> 200		> 200		> 200		200	20.86
14b	C(CH <sub>3</sub> ) <sub>3</sub>	OMe	<i>t</i> -Boc	100	4.15	200	20.14	> 200		> 200		> 200		> 200	
14c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	OMe	<i>t</i> -Boc	> 200		> 200		200	18.12	200	14.97	> 200		> 200	
14d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	OMe	<i>t</i> -Boc	200	10.08	100	10.00	100	4.93	> 200		200	27.53	100	9.44
14e	adamantan-1-yl	OMe	<i>t</i> -Boc	50	9.77	50	6.87	100	8.33	200	31.47	100	25.71	200	16.11
14f	H	OMe	<i>t</i> -Boc	> 200		100	5.07	> 200		> 200		> 200		> 200	
15a	CH(CH <sub>3</sub> ) <sub>2</sub>	OMe	H	50	4.82	50	5.42	200	9.98	> 200		> 200		> 200	
15b	C(CH <sub>3</sub> ) <sub>3</sub>	OMe	H	20	5.39	20	2.53	200	18.48	200	10.52	> 200		100	9.65
15c	<i>c</i> -C <sub>4</sub> H <sub>7</sub>	OMe	H	200	9.28	100	6.66	200	15.31	> 200		> 200		200	13.96
15d	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	OMe	H	50	20.95	10	1.92	> 200		> 200		> 200		200	27.55
15e	adamantan-1-yl	OMe	H	10	3.67	20	4.28	> 200		200	19.38	200	35.48	> 200	
15f	H	OMe	H	100	4.88	50	2.72	> 200		> 200		> 200		> 200	

<sup>a</sup> In  $\mu\text{g/mL}$ . Standard used: *C. albicans* (amphotericin B, IC<sub>50</sub> = 0.34  $\mu\text{g/mL}$ ), MRSA (vancomycin, IC<sub>50</sub> = 0.14  $\mu\text{g/mL}$ ), MRSE (vancomycin, IC<sub>50</sub> = 0.14  $\mu\text{g/mL}$ ), *E. coli* (streptomycin, IC<sub>50</sub> = 0.73  $\mu\text{g/mL}$ ), *K. pneumoniae* (neomycin, IC<sub>50</sub> = 0.6  $\mu\text{g/mL}$ ), *P. aeruginosa* (ciprofloxacin, IC<sub>50</sub> = 1.18  $\mu\text{g/mL}$ ).

the imidazole ring affects the antimicrobial activity against MRSA in a positive way. As all but one of the above stated active peptides have a methyl ester group at the C-terminus, it can also be stated that OMe is more effective as compared to the NHBzl group for increasing the activity against MRSA for such Trp-His peptides. Also, it was observed that analogues **4a**, **4c**, **4d**, **4e**, **5c**, **5e**, **5f**, **8d**, **8f**, **9a**, and **9d** were found to be moderately effective, with IC<sub>50</sub> values ranged between 7 and 32  $\mu\text{g/mL}$  against MRSA. Whereas against methicillin resistant *S. epidermidis* (MRSE), analogues **8e**, **9e**, and **5e** were found to be most potent with IC<sub>50</sub> values of 4.03, 4.59, and 4.70  $\mu\text{g/mL}$ , respectively. The importance of bulk at position R<sub>1</sub> has been underlined with the fact that all these active peptides contain adamantan-1-yl at this position. Some other analogues like **5b**, **5c**, **5d**, **8b**, **8c**, **8d**, **9b**, **9c**, and **9d** also displayed encouraging activities with IC<sub>50</sub> values in the range between 5 and 15  $\mu\text{g/mL}$ . Coming to Gram-negative bacterial strains, some compounds were found to be active against *E. coli*. Analogue **8e** was found to be most potent against *E. coli* with an IC<sub>50</sub> value of 6.59  $\mu\text{g/mL}$ . Some other analogues like **9c**, **9a**, and **8a** also displayed encouraging activities with IC<sub>50</sub> values of 9.48, 12.19, and 16.27  $\mu\text{g/mL}$ , respectively. Interestingly, only peptides having a Boc group at the N-terminus were found most active against *E. coli*.

This can be interpreted in a way that the presence of hydrophobicity and/or the deficiency of positively ionizable features at the N-terminus of His-Trp peptides is essential for activity against *E. coli*. However, the importance of a bulkier residue at position R<sub>1</sub> seems to be equally important as discussed earlier. In the case of *Klebsiella pneumoniae*, very few peptides exhibited moderate activities such as analogues **5d** (IC<sub>50</sub> = 17.37  $\mu\text{g/mL}$ ), **5e** (IC<sub>50</sub> = 31.99  $\mu\text{g/mL}$ ), and **4f** (IC<sub>50</sub> = 21.24  $\mu\text{g/mL}$ ). However, except **4f** (IC<sub>50</sub> = 40.70  $\mu\text{g/mL}$ ), no analogue of this series was found to be active against *Pseudomonas aeruginosa*. Analogues **5e** and **5d** with adamantan-1-yl and cyclohexyl groups at the imidazole ring and an NHBzl group at the C-terminus were most potent against the fungus *Candida albicans* with IC<sub>50</sub> values of 8.25 and 8.41  $\mu\text{g/mL}$ , respectively. This leads to the conclusion that the presence of a bulky group at the C-terminus as well as at the C-2 position of the imidazole ring increases the activity of Trp-His peptides against *C. albicans*. Also, the presence of a free amino group at the N-terminus, which increases the overall cationicity of the peptide, seems to be important for a good activity against the same strain. Apart from these peptides, many other analogues such as **4c**, **4e**, **5c**, **5f**, **8d**, **8e**, **8f**, **9a**, **9c**, and **9f** produced modest activities with IC<sub>50</sub> values in the range of

10–35  $\mu\text{g/mL}$  for the same strain. Thus, broadly, it is observed that out of all synthesized Trp-His peptides, **5e**, which is substituted with a bulky adamantan-1-yl group at the C-2-position in the imidazole ring and has a NHBzl group at the C-terminus, was active against five strains exhibiting a broad spectrum of antimicrobial activities. Analogues **4f**, **5c**, **5d**, **8e**, and **9c** displayed promising antimicrobial activity against four tested strains.

The antimicrobial results for the His-Arg peptides (series 2) are shown in Table 2. These peptides showed high activities against MRSA. Analogue **12d** exhibited the most potent activity with an  $\text{IC}_{50}$  of 2.06  $\mu\text{g/mL}$ . In total, the eight compounds **12a**, **12b**, **12d**, **12e**, **14b**, **15a**, **15e**, and **15f** exhibited  $\text{IC}_{50}$  values  $\leq 5 \mu\text{g/mL}$ . In the case of the His-Arg peptides, the peptides with a Boc group at the N-terminus are more active as compared to the unprotected ones. A possible explanation for this observation is that the reduction in hydrophobicity that is caused by the substitution of tryptophan with histidine is compensated by the bulkier Boc group, whereas the cationicity imparted by the guanidinium group of the arginine side chain plays an important role in the activity. Some of the tested peptides also showed encouraging activities against MRSE. Analogues **12c**, **15d**, **15b**, and **15f** were highly potent with  $\text{IC}_{50}$  values of 1.00, 1.92, 2.53, and 2.72  $\mu\text{g/mL}$ , respectively. Apart from these analogues, at least 10 compounds from series 2 exhibited  $\text{IC}_{50}$  values in the range of 2–10  $\mu\text{g/mL}$ . Contrary to the observation made in case of MRSA, peptides with a free amino group at the N-terminus are most active against MRSE. Very few compounds were active against *E. coli*. Analogue **14d** was found to be the most potent peptide against *E. coli* ( $\text{IC}_{50} = 4.93 \mu\text{g/mL}$ ). Analogues **14e**, **12e**, and **15a** exhibited modest activity with  $\text{IC}_{50}$  values of 8.33, 9.29, and 9.98  $\mu\text{g/mL}$ , respectively. Similarly, analogues **15b** ( $\text{IC}_{50} = 10.52 \mu\text{g/mL}$ ), **14c** ( $\text{IC}_{50} = 14.97 \mu\text{g/mL}$ ), **11e** ( $\text{IC}_{50} = 15.08 \mu\text{g/mL}$ ), **12e** ( $\text{IC}_{50} = 16.15 \mu\text{g/mL}$ ), and **11c** ( $\text{IC}_{50} = 20.81 \mu\text{g/mL}$ ) were the only compounds exhibiting moderate activities against *K. pneumoniae*. However, except of a few peptides that were moderately active, most peptides belonging to series 2 were inactive against *P. aeruginosa*. Analogues **12e**, **14d**, **15b**, and **12c** were found to be most potent against the fungus *C. albicans* with  $\text{IC}_{50}$  values of 7.46, 9.44, 9.65, and 9.95  $\mu\text{g/mL}$ , respectively. Many other analogues produced modest activities with  $\text{IC}_{50}$  values in the range of 16–34  $\mu\text{g/mL}$  for the same pathogen. It seems that the presence of a bulky group at position  $R_1$  is much more important than the variations tried at positions  $R_2$  and  $R_3$  for the His-Arg peptides concerning activity against *C. albicans*.

As a conclusion, it can be interpreted that out of all synthesized His-Arg analogues, compound **12e** (substituted with a bulky and hydrophobic adamantan-1-yl group at the C-2-position of the imidazole ring with NHBzl at the C-terminus) and **14d** and **14e** substituted with bulky cyclohexyl and adamantan-1-yl groups, respectively, were active against five strains exhibiting a broad spectrum of antimicrobial activities. Analogue **15d** displayed promising antimicrobial activity against four tested strains.

**Cytotoxic Experiments.** Upon the basis of the results from the antimicrobial testing, the most active peptides were screened for cytotoxicity using a MTT assay with mouse fibroblasts. All of the tested dipeptides, belonging to both series 1 and 2, showed no cytotoxic effect up to the highest test concentration of 200  $\mu\text{g/mL}$ .

## Conclusions

Our experiments have resulted in the identification of novel dipeptide analogues composed of the modified natural amino acids His, Trp, and Arg, which will help in contributing to increase the enzymatic stability of the peptides. Apart from containing synthetic residues, these peptides have an extremely small size, allowing the possibility of designing active peptides far smaller than previously described. Also, with the help of this study, we have established that His and its derivatives can replace both Trp and Arg to different extents in the Trp-Arg class of AMPs, thereby introducing His-Arg and Trp-His classes of AMPs. It has also been noted that variation of hydrophobicity, particularly as a substituent of the imidazole ring of histidine, plays an important role in effecting antimicrobial activity of compounds of both classes. In general, the His-Arg peptides were more active against the tested strains. This is mainly attributed to enhanced cationic character as compared to the other class.

## Experimental Section

The synthesized amino acids and dipeptides were checked for their purity on precoated silica gel G<sub>254</sub> TLC plates (Merck), and the spots were visualized under UV light and by exposing them to iodine vapors. Column chromatographic purification was carried out on Merck silica gel (230–400 mesh) or neutral alumina. Melting points were recorded on a capillary melting point apparatus and are uncorrected. All solvents used for synthesis were of analytical grade and used without any further purification unless otherwise stated.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded on a 300 MHz Bruker FT-NMR (Avance DPX 300) spectrometer using tetramethylsilane as internal standard, and the chemical shifts are reported in  $\delta$  units. Mass spectra were recorded on a Finnigan Mat LCQ spectrometer (APCI/ESI). Optical rotations were recorded on a Perkin-Elmer 241MC polarimeter. Elemental analyses were recorded on Elementar Vario EL spectrometer. The elemental analyses of all final peptides were within  $\pm 0.4\%$  of the expected values unless otherwise stated. All final peptides were checked for their homogeneity on a Shimadzu SPD-M20A HPLC system using a Supelcosil LC-8, 5  $\mu\text{m}$  (25 cm  $\times$  4.6 mm ID) column. The peptides were analyzed by using a solvent system of  $\text{CH}_3\text{CN}-\text{H}_2\text{O}$  (0.1% TFA) in the form of a 20 min gradient: 5–65%  $\text{CH}_3\text{CN}$  in 12 min, 65–95%  $\text{CH}_3\text{CN}$  in 3 min, and 100%  $\text{CH}_3\text{CN}$  in 5 min. The peptides were  $\geq 95\%$  pure on HPLC analysis. Amino acids, and coupling reagents, DMF, DCC, DIC, and TFA, were purchased from either Chem-Impex International or NovaBiochem (Merck Ltd.).

**General Method for the Synthesis of *N*- $\alpha$ -Boc-L-Trp-L-His(2-alkyl)-NHBzl (**4a–f**, Scheme 1).** HONB (1.2 mmol) was added to *N*- $\alpha$ -Boc-L-Trp (1 mmol) in water-free DMF (10 mL). At  $-10^\circ\text{C}$ , DIC (1.2 mmol) was added, and after 5 min stirring, L-His(2-alkyl)-NHBzl (**3a–f**, 1 mmol) was added. After stirring for 36 h at  $4^\circ\text{C}$ , the solvent was removed and the crude product chromatographed on neutral alumina using  $\text{CH}_3\text{OH}:\text{CHCl}_3$  (7:93) to afford *N*- $\alpha$ -Boc-L-Trp-L-His(2-alkyl)-NHBzl (**4a–f**).

***N*- $\alpha$ -Boc-L-Trp-L-His(2-isopropyl)-NHBzl (**4a**).** Yield: 52%, light-yellowish solid; mp 209–210  $^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.17 (d, 6H,  $J = 7.1$  Hz), 1.28 (s, 9H), 2.88 (m, 2H), 3.07 (m, 1H), 3.09 (m, 2H), 3.88 (t, 1H,  $J = 7.3$  Hz), 4.37 (t, 1H,  $J = 6.8$  Hz), 4.69 (m, 1H), 4.87 (m, 1H), 6.66 (s, 1H), 7.08–7.11 (m, 3H), 7.30–7.38 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  21.53, 27.94, 28.29, 28.60, 29.83, 42.71, 52.45, 56.95, 80.32, 107.29, 111.28, 117.40, 118.59, 120.32, 122.46, 123.85, 126.15, 128.59, 128.91, 129.69, 134.81, 138.34, 155.37, 170.79, 171.40. MS (APCI):  $m/z$  573.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His(2-tert-butyl)-NHBzl (**4b**).** Yield: 62%, colorless solid; mp 214–216  $^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.24 (s,

9H), 1.31 (s, 9H), 2.71 (m, 2H), 3.14 (m, 2H), 3.68 (t, 1H,  $J = 7.0$  Hz), 4.26 (t, 1H,  $J = 6.9$  Hz), 4.70 (m, 1H), 4.92 (m, 1H), 6.89 (s, 1H), 7.10–7.13 (m, 3H), 7.20–7.22 (m, 2H), 7.35–7.40 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  28.27, 28.91, 29.37, 30.17, 31.82, 37.29, 42.28, 52.37, 55.93, 79.80, 109.71, 111.48, 116.82, 118.38, 119.38, 123.60, 124.31, 127.74, 128.09, 129.51, 137.11, 138.91, 156.27, 172.52, 174.51. MS (APCI):  $m/z$  587.2  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His(2-cyclobutyl)-NHBzl (4c).** Yield: 48%, colorless solid; mp 224–227 °C (dec.);  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.27 (s, 9H), 1.86 (m, 2H), 2.29 (m, 4H), 2.89 (m, 2H), 3.06 (m, 1H), 3.17 (m, 2H,  $\text{CH}_2$ ), 3.57 (t, 1H,  $J = 6.9$  Hz), 4.27 (t, 1H,  $J = 6.8$  Hz), 4.48 (m, 1H), 4.71 (m, 1H), 6.58 (s, 1H), 7.05–7.09 (m, 3H), 7.21–7.32 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  19.37, 25.73, 27.91, 28.39, 28.90, 29.81, 31.81, 43.82, 52.17, 54.71, 78.96, 110.15, 111.63, 116.87, 118.71, 119.35, 123.04, 124.10, 126.19, 127.89, 129.00, 130.46, 136.51, 139.90, 158.27, 173.80, 181.61. MS (APCI):  $m/z$  585.1  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His(2-cyclohexyl)-NHBzl (4d).** Yield: 60%, light-yellowish solid; mp 241–242 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.23 (m, 6H), 1.35 (s, 9H), 1.71 (m, 4H), 1.96 (m, 1H), 2.97 (m, 2H), 3.24 (m, 2H), 4.04 (t, 1H,  $J = 7.0$  Hz), 4.35 (t, 1H,  $J = 6.6$  Hz), 4.63 (m, 1H), 4.88 (m, 1H), 6.83 (s, 1H), 7.02–7.10 (m, 3H), 7.16–7.21 (m, 2H), 7.25–7.30 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  24.90, 26.41, 26.84, 28.05, 29.17, 30.37, 32.77, 38.27, 44.10, 52.78, 55.38, 77.71, 111.25, 111.34, 115.72, 119.10, 120.73, 123.38, 124.29, 126.16, 127.90, 128.15, 129.70, 137.33, 142.51, 156.21, 171.92, 177.52. MS (APCI):  $m/z$  613.2  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His[2-(adamantan-1-yl)]-NHBzl (4e).** Yield: 63%, colorless solid; mp 254–256 °C.;  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.24 (s, 9H), 1.51–1.54 (m, 12H), 1.97 (m, 3H), 3.00 (m, 2H), 3.22 (m, 2H), 4.09 (t, 1H,  $J = 7.2$  Hz), 4.28 (t, 1H,  $J = 6.8$  Hz), 4.61 (m, 1H), 4.85 (m, 1H), 6.89 (s, 1H), 7.01–7.09 (m, 3H), 7.17–7.30 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  26.34, 27.05, 27.70, 28.17, 34.69, 38.46, 41.23, 44.57, 53.82, 55.38, 80.02, 110.14, 111.61, 115.41, 118.60, 120.51, 122.28, 124.55, 126.81, 127.68, 128.73, 129.12, 136.07, 144.28, 159.61, 172.28, 175.04. MS (APCI):  $m/z$  665.2  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His-NHBzl (4f).** Yield: 65%, colorless solid; mp 194–196 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.37 (s, 9H), 2.77 (m, 2H), 3.25 (m, 2H), 3.71 (t, 1H,  $J = 6.9$  Hz), 4.29 (t, 1H,  $J = 6.8$  Hz), 4.51 (m, 1H), 4.72 (m, 1H), 6.79 (s, 1H), 7.03–7.09 (m, 3H), 7.23–7.34 (m, 7H, Ar-H), 7.41 (s, 1H, CH).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  26.61, 28.56, 29.18, 42.26, 53.75, 59.02, 73.29, 109.18, 111.39, 117.56, 118.12, 119.19, 121.89, 123.48, 128.76, 129.19, 129.82, 135.80, 137.92, 153.21, 170.74, 173.96. MS (APCI):  $m/z$  531.0  $[\text{MH}]^+$ .

**General Method for the Synthesis of L-Trp-L-His(2-alkyl)-NHBzl·2HCl (5a–f, Scheme 1).** A solution of *N*- $\alpha$ -Boc-L-Trp-L-His(2-alkyl)-NHBzl (4a–f, 1 mmol) in 3 N HCl in 1,4-dioxane (15 mL) was stirred at ambient temperature for 15 min. The solvent was evaporated under reduced pressure to produce L-Trp-L-His(2-alkyl)-NHBzl·2HCl (5a–f).

**L-Trp-L-His(2-isopropyl)-NHBzl·2HCl (5a).** Yield: 90%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.27 (d, 6H), 2.96 (m, 2H), 3.15 (m, 1H), 3.20 (m, 2H), 3.82 (t, 1H,  $J = 7.0$  Hz), 4.42 (t, 1H,  $J = 6.9$  Hz), 4.68 (m, 1H), 4.89 (m, 1H), 6.91 (s, 1H), 7.11–7.18 (m, 3H), 7.34–7.41 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  20.59, 26.28, 28.02, 29.61, 41.38, 52.70, 58.93, 109.22, 112.65, 117.42, 118.97, 121.64, 122.86, 125.32, 127.78, 128.81, 129.10, 129.93, 138.13, 145.34, 168.58, 173.46. MS (APCI):  $m/z$  473.0  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -14.9^\circ$  ( $c = 1.0$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His(2-tert-butyl)-NHBzl·2HCl (5b).** Yield: 94%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.26 (s, 9H), 2.61 (m, 2H), 2.99 (m, 2H), 3.16 (t, 1H,  $J = 7.2$  Hz), 4.32 (t, 1H,  $J = 6.9$  Hz), 4.57 (m, 1H), 4.79 (m, 1H), 6.75 (s, 1H), 7.03 (m, 3H), 7.26–7.35 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  28.70, 28.66, 29.58, 30.48, 32.51, 36.51, 43.02, 53.16, 57.81, 111.19, 113.34, 117.41, 118.30, 119.83, 122.65, 124.00, 127.67, 128.89, 129.26, 135.31, 138.35, 170.57, 178.13. MS (APCI):  $m/z$  487.1  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -19.2^\circ$  ( $c = 1.8$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His(2-cyclobutyl)-NHBzl·2HCl (5c).** Yield: 93%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.73 (m, 2H), 2.18 (m, 4H), 2.83 (m, 2H), 3.08 (m, 1H), 3.21 (m, 2H), 3.71 (t, 1H,  $J = 6.8$  Hz), 4.37 (t, 1H,  $J = 6.7$  Hz), 4.56 (m, 1H), 4.87 (m, 1H), 6.80 (s, 1H), 7.09–7.14 (m, 3H), 7.29–7.37 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  21.32, 25.81, 27.45, 28.79, 30.37, 31.94, 44.18, 53.78, 54.61, 109.19, 111.61, 118.04, 118.96, 120.24, 122.40, 124.37, 127.02, 128.83, 129.55, 130.68, 133.83, 138.28, 171.83, 176.91. MS (APCI):  $m/z$  485.2  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -15.0^\circ$  ( $c = 1.5$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His(2-cyclohexyl)-NHBzl·2HCl (5d).** Yield: 90%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.19–1.23 (m, 6H), 1.66 (m, 4H), 2.07 (m, 1H), 2.89 (m, 2H), 3.18 (m), 3.94 (t, 1H,  $J = 7.1$  Hz), 4.27 (t, 1H,  $J = 6.7$  Hz), 4.60 (m, 1H), 4.98 (m, 1H), 6.68 (s, 1H), 6.99–7.06 (m, 3H), 7.21 (m, 2H), 7.29–7.35 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  25.19, 26.79, 27.44, 28.10, 30.07, 32.77, 36.12, 43.72, 52.76, 56.31, 109.25, 111.08, 116.51, 119.89, 121.76, 124.35, 125.29, 127.21, 128.54, 129.80, 136.67, 145.14, 173.27, 175.55. MS (APCI):  $m/z$  513.3  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -22.6^\circ$  ( $c = 1.6$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His[2-(adamantan-1-yl)]-NHBzl·2HCl (5e).** Yield: 95%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.49 (m, 12H), 2.07 (m, 3H), 2.95 (m, 2H), 3.13 (m, 2H), 3.95 (t, 1H,  $J = 6.8$  Hz), 4.26 (t, 1H,  $J = 6.9$  Hz), 4.72 (m, 1H), 4.88 (m, 1H), 6.95 (s, 1H), 7.07–7.14 (m, 3H), 7.26–7.33 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  25.94, 26.89, 27.41, 28.07, 35.16, 39.30, 42.27, 44.36, 52.63, 57.17, 110.19, 111.99, 116.80, 118.68, 122.00, 122.93, 124.51, 126.58, 127.37, 128.56, 129.33, 135.88, 143.28, 170.98, 174.49. MS (APCI):  $m/z$  565.3  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -17.0^\circ$  ( $c = 1.5$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His-NHBzl·2HCl (5f).** Yield: 95%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  2.69 (m, 2H), 3.16 (m, 2H), 3.83 (t, 1H,  $J = 6.7$  Hz), 4.37 (t, 1H,  $J = 6.8$  Hz), 4.69 (m, 1H), 4.95 (m, 1H), 6.86 (s, 1H), 7.06–7.10 (m, 3H), 7.28–7.37 (m, 7H), 7.55 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  27.51, 29.66, 43.76, 52.42, 57.81, 107.61, 112.25, 118.28, 119.69, 119.99, 120.56, 123.14, 127.43, 128.71, 129.35, 137.90, 139.28, 172.40, 177.22. MS (APCI):  $m/z$  431.2  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -11.6^\circ$  ( $c = 1.2$ ,  $\text{CH}_3\text{OH}$ ).

**General Method for the Synthesis of *N*- $\alpha$ -Boc-L-Trp-L-His(2-alkyl)-OMe (8a–f, Scheme 2).** To a solution of L-His(2-alkyl)-OMe·2HCl (7a–f, 1 mmol) was added ammonia (7 N solution in  $\text{CH}_3\text{OH}$ ) at 0 °C. The reaction mixture was stirred for 10 min, and the solvent was removed under reduced pressure to produce L-His(2-alkyl)-OMe. HONB (1.2 mmol) was added to *N*- $\alpha$ -Boc-L-Trp (1 mmol) in water-free DMF (10 mL). At –10 °C, DIC (1.2 mmol) was added, and after 5 min stirring, L-His(2-alkyl)-OMe (1 mmol) was added. After stirring for 36 h at 4 °C, the solvent was removed and the crude product chromatographed on neutral alumina using  $\text{CH}_3\text{OH}:\text{CHCl}_3$  (5:95) to afford *N*- $\alpha$ -Boc-L-Trp-L-His(2-alkyl)-OMe (8a–f).

***N*- $\alpha$ -Boc-L-Trp-L-His(2-isopropyl)-OMe (8a).** Yield: 70%, colorless solid; mp 188–190 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.26 (d, 6H,  $J = 7.1$  Hz), 1.35 (s, 9H), 2.96 (m, 2H), 3.17 (m, 1H), 3.34 (m, 2H), 3.63 (s, 3H), 3.66 (t, 1H,  $J = 7.2$  Hz), 4.35 (t, 1H,  $J = 6.9$  Hz), 6.71 (s, 1H), 7.02–7.05 (m, 3H), 7.31–7.33 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  21.26, 28.10, 28.54, 30.17, 43.21, 51.68, 56.43, 80.12, 110.32, 111.74, 117.46, 118.77, 119.21, 121.81, 124.13, 128.28, 129.32, 134.85, 154.18, 157.01, 172.15, 175.12. MS (APCI):  $m/z$  498.8  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His(2-tert-butyl)-OMe (8b).** Yield: 67%, colorless solid; mp 198 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.30 (s, 9H), 1.34 (s, 9H), 2.96 (m, 2H), 3.17 (m, 2H), 3.66 (s, 3H), 3.78 (t, 1H,  $J = 7.1$  Hz), 4.34 (t, 1H,  $J = 6.9$  Hz), 6.68 (s, 1H), 7.01–7.08 (m, 3H), 7.30–7.33 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  28.07, 28.66, 29.19, 29.86, 30.16, 33.19, 43.20, 51.66, 52.19, 56.40, 80.08, 110.31, 111.69, 117.65, 118.76, 119.18, 121.78, 124.08, 128.28, 131.73, 134.85, 137.43, 156.60, 172.28, 175.64; MS (APCI):  $m/z$  512.2  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His(2-cyclobutyl)-OMe (8c).** Yield: 50%, light-yellowish solid; mp 194–196 °C.;  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$

1.35 (s, 9H), 1.50 (m, 2H), 2.31 (m, 4H), 2.95 (m, 2H), 3.04 (m, 1H), 3.19 (m, 2H), 3.61 (s, 3H), 3.81 (t, 1H,  $J = 6.8$  Hz), 4.35 (t, 1H,  $J = 6.9$  Hz), 6.73 (s, 1H), 7.04–7.08 (m, 3H), 7.31–7.33 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  18.91, 25.23, 27.61, 28.02, 28.47, 29.67, 31.98, 42.18, 51.92, 52.31, 54.89, 79.83, 107.67, 111.72, 112.56, 117.34, 118.70, 119.89, 120.56, 125.23, 128.13, 134.39, 141.90, 157.36, 171.67, 176.28. MS (APCI):  $m/z$  510.9  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His(2-cyclohexyl)-OMe (8d).** Yield: 61%, colorless solid; mp 211–214 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.25 (m, 6H), 1.39 (s, 9H), 1.65 (m, 4H), 1.90 (m, 1H), 2.93 (m, 2H), 3.20 (m, 2H), 3.59 (s, 3H), 3.69 (t, 1H,  $J = 7.1$  Hz), 4.35 (t, 1H,  $J = 6.8$  Hz), 6.49 (s, 1H), 7.05–7.10 (m, 3H), 7.33–7.35 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  23.94, 26.14, 26.45, 28.33, 28.79, 30.21, 31.98, 37.10, 43.18, 51.91, 53.01, 56.51, 80.83, 109.97, 112.09, 118.90, 119.81, 122.36, 124.15, 128.02, 128.86, 134.77, 153.48, 156.47, 171.60, 172.75. MS (APCI):  $m/z$  538.2  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His[2-(adamantan-1-yl)]-OMe (8e).** Yield: 63%, colorless solid; mp 233–235 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.36 (s, 9H), 1.52–1.55 (m, 12H), 1.99 (m, 3H), 3.01 (m, 2H), 3.22 (m, 2H), 3.65 (s, 3H), 3.73 (t, 1H,  $J = 7.0$  Hz), 4.32 (t, 1H,  $J = 6.8$  Hz), 6.83 (s, 1H), 7.01–7.08 (m, 3H), 7.31–7.33 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  26.53, 26.62, 27.35, 28.91, 35.15, 39.63, 41.66, 43.47, 51.23, 54.89, 78.52, 108.71, 110.15, 115.95, 117.16, 119.89, 120.23, 122.51, 126.68, 129.14, 133.30, 135.85, 155.38, 158.19, 170.34, 173.63. MS (APCI):  $m/z$  591.0  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-Trp-L-His-OMe (8f).** Yield: 67%, white solid; mp 171–173 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  1.36 (s, 9H), 3.05 (m, 2H), 3.23 (m, 2H), 3.64 (s, 3H), 3.77 (t, 1H,  $J = 7.1$  Hz), 4.35 (t, 1H,  $J = 6.9$  Hz), 6.85 (s, 1H), 7.07 (m, 3H), 7.31 (m, 2H), 7.59 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  22.97, 27.69, 29.09, 51.68, 53.20, 56.36, 110.31, 111.69, 118.03, 119.19, 121.54, 124.11, 128.25, 134.88, 155.62, 172.14, 175.37. MS (APCI):  $m/z$  455.9  $[\text{MH}]^+$ .

**General Method for the Synthesis of L-Trp-2-alkyl-L-His-OMe·2HCl (9a–f, Scheme 2).** A solution of *N*- $\alpha$ -Boc-L-Trp-L-His(2-alkyl)-OMe (8a–f, 1 mmol) in 3 N HCl in 1,4-dioxane (15 mL) was stirred at ambient temperature for 15 min. The solvent was evaporated under reduced pressure to afford L-Trp-2-alkyl-L-His-OMe·2HCl (9a–f).

**L-Trp-L-His(2-isopropyl)-OMe·2HCl (9a).** Yield: 90%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.32 (d, 6H), 2.93 (m, 2H), 3.24 (m, 1H), 3.34 (m, 2H), 3.57 (s, 3H), 3.76 (t, 1H,  $J = 6.9$  Hz), 4.07 (t, 1H,  $J = 6.6$  Hz), 6.88 (s, 1H), 7.02–7.04 (m, 3H), 7.22 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  20.11, 27.43, 28.56, 30.13, 51.67, 54.40, 63.69, 114.58, 117.20, 118.57, 120.41, 122.23, 124.53, 127.52, 134.92, 170.50, 174.76. MS (APCI):  $m/z$  398.2  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -13.0^\circ$  ( $c = 1.0$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His(2-tert-butyl)-OMe·2HCl (9b).** Yield: 92%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.44 (s, 9H), 2.76 (m, 2H), 3.16 (m, 2H), 3.67 (s, 3H), 3.73 (t, 1H,  $J = 7.0$  Hz), 4.43 (t, 1H,  $J = 6.7$  Hz), 6.91 (s, 1H), 7.09–7.12 (m, 3H), 7.21–7.23 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  27.16, 28.17, 29.53, 30.13, 43.26, 51.66, 52.69, 54.36, 111.80, 117.24, 118.58, 119.67, 120.04, 122.24, 125.22, 129.48, 134.92, 170.89, 174.75. MS (APCI):  $m/z$  412.2  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -22.9^\circ$  ( $c = 1.7$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His(2-cyclobutyl)-OMe·2HCl (9c).** Yield: 86%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.49 (m, 2H), 2.33 (m, 4H), 2.87 (m, 2H), 3.21 (m, 1H), 3.29 (m, 2H), 3.54 (s, 3H), 3.73 (t, 1H,  $J = 6.7$  Hz), 4.32 (t, 1H,  $J = 6.8$  Hz), 6.94 (s, 1H), 7.10–7.13 (m, 3H), 7.28 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  18.74, 27.17, 28.18, 29.46, 31.43, 52.55, 52.78, 54.44, 107.19, 112.014, 117.25, 118.64, 119.65, 119.99, 122.22, 125.15, 128.56, 134.93, 169.67, 170.82. MS (APCI):  $m/z$  410.3  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -16.7^\circ$  ( $c = 1.0$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His(2-cyclohexyl)-OMe·2HCl (9d).** Yield: 95%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.28 (m, 6H), 1.74 (m, 4H), 2.02 (m, 1H), 2.98 (m, 2H), 3.30 (m, 2H), 3.64 (s, 3H), 3.73 (t, 1H,  $J = 6.9$  Hz), 4.24 (t, 1H,  $J = 6.8$  Hz), 6.88 (s, 1H), 7.03–7.08 (m, 3H), 7.25 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$

25.71, 25.98, 27.19, 28.86, 30.12, 31.24, 36.44, 43.29, 51.70, 52.78, 63.70, 107.18, 111.80, 117.16, 118.10, 119.66, 120.04, 122.24, 125.26, 129.03, 134.94, 170.82, 170.75. MS (APCI):  $m/z$  438.4  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -13.9^\circ$  ( $c = 1.5$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His[2-(adamantan-1-yl)]-OMe·2HCl (9e).** Yield: 85%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.58–1.65 (m, 12H), 2.07 (m, 3H), 2.89 (m, 2H), 3.31 (m, 2H), 3.65 (s, 3H), 3.74 (t, 1H,  $J = 7.1$  Hz), 4.17 (t, 1H,  $J = 6.8$  Hz), 6.78 (s, 1H), 7.07–7.13 (m, 3H), 7.28–7.30 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  25.39, 28.58, 28.91, 30.12, 31.34, 36.29, 43.27, 51.67, 52.02, 67.54, 108.43, 111.88, 116.86, 118.30, 119.82, 121.29, 122.61, 125.92, 128.42, 134.92, 171.02, 174.75. MS (APCI):  $m/z$  490.3  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -25.4^\circ$  ( $c = 2.0$ ,  $\text{CH}_3\text{OH}$ ).

**L-Trp-L-His-OMe·2HCl (9f).** Yield: 88%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  2.90 (m, 2H), 3.21 (m, 2H), 3.55 (s, 3H), 3.64 (t, 1H,  $J = 7.0$  Hz), 3.74 (t, 1H,  $J = 6.8$  Hz), 6.93 (s, 1H), 7.12–7.15 (m, 3H), 7.27 (m, 2H), 7.42 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  27.12, 30.12, 52.58, 63.67, 67.55, 111.81, 118.19, 118.60, 119.72, 120.04, 124.53, 128.69, 134.94, 171.75, 174.97. MS (APCI):  $m/z$  356.1  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -18.4^\circ$  ( $c = 1.3$ ,  $\text{CH}_3\text{OH}$ ).

**General Method for the Synthesis of *N*- $\alpha$ -Boc-L-His(2-alkyl)-L-Arg-NHBzl (11a–f, Scheme 3).** HONB (1.2 mmol) was added to *N*- $\alpha$ -Boc-L-His(2-alkyl)-OH (1a–f, 1 mmol) in water-free DMF (10 mL). At  $-10$  °C, DIC (1.2 mmol) was added, and after 5 min stirring, L-Arg-NHBzl (10, 1 mmol) was added. After stirring for 36 h at 4 °C, the solvent was removed and the crude product chromatographed on neutral alumina using  $\text{CH}_3\text{OH}:\text{CHCl}_3$  (1:9) to afford *N*- $\alpha$ -Boc-L-His(2-alkyl)-L-Arg-NHBzl (11a–f).

***N*- $\alpha$ -Boc-L-His(2-isopropyl)-L-Arg-NHBzl (11a).** Yield: 58%, colorless solid; mp 182–183 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.18 (d, 6H,  $J = 7.0$  Hz), 1.37 (s, 9H), 1.63 (m, 2H), 1.84 (m, 2H), 2.67 (t, 2H,  $J = 7.1$  Hz), 3.03 (m, 2H), 3.29 (m, 1H), 4.46 (s, 2H), 4.68 (t, 1H,  $J = 6.7$  Hz), 4.93 (t, 1H,  $J = 6.9$  Hz), 6.90 (s, 1H), 7.26–7.31 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  21.4, 23.11, 28.08, 29.5, 29.84, 30.12, 41.89, 44.81, 54.27, 58.1, 75.7, 116.06, 124.88, 126.12, 127.8, 128.95, 129.26, 131.83, 139.53, 151.37, 157.99, 172.92. MS (APCI):  $m/z$  543.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His(2-tert-butyl)-L-Arg-NHBzl (11b).** Yield: 53%, colorless solid; mp 197–200 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.35 (s, 9H), 1.46 (s, 9H), 1.70 (m, 2H), 1.84 (m, 2H), 2.71 (t, 2H,  $J = 6.9$  Hz), 3.04 (m, 2H), 4.34 (s, 2H), 4.51 (t, 1H,  $J = 6.7$  Hz), 4.85 (t, 1H,  $J = 6.8$  Hz), 6.78 (s, 1H), 7.24–7.27 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  23.78, 28.72, 29.36, 29.79, 40.12, 43.92, 55.7, 58.86, 73.2, 119.61, 125.11, 126.6, 127.44, 128.72, 137.85, 148.17, 154.31, 158.63, 174.8. MS (APCI):  $m/z$  557.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His(2-cyclobutyl)-L-Arg-NHBzl (11c).** Yield: 50%, colorless solid; mp 213–215 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.34 (s, 9H), 1.57 (m, 2H), 1.87 (m, 2H), 1.93 (m, 2H), 2.14 (m, 4H), 2.56 (t, 2H,  $J = 7.1$  Hz), 3.12 (m, 2H), 3.27 (m, 1H), 4.37 (s, 2H), 4.58 (t, 1H,  $J = 6.9$  Hz), 4.80 (t, 1H,  $J = 6.8$  Hz), 6.94 (s, 1H), 7.27–7.32 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  19.57, 24.73, 26.2, 27.84, 28.3, 29.45, 31.92, 35.56, 40.9, 44.6, 56.76, 59.44, 78.3, 119.52, 125.4, 126.55, 127.78, 129.6, 136.48, 146.2, 155.0, 158.28, 178.4. MS (APCI):  $m/z$  555.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His(2-cyclohexyl)-L-Arg-NHBzl (11d).** Yield: 52%, colorless solid; mp 220 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.33–1.36 (m, 6H), 1.40 (s, 9H), 1.73 (m, 2H), 1.80 (m, 4H), 1.92 (m, 2H), 2.67 (t, 2H,  $J = 7.2$  Hz), 3.03 (m, 2H), 4.39 (s, 2H), 4.55 (t, 1H,  $J = 6.9$  Hz), 4.81 (t, 1H,  $J = 6.8$  Hz), 6.68 (s, 1H), 7.28–7.31 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  25.57, 26.42, 26.66, 28.10, 29.86, 30.14, 32.47, 38.60, 41.29, 43.47, 53.76, 56.72, 72.13, 119.80, 126.18, 127.61, 128.93, 130.16, 137.71, 147.74, 154.62, 158.63, 177.52. MS (APCI):  $m/z$  583.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His[2-(adamantan-1-yl)]-L-Arg-NHBzl (11e).** Yield: 45%, colorless solid; mp 268–270 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.31 (s, 9H), 1.58–1.62 (m, 12H), 1.72 (m, 2H), 1.85 (m, 2H), 2.01 (m, 3H), 2.59 (t, 2H,  $J = 7.0$  Hz), 3.08 (m, 2H), 4.30 (s, 2H), 4.54 (t, 1H,  $J = 6.8$  Hz), 4.86 (t, 1H,  $J = 6.7$  Hz), 6.67 (s, 1H), 7.24–7.28



(m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  24.12, 27.81, 28.3, 29.66, 30.82, 32.67, 35.97, 40.94, 43.45, 44.68, 53.56, 59.45, 73.1, 119.38, 125.85, 126.34, 128.04, 137.91, 138.73, 146.51, 152.15, 157.8, 176.17; MS (APCI):  $m/z$  635.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His-L-Arg-NHBzl (11f).** Yield: 62%, colorless solid; mp 161–163 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.40 (s, 9H), 1.61 (m, 2H), 1.89 (m, 2H), 2.71 (t, 2H,  $J = 7.0$  Hz), 3.12 (m, 2H), 4.29 (s, 2H), 4.57 (t, 1H,  $J = 6.7$  Hz), 4.82 (t, 1H,  $J = 6.7$  Hz), 6.86 (s, 1H), 7.27–7.30 (m, 5H), 7.65 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  23.17, 28.08, 29.57, 30.62, 39.96, 43.52, 54.04, 59.83, 77.63, 119.42, 124.21, 124.82, 126.18, 126.92, 127.91, 131.7, 136.0, 139.8, 153.15, 157.82, 177.1. MS (APCI):  $m/z$  501.2  $[\text{MH}]^+$ .

**General Method for the Synthesis of L-His(2-alkyl)-L-Arg-NHBzl·2HCl (12a–f, Scheme 3).** A solution of *N*- $\alpha$ -Boc-L-His(2-alkyl)-L-Arg-NHBzl (11a–f, 1 mmol) in 3 N HCl in 1,4-dioxane (15 mL) was stirred at ambient temperature for 15 min. The solvent was evaporated under reduced pressure to produce L-His(2-alkyl)-L-Arg-NHBzl·2HCl (12a–f).

**L-His(2-isopropyl)-L-Arg-NHBzl·2HCl (12a).** Yield: 87%, light-yellowish semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.12 (d, 6H,  $J = 7.1$  Hz), 1.67 (m, 2H), 1.93 (m, 2H), 2.87 (t, 2H,  $J = 7.2$  Hz), 3.09 (m, 2H), 3.22 (m, 1H), 3.74 (t, 1H,  $J = 6.8$  Hz), 4.45 (m, 2H), 4.62 (t, 1H,  $J = 6.7$  Hz), 6.97 (s, 1H), 7.28–7.33 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  22.39, 24.54, 28.7, 29.91, 30.67, 39.96, 45.41, 55.7, 59.71, 119.52, 125.4, 126.88, 127.2, 127.92, 128.19, 132.07, 136.5, 138.74, 153.82, 176.89. MS (APCI):  $m/z$  443.9  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -18.6^\circ$  ( $c = 1.5$ ,  $\text{H}_2\text{O}$ ).

**L-His(2-*tert*-butyl)-L-Arg-NHBzl·2HCl (12b).** Yield: 89%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.43 (s, 9H), 1.79 (m, 2H), 1.86 (m, 2H), 2.77 (t, 2H,  $J = 7.1$  Hz), 3.14 (m, 2H), 3.85 (t, 1H,  $J = 6.8$  Hz), 4.45 (s, 2H), 4.57 (t, 1H,  $J = 6.7$  Hz), 6.91 (s, 1H), 7.30–7.35 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  24.08, 25.74, 29.17, 30.25, 33.93, 41.22, 42.72, 53.79, 58.24, 120.61, 126.21, 126.6, 127.32, 128.47, 133.35, 146.7, 159.46, 173.58. MS (APCI):  $m/z$  457.9  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -16.5^\circ$  ( $c = 1$ ,  $\text{H}_2\text{O}$ ).

**L-His(2-cyclobutyl)-L-Arg-NHBzl·2HCl (12c).** Yield: 93%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.64 (m, 2H), 1.82 (m, 2H), 1.98 (m, 2H), 2.23 (m, 4H), 2.68 (t, 2H,  $J = 7.0$  Hz), 3.05 (m, 2H), 3.31 (m, 1H), 3.72 (t, 1H,  $J = 6.7$  Hz), 4.33 (s, 2H), 4.51 (t, 1H,  $J = 6.8$  Hz), 6.78 (s, 1H), 7.25–7.28 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  20.31, 24.51, 25.32, 28.2, 31.92, 36.73, 41.82, 44.31, 54.47, 59.04, 122.24, 126.4, 126.9, 127.62, 128.71, 134.67, 147.22, 157.63, 174.72. MS (APCI):  $m/z$  455.7  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -11.7^\circ$  ( $c = 2.0$ ,  $\text{H}_2\text{O}$ ).

**L-His(2-cyclohexyl)-L-Arg-NHBzl·2HCl (12d).** Yield: 95%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.39–1.41 (m, 6H), 1.78 (m, 2H), 1.85 (m, 4H), 1.95 (m, 2H), 2.56 (t, 2H,  $J = 7.1$  Hz), 3.10 (m, 2H), 3.88 (t, 1H,  $J = 6.9$  Hz), 4.30 (s, 2H), 4.47 (t, 1H,  $J = 6.7$  Hz), 6.87 (s, 1H), 7.29–7.34 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  24.1, 25.2, 26.42, 28.32, 31.78, 33.5, 38.95, 41.67, 44.0, 55.27, 58.84, 121.52, 126.1, 126.78, 128.62, 128.9, 130.16, 138.3, 145.42, 159.36, 175.5. MS (APCI):  $m/z$  483.8  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -12.2^\circ$  ( $c = 1.8$ ,  $\text{H}_2\text{O}$ ).

**L-His[2-(adamantan-1-yl)]-L-Arg-NHBzl·2HCl (12e).** Yield: 90%, colorless semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.63–1.68 (m, 12H), 1.76 (m, 2H), 1.90 (m, 2H), 2.06 (m, 3H), 2.66 (t, 2H,  $J = 7.2$  Hz), 3.18 (m, 2H), 3.98 (t, 1H,  $J = 6.8$  Hz), 4.44 (s, 2H), 4.61 (t, 1H,  $J = 6.7$  Hz), 6.79 (s, 1H), 7.28–7.31 (m, 5H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  23.42, 28.47, 28.93, 30.5, 32.82, 36.32, 41.24, 43.32, 44.06, 52.77, 58.61, 120.43, 126.19, 126.8, 128.41, 138.16, 138.3, 147.66, 156.13, 173.4. MS (APCI):  $m/z$  535.9  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -15.8^\circ$  ( $c = 1.4$ ,  $\text{H}_2\text{O}$ ).

**L-His-L-Arg-NHBzl·2HCl (12f).** Yield: 95%, light-yellowish semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.59 (m, 2H), 1.86 (m, 2H), 2.79 (t, 2H,  $J = 7.2$  Hz), 3.05 (m, 2H), 3.81 (t, 1H,  $J = 6.8$  Hz), 4.36 (s, 2H), 4.51 (t, 1H,  $J = 6.7$  Hz), 6.87 (s, 1H), 7.29–7.34 (m, 5H), 7.55 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  24.78, 29.23, 30.12, 40.56, 44.18, 53.67, 61.20, 116.28, 123.85, 124.34, 124.78, 125.82, 126.52, 128.11, 130.72, 135.25, 138.13, 155.27, 178.91. MS (APCI):  $m/z$  401.5  $[\text{MH}]^+$ .  $[\alpha]_{\text{D}}^{25} = -8.8^\circ$  ( $c = 1$ ,  $\text{H}_2\text{O}$ ). H

**General Method for the Synthesis of *N*- $\alpha$ -Boc-L-His(2-alkyl)-L-Arg-OMe (14a–f, Scheme 4).** HONB (1.2 mmol) was added to *N*- $\alpha$ -Boc-L-His(2-alkyl)-OH (1a–e, 1 mmol) in water-free DMF (10 mL). At  $-10^\circ\text{C}$ , DIC (1.2 mmol) was added, and after 5 min stirring, L-Arg-OMe (13, 1 mmol) was added. After stirring for 36 h at  $4^\circ\text{C}$ , the solvent was removed and the crude product chromatographed on neutral alumina using  $\text{CH}_3\text{OH}:\text{CHCl}_3$  (1:9) to afford *N*- $\alpha$ -Boc-L-His(2-alkyl)-L-Arg-OMe (14a–f).

***N*- $\alpha$ -Boc-L-His(2-isopropyl)-L-Arg-OMe (14a).** Yield: 61%, white solid; mp 167–168 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.23 (d, 6H,  $J = 7.0$  Hz), 1.37 (s, 9H), 1.55 (m, 2H), 1.81 (m, 2H), 2.45 (t, 2H,  $J = 6.7$  Hz), 3.12 (m, 2H), 3.25 (m, 1H), 3.68 (s, 3H), 3.91 (t, 1H,  $J = 6.8$  Hz), 4.36 (t, 1H,  $J = 6.5$  Hz), 6.99 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  22.80, 25.15, 26.15, 29.27, 31.82, 32.55, 45.29, 51.86, 52.71, 57.81, 83.76, 118.45, 128.19, 144.23, 157.96, 160.73, 176.64, 180.02. MS (APCI):  $m/z$  468.2  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His(2-*tert*-butyl)-L-Arg-OMe (14b).** Yield: 58%, colorless solid; mp 174 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.38 (s, 9H), 1.41 (s, 9H), 1.68 (m, 2H), 1.91 (m, 2H), 3.00 (t, 2H,  $J = 6.8$  Hz), 3.18 (m, 2H), 3.69 (s, 3H), 3.83 (t, 1H,  $J = 6.9$  Hz), 4.40 (t, 1H,  $J = 6.7$  Hz), 6.72 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  25.62, 28.29, 29.17, 29.51, 31.90, 33.24, 41.66, 52.52, 52.67, 54.16, 80.15, 118.02, 132.49, 156.69, 157.06, 158.08, 172.99, 176.37. MS (APCI):  $m/z$  482.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His(2-cyclobutyl)-L-Arg-OMe (14c).** Yield: 55%, colorless solid; mp 180–182 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.28 (m, 2H), 1.40 (s, 9H), 1.68 (m, 2H), 1.92 (m, 2H), 2.34 (m, 4H), 2.51 (t, 2H,  $J = 6.8$  Hz), 3.22 (m, 2H), 3.29 (m, 1H), 3.36 (s, 3H), 3.93 (t, 1H,  $J = 6.9$  Hz), 4.44 (t, 1H,  $J = 6.6$  Hz), 6.75 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  18.80, 25.62, 28.31, 28.88, 29.14, 30.25, 34.33, 41.68, 52.55, 54.56, 62.97, 78.39, 117.89, 132.82, 139.43, 155.47, 158.09, 173.03, 176.33. MS (APCI):  $m/z$  480.2  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His(2-cyclohexyl)-L-Arg-OMe (14d).** Yield: 55%, colorless solid; mp 187 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.30–1.32 (m, 6H), 1.39 (s, 9H), 1.72 (m, 2H), 1.82 (m, 4H), 1.93 (m, 2H), 2.51 (t, 2H,  $J = 7.0$  Hz), 2.83 (m, 1H), 3.20 (m, 2H), 3.75 (s, 3H), 3.85 (t, 1H,  $J = 6.8$  Hz), 4.41 (t, 1H,  $J = 6.6$  Hz), 7.04 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  18.94, 25.97, 26.18, 26.40, 28.64, 33.83, 36.87, 39.45, 41.38, 52.64, 53.13, 58.70, 119.53, 128.29, 140.48, 156.43, 158.09, 173.14, 174.99. MS (APCI):  $m/z$  508.4  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His[2-(adamantan-1-yl)]-L-Arg-OMe (14e).** Yield: 56%, colorless solid; mp 202–204 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.39 (s, 9H), 1.49–1.53 (m, 12H), 1.56 (m, 2H), 1.82 (m, 2H), 2.04 (m, 3H), 2.43 (t, 2H,  $J = 7.2$  Hz), 3.21 (m, 2H), 3.72 (s, 3H), 3.97 (t, 1H,  $J = 6.9$  Hz), 4.53 (t, 1H,  $J = 6.7$  Hz), 6.84 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  21.22, 24.30, 27.49, 28.40, 29.02, 30.15, 33.90, 38.28, 41.41, 46.30, 52.23, 56.95, 60.26, 120.81, 125.10, 128.11, 129.28, 141.62, 155.17, 158.08, 172.70, 177.52. MS (APCI):  $m/z$  560.3  $[\text{MH}]^+$ .

***N*- $\alpha$ -Boc-L-His-L-Arg-OMe (14f).** Yield: 65%, colorless solid; mp 153–155 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.40 (s, 9H), 1.48 (m, 2H), 1.67 (m, 2H), 2.51 (t, 2H,  $J = 6.9$  Hz), 3.11 (m, 2H), 3.72 (s, 3H), 3.84 (t, 1H,  $J = 6.9$  Hz), 4.41 (t, 1H,  $J = 6.5$  Hz), 6.78 (s, 1H), 7.08 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  25.71, 28.18, 28.97, 29.49, 41.51, 52.56, 53.26, 55.32, 80.34, 118.23, 132.83, 135.41, 156.99, 158.07, 171.08, 173.72. MS (APCI):  $m/z$  426.1  $[\text{MH}]^+$ .

**General Method for the Synthesis of L-His(2-alkyl)-L-Arg-OMe·2HCl (15a–f, Scheme 4).** A solution of *N*- $\alpha$ -Boc-L-His(2-alkyl)-L-Arg-OMe (14a–f, 1 mmol) in 3 N HCl in 1,4-dioxane (15 mL) was stirred at ambient temperature for 15 min. The solvent was evaporated under reduced pressure to produce L-His(2-alkyl)-L-Arg-OMe·2HCl (15a–f).

**L-His(2-isopropyl)-L-Arg-OMe·2HCl (15a).** Yield: 90%, light-yellowish semisolid.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.28 (d, 6H), 1.57 (m, 2H), 1.87 (m, 2H), 2.65 (t, 2H,  $J = 6.9$  Hz), 3.07 (m, 2H), 3.31 (m, 1H), 3.75 (s, 3H), 3.86 (t, 1H,  $J = 6.9$  Hz), 4.43 (t, 1H,  $J = 6.6$  Hz), 6.88 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  20.22, 25.15, 27.63, 28.09, 28.63, 41.41, 52.74, 53.36, 58.34, 118.67,

126.39, 148.2, 159.74, 170.48, 177.51. MS (APCI):  $m/z$  368.3 [MH]<sup>+</sup>. [ $\alpha$ ]<sub>D</sub><sup>25</sup> = -21.0° ( $c$  = 1.8, H<sub>2</sub>O).

**L-His(2-*tert*-butyl)-L-Arg-OMe·2HCl (15b).** Yield: 95%, light-yellowish semisolid. <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  1.51 (s, 9H), 1.75 (m, 2H), 1.98 (m, 2H), 2.72 (t, 2H,  $J$  = 6.9 Hz), 3.12 (m, 2H), 3.40 (s, 3H), 3.73 (t, 1H,  $J$  = 6.9 Hz), 4.47 (t, 1H,  $J$  = 6.6 Hz), 7.41 (s, 1H). <sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD):  $\delta$  24.29, 25.78, 28.21, 28.57, 33.55, 41.4, 53.07, 56.73, 118.79, 126.57, 141.23, 158.09, 174.01, 176.12. MS (APCI):  $m/z$  382.3 [MH]<sup>+</sup>. [ $\alpha$ ]<sub>D</sub><sup>25</sup> = -10.7° ( $c$  = 2.0, H<sub>2</sub>O).

**L-His(2-cyclobutyl)-L-Arg-OMe·2HCl (15c).** Yield: 86%, light-yellowish semisolid. <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  1.46 (m, 2H), 1.75 (m, 2H), 1.99 (m, 2H), 2.46 (m, 4H), 2.73 (t, 2H,  $J$  = 6.8 Hz), 3.34 (m, 2H), 3.43 (m, 1H), 3.74 (s, 3H), 3.97 (t, 1H,  $J$  = 6.7 Hz), 4.40 (t, 1H,  $J$  = 6.5 Hz), 7.55 (s, 1H). <sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD):  $\delta$  18.78, 25.87, 27.1, 28.06, 30.07, 35.17, 41.4, 52.47, 53.66, 58.32, 119.28, 127.75, 135.32, 158.07, 168.77, 173.14. MS (APCI):  $m/z$  380.3 [MH]<sup>+</sup>. [ $\alpha$ ]<sub>D</sub><sup>25</sup> = -9.0° ( $c$  = 1.0, H<sub>2</sub>O).

**L-His(2-cyclohexyl)-L-Arg-OMe·2HCl (15d).** Yield: 90%, light-yellowish semisolid. <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  1.46–1.49 (m, 6H), 1.77 (m, 2H), 1.83 (m, 4H), 1.92 (m, 2H), 2.59 (t, 2H,  $J$  = 7.0 Hz), 2.98 (m, 1H), 3.11 (m, 2H), 3.63 (s, 3H), 3.89 (t, 1H,  $J$  = 6.8 Hz), 4.44 (t, 1H,  $J$  = 6.6 Hz), 6.98 (s, 1H). <sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD):  $\delta$  24.86, 25.72, 26.56, 28.96, 30.63, 34.13, 37.84, 41.49, 53.44, 54.71, 58.81, 118.12, 128.93, 137.63, 157.88, 170.71, 176.7. MS (APCI):  $m/z$  408.3 [MH]<sup>+</sup>. [ $\alpha$ ]<sub>D</sub><sup>25</sup> = -15.6° ( $c$  = 1.1, H<sub>2</sub>O).

**L-His[2-(adamantan-1-yl)]-L-Arg-OMe·2HCl (15e).** Yield: 82%, light-yellowish semisolid. <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  1.57 (m, 2H), 1.66–1.69 (m, 12H), 1.82 (m, 2H), 2.08 (m, 3H), 2.54 (t, 2H,  $J$  = 7.0 Hz), 3.09 (m, 2H), 3.62 (s, 3H), 3.88 (t, 1H,  $J$  = 6.7 Hz), 4.48 (t, 1H,  $J$  = 6.9 Hz), 6.93 (s, 1H). <sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD):  $\delta$  24.34, 26.25, 27.71, 28.56, 30.89, 32.14, 37.67, 41.72, 43.61, 52.74, 55.31, 59.49, 119.81, 128.75, 139.67, 159.08, 171.17, 174.67. MS (APCI):  $m/z$  460.4 [MH]<sup>+</sup>. [ $\alpha$ ]<sub>D</sub><sup>25</sup> = -21.3° ( $c$  = 1.0, H<sub>2</sub>O).

**L-His-L-Arg-OMe·2HCl (15f).** Yield: 92%, light-yellowish semisolid. <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  1.66 (m, 2H), 1.93 (m, 2H), 2.71 (t, 2H,  $J$  = 7.1 Hz), 3.11 (m, 2H), 3.66 (s, 3H), 3.75 (t, 1H,  $J$  = 6.8 Hz), 4.38 (t, 1H,  $J$  = 6.6 Hz), 6.78 (s, 1H), 7.58 (s, 1H). <sup>13</sup>C NMR (75 MHz, CD<sub>3</sub>OD):  $\delta$  28.84, 29.42, 30.73, 45.12, 51.61, 52.75, 56.43, 123.24, 130.94, 138.92, 161.76, 171.88, 176.53. MS (APCI):  $m/z$  326.2 [MH]<sup>+</sup>. [ $\alpha$ ]<sub>D</sub><sup>25</sup> = -12.7° ( $c$  = 1.2, H<sub>2</sub>O).

**Antimicrobial Activity Determination.** The synthesized dipeptides as well as their Boc protected derivatives were evaluated for antibacterial and antifungal activities against three Gram-negative bacteria [*E. coli* (ATCC 35218), *P. aeruginosa* (ATCC 9027), and *K. pneumoniae* (ATCC 700603)], two Gram-positive bacteria [MRSA (DSM 50128509) and MRSE (DSM 50160384)], and one fungal strain [*C. albicans* (ATCC 10231)], which were obtained from the German Collection of Microorganisms and Cell Cultures (DSMZ). Minimum inhibition concentration (MIC), defined as the lowest test concentration that completely suppresses growth of microorganism, was measured using a modified broth microdilution method<sup>25</sup> whereby the assay was performed in sterile 96-well, flat-bottom polypropylene microtiter plates (Nunc, Denmark) with inocula of  $5 \times 10^5$  bacteria or fungi per mL. The yeast extract/peptone/dextrose (YPD) medium was used in the case of the fungal strain, whereas a modification of a reported medium, with addition of a buffer substance (MOPS) to stabilize the pH for the addition of acidic peptides, was used for the bacterial strains.<sup>26</sup> After aerobic incubation for 15–18 h at 37 °C, inhibition of bacterial and fungal growth was determined by measuring absorbance at 600 nm with a Fusion universal microplate analyzer (Perkin-Elmer, MA). Half-maximal inhibitory concentration (IC<sub>50</sub>) is expressed as the concentration that affords 50% inhibition of microbial growth. IC<sub>50</sub> values were obtained, in triplicate per assay, from nonlinear regression analysis of plots of percentage inhibition versus log [concentration of peptide].

**Cytotoxicity Experiments.** Murine L-929 fibroblasts were obtained from the German Collection of Microorganisms and Cell Cultures (DSMZ) and cultivated in the media recommended by the supplier at 37 °C and 10% CO<sub>2</sub>. MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide] was used to measure growth and viability of cells, which are capable of reducing it by dehydrogenases of the mitochondria to a violet formazan product. Serial dilutions of the test compounds (60  $\mu$ L) were added to 120  $\mu$ L aliquots of a cell suspension ( $5 \times 10^4$ /mL) in a 96-well microplate. Blank and solvent controls were incubated under identical conditions. MTT in phosphate buffered saline (PBS) (20  $\mu$ L) was added to a final concentration of 0.5 mg/mL after 5 days. The precipitate of formazan crystals was centrifuged after 2 h and the supernatant discarded. The precipitate was washed with 100  $\mu$ L of PBS and dissolved in 100  $\mu$ L of isopropyl alcohol containing 0.4% hydrochloric acid. The microplates were gently shaken for 20 min to ensure a complete dissolution of the formazan and finally measured at 595 nm using an ELISA plate reader. All studies were carried out in two parallel experiments; the percentage of viable cells was calculated as the mean with respect to the controls set to 100%.

**Acknowledgment.** Rohit K. Sharma thanks the Council of Scientific and Industrial Research (CSIR), New Delhi for the award of Senior Research Fellowship. The authors would like to thank Dr. Florenz Sasse, Helmholtz Centre for Infection Research for his support in carrying out the cytotoxicity experiments.

**Supporting Information Available:** Detailed experimental procedures and spectral data for the intermediates **2a–f**, **3a–f**, and **7a–e**. The HPLC analysis results of all final peptides. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- Walsh, C. Where will the new antibiotics come from?. *Nat. Rev. Microbiol.* **2003**, *1*, 65–70.
- Finch, R. Bacterial resistance—the clinical challenge. *Clin. Microbiol. Infect.* **2002**, *8*, 21–32.
- Shain, C. S. Telithromycin: the first of the ketolides. *Ann. Pharmacother.* **2002**, *36*, 452–464.
- Daptomycin (cubicin) for skin and soft tissue infections. *Med. Lett. Drugs Ther.* **2004**, *46*, 11–12.
- Papagianni, M. Ribosomally synthesized peptides with antimicrobial properties: biosynthesis, structure, function, and applications. *Biotechnol. Adv.* **2003**, *21*, 465–499.
- Hancock, R. E. W.; Scott, M. G. The role of antimicrobial peptides in animal defences. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 8856–8861.
- Tossi, A.; Sandri, L.; Giangaspero, A. Amphipatic, alpha-helical antimicrobial peptides. *Biopolymers* **2000**, *55*, 4–30.
- Huang, H. W. Action of antimicrobial peptides: two-state model. *Biochemistry* **2000**, *39*, 8347–8352.
- Hancock, R. E. W.; Lehrer, R. I. Cationic peptides: a new source of antibiotics. *Trends. Biotechnol.* **1998**, *16*, 82–88.
- Sengupta, D.; Leontiadou, H.; Mark, A. E.; Marrink, S. J. Toroidal pores formed by antimicrobial peptides show significant disorder. *Biochim. Biophys. Acta* **2008**, *1778*, 2308–2317.
- Reddy, K. V. R.; Yedery, R. D.; Aranha, C. Antimicrobial peptides: premisses and promises. *Int. J. Antimicrob. Agents* **2004**, *24*, 536–547.
- Yeaman, M. R.; Yount, N. Y. Mechanisms of Antimicrobial Peptide Action and Resistance. *Pharmacol. Rev.* **2003**, *55*, 27–55.
- Sundriyal, S.; Sharma, R. K.; Jain, R.; Bharatam, P. V. Minimum requirements of hydrophobic and hydrophilic features in cationic peptide antibiotics (CPAs): pharmacophore generation and validation with cationic steroid antibiotics (CSAs). *J. Mol. Model.* **2008**, *14*, 265–278.
- Kurogi, Y.; Guner, O. F. Pharmacophore modeling and three-dimensional database searching for drug design using catalyst. *Curr. Med. Chem.* **2001**, *8*, 1035–1055.

- (15) Strøm, M. B.; Haug, B. E.; Skar, M. L.; Stensen, W.; Stiberg, T.; Svendsen, J. S. The pharmacophore of short cationic antibacterial peptides. *J. Med. Chem.* **2003**, *46*, 1567–1570.
- (16) Haug, B. E.; M. L.; Stensen, W.; Stiberg, T.; Svendsen, J. S. Bulky nonproteinogenic amino acids permit the design of very small and effective cationic antibacterial peptides. *J. Med. Chem.* **2004**, *47*, 4159–4162.
- (17) Chan, D. I.; Prenner, E. J.; Vogel, H. J. Tryptophan- and arginine-rich antimicrobial peptides: structures and mechanisms of action. *Biochim. Biophys. Acta* **2006**, *1758*, 1184–1202.
- (18) Kacprzyk, L.; Rydengård, V.; Morgelin, M.; Davoudi, M.; Pasupuleti, M.; Malmsten, M.; Schmidtchen, A. Antimicrobial activity of histidine-rich peptides is dependent on acidic conditions. *Biochim. Biophys. Acta* **2007**, *1768*, 2667–2680.
- (19) Jain, R.; Cohen, L. A. Regiospecific alkylation of histidine and histamine at C-2. *Tetrahedron* **1997**, *53*, 2365–2370.
- (20) Jain, R.; Cohen, L. A.; King, M. M. Synthesis of novel ring-substituted histidines and histamines. *Tetrahedron* **1997**, *53*, 4539–4548.
- (21) Kaur, N.; Lu, X.; Gershengorn, G. C.; Jain, R. Thyrotropin-releasing hormone (TRH) analogues that exhibit selectivity to TRH receptor subtype 2. *J. Med. Chem.* **2005**, *48*, 6162–6165.
- (22) Minisci, F. Novel applications of free-radical reactions in preparative organic chemistry. *Synthesis* **1973**, 1–24.
- (23) Minisci, F.; Vismara, E.; Fontana, F. Recent developments of free-radical substitutions of heteroaromatic bases. *Heterocycles* **1989**, *28*, 489–519.
- (24) Sharma, R. K.; Jain, R. Unprecedented 1,1'-carbonyldiimidazole mediated amidation of unprotected  $\alpha$ -amino acids in water. *Synlett* **2007**, 603–606.
- (25) Saginur, R.; Melissa, S.; Ferris, W.; Aaron, S. D.; Chan, F.; Lee, C.; Ramotar, K. Multiple combination bactericidal testing of staphylococcal biofilms from implant-associated infections. *Antimicrob. Agents Chemother.* **2006**, *50*, 55–61.
- (26) Stepanovic, S.; Vukovic, D.; Dakic, I.; Savic, B.; Svabic-Vlahovic, M. A modified microtiter-plate test for quantification of staphylococcal biofilm formation. *J. Microbiol. Methods* **2000**, *40*, 175–179.