

Correction to 3'-Deoxy Phosphoramidate Dinucleosides as Improved Inhibitors of Hepatitis C Virus Subgenomic Replicon and NS5B Polymerase Activity [*Journal of Medicinal Chemistry* 2010, 53, 6608–6617 DOI: 10.1021/jm100102v].
 Stephane Priet, Ivan Zlatev, Ivan Barvik, Jr., Katrien Geerts, Pieter Leyssen, Johan Neyts, Helene Dutartre, Bruno Canard, Jean-Jacques Vasseur, François Morvan,* and Karine Alvarez*

Page 6612. Table 2 contains errors. The list of compounds and IC₅₀ values should be as follows.

Table 2. Except for Compound 1, Separation of Diastereoisomers Does Not Modify the Inhibition Potency of 3'-Deoxy Phosphoramidate Dinucleosides^a

| compd | IC ₅₀ (μM) |
|-------|-----------------------|
| 1-Rp | 72 ± 25 |
| 1-Sp | 7 ± 3 |
| 1'-Rp | 121 ± 11 |
| 1'-Sp | 57 ± 18 |
| 2-Rp | 28 ± 6 |
| 2-Sp | 37 ± 6 |
| 2'-Rp | 64 ± 19 |
| 2'-Sp | 42 ± 7 |
| 6-Rp | 9 ± 1 |
| 6-Sp | 13 ± 6 |
| 7-Rp | 26 ± 6 |
| 7-Sp | 16 ± 4 |
| 8-Rp | 20.3 ± 1 |
| 8-Sp | 18.5 ± 8 |

^a IC₅₀ values of separated diastereoisomers of neutral and negatively charged (1, 1', 2, 2', 6, 7, and 8) phosphoramidate dinucleosides. All data represent average values for at least three separated experiments.

DOI: 10.1021/jm200628q
 Published on Web 06/14/2011