

Fluorophosphonylated Nucleoside Derivatives as New Series of Thymidine Phosphorylase Multisubstrate Inhibitors


Sonia Amel Diab,[†] Coralie De Schutter,[†] Murielle Muzard,[‡] Richard Plantier-Royon,[‡] Emmanuel Pfund,[†] and Thierry Lequeux^{*†}

[†]Laboratoire de Chimie Moléculaire et Thioorganique, UMR CNRS 6507 and FR3038, ENSICAEN, Université de Caen Basse-Normandie, 6 Boulevard du Maréchal Juin, 14050 Caen Cedex, France

[‡]Institut de Chimie Moléculaire de Reims, UMR CNRS 6229, UFR des Sciences Exactes et Naturelles, BP 1039, 51687 Reims Cedex 2, France

S Supporting Information

ABSTRACT: The synthesis of new class of potential TPase inhibitors containing a difluoromethylphosphonate function as phosphate mimic is reported. This new series was prepared from a readily available fluorinated building block in few steps. Two series were evaluated as potential inhibitors: a linear series and a conformational constrained series. The activity of these multisubstrate inhibitors depends on the size of the spacer introduced between the pyrimidine ring and the phosphonate function. Best results were observed from propargylthymine and corresponding azides.



INTRODUCTION

Platelet-derived endothelial cell growth factor (PD-ECGF) is a protein involved in tumor angiogenesis. This protein at low abundance is overexpressed in many human solid tumors and might be an attractive cancer chemotherapy target for inhibition of tumor angiogenesis, subsequent tumor growth, and metastasis. Because of a structural and genetic similarity between the *Escherichia coli* thymidine phosphorylase (EC 2.4.2.4) and the human recombinant PD-ECGF, it has been suggested that PD-ECGF could be identified as the human thymidine phosphorylase.^{1,2} Thymidine phosphorylase (TPase) catalyzes the reversible phosphorolysis of pyrimidine 2'-deoxynucleosides (Figure 1). The




Figure 1. Thymidine phosphorylase EC 2.4.2.4.

main product of the conversion, the 2-deoxyribose-1-phosphate, is transformed into 2-deoxyribose (2dR), which stimulates endothelial cell migration and angiogenesis.³

An important effect on angiogenesis and apoptosis in tumors has been observed by using thymidine phosphorylase inhibitors.^{4,5} The known TPase inhibitors are commonly structurally close to modified nucleosides or acyclonucleosides (Figure 2), and some of them are well-known as antiviral agents.⁶ Among these inhibitors, the most efficient on the tumor

angiogenesis is a chloropyrrolidinyl uracil derivative called TPI (Figure 2).^{4a,5}

Multisubstrate inhibitors, simultaneously bound to the nucleoside and phosphate binding sites, are less efficient than TPI. Their moderate activities could be attributed to the low acidic character of the phosphonate function (pK_a (phosphonate) = 7.5–8). Indeed, high difference in activities between naturally occurring phosphates and phosphonates has been already reported for purine nucleoside phosphorylase and tyrosine kinase inhibitors.⁷ As difluoromethylphosphonates are the best surrogate mimic of naturally occurring phosphates,⁸ our work is focused on the synthesis and the evaluation of the inhibitory activities of new fluorinated multisubstrate compounds. Taking into account that the distance between the phosphate oxygen atom and the thymidine N¹ atom lies between 4 and 10 Å when the enzymatic substitution occurred,⁹ the present study deals with the synthesis of fluorinated phosphonates as multisubstrate inhibitors of TPase containing different spacers such as alkyl chains and heterocycles (Figure 3).

RESULTS AND DISCUSSION

Chemistry. In connection with our previous studies regarding the synthesis of fluorophosphonate derivatives,¹⁰ the preparation of aliphatic series 7–10 and 14–16 was realized from the readily available tosylate derivatives 1 and 2 (Scheme 1).¹¹

We recently reported that 6-chloropurine or N³-benzoylthymine alkylation with fluorophosphonylated tosylates 1 and 2 affording compounds 3, 4, and 12 proceeded in good yields and

Received: December 15, 2011

Published: February 28, 2012





Figure 2. Thymidine phosphorylase inhibitors.^{6a,b,e,f}




□ = alkyl chain, heterocycle

X = alkyl, H or Br

Figure 3. Multisubstrate inhibitors.

Scheme 1. Synthesis of Phosphonic Acids 7–10 and 14–16^a




^aReagents and conditions: (a) (i) *N*³-benzoylthymine, TMG, DMSO, rt, 15 h; (ii) MeNH₂, MeOH, rt, 15 h to give 3 (75%) and 4 (81%); (i) *N*³-benzoyluracil, TMG, DMSO, rt, 15 h; (ii) MeNH₂, MeOH, rt, 15 h to give 5 (78%); (b) NBS, AIBN, THF, 60 °C, 1.5 h (75%); (c) (i) TMSBr, CH₂Cl₂, 0 °C to rt, 72 h; (ii) MeOH, rt, 2 h to give 7 (67%), 8 (75%), 9 (70%), and 10 (74%); (d) 6-chloropurine, TMG, DMSO, rt, 15 h to give 12 (67%); 2-amino-6-chloropurine, TMG, DMSO, rt, 15 h to give 11 (60%) and 13 (68%); (e) (i) TMSBr, CH₂Cl₂, 0 °C to rt, 72 h; (ii) H₂O, rt, 16 h to give 14 (77%), 15 (75%), and 16 (64%).

excellent regioselectivities when the reaction was conducted in the presence of 1,1,3,3-tetramethylguanidine (TMG).¹¹ These coupling conditions were extended to other nucleic bases. Indeed, tosylates 1 and 2 were treated with 2-amino-6-chloropurine in the presence of TMG. After 15 h of stirring, corresponding alkylation products 11 and 13 were isolated in 60–68% yields. From *N*³-benzoyluracil and tosylate 2, the reaction was also very efficient and uracil derivative 5 was

obtained in 78% yield after deprotection of the benzoyl group. Bromination of the 5-position of uracil was realized from 5 to afford fluorophosphonylated 5-bromouracil 6 in 75% yield. Cleavage of phosphonic alkyl esters was carried out under standard conditions. Fluorinated phosphonates 3–6 were treated with TMSBr (5 equiv) followed by addition of methanol, leading to phosphonic acids 7–10 in 67–75% yields after precipitation. From 6-chloropurine derivatives 11–13, an additional step allowing the chlorine atom substitution by a hydroxyl group led to phosphonic acids 14–16 in 64–77% yields. Fluorophosphonylated acyclonucleosides 27–29, containing longer alkyl chain spacers (8–11 carbon atoms), were prepared from 17 (Scheme 2).^{10c}

Scheme 2. Synthesis of Phosphonic Acids 27–29^a




^aReagents and conditions: (a) (i) ^tBuLi, THF, –78 °C, 10 min; (ii) 1,8-dibromooctane, THF, 30 min at –78 °C, then 1 h at –10 °C to give 18 (52%); 1,9-dibromononane, THF, 30 min at –78 °C, then 1 h at –10 °C to give 19 (51%); 1,11-dibromoundecane, THF, 30 min at –78 °C, then 1 h at –10 °C to give 20 (50%); (b) *N*³-benzoylthymine, TMG, DMSO, rt, 15 h to give 21 (64%), 22 (77%), and 23 (76%); (c) MeNH₂, MeOH, rt, 18 h to give 24 (94%), 25 (76%), and 26 (90%); (d) (i) TMSBr, CH₂Cl₂, 0 °C to rt, 72 h; (ii) MeOH, rt, 2 h to give 27 (99%), 28 (88%), and 29 (97%).

Alkylation reactions proceeded smoothly when the carbanion was formed from 17. In this case, the anion trapped with 1,8-dibromooctane, 1,9-dibromononane, and 1,11-dibromoundecane afforded corresponding difluorophosphonylated bromoalkanes 18–20 in 50–52% yields. Products 18–20 were reacted with *N*³-benzoylthymine, and nucleotide analogues 24–26 were obtained after nucleobase deprotection. Deprotection of phosphonic esters with TMSBr/MeOH afforded analogues 27–29

isolated by precipitation in 88–99% yields. In this series, better yields were obtained when intermediates **21**–**23** were isolated prior to debenzoylation reaction. Surprisingly, when the alkylation step was conducted from an anion formed by deprotonation of the diethyl difluoromethylphosphonate by LDA, less than 10% of products were detected in the crude.¹²

In order to modify the conformation of the alkyl chain and the enzymatic activity, the introduction of a sulfur atom was realized (Scheme 3). The presence of a sulfur atom has already

Scheme 3. Synthesis of Sulfur-Containing Nucleotide Analogues **39–**42**^a**



^aReagents and conditions. (a) 2-mercaptoethanol, TMG, MeCN, rt, 15 h to give **30** (86%); 3-mercaptoopropanol, TMG, MeCN, rt, 15 h to give **31** (77%); 4-mercaptobutanol, TMG, MeCN, rt, 15 h to give **32** (89%); 6-mercaptohexanol, TMG, MeCN, rt, 15 h to give **33** (79%); (b) (i) *N*³-benzoylthymine, PPh₃, DIAD, THF, rt, 15 h; (ii) MeNH₂, MeOH, rt, 18 h to give **35** (64%), **36** (55%), and **37** (64%); (c) (i) TMSBr, CH₂Cl₂, rt, 72 h; (ii) MeOH, rt, 2 h to give **39** (50%), **40** (86%), **41** (94%), and **42** (83%); (d) TsCl, NEt₃, CH₂Cl₂, rt, 15 h (79%); (e) (i) *N*³-benzoylthymine, TMG, rt, 15 h; (ii) MeNH₂, MeOH, rt, 15 h (59%).


been applied with success to improve the activity of PNP inhibitors.¹³

Sulfur-containing spacers of variable length were prepared by alkylation of thiols using tosylate **2**. This latter was treated with 2 equiv of thiol (2-mercaptoethanol, 3-mercaptoopropanol, 4-mercaptobutanol, or 6-mercaptohexanol) in the presence of 1.5 equiv of TMG. After 15 h of stirring in acetonitrile at room temperature, corresponding fluorophosphonylated sulfanyl alcohols **30**–**33** were obtained in 77–89% yields. *N*³-Benzoylthymine was introduced after activation of alcohols **30**–**33** from the corresponding tosylate or by using the Mitsunobu conditions. From fluorinated hydroxyphosphonate **33**, corresponding tosylate **34** was formed in 79% yield, while tosylation of **30**–**32** was unsuccessful. Alternatively, alcohols **30**–**32** were treated with *N*³-benzoylthymine in the presence of PPh₃ and DIAD to afford corresponding thymine derivatives **35**–**37** in 55–64% yields after deprotection. In this later case, no *O*-alkylation product was detected in the crude. Indeed, the observed chemical shift was in agreement with the formation of a methylene–nitrogen bond when compared to the starting alcohols **30**–**32** (¹³C NMR analysis). Furthermore, tosylate **34** was easily transformed into its corresponding alkylated thymine **38**. Finally, deprotection of phosphonate esters **35**–**38** with

TMSBr followed by methanolysis and precipitation yielded the targeted phosphonic acids **39**–**42** with good efficiency.

Functionalization of the spacer by introduction of an iodine atom was explored from iododifluoromethylphosphonate **43** and alkenes.¹⁴ Free radical addition onto allylthymine was realized (Scheme 4).

Scheme 4. Synthesis of Phosphonic Acid **45^a**




^aReagents and conditions. (a) (i) ^tBuLi, THF, –78 °C, 5 min; (ii) I₂, THF, –78 °C, 1 h (81%). (b) Method A: allylthymine, dilauroyl peroxide, C₂H₄Cl₂, 80 °C, 3 h (71%). Method B: allylthymine, Na₂S₂O₄, NaHCO₃, MeCN, H₂O, rt, 20 h (17%). Method C: allylthymine, Et₃B, CH₂Cl₂, rt, 3h (58%). (c) (i) TMSBr, CH₂Cl₂, rt, 72 h; (ii) MeOH, rt, 2 h (76%).

Iododifluoromethylphosphonate **43** was reacted with allylthymine in the presence of three different initiators (sodium dithionite, triethylborane, dilauroyl peroxide). Best results were observed in the presence of 0.3 equiv of dilauroyl peroxide. After 3 h of stirring under refluxed dichloroethane, fluorinated iodophosphonate **44** was isolated in 71% yield. It is worthy of note that no protection of the nucleobase is required for the reaction to proceed. After deprotection the corresponding phosphonic acid **45** was obtained in 76% yield.


Finally, phosphonic acids **56**–**62** containing a triazolyl heterocycle were synthesized from fluorophosphonylated azides **46**–**48** and propargyl nucleic bases (Scheme 5).¹⁵ Azido compounds **46**–**48** were easily obtained from their corresponding tosylates and sodium azide, as previously reported.¹⁵ They were reacted with propargyl nucleic bases (1.1 equiv) in the presence of sodium ascorbate (10 mol %) and copper sulfate (5 mol %). From propargylthymine, uracil, 6-chloropurine, and 2-amino-6-chloropurine, corresponding triazolyl derivatives **49**–**55** were isolated in 71–96% yields after 24 h of stirring at room temperature in ^tBuOH/H₂O (1/1). Phosphonic ester hydrolysis and chlorine substitution by a hydroxyl group, carried out with TMSBr followed by hydrolysis steps with MeOH and/or H₂O, afforded pure phosphonic acids **56**–**62** in 34–87% yields after purification by precipitation.

To confirm the effect of the fluorine atoms in the series of targeted acyclic nucleotides, phosphonic acid **65**, as non-fluorinated analogue of **7**, was also prepared. The synthesis was performed in three steps starting from commercially available 1,4-dibromobutane as depicted (Scheme 6).

Biological Studies. The magnitude of the size of the spacer was evaluated for compound **49** by X-ray analysis¹⁶ and compared to the value obtained by molecular modeling. In this case, similar values were observed by both methods (Figure 4), and the molecular modeling approach was extended to all the prepared compounds to correlate the enzymatic inhibition percentage with the phosphorus–nitrogen atoms distance. As expected, the different spacers introduced in the series permitted

Scheme 5. Synthesis of Phosphonic Acids 56–62^a

^aReagents and conditions. (a) propargylthymine, $CuSO_4$, sodium ascorbate, $tBuOH, H_2O$, rt, 24 h to give **49** (71%), **50** (96%), and **51** (93%); propargyluracil, $CuSO_4$, sodium ascorbate, $tBuOH, H_2O$, rt, 24 h to give **52** (85%); propargyl-6-chloropurine, $CuSO_4$, sodium ascorbate, $tBuOH, H_2O$, rt, 24 h to give **53** (85%) and **54** (82%); propargyl-2-amino-6-chloropurine, $CuSO_4$, sodium ascorbate, $tBuOH, H_2O$, rt, 24 h to give **55** (75%); (b) (i) $TMSBr, CH_2Cl_2$, rt, 72 h; (ii) $MeOH$, rt, 2 h to give **56** (34%), **57** (87%), **58** (73%), and **59** (75%); (c) (i) $TMSBr, CH_2Cl_2, 0^\circ C$ to rt, 72 h; (ii) $MeOH$, rt, 2 h; (iii) H_2O , rt, 16 h to give **60** (56%), **61** (63%), and **62** (81%).

Scheme 6. Synthesis of Phosphonic Acid 65^a

^aReagents and conditions: (a) $P(O^iPr)_3$, $130^\circ C$, 24 h (81%); (b) (i) N^3 -benzoylthymine, $TMG, DMSO$, rt, 16 h; (ii) $MeNH_2, MeOH$, rt, 16 h (92%); (c) (i) $TMSBr, CH_2Cl_2$, rt, 35 h; (ii) $MeOH$, rt, 2 h (88%).




Figure 4. (a) X-ray structure of difluorophosphonate **49**. (b) Computational modeling of **49**.

coverage of a large range of distance between 6.5 and 16.3 Å (Table 1). These distances should cover the two possible opened and closed conformations adopted by the enzyme.

Table 1. TP Inhibition Studies with Phosphonic Acids Containing a Linear Spacer


entry	compd	phosphorus–nitrogen distance (Å)	TP inhibition (%) at 1 mM	TP inhibition (%) at 100 μM
1	6ASBU		100 (100) ^{6d}	77 (71) ^{6d}
2	POT		76 (75) ^{6d}	15 (29) ^{6d}
3	7	6.60	51	4
4	8	7.38	68	15
5	9	7.38	44	15
6	10	7.38	61	20
7	14	7.81	25	4
8	15	7.82	25	9
9	16	7.82	74	34
10	27	12.57	85	55
11	28	13.84	86	55
12	29	16.33	88	56
13	39	10.23	36	0
14	40	11.72	90	43
15	41	7.82	78	40
16	42	15.22	86	38
17	45	5.36	95	52
18	65	6.56	23	0

Inhibitory activity of phosphonic acids **7–10**, **14–16**, **27–29**, **39–42**, **45**, **56–62**, and **65** against thymidine phosphor-ylase from *E. coli* was evaluated using the assay reported by Pérez-Pérez (Tables 1 and 2).^{6d} Typically, to a mixture of thymidine, orthophosphate, and TPase in Tris buffer was added phosphonic acid at two concentrations (1 mM, 100 μM). After 20 min of incubation at room temperature followed by 5 min at 90 °C, percentage of inhibition was determined by HPLC after separation of thymidine from thymine. Results were compared to those observed from 6-amino-5-bromouracil (6ASBU), a well-established TPase inhibitor, and 1-phosphonoctylthymine (POT), a moderate inhibitor.^{6d} Percentages of inhibition of phosphonic acids containing a linear spacer are listed in Table 1. As expected, introduction of fluorine atoms increased the activity

Table 2. TP Inhibition Studies with Conformational Constrained Phosphonic Acids

entry	compd	phosphorus–nitrogen distance (Å)	TP inhibition (%) at 1 mM	TP inhibition (%) at 100 μ M
1	6ASBU		100 (100) ^{6d}	77 (71) ^{6d}
2	POT		76 (75) ^{6d}	15 (29) ^{6d}
3	56	7.82	29	9
4	57	10.86	79	36
5	58	10.86	90	68
6	59	10.86	79	22
7	60	9.95	69	29
8	61	10.83	76	35
9	62	10.83	76	35

as already observed for the inhibition of PNP.^{7a} Indeed, compound 7 containing a short spacer was found to be twice more active at 1 mM than its corresponding nonfluorinated analogue 65 (entries 3 and 18). This difference in activity was confirmed when the inhibition percentages of compounds 27 and POT were compared. In addition, the substitution of the pyrimidine ring appeared essential to improve the inhibition of TPase, as observed for TP64 and POT (Figure 5). Unfortunately,

**Figure 5.** Influence of the presence of fluorine atoms and the pyrimidine ring onto the inhibition percentage of TPase at 100 μ M.^{6d}

the synthesis of the 6ASBU derivatives was attempted but not successful in our hands.

Surprisingly, phosphonic acid 45, containing an iodine atom in the same size spacer, inhibited TPase at 95% inhibition at 1 mM and 52% at 100 μ M (entry 17). The introduction of a supplementary halogen atom into the spacer had a remarkable positive effect on the inhibition compared to 7 (entry 3). This effect is difficult to rationalize; we presume that the iodine atom might occupy the pentose ring pocket of the natural substrate. This later case suggests that TPase would be inhibited in its closed active conformation.¹⁷

Substitution of thymine by uracil decreased the inhibitory effect, and compound 9 was consequently less active than 8 at 1 mM (entries 5 and 4). However, phosphonic acid 10 in which a bromine atom was introduced onto the 5-position of uracil displayed roughly the same activity as its corresponding thymine derivative 8 (entries 6 and 4), confirming the presence of lipophilic pocket around the nucleic base.^{6c} Substitution of thymine by a purine nucleic base such as hypoxanthine and guanine induced a decrease of the activity. Compounds 14 and 15 were less active than their thymine analogues 7 and 8 (entries 3, 4, 7, and 8). In contrast, only phosphonic acid 16 showed a marginal inhibitory effect of 74% at 1 mM (entry 9).

We also noticed that the activity increased with the size of the spacer. Nucleotide analogue 7 exhibited 50% inhibition at

1 mM, while 70% was observed for compound 8 and 85% for compound 27, both containing, respectively, one and seven additional carbon atoms in the spacer (entries 3, 4, and 10). This pattern was even more pronounced at 100 μ M (5% for 7, 15% for 8, and 55% for 27). In this series, results were found to be optimum with 27 and the activity remained unchanged when more than eight carbon atoms were introduced in the spacer. Indeed, compounds 28 and 29 displayed about 85% inhibition at 1 mM and 55% at 100 μ M (entries 10, 11, and 12). These results seem to indicate that thymidine phosphorylase is locked in its open, inactive conformation, as already demonstrated by Balzarini and co-workers in a previous study.^{6b}

Substitution of a methylene group in the spacer by a sulfur atom did not induce any change in the activity in almost every case. In fact, sulfur-containing phosphonic acids 40–42 showed the same activity as their corresponding analogues 27–29 (entries 10–12 and 14–16). Surprisingly, the difluorophosphonylated nucleoside 39, in which the sulfur atom and the thymine are separated by only two carbon atoms, was found to be inactive even at high concentration (entry 13).

PNP inhibition studies realized in the late 1990s revealed that the distance between the nitrogen atom of the nucleic base and the phosphorus atom of the difluoromethylphosphonate function was not the only important factor required for the design of potent inhibitors. In fact, it was also reported that introduction of a cycle (phenyl, cyclopropane, or THF) in the spacer allowed an increase in the activity.¹⁸ In addition, to explore if a click enzymatic reaction could be possible,¹⁹ the tolerance of a triazolyl ring was evaluated. This was confirmed by the difluorophosphonylated nucleoside analogues 56–60, containing a triazolyl ring (Table 2). The presence of a triazolyl ring is tolerated by the enzyme, and phosphonic acids 56–60 exhibited moderate to excellent activities at 1 mM (entries 3–9). Compound 56 with only one carbon atom between the difluorophosphonate group and the triazolyl ring displayed a weak inhibition activity (entry 3). This result indicates that the distance between the fluorinated phosphonate and the nucleic base is crucial and reinforces the idea of an inhibition of the enzyme in its opened conformation. Furthermore, we noticed a drastic improvement of the activity when the triazolyl ring was introduced onto a larger spacer. Indeed, nucleotide analogues 57 and 58 were found to be 3 times more active than 56 at 1 mM and up to 7 times at 100 μ M (entries 3–5). As a high difference in activity was observed from nonfluorinated (Table 1, POT, 65) and the fluorinated series (Table 1, 27, 7), the synthesis of nonfluorinated triazolyl derivatives was not explored. As previously discussed, substitution of the thymine nucleobase by uracil, hypoxanthine, or guanine had a negative effect. Derivative 58 presented 68% inhibition at 100 μ M, while compounds 59, 61, and 62 did not exceed 35% at the same concentration (entries 5, 6, 8, 9). The same pattern was observed with 57 and 60 (entries 4 and 7).

Difluorophosphonylated analogue 58 combining the triazolyl ring and a large size spacer appears to be the best inhibitor revealed by this study. Thus, its inhibition constant was evaluated ($K_i = 58 \mu$ M), and 58 acted as a competitive inhibitor (Figure 6). Note that the other competitive inhibitors TP64 and TP65, evaluated on *E. coli* TPase, presented 142 and 54 μ M respectively,^{6b} for a similar assay, showing again the superiority of the difluoromethylphosphonates as phosphate surrogates.




Figure 6. (A) Lineweaver–Burk plot and (B) replot of slope versus [58]: (■) 58, 100 μM ; (▲) 58, 50 μM ; (×) 58, 20 μM ; (◆) 58, 0 μM .

CONCLUSION

Novel fluorinated acyclic nucleotides have been synthesized to develop new multisubstrate inhibitors of *E. coli* thymidine phosphorylase as potent antiangiogenic factors. The first series contained a difluoromethylphosphonate group and a nucleic base that were separated by linear spacers from different size. A second series of inhibitors functionalized by a triazolyl ring was developed, and their biological activities were evaluated.

For both series, thymine derivatives were much more active than their analogues bearing uracil, guanine, and hypoxanthine nucleic base. It has been confirmed that difluoromethylphosphonate derivatives are more active than their nonfluorinated analogues (65 vs 7). However in the fluorinated series, compounds having a short spacer (6–8 Å) are less active except when the iodine atom was present on the alkyl chain (compound 45). In contrast, a long length spacer up to 16 Å dramatically enhanced the inhibition properties of the compounds. In this series the optimum activity for a chain length of 14–15 Å suggests an inhibition of the enzyme in its opened, inactive conformation, as previously observed in the literature. Substitution of a methylene group in the spacer by a sulfur atom induced no change of the activity. The presence of the triazolyl ring in the spacer is well tolerated by the enzyme, and phosphonic acids 56–60 exhibited moderate to good activities at 1 mM. The best inhibition was obtained with 58 (90% at 1 mM and 68% at 100 μM) that combined the triazolyl ring and a long spacer. Because of its large size, it is reasonable to assume that 58 inhibits the thymidine phosphorylase in its opened inactive conformation. Finally, these fluorinated acyclic nucleosides are the first examples of multisubstrate TPase inhibitors bearing a difluoromethylphosphonate group as phosphate mimic. These new series will be further exploited in order to improve the activity against thymidine phosphorylase to design new antiangiogenic agents after derivation into their corresponding prodrugs.^{61–k,20} In addition, their activity toward viruses will be evaluated and reported elsewhere.

EXPERIMENTAL SECTION

General. All commercially available reagents were bought from Aldrich and used as received. For anhydrous conditions, the glassware was dried in the oven at 120 °C and cooled to room temperature under a continuous nitrogen flow. THF, CH_2Cl_2 , Et_2O , and CH_3CN were dried in a solvent generator from Innovative Technologies Inc., which uses an activated alumina column to remove water. DMF and NEt_3 were distilled under CaH_2 or 4 Å molecular sieves. Flash column chromatography was realized on silica gel 60 (40–63 μm) from Merck

with air pressure, and products were detected by thin layer chromatography, in which the spots were visualized by UV irradiation and/or KMnO_4 solution. NMR spectra were recorded on a 250 or 400 MHz apparatus in deuterated solvent at 25 °C. ^{31}P and ^{19}F NMR spectral lines are with respect to the internal references H_3PO_4 (capillary) and CFCl_3 . All chemical shifts are reported in δ parts per million (ppm), and coupling constants are in hertz (Hz). High-resolution mass data were recorded on a high-resolution mass spectrometer in the EI or ESI mode. IR spectra were recorded on a Perkin-Elmer ATR IR instrument. Analytical HPLC was performed on a Waters systems (model 600 controller, model 717plus autosampler, model 996 photodiode array detector) using reverse phase Phenomenex Gemini C18 110A (5 μm , 4.6 mm \times 250 mm). The mobile phase was 70% methanol, 30% H_2O with 0.05% of TFA, and the flow rate was 1.0 mL/min. UV spectra were recorded using a Waters 996 photodiode array detector. Data were integrated and reported using Waters Empower software. All compounds submitted to enzymatic assays displayed purity of >95% as determined by this method, unless stated otherwise.

Diisopropyl 1,1-Difluoro-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pentylphosphonate (5). *Typical Procedure for the Preparation of Alkyl Spacer from Tosylate Derivatives.* To a stirred solution of tosylate 2 (442.1 mg, 1.0 mmol) in DMSO (5 mL), N^3 -protected pyrimidine (324 mg, 1.50 mmol) was added followed by dropwise addition of TMG (0.19 mL, 1.50 mmol). The mixture was stirred and monitored by TLC. After completion (15 h), DMSO was evaporated and the residue diluted in CH_2Cl_2 (15 mL), washed with water (5 mL) and brine (5 mL). The aqueous layers were extracted with CH_2Cl_2 (2 \times 10 mL) and the combined organic layers washed with brine (5 mL) and dried over MgSO_4 . Solvents were evaporated under reduced pressure. To a stirred solution of crude product (382.8 mg, 0.46 mmol) in MeOH (5 mL) was added 40% *N*-methylamine in H_2O (5 mL, 20.0 mmol), and the solution was stirred for 15 h at room temperature. The methanol was evaporated under reduced pressure. The residue was purified by flash column chromatography on silica using EtOAc–pentane (8/2) as eluent to give 5 as a light yellow syrup (298.6 mg, 78%). ^1H NMR (CDCl_3 , 400 MHz) δ 9.75 (s, 1H), 7.15 (d, J = 8.0 Hz, 1H), 5.66 (d, J = 8.0 Hz, 1H), 4.80 (dsept, J = 6.0 Hz, 2H), 3.71 (t, J = 6.8 Hz, 2H), 1.95–2.15 (m, 2H), 1.68–1.78 (m, 2H), 1.49–1.65 (m, 2H), 1.32 (dd, J = 6.1, 3.1 Hz, 12H); ^{19}F NMR (CDCl_3 , 376 MHz) δ -112.83 (dt, J = 108.5, 19.8 Hz, 2F); ^{31}P NMR (CDCl_3 , 162 MHz) δ 5.1 (t, J = 108.5 Hz, 1P); ^{13}C NMR (CDCl_3 , 63 MHz) δ 164.3, 151.1, 144.7, 121.6 (dt, J = 260.6, 218.1 Hz), 102.4, 73.9 (d, J = 7.1 Hz), 48.8, 33.6 (dt, J = 21.2, 15.1 Hz), 29.9, 24.3 (d, J = 3.4 Hz), 23.9 (d, J = 4.8 Hz), 18.13 (dt, J = 4.9 Hz). LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 383 (31), 341 (73), 299 (100). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{15}\text{H}_{26}\text{F}_2\text{N}_2\text{O}_5\text{P}$ 383.1547, found 383.1532.

Diisopropyl 1,1-Difluoro-5-[5-bromo-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl]pentylphosphonate (6). To a suspension of

5 (250.0 mg, 0.65 mmol) in dry THF (5 mL) were added *N*-bromosuccinimide (115.7 mg, 0.65 mmol) and AIBN (a pinch). The mixture was heated at 60 °C for 90 min. Then it was diluted with EtOH (10 mL) and filtered through Celite. The filtrate was evaporated and purified by flash column chromatography on silica using CH₂Cl₂–MeOH (20/1) as eluent to give **6** (224.8 mg, 75%) as a yellow syrup. ¹H NMR (CDCl₃, 400 MHz) δ 10.09 (s, 1H), 7.56 (s, 1H), 4.80 (2H, dsept, *J* = 6.0 Hz), 3.73 (2H, t, *J* = 7.2 Hz), 1.99–2.16 (m, 2H), 1.64–1.80 (m, 2H), 1.50–1.62 (m, 2H), 1.31 (dd, *J* = 6.2, 3.5 Hz, 12H); ¹⁹F NMR (CDCl₃, 376 MHz) δ –112.75 (dt, *J* = 108.6, 19.6 Hz, 2F); ³¹P NMR (CDCl₃, 162 MHz) δ 4.9 (t, *J* = 108.6 Hz, 1P); ¹³C NMR (CDCl₃, 101 MHz) δ 160.1, 150.6, 144.2, 120.4 (dt, *J* = 260.3, 218.4 Hz), 96.5, 74.0 (d, *J* = 7.1 Hz), 49.1, 33.5 (dt, *J* = 21.2, 14.7 Hz), 28.8, 24.1 (d, *J* = 3.5 Hz), 23.9 (d, *J* = 4.8 Hz), 18.1 (dt, *J* = 4.9 Hz). LRMS-ESI (*m/z*): [M + H]⁺ 461 (58), 419 (100), 377 (75), 229 (2). HRMS-ESI (*m/z*): [M + H]⁺ calcd for C₁₅H₂₅BrF₂N₂O₃P 461.0653, found 461.0633.

1,1-Difluoro-5-(6-hydroxy-9H-purin-9-yl)pentylphosphonic Acid (9). *General Procedure for the Hydrolysis of Difluorophosphonylated Pyrimidine Derivatives.* To a cooled solution of **5** (84.1 mg, 0.22 mmol) in anhydrous CH₂Cl₂ (5 mL) was added TMSBr (0.174 mL, 1.32 mmol). The mixture was stirred at room temperature for 72 h and then concentrated under vacuum. The residue was diluted in MeOH (2 mL) and stirred for 2 h at room temperature. The solvent was removed, and the crude was diluted in MeOH (0.5 mL). Compound **9** (45.9 mg, 70%) was obtained as a white powder after precipitation in Et₂O–CH₂Cl₂ (3/1): mp 193 °C; ¹H NMR (MeOD, 500 MHz) δ 7.41 (d, *J* = 9.5 Hz, 1H), 5.62 (d, *J* = 9.5 Hz, 1H), 3.79 (t, *J* = 9.0 Hz, 2H), 2.02–2.22 (m, 2H), 1.71–1.83 (m, 2H), 1.57–1.70 (m, 2H); ³¹P NMR (MeOD, 202 MHz) δ 5.7 (t, *J* = 110.1 Hz, 1P); ¹⁹F NMR (MeOD, 470 MHz) δ –118.65 (dt, *J* = 110.1, 23.5 Hz, 2F); ¹³C NMR (DMSO, 101 MHz) δ 163.9, 152.3, 146.8, 121.3 (dt, *J* = 260.8, 194.2 Hz), 106.5, 43.9, 31.2 (m), 23.5, 17.0 (m). LRMS-ESI (*m/z*): [M – H][–] 297 (90), 277 (6), 254 (100), 234 (8), 185 (32), 111 (36). HRMS-ESI (*m/z*): [M – H][–] calcd for C₉H₁₂F₂N₂O₃P 297.0452, found 297.0443.

Diisopropyl 4-(2-Amino-6-chloro-9H-purin-9-yl)-1,1-difluorobutylphosphonate (11). *General Procedure for the Introduction of Purine Nucleic Bases.* To a stirred solution of tosylate **1** (200.0 mg, 0.47 mmol) and 2-amino-6-chloropurine (119.5 mg, 0.71 mmol) in DMSO (5 mL) was added dropwise TMG (0.089 mL, 0.71 mmol). The mixture was stirred at 20 °C and monitored by TLC. After completion (15 h), DMSO was evaporated and the residue diluted in CH₂Cl₂ (15 mL), washed with water (5 mL), and brine (5 mL). The aqueous layer was extracted with CH₂Cl₂ (2 × 10 mL), and the combined organic layers were washed with brine (10 mL), dried over MgSO₄, and filtered. Solvents were evaporated under reduced pressure and crude product was purified by flash column chromatography on silica using ethyl acetate–pentane (7/3) as eluent to give **11** (120.1 mg, 60%) as a light yellow syrup. ¹H NMR (CDCl₃, 250 MHz) δ 7.78 (s, 1H), 5.28 (sbr, 2H), 4.83 (dsept, *J* = 6.3 Hz, 2H), 4.14 (t, *J* = 6.7 Hz, 2H), 2.02–2.40 (m, 4H), 1.33 (dd, *J* = 5.9, 3.3 Hz, 12H); ¹⁹F NMR (CDCl₃, 235 MHz) δ –112.26 (dt, *J* = 106.9, 19.0 Hz, 2F); ³¹P NMR (CDCl₃, 101 MHz) δ 4.8 (t, *J* = 106.9 Hz, 1P); ¹³C NMR (CDCl₃, 101 MHz) δ 160.1, 154.4, 153.0, 146.7, 131.9, 120.1 (dt, *J* = 259.8, 216.9 Hz), 73.8 (d, *J* = 7.0 Hz), 48.1, 32.6 (dt, *J* = 21.0, 14.9 Hz), 24.4 (d, *J* = 3.3 Hz), 23.9 (d, *J* = 4.7 Hz), 22.2 (dt, *J* = 5.0 Hz). LRMS-ESI (*m/z*): [M + H]⁺ 427 (62), 385 (100), 342 (23). HRMS-ESI (*m/z*): [M + H]⁺ calcd for C₁₅H₂₄ClF₂N₅O₃P 426.1273, found 426.1285.

1,1-Difluoro-4-(2-amino-6-hydroxy-9H-purin-9-yl)pentylphosphonic Acid (14).²¹ *General Procedure for the Hydrolysis of Difluorophosphonylated Purine Derivatives.* To a solution of **11** (93.7 mg, 0.22 mmol) in anhydrous CH₂Cl₂ (3 mL) cooled at 0 °C was added TMSBr (0.174 mL, 1.32 mmol). The mixture was stirred at room temperature for 72 h. Volatiles were removed, and the residue was stirred in H₂O (1 mL) for

16 h and dried under reduced pressure. The residue was diluted in MeOH, and compound **14** (54.75 mg, 77%) was obtained as a white solid after precipitation in Et₂O–CH₂Cl₂: mp 230 °C; ¹H NMR (D₂O, 400 MHz) δ 8.53 (s, 1H), 4.21 (t, *J* = 7.5 Hz, 2H), 1.89–2.22 (m, 4H); ¹⁹F NMR (D₂O, 376 MHz) δ –112.43 (dt, *J* = 112.3, 21.5 Hz, 2F); ³¹P NMR (D₂O, 162 MHz) δ 3.6 (t, *J* = 112.3 Hz, 1P); ¹³C NMR (D₂O, 101 MHz) δ 166.2, 163.5, 148.6, 141.2, 128.3, 122.4 (m), 46.0, 29.9 (m), 23.9 (m). LRMS-ESI (*m/z*): [M – H][–] 322 (100), 279 (25), 222 (10). HRMS-ESI (*m/z*): [M – H][–] calcd for C₉H₁₁F₂N₅O₄P 322.0517, found 322.0512.

Diisopropyl 1,1-Difluoro-9-bromononylphosphonate (18). *General Procedure for the Preparation of Diisopropyl 1,1-Difluorobromoalkylphosphonate.* To a solution of *tert*-butyllithium (2.17 mL 1 M in pentane, 3.47 mmol, 1.3 equiv) in anhydrous THF (40 mL) at –78 °C, difluoromethylphosphonate **17** (700 mg, 2.67 mmol, 1 equiv) was added dropwise. The mixture was stirred for 10 min at –78 °C, and 1,8-dibromooctane (0.98 mL, 5.34 mmol, 2 equiv) was added slowly. The mixture was stirred for 30 min at –78 °C and warmed from –78 to –10 °C over 1 h. The reaction mixture was quenched by addition of saturated NH₄Cl_{aq} (4 mL), and the mixture was extracted with CH₂Cl₂ (10 mL), washed with saturated aqueous solution of NaHCO₃, dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography using EtOAc/pentane (1/9 and 2/8) as eluent to afford compound **18** (565 mg, 52%) as a colorless oil. ¹H NMR (CDCl₃, 400 MHz) δ 4.84 (dsept, *J* = 6.8 Hz, 2H), 3.40 (t, *J* = 6.8 Hz, 2H), 2.10–1.95 (m, 2H), 1.85 (quint, *J* = 6.9 Hz, 2H), 1.60–1.38 (m, 4H), 1.37 (dd, *J* = 6.1, 4.1 Hz, 12H), 1.35–1.30 (m, 6H); ¹⁹F NMR (CDCl₃, 376 MHz) δ –112.83 (dt, *J* = 20.0, 109.7 Hz, 2F); ³¹P NMR (CDCl₃, 161 MHz) δ 5.7 (t, *J* = 109.7 Hz, 1P); ¹³C NMR (CDCl₃, 101 MHz) δ 120.6 (dt, *J* = 260.0, 217.1 Hz), 73.3 (d, *J* = 7.1 Hz), 33.8 (dt, *J* = 20.9, 14.2 Hz), 33.9, 32.7, 29.1, 29.0, 28.5, 28.0, 24.1 (d, *J* = 3.5 Hz), 23.7 (d, *J* = 4.9 Hz), 20.5 (dd, *J* = 4.7 Hz). LRMS-ESI (*m/z*): [M + H]⁺ 407 (30), 365 (62), 323 (100). HRMS-ESI (*m/z*): [M + H]⁺ calcd for C₁₅H₃₁O₃F₂PBr 407.1162, found 407.1161.

Diisopropyl 1,1-Difluoro-9-(5-methyl-2,4-dihydropyrimidin-1,3(2H)-yl)nonylphosphonate (21). *General Procedure for Thymine Introduction from Bromoalkyl Derivatives.* To a solution of **18** (285 mg, 0.70 mmol) and *N*³-benzoylthymine (320 mg, 1.39 mmol) in CH₃CN (6 mL) and DMSO (0.5 mL), was added dropwise TMG (0.191 mL, 1.53 mmol). After 15 h of stirring at 20 °C the mixture was poured in Et₂O (10 mL). The organic layer was washed with water (2 × 2 mL), brine (5 mL), dried over MgSO₄, filtered, and concentrated under reduced pressure. Compound **21** (249 mg, 64%) was isolated as colorless oil by flash column chromatography using EtOAc/pentane (6/4) as eluent. ¹H NMR (CDCl₃, 400 MHz) δ 7.89 (dd, *J* = 8.0, 0.8 Hz, 2H), 7.62 (dd, *J* = 7.4 Hz, 1H), 7.47 (dd, *J* = 7.7 Hz, 2H), 7.09 (s, 1H), 4.82 (dsept, *J* = 6.4 Hz, 2H), 3.70 (t, *J* = 7.3 Hz, 2H), 2.20–1.99 (m, 2H), 1.94 (s, 3H), 1.70–1.50 (m, 4H), 1.35 (dd, *J* = 6.1, 4.5 Hz, 12H), 1.35–1.30 (m, 8H); ¹⁹F NMR (CDCl₃, 376 MHz) δ –112.79 (dt, *J* = 20.0, 109.6 Hz, 2F); ³¹P NMR (CDCl₃, 162 MHz) δ 5.6 (t, *J* = 109.6 Hz, 1P); ¹³C NMR (CDCl₃, 101 MHz) δ 169.2, 163.2, 149.8, 140.2, 134.9, 131.7, 130.4, 129.1, 120.6 (dt, *J* = 259.0, 217.1 Hz), 110.6, 73.93 (d, *J* = 7.1 Hz), 48.8, 33.8 (dt, *J* = 20.9, 14.3 Hz), 29.2, 29.1, 29.0, 28.9, 26.3, 24.1 (d, *J* = 3.5 Hz), 23.7 (d, *J* = 4.8 Hz), 20.5 (dd, *J* = 4.6 Hz, C₃), 12.4. LRMS-ESI (*m/z*): [M + H]⁺ 557 (61), 515 (64), 473 (100), 393 (8), 351 (16). HRMS-ESI (*m/z*): [M + H]⁺ calcd for C₂₇H₄₀N₂O₆F₂P 557.2592, found 557.2607.

Diisopropyl 1,1-Difluoro-9-(5-methyl-2,4-dihydropyrimidin-1,3(2H)-yl)nonylphosphonate (24). *General Procedure for the Deprotection of *N*³-Benzoylthymine.* To a stirred solution of **21** (256 mg, 0.46 mmol) in MeOH (5 mL) was added *N*-methylamine (5 mL, 40% aqueous solution). After 18 h, the mixture was concentrated under reduced pressure. Flash column chromatography using EtOAc/pentane (7/3) afforded **24** (195 mg,

94%) as a colorless oil. ^1H NMR (CDCl_3 , 400 MHz) δ 9.77 (sbr, 1H), 6.94 (d, $J = 1.2$ Hz, 1H), 4.77 (dsept, $J = 6.2$ Hz, 2H), 3.62 (t, $J = 7.4$ Hz, 2H), 2.05–1.92 (m, 2H), 1.85 (d, $J = 1.1$ Hz, 3H), 1.65–1.50 (m, 4H), 1.30 (dd, $J = 6.2$, 4.6 Hz, 12H), 1.27–1.20 (m, 8H); ^{19}F NMR (CDCl_3 , 376 Mz) δ –112.81 (dt, $J = 20.0$, 109.7 Hz, 2F); ^{31}P NMR (CDCl_3 , 162 MHz) δ 5.6 (t, $J = 109.7$ Hz, 1P); ^{13}C NMR (CDCl_3 , 101 MHz) δ 164.7, 151.1, 140.4, 120.6 (dt, $J = 259.0$, 217.0 Hz), 110.5, 73.3 (d, $J = 7.1$ Hz), 48.4, 33.7 (dt, $J = 20.8$, 14.2 Hz), 29.1, 29.1, 29.0, 28.9, 26.3, 24.1 (d, $J = 3.4$ Hz), 23.7 (d, $J = 4.9$ Hz), 20.5 (dd, $J = 4.6$ Hz), 12.3. LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 453 (92), 411 (100), 369 (51). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{36}\text{N}_2\text{O}_5\text{F}_2\text{P}$ 453.2330, found 453.2333.

1,1-Difluoro-9-(5-methyl-2,4-dihydropyrimidin-1,3(2H)-yl)-nonylphosphonic Acid (27). Following the general procedure used for **9** from difluorophosphonate **24** (99 mg, 0.22 mmol), TMSBr (0.174 mL, 1.32 mmol), and CH_2Cl_2 (5 mL), compound **27** was isolated as a white solid (80 mg, 99%): mp 175–180 °C; ^1H NMR (DMSO, 400 MHz) δ 11.19 (s, 1H), 7.53 (d, $J = 1.2$ Hz, 1H), 3.60 (t, $J = 7.2$ Hz, 2H), 2.00–1.85 (m, 2H), 1.75 (d, $J = 1.0$ Hz, 3H), 1.60–1.40 (m, 4H), 1.30–1.20 (m, 8H); ^{19}F NMR (DMSO, 376 Mz) δ –112.41 (dt, $J = 20.1$, 101.1 Hz, 2F); ^{31}P NMR (DMSO, 162 MHz) δ 4.2 (t, $J = 101.1$ Hz, 1P); ^{13}C NMR (DMSO, 101 MHz) δ 164.4, 151.0, 141.6, 121.5 (dt, $J = 257.9$, 204.5 Hz), 108.6, 48.7, 47.3, 33.5 (dt, $J = 21.0$, 14.2 Hz), 28.8, 28.6, 25.9, 20.6 (dt, $J = 4.4$ Hz), 12.0. LRMS-ESI (m/z) $[\text{M} - \text{H}]^-$ 367 (7), 347 (10), 324 (100), 304 (15). HRMS-ESI (m/z): $[\text{M} - \text{H}]^-$ calcd for $\text{C}_{14}\text{H}_{22}\text{N}_2\text{O}_5\text{F}_2\text{P}$ 367.1234, found 367.1231.

Diisopropyl 1,1-Difluoro-5-(2-hydroxy-1-ethylsulfanyl)-pentylphosphonate (30). *General Procedure for the Preparation of Difluorophosphonylated Thioethers.* To a solution of **2** (845 mg, 1.91 mmol) and 2-mercaptoethanol (0.270 μL , 3.82 mmol) in anhydrous CH_3CN (10 mL) was added TMG (0.350 μL , 2.86 mmol). The mixture was stirred 15 h at room temperature. The solution was diluted in Et_2O (10 mL) and washed with water (2×5 mL). The aqueous layers were combined and extracted with CH_2Cl_2 (10 mL). The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and concentrated under reduced pressure. The crude product was purified by flash column chromatography using EtOAc/pentane (7/3) as eluent to give **30** (572 mg, 86%) as a colorless oil; ^1H NMR (CDCl_3 , 400 MHz) δ 4.81 (dsept, $J = 6.4$ Hz, 2H), 3.69 (t, $J = 6.0$ Hz, 2H), 2.70 (t, $J = 6.0$ Hz, 2H), 2.52 (t, $J = 6.8$ Hz, 2H), 2.10–1.95 (m, 2H), 1.70–1.60 (m, 4H), 1.35 (dd, $J = 6.0$, 3.5 Hz, 12H); ^{19}F NMR (CDCl_3 , 376 Mz) δ –112.70 (dt, $J = 19.7$, 108.8 Hz, 2F); ^{31}P NMR (CDCl_3 , 161 Mz) δ 5.4 (t, $J = 108.8$ Hz, 1P); ^{13}C NMR (CDCl_3 , 101 MHz) δ 120.4 (dt, $J = 259.3$, 217.4 Hz), 73.6 (d, $J = 7.1$ Hz), 60.4, 35.2, 33.4 (dt, $J = 35.6$, 14.5 Hz), 31.3, 29.3, 24.1 (d, $J = 3.5$ Hz), 23.7 (d, $J = 4.9$ Hz), 20.0 (dd, $J = 4.9$ Hz). LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 349 (15), 331 (100), 289 (38), 271 (2), 247 (10), 229 (2). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{13}\text{H}_{28}\text{O}_4\text{F}_2\text{PS}$ 349.1414, found 349.1412.

Diisopropyl 1,1-Difluoro-5-(2-(3-benzoyl-5-methyl-2,4-dihydropyrimidin-1(2H)-yl)-1-ethanesulfure)pentylphosphonate (35). *General Procedure for Thymine Introduction from Fluorinated Hydroxyphosphonates.* Alcohol **30** (570 mg, 1.64 mmol), N^3 -benzoylthymine (453 mg, 1.97 mmol), and Ph_3P (515 mg, 1.97 mmol) in anhydrous THF (12 mL) were cooled at 0 °C. DIAD (0.485 μL , 2.46 mmol) was introduced dropwise, and the reaction mixture was stirred for 15 h at room temperature. The solvent was evaporated under reduced pressure, and the crude was filtered through a pad of silica gel using EtOAc/pentane (6/4 and 7/3). After concentration, the crude product was diluted with MeOH (17 mL), and N -methylamine (17 mL, 40% aqueous solution) was added slowly. After 18 h, the mixture was concentrated under reduced pressure. Flash column chromatography using EtOAc/pentane (8/2) afforded **35** (479.1 mg, 64%) as a colorless oil; ^1H NMR (CDCl_3 , 400 MHz) δ 8.80 (s, 1H), 7.03 (d, $J = 1.2$ Hz, 1H), 4.84 (dsept, $J = 6.2$ Hz, 2H), 3.85

(t, $J = 6.8$ Hz, 2H), 2.80 (t, $J = 6.8$ Hz, 2H), 2.55 (t, $J = 6.9$ Hz, 2H), 2.10–1.97 (m, 2H), 1.92 (d, $J = 1.1$ Hz, 3H), 1.67–1.65 (m, 4H), 1.36 (dd, $J = 6.2$, 4.0 Hz, 12H); ^{19}F NMR (CDCl_3 , 376 Mz) δ –112.81 (dt, $J = 19.7$, 109.0 Hz, 2F); ^{31}P NMR (CDCl_3 , 162 MHz) δ 5.4 (t, $J = 109.0$ Hz, 1P); ^{13}C NMR (CDCl_3 , 101 MHz) δ 164.4, 150.9, 140.8, 120.4 (dt, $J = 259.5$, $J = 217.3$ Hz), 110.4, 73.6 (d, $J = 7.1$ Hz), 48.6, 33.4 (dt, $J = 20.9$, 14.5 Hz), 32.1, 30.8, 29.1, 24.1 (d, $J = 3.5$ Hz), 23.8 (d, $J = 4.9$ Hz), 19.9 (dd, $J = 4.9$ Hz), 12.3. LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 457 (75), 415 (100), 373 (70), 331 (7). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{32}\text{N}_2\text{O}_5\text{F}_2\text{PS}$ 457.1738, found 457.1752.

1,1-Difluoro-5-(2-(5-methyl-2,4-dihydropyrimidin-1,3(2H)-yl)-1-ethanesulfure)pentylphosphonic Acid (39). Following the general procedure used for **9** from difluorophosphonate **35** (100 mg, 0.22 mmol), TMSBr (0.174 mL, 1.32 mmol), and CH_2Cl_2 (5 mL), compound **39** was isolated as a white solid (41 mg, 50%): mp 155 °C; ^1H NMR (DMSO, 400 MHz) δ 11.29 (s, 1H), 7.61 (d, $J = 1.2$ Hz, 1H), 3.85 (t, $J = 6.8$ Hz, 2H), 2.79 (t, $J = 7.3$ Hz, 2H), 2.61 (t, $J = 6.7$ Hz, 2H), 2.10–1.97 (m, 2H), 1.81 (d, $J = 1.0$ Hz, 3H), 1.67–1.60 (m, 4H); ^{19}F NMR (DMSO, 376 MHz) δ –112.35 (dt, $J = 20.0$, 100.8 Hz, 2F); ^{31}P NMR (DMSO, 162 MHz) δ 4.1 (t, $J = 100.8$ Hz, 1P); ^{13}C NMR (MeOD, 101 MHz) δ 166.9, 152.8, 143.5, 121.9 (dt, $J = 257.1$, 210.5 Hz), 110.8, 34.4 (dt, $J = 21.2$, 14.8 Hz), 32.4, 31.2, 30.3 (4C), 21.2 (dd, $J = 4.7$ Hz), 12.2. LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 373 (19), 247 (100), 187 (43). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{12}\text{H}_{20}\text{N}_2\text{O}_5\text{F}_2\text{PS}$ 373.0799, found 373.0800.

Diisopropyl 1,1-Difluoro-3-iodo-4-(5-methyl-2,4-dioxo-3,4-dihydropyridin-1(2H)-yl)butylphosphonate (44). To a refluxed solution of **43** (100.0 mg, 0.29 mmol) and allylthymine (54.0 mg, 0.32 mmol) in 1,2-dichloroethane (2 mL) was added over a period of 1 h via a syringe pump dilauroyl peroxide (35.0 mg, 0.09 mmol) in 1,2-dichloroethane (1 mL). After 3 h, the mixture was cooled, concentrated and the crude product was purified by flash column chromatography using EtOAc/pentane (6/1) as eluent to give **44** (106.0 mg, 71%) as a white solid: mp 110.5 °C; ^1H NMR (CDCl_3 , 400 MHz) δ 8.87 (s, 1H), 6.98 (d, $J = 1.1$ Hz, 1H), 4.79 (dsept, $J = 6.2$ Hz, 2H), 4.59–4.71 (m, 1H), 4.11 (dd, $J = 14.5$, 10.0 Hz, 1H), 3.80 (dd, $J = 14.5$, 10.0 Hz, 1H), 2.67–2.96 (m, 2H), 1.88 (d, $J = 1.0$ Hz, 3H), 1.32 (dd, $J = 6.2$, 3.0 Hz, 12H); ^{19}F NMR (CDCl_3 , 376 MHz) δ –112.29 (dddd, $J = 296.1$, 104.0, 25.0, 13.0 Hz, 1F), –109.8 (dddd, $J = 296.1$, 103.0, 27.1, 11.2 Hz, 1F); ^{31}P NMR (CDCl_3 , 162 MHz) δ 3.5 (dd, $J = 104.0$, 103.0 Hz, 1P); ^{13}C NMR (CDCl_3 , 101 MHz) δ 164.7, 149.9, 140.1, 119.2 (dt, $J = 262.8$, 216.8 Hz), 110.4, 74.2 (d, $J = 7.0$ Hz), 55.3, 41.9 (ddd, $J = 20.6$, 15.6 Hz), 24.0 (d, $J = 3.5$ Hz), 23.6 (d, $J = 4.6$ Hz), 16.6, 12.2. LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 509 (28), 466 (87), 425 (100). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{15}\text{H}_{23}\text{F}_2\text{IN}_2\text{O}_5\text{P}$ 509.0514, found 509.0519.

1,1-Difluoro-3-iodo-4-(5-methyl-2,4-dioxo-3,4-dihydropyridin-1(2H)-yl)butylphosphonic Acid (45). Following the general procedure used for **9**, from **44** (112 mg, 0.22 mmol), TMSBr (0.174 mL, 1.32 mmol), and CH_2Cl_2 (5 mL), compound **45** was obtained as a white powder (71 mg, 76%): mp 168 °C; ^1H NMR (DMSO, 400 MHz) δ 10.51 (s, 1H), 7.19 (d, $J = 1.2$ Hz, 1H), 4.25–4.73 (m, 1H), 3.85–4.10 (m, 2H), 2.58–2.89 (m, 2H), 1.96 (d, $J = 1.0$ Hz, 3H); ^{31}P NMR (DMSO, 162 MHz) δ 3.6 (dd, $J = 104.0$, 103.0 Hz, 1P); ^{19}F NMR (DMSO, 376 MHz) δ –113.05 (dddd, $J = 297.0$, 104.0, 25.2, 12.6 Hz, 1F), –111.36 (dddd, $J = 297.0$, 103.0, 26.7, 11.2 Hz, 1F); ^{13}C NMR (DMSO, 101 MHz) δ 163.7, 150.3, 147.5, 121.2 (dt, $J = 263.9$, 217.6 Hz), 110.9, 57.2, 41.9 (m), 17.1 (m), 14.2. LRMS-ESI (m/z): $[\text{M} - \text{H}]^-$ 422 (64), 295 (100), 126 (2). HRMS-ESI (m/z): $[\text{M} - \text{H}]^-$ calcd for $\text{C}_9\text{H}_11\text{F}_2\text{IN}_2\text{O}_5\text{P}$ 422.9418, found 422.9423.

1-[1-(5-Diisopropoxyphosphono-5,5-difluoropentyl)-1,2,3-triazolo-4-methyl]uracil (52). *General Procedure for the Preparation of Triazolyl Derivatives.* To a stirred solution of **48** (320.0 mg, 1.02 mmol) in $^t\text{BuOH}$ (2.5 mL) and H_2O (2.5 mL) were added propargyl uracil (168.1 mg, 1.12 mmol), sodium ascorbate (19.8 mg, 0.10 mmol), and $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (8.00 mg, 0.05 mmol). The mixture was stirred for 24 h at room temperature, then concentrated under reduced pressure. The residue was diluted in CH_2Cl_2 (10 mL) and washed with water

(10 mL). The aqueous layer was extracted with CH_2Cl_2 (2×10 mL), and combined organic layers were washed with a saturated solution of NaCl (5 mL), dried over MgSO_4 , filtered, and concentrated. The crude product was purified by flash column chromatography using $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ (10/1) as eluent to give **52** (401.8 mg, 85%) as a white solid: mp 105 °C; ^1H NMR (CDCl_3 , 400 MHz) δ 8.69 (s, 1H), 7.59 (s, 1H), 7.23 (d, $J = 1.1$ Hz, 1H), 5.60 (s, 1H), 4.80 (s, 2H), 4.76 (dsept, $J = 6.2$ Hz, 2H), 4.16 (t, $J = 6.9$ Hz, 2H), 1.81–2.06 (m, 4H), 1.52–1.68 (m, 2H), 1.33 (dd, $J = 6.1, 3.1$ Hz, 12H); ^{19}F NMR (CDCl_3 , 376 MHz) δ -112.90 (dt, $J = 106.9, 19.0$ Hz, 2F); ^{31}P NMR (CDCl_3 , 162 MHz) δ 5.0 (t, $J = 106.9$ Hz, 1P); ^{13}C NMR (CDCl_3 , 101 MHz) δ 164.4, 151.3, 142.1, 139.9, 124.0, 119.6 (dt, $J = 261.3, 218.0$ Hz), 110.9, 74.2 (d, $J = 7.1$ Hz), 50.1, 43.0, 33.2 (dt, $J = 20.1, 14.9$ Hz), 29.7, 24.3 (d, $J = 3.6$ Hz), 23.7 (d, $J = 5.1$ Hz), 18.0 (dt, $J = 4.7$ Hz). LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 464 (46), 422 (100), 380 (72). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{20}\text{F}_2\text{N}_5\text{O}_5\text{P}$ 464.1874, found 464.1868.

9-[1-(4-Diisopropoxyphosphono-4,4-difluorobutyl)-1,2,3-triazolo-4-methyl]-6-chloropurine (53). General procedure used for **52** was followed with **47** (300.0 mg, 1.00 mmol) and propargyl-6-chloropurine (211.9 mg, 1.10 mmol) to afford **53** (418.1 mg, 85%) as a white solid: mp 108 °C; ^1H NMR (CDCl_3 , 400 MHz) δ 8.66 (s, 1H), 8.31 (s, 1H), 7.72 (s, 1H), 5.51 (s, 2H), 4.73 (dsept, $J = 6.0$ Hz, 2H), 4.35 (t, $J = 7.2$ Hz, 2H), 2.05–2.18 (m, 2H), 1.89–2.05 (m, 2H), 1.25 (dd, $J = 6.4, 4.0$ Hz, 12H); ^{31}P NMR (CDCl_3 , 162 MHz) δ 4.4 (t, $J = 106.4$ Hz, 1P); ^{19}F NMR (CDCl_3 , 376 MHz) δ -112.34 (dt, $J = 106.4, 18.8$ Hz, 2F); ^{13}C NMR (CDCl_3 , 63 MHz) δ 156.5, 151.0, 145.4, 131.6, 129.6, 126.2, 123.4, 119.5 (dt, $J = 260.8, 218.1$ Hz), 74.0 (d, $J = 7.2$ Hz, 2C), 49.8, 39.1, 31.00 (dt, $J = 21.4, 15.1$ Hz), 24.1 (d, $J = 3.5$ Hz, 2C), 23.8 (d, $J = 4.7$ Hz, 2C), 22.2 (dt, $J = 4.8$ Hz). LRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ 492 (100), 450 (82), 408 (22), 226 (15), 173 (18). HRMS-ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{26}\text{ClF}_2\text{N}_7\text{O}_3\text{P}$ 492.1491, found 492.1470.

1,1-Difluoro-5-(4-((2,4-dioxo-3,4-dihydropyrimidin-2(1H-yl)methyl)-1H-1,2,3-triazol-1-yl)pentylphosphonic Acid (59). General procedure used for **9** was followed with **52** (101.9 mg, 0.22 mmol), TMSBr (0.174 mL, 1.32 mmol), and CH_2Cl_2 (5 mL) to afford **59** (62.6 mg, 75%) as a white solid: mp 184 °C; ^1H NMR (DMSO, 400 MHz) δ 8.15 (s, 1H), 8.07 (s, 1H), 7.87 (s, 1H), 5.80 (s, 2H), 4.54 (t, $J = 7.2$ Hz, 2H), 1.90–2.15 (m, 2H), 1.70–1.85 (m, 2H), 1.43–1.55 (m, 2H); ^{31}P NMR (DMSO, 162 MHz) δ 4.3 (t, $J = 97.0$ Hz, 1P); ^{19}F NMR (DMSO, 376 MHz) δ -112.82 (dt, $J = 97.0, 19.9$ Hz, 2F); ^{13}C NMR (DMSO, 101 MHz) δ 164.9, 151.4, 141.9, 139.3, 129.6, 119.9 (m), 104.1, 55.3, 47.8, 33.5 (m), 28.9, 25.3 (m). LRMS-ESI (m/z): $[\text{M} - \text{H}]^-$ 378 (100), 358 (15), 335 (70), 315 (45). HRMS-ESI (m/z): $[\text{M} - \text{H}]^-$ calcd for $\text{C}_{12}\text{H}_{13}\text{F}_2\text{N}_5\text{O}_5\text{P}$ 378.1235, found 378.1246.

1,1-Difluoro-4-(4-((6-hydroxy-9H-purin-9-yl)methyl)-1H-1,2,3-triazol-1-yl)butylphosphonic Acid (60). General procedure used for **14** was followed with **53** (108.2 mg, 0.22 mmol), TMSBr (202.1 mg, 1.32 mmol), and CH_2Cl_2 (5 mL) to afford **60** (47.9 mg, 56%) as a white solid: mp 185 °C; ^1H NMR (D_2O , 400 MHz) δ 8.45 (s, 1H), 8.23 (s, 1H), 8.12 (s, 1H), 5.62 (s, 2H), 4.49 (t, $J = 6.8$ Hz, 2H), 2.06–2.25 (m, 2H), 1.85–2.05 (m, 2H); ^{31}P NMR (D_2O , 162 MHz) δ 4.5 (t, $J = 97.2$ Hz, 1P); ^{19}F NMR (D_2O , 376 MHz) δ -112.23 (dt, $J = 97.2, 19.9$ Hz, 2F); ^{13}C NMR (D_2O , 101 MHz) δ 155.8, 155.2, 146.0, 144.4, 142.5, 123.3, 121.3, 114.89 (m), 50.1, 42.1, 32.4 (m), 17.3 (m). LRMS-ESI (m/z): $[\text{M} - \text{H}]^-$ 388 (100), 368 (8), 216 (20), 171 (62), 135 (26). HRMS-ESI (m/z): $[\text{M} - \text{H}]^-$ calcd for $\text{C}_{12}\text{H}_{13}\text{F}_2\text{N}_7\text{O}_4\text{P}$ 388.0735, found 388.0718.

■ ASSOCIATED CONTENT

Supporting Information

Experimental details for the preparation of compounds **7–65** not described in the manuscript, HPLC data of all tested compounds (POT, SBRU, **7–10**, **14–16**, **27–29**, **39–42**, **45**, **56–61**, **65**), and ^1H and ^{13}C NMR spectra for compounds **5–65**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Phone: (33)231452854. E-mail: Thierry.Lequeux@ensicaen.fr.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the CNRS and the Crunch (Interregional Organic Chemistry Network). Dr. J. F. Lohier is gratefully thanked for providing crystallographic data.

■ ABBREVIATIONS USED

TPase, thymidine phosphorylase; PD-ECGF, platelet-derived endothelial cell growth factor; 2dR, 2-deoxyribose; TPI, thymidine phosphorylase inhibitor; TMG, tetramethylguanidine; LDA, lithium diisopropylamide; DIAD, diethyl diazodicarboxylate; 6ASBU, 6-amino-5-bromouracyl; POT, 1-phosphonocytlythymine; PNP, purine nucleoside phosphorylase; K_i , inhibition constant; TMSBr, trimethylsilyl bromide

■ REFERENCES

- (1) Moghaddam, A.; Bicknell, R. Expression of Platelet-Derived Endothelial Cell Growth Factor in *Escherichia coli* and Confirmation of Its Thymidine Phosphorylase Activity. *Biochemistry* **1992**, *31*, 12141–12146.
- (2) Barton, G. J.; Ponting, C. P.; Spraggon, G.; Finnis, C.; Sleep, D. Human Platelet-Derived Endothelial Cell Growth Factor Is Homologous to *Escherichia coli* Thymidine Phosphorylase. *Protein Sci.* **1992**, *1*, 688–690.
- (3) Schwartz, E. L.; Hotchkiss, K. A.; Ashton, A. W. Thymidine Phosphorylase and 2-Deoxyribose Stimulate Human Endothelial Cell Migration by Specific Activation of the Integrins $\alpha_5\beta_1$ and $\alpha_v\beta_3$. *J. Biol. Chem.* **2003**, *278*, 19272–19279.
- (4) (a) Matsushita, S.; Nitanda, T.; Furukawa, T.; Sumizawa, T.; Tani, A.; Nishimoto, K.; Akiba, S.; Miyadera, K.; Fukushima, M.; Yamada, Y.; Yoshida, H.; Kanzaki, T.; Akiyama, S. The Effect of a Thymidine Phosphorylase Inhibitor on Angiogenesis and Apoptosis in Tumors. *Cancer Res.* **1999**, *59*, 1911–1916. (b) Takao, S.; Akiyama, S.; Nakajo, A.; Yoh, H.; Kitazono, M.; Natsgoe, S.; Miyadera, K.; Fukushima, M.; Yamada, Y.; Aikou, T. Suppression of Metastasis by Thymidine Phosphorylase Inhibitor. *Cancer Res.* **2000**, *60*, 5345–5348. (c) Sengupta, S.; Sellers, L. A.; Matheson, H. B.; Fan, T. P. D. Thymidine Phosphorylase Induces Angiogenesis in Vivo and in Vitro: An Evaluation of Possible Mechanisms. *Br. J. Pharmacol.* **2003**, *139*, 219–231.
- (5) (a) Uracil Derivatives, and Antitumor Effect Potentiator and Antitumor Agent Containing the Same. Patent US5744475, 1998; Taiho Pharmaceutical Co. Ltd.; *Chem. Abstr.* **1998**, *125*, 328727. (b) Angiogenesis Inhibitors. Patent EP1354589, 2003; Japan Science and Tech Corp.; *Chem. Abstr.* **2003**, *137*, 41733.
- (6) (a) Foche, F.; Spadari, S. Thymidine Phosphorylase: A Two-Face Janus in Anticancer Chemotherapy. *Curr. Cancer Drug Targets* **2001**, *1*, 141–153. (b) Balzarini, J.; Degève, B.; Esteban-Gamboa, A.; Esnouf, R.; De Clercq, E.; Engelborghs, Y.; Camarasa, M. J.; Pérez-Pérez, M. J. Kinetic Analysis of Novel Multisubstrate Analogue Inhibitors of Thymidine Phosphorylase. *FEBS Lett.* **2000**, *483*, 181–185. (c) Langen, P.; Etzold, G.; Bärwolff, D.; Preussel, B. Inhibition of Thymidine Phosphorylase by 6-Aminothymine and Derivatives of 6-Aminouracil. *Biochem. Pharmacol.* **1967**, *16*, 1833–1837. (d) Esteban-Gamboa, A.; Balzarini, J.; Esnouf, R.; De Clercq, E.; Camarasa, M. J.; Pérez-Pérez, M. Design, Synthesis, and Enzymatic Evaluation of Multisubstrate Analogue Inhibitors of *Escherichia coli* Thymidine Phosphorylase. *J. Med. Chem.* **2000**, *43*, 971–983. (e) Votruba, I.; Pomeisl, K.; Tloustova, E.; Holy, A.; Otova, B. Inhibition of Thymidine Phosphorylase (PD-ECGF) from SD-Lymphoma by Phosphonomethoxyalkyl Thymines. *Biochem. Pharmacol.* **2005**, *69*,

- 1517–1521. (f) Allan, A. L.; Gladstone, P. L.; Price, M. L. P.; Hopkins, S. A.; Juarez, J. C.; Donate, F.; Fernansky, R. J.; Shaw, D. E.; Ganem, B.; Lee, Y.; Wang, W.; Ealick, S. Synthesis and Evaluation of Multisubstrate Bicyclic Pyrimidine Nucleoside Inhibitors of Human Thymidine Phosphorylase. *J. Med. Chem.* **2006**, *49*, 7807–7815. (g) Casanova, E.; Hernandez, A. I.; Priego, E. M.; Liekens, S.; Camarasa, M. J.; Balzarini, J.; Pérez-Pérez, M. J. 5'-O-Tritylinosine and Analogues as Allosteric Inhibitors of Human Thymidine Phosphorylase. *J. Med. Chem.* **2006**, *49*, 5562–5570. (h) Reigan, P.; Edwards, P. N.; Gbaj, A.; Cole, C.; Barry, S. T.; Page, K. M.; Ashton, S. E.; Luke, R. W. A.; Douglas, K. T.; Stratford, I. J.; Jaffar, M.; Bryce, R. A.; Freeman, S. Aminoimidazolymethyluracil Analogues as Potent Inhibitors of Thymidine Phosphorylase and Their Bioreductive Nitroimidazolyl Prodrugs. *J. Med. Chem.* **2005**, *48*, 392–402. (i) Cole, C.; Reigan, P.; Gbaj, A.; Edwards, P. N.; Douglas, K. T.; Stratford, I. J.; Freeman, S.; Jaffar, M. Potential Tumor-Selective Nitroimidazolymethyluracil Prodrug Derivatives: Inhibitors of the Angiogenic Enzyme Thymidine Phosphorylase. *J. Med. Chem.* **2003**, *46*, 207–209. (j) Pradère, U.; Clavier, H.; Roy, V.; Nolan, S. P.; Agrofoglio, L. A. The Shortest Strategy for Generating Phosphonate Prodrugs by Olefin Cross-Metathesis: Application to Acyclonucleoside Phosphonates. *Eur. J. Org. Chem.* **2011**, 7324–7330. (k) Montagu, A.; Pradère, U.; Roy, V.; Nolan, S. P.; Agrofoglio, L. A. Expedient Convergent Procedure for the Preparation of Bis(POC) Prodrugs of New (E)-4-Phosphonobut-2-en-1-yl Nucleosides. *Tetrahedron* **2011**, *67*, 5319–5328. (l) Barral, K.; Weck, C.; Payrot, N.; Roux, L.; Durafour, C.; Zoulim, F.; Neyts, J.; Balzarini, J.; Canard, B.; Priet, S.; Alvarez, K. Acyclic Nucleoside Thiophosphonates as Potent Inhibitors of HIV and HBV Replication. *Eur. J. Med. Chem.* **2011**, *46*, 4281–4288. (m) Vakiaeva, N.; Wyles, D. L.; Schooley, R. T.; Hwu, J. B.; Beadle, J. R.; Prichard, M. N.; Hostetler, K. Y. Synthesis and Antiviral Evaluation of 9-(S)-[3-Alkoxy-2-(phosphonomethoxy)propyl]nucleoside Alkoxyalkyl Esters: Inhibitors of Hepatitis C Virus and HIV-1 Replication. *Bioorg. Med. Chem.* **2011**, *19*, 4616–4625. (7) (a) Halazy, S.; Ehrhard, A.; Danzin, C. 9-(Difluorophosphonoalkyl)guanines as a New Class of Multisubstrate Analog Inhibitors of Purine Nucleoside Phosphorylase. *J. Am. Chem. Soc.* **1991**, *113*, 315–317. (b) Burke, T. R.; Smyth, M. S.; Otaka, A.; Nomizu, M.; Roller, P. P.; Wolf, G.; Case, R.; Shoelson, S. E. Nonhydrolyzable Phosphotyrosyl Mimetics for the Preparation of Phosphatase-Resistant SH2 Domain Inhibitors. *Biochemistry* **1994**, *33*, 6490–6494. (c) Berkowitz, D. B.; Bose, M. α -Monofluoroalkylphosphonates: A Class of Isoacidic and “Tunable” Mimics of Biological Phosphates. *J. Fluorine Chem.* **2001**, *112*, 13–33. (8) Romanenko, V. D.; Kukhar, V. P. Fluorinated Phosphonates: Synthesis and Biomedical Application. *Chem. Rev.* **2006**, *106*, 3868–3935. (9) (a) Birck, M. R.; Schramm, V. L. Nucleophilic Participation in the Transition State for Human Thymidine Phosphorylase. *J. Am. Chem. Soc.* **2004**, *126*, 2447–2453. (b) Birck, M. R.; Schramm, V. L. Binding Causes the Remote [^3H]Thymidine Kinetic Isotope Effect in Human Thymidine Phosphorylase. *J. Am. Chem. Soc.* **2004**, *126*, 6882–6883. (10) (a) Pfund, E.; Lequeux, T.; Vazeux, M.; Masson, S. Synthesis of Thiapyranoside Precursors Using the Building-Block Approach from a Phosphonodifluorodithioacetate. *Tetrahedron Lett.* **2002**, *43*, 2033–2036. (b) Pfund, E.; Lequeux, T.; Vazeux, M.; Masson, S. Synthesis of Thiazolines Linked to a Difluoromethylphosphonate Diester via Dithioester Chemistry. *Org. Lett.* **2002**, *4*, 843–846. (c) Henry-dit-Quesnel, A.; Toupet, L.; Pommelet, J. C.; Lequeux, T. A Difluorosulfide as a Freon-Free Source of Phosphonodifluoromethyl Carbanion. *Org. Biomol. Chem.* **2003**, *1*, 2486–2491. (d) Pfund, E.; Masson, S.; Vazeux, M.; Lequeux, T. Syntheses of α -Fluoro- α,β -unsaturated Thioamides and Thiazolines from a Fluorophosphonodithioacetate. *J. Org. Chem.* **2004**, *69*, 4670–4676. (e) Ozouf, P.; Binot, G.; Pommelet, J. P.; Lequeux, T. Syntheses of ω -Hydroxy- α,α -difluoromethylphosphonates by Oxacycle Ring-Opening Reactions. *Org. Lett.* **2004**, *6*, 3747–3750. (f) Pfund, E.; Lequeux, T.; Masson, S.; Cordi, A.; Pierre, A.; Hervé, G.; Serre, V. Efficient Synthesis of Fluorothiosparfosic Acid Analogues with Potential Antitumoral Activity. *Bioorg. Med. Chem.* **2005**, *13*, 4921–4928. (11) Diab, S. A.; Sene, A.; Pfund, E.; Lequeux, T. Efficient Synthesis of Fluorophosphonylated Alkyls by Ring-Opening Reaction of Cyclic Sulfates. *Org. Lett.* **2008**, *10*, 3895–3898. (12) Jakeman, D. L.; Ivory, A. J.; Williamson, M. P.; Blackburn, G. M. Highly Potent Bisphosphonate Ligands for Phosphoglycerate Kinase. *J. Med. Chem.* **1998**, *41*, 4439–4452. (13) Kelly, J. L.; Linn, J. A.; McLean, E. W.; Tuttle, J. V. 9-[(Phosphonoalkyl)benzyl]guanines. Multisubstrate Analogue Inhibitors of Human Erythrocyte Purine Nucleoside Phosphorylase. *J. Med. Chem.* **1993**, *36*, 3455–3463. (14) (a) Sene, A.; Diab, S. A.; Hienzsch, A.; Cahard, D.; Lequeux, T. Radical Conjugated Addition: Addition of Dialkyl Phosphonodifluoromethyl Radical onto Unsaturated Ketones. *Synlett* **2009**, 981–985. (b) Yang, Z.; Burton, D. J. A Novel and Practical Preparation of α,α -Difluoro Functionalized Phosphonates from Iododifluoromethylphosphonate. *J. Org. Chem.* **1992**, *57*, 4676–4683. (15) Diab, S. A.; Hienzsch, A.; Lebargy, C.; Guillaume, S.; Pfund, E.; Lequeux, T. Synthesis of Fluorophosphonylated Acyclic Nucleotide Analogues via Copper(I)-Catalyzed Huisgen 1–3 Dipolar Cycloaddition. *Org. Biomol. Chem.* **2009**, *7*, 4481–4490. (16) The following crystal structure has been deposited at the Cambridge Crystallographic Data Centre and allocated the deposition number CCDC 850861. (17) Birck, M. R.; Schramm, V. L. Nucleophilic Participation in the Transition State for Human Thymidine Phosphorylase. *J. Am. Chem. Soc.* **2004**, *126*, 2447–2453. (18) (a) Yokomatsu, T.; Hayakawa, Y.; Suemune, K.; Kihara, T.; Soeda, S.; Shimeno, H.; Shibuya, S. Synthesis and Biological Evaluation of 1,1-Difluoro-2-(tetrahydro-3-furanyl)ethylphosphonic Acids Possessing a N^9 -Purinylmethyl Functional Group at the Ring. A New Class of Inhibitors for Purine Nucleoside Phosphorylases. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 2833–2836. (b) Yokomatsu, T.; Abe, H.; Sato, M.; Suemune, K.; Kihara, T.; Soeda, S.; Shimeno, H.; Shibuya, S. Selective α -1A Adrenergic Receptor Antagonists. Effects of Pharmacophore Regio- and Stereochemistry on Potency and Selectivity. *Bioorg. Med. Chem.* **1998**, *6*, 2495–2505. (c) Yokomatsu, T.; Hayakawa, Y.; Kihara, T.; Koyanagi, S.; Soeda, S.; Shimeno, H.; Shibuya, S. Synthesis of the Novel Analogues of Dysidiolide and Their Structure–Activity Relationship. *Bioorg. Med. Chem.* **2000**, *8*, 2571–2579. (d) Halazy, S.; Ehrhard, A.; Eggenpiller, A.; Berges-Gross, V.; Danzin, C. Fluorophosphonate Derivatives of N^9 -Benzylguanine as Potent, Slow-Binding Multisubstrate Analogue Inhibitors of Purine Nucleoside Phosphorylase. *Tetrahedron* **1996**, *52*, 177–184. (19) Lewis, W. G.; Green, L. G.; Grynszpan, F.; Radic, Z.; Carlier, P. R.; Taylor, P.; Finn, M. G.; Sharpless, K. B. Click Chemistry in Situ: Acetylcholinesterase as a Reaction Vessel for the Selective Assembly of a Femtomolar Inhibitor from an Array of Building Blocks. *Angew. Chem., Int. Ed.* **2002**, 1053–1057. (20) (a) Siddiqui, A.; Ballatore, C.; McGuigan, C.; De Clercq, E.; Balzarini, J. The Presence of Substituents on the Aryl Moiety of the Aryl Phosphoramidate Derivative of d4T Enhances Anti-HIV Efficacy in Cell Culture: A Structure–Activity Relationship. *J. Med. Chem.* **1999**, *42*, 393–399. (b) Mathé, C.; Périgaud, C.; Gosselin, G.; Imbach, J. L. Phosphopeptide Prodrug Bearing an S-Acyl-2-thioethyl Enzyme-Labile Phosphate Protection. *J. Org. Chem.* **1998**, *63*, 8547–8550. (c) Valette, G.; Pompon, A.; Girardet, J. L.; Cappellacci, L.; Franchetti, P.; Grifantini, M.; La Colla, P.; Loi, A. G.; Périgaud, C.; Gosselin, G.; Imbach, J. L. Decomposition Pathways and in Vitro HIV Inhibitory Effects of IsoddA Pronucleotides: Toward a Rational Approach for Intracellular Delivery of Nucleoside 5'-Monophosphates. *J. Med. Chem.* **1996**, *39*, 1981–1990. (d) Peyrottes, S.; Coussot, G.; Lefebvre, I.; Imbach, J. L.; Gosselin, G.; Aubertin, A. M.; Périgaud, C. S-Acyl-2-Thioethyl Aryl Phosphotriester Derivatives of AZT: Synthesis, Antiviral Activity, and Stability Study. *J. Med. Chem.* **2003**, *46*, 782–793. (e) Egron, D.; Imbach, J. L.; Gosselin, G.; Aubertin, A. M.; Périgaud, C. S-Acyl-2-thioethyl Phosphoramidate

Diester Derivatives as Mononucleotide Prodrugs. *J. Med. Chem.* **2003**, 46, 4564–4571.

(21) Kim, C. U.; Luh, B. Y.; Misco, P. F.; Bronson, J. J.; Hitchcock, M. J. M.; Ghazzouli, I.; Martin, J. C. Acyclic Purine Phosphonate Analogues as Antiviral Agents. Synthesis and Structure–Activity Relationships. *J. Med. Chem.* **1990**, 33, 1207–1213.