

NOTE

The *Helicobacter pylori* Ferric Uptake Regulator (Fur) Is Essential for Growth Under Sodium Chloride Stress

Hanan Gancz and D. Scott Merrell*

Department of Microbiology and Immunology, Uniformed Services University of the Health Sciences,
4301 Jones Bridge Rd., Bethesda, MD 20814, USA

(Received September 29, 2010 / Accepted November 22, 2010)

Epidemiological data and animal models indicate that *Helicobacter pylori* and dietary NaCl have a synergistic ill effect on gastric maladies. Here we show that the Ferric Uptake Regulator (Fur), which is a crucial regulatory factor required for *H. pylori* colonization, is essential for growth in the presence of high NaCl concentrations. Moreover, we demonstrate that the transcriptional response induced by sodium chloride stress exhibits similarities to that seen under iron depletion.

Keywords: iron, regulation, Fur, *Helicobacter pylori*

Helicobacter pylori can be found in the gastric mucosa of more than half of the human population (Pounder and Ng, 1995). Long-term colonization results in gastritis and may lead to gastric or duodenal ulcers (Malaty, 2007). Infection with *H. pylori* is also a risk factor for development of gastric cancer (Matysiak-Budnik and Megraud, 2006; Mbulaiteye *et al.*, 2009); to date, *H. pylori* is the only bacterium classified as a carcinogen (Anon, 1994). For colonized individuals, epidemiological data indicate a synergistic ill effect between diets rich in sodium chloride and *H. pylori* infection (Tsugane and Sasazuki, 2007; Wang *et al.*, 2009). The role of a high salt diet on *H. pylori* infection was also previously examined in several animal models. In a mouse model of infection, an increase in bacterial load was observed in animals consuming a high salt diet (Rogers *et al.*, 2005). Whereas, in a gastric chemical carcinogenesis model of *H. pylori*-infected Mongolian gerbils, a high salt diet resulted in the dose-dependent promotion of cancer initiation that was accompanied by a shift in mucin production from the glandular to the surface mucous cells (Kato *et al.*, 2006). Taken together, the epidemiological and animal based evidence suggest dietary mediated increases in the severity of *H. pylori*-induced disease outcomes.

Based on the above observations, we and others postulated and subsequently showed that salt concentration contributes to changes in expression of *H. pylori* virulence factors such as *cagA* (Loh *et al.*, 2007) and *vacA* (Gancz *et al.*, 2008) *in vitro*. However, the mechanisms of regulation of the salt stress response and the factors required for salt survival are not yet characterized for this organism.

In order to identify genes that contribute to *H. pylori*'s ability to withstand salt stress, we took a targeted approach and screened defined *H. pylori* mutants that were constructed in

the G27 strain background and already existed in our strain collection (Table 1). Specifically, we targeted strains containing mutations in known regulatory genes (*flgR*, *fur*, *luxS*, *spoT*) or known virulence genes (*cagA*, *ureB*, *vacA*). All *H. pylori* strains were maintained as frozen stocks at 80°C as previously described (Carpenter *et al.*, 2010). Prior to each experiment, strains were revived on horse blood agar (HBA) plates as previously described (Carpenter *et al.*, 2010) and used to inoculate liquid cultures that consisted of Brucella broth (Neogen Corporation) supplemented with 10% fetal bovine serum and 10 µg/ml vancomycin (BB). These cultures were grown at 37°C with shaking at 110 rpm. Both liquid and plate cultures were grown under microaerobic conditions (5% O₂, 10% CO₂, and 85% N₂) generated with an Anoxomat gas evacuation and replacement system (Spiral Biotech) in gas evacuation jars. When reviving mutant bacterial strains from the freezer only, plates were supplemented with 8 µg/ml chloramphenicol (Cm) (EMD Chemicals, Inc.) and/or 25 µg/ml kanamycin (Kan) (Gibco) where appropriate (Table 1).

To determine whether mutation of any of the *H. pylori* regulatory or virulence genes affected growth in the presence of elevated salt concentrations, we calculated the plating efficiency of each mutant strain and the wildtype G27 control strain by determining the number of colonies obtained after spotting on solidified (1.3% agar final) BB supplemented with various concentrations of NaCl. The normal solidified BB medium contains 5 g/L of NaCl (BB5), and we achieved increased salt concentration by use of BB containing 13.5 g/L of NaCl (BB13.5). This elevated concentration of sodium chloride is lower than the dramatic growth inhibitory condition we previously described for this strain (BB15) (Gancz *et al.*, 2008). The wildtype strain and each of the defined isogenic mutant strains were grown to mid-exponential phase in liquid media and then OD equilibrated to give identical cell densities. Cells were then serially diluted and plated side-by-side on

* For correspondence. E-mail: dmerrell@usuhs.mil; Tel: +1-301-295-1584; Fax: +1-301-295-3773

Table 1. Plasmids and strains used in this study

Plasmid or strain	Description	Reference
Plasmids		
pDSM142	pΔHP1027-K7	Gancz <i>et al.</i> (2006)
pDSM340	<i>fur</i> ::pTM117 complementation vector	Carpenter <i>et al.</i> (2007)
<i>H. pylori</i> strains		
G27	WT <i>H. pylori</i>	Covacci <i>et al.</i> (1993)
26695	WT <i>H. pylori</i>	Tomb <i>et al.</i> (1997)
43504	WT <i>H. pylori</i>	Alamuri <i>et al.</i> (2006)
DSM43	G27 <i>ureB</i> :: <i>aphA</i> , Kan ^r	Joyce <i>et al.</i> (2001)
DSM90	G27 <i>flgR</i> :: <i>cat</i> , Cm ^r	This study
DSM205	G27 <i>vacA</i> :: <i>aphA</i> , Kan ^r	Amieva <i>et al.</i> (2002)
DSM207	G27 <i>cagA</i> :: <i>cat</i> , Cm ^r	Amieva <i>et al.</i> (2002)
DSM211	G27 <i>luxS</i> :: <i>aphA</i> , Kan ^r	Kind gift from J.V. Solnick laboratory
DSM300	G27 <i>Δfur</i> :: <i>cat</i> , Cm ^r	Carpenter <i>et al.</i> (2007)
DSM343	DSM300 (pDSM340), Kan ^r Cm ^r	Carpenter <i>et al.</i> (2007)
DSM357	26695 <i>Δfur</i> :: <i>cat</i> , Cm ^r	Carpenter <i>et al.</i> (2009)
DSM388	43504 <i>Δfur</i> :: <i>aphA</i> , Kan ^r	This study
DSM594	G27 <i>spoT</i> :: <i>aphA</i> , Kan ^r	Zhou <i>et al.</i> (2008)

BB5 and BB13.5 plates (Fig. 1) as well as on HBA plates (data not shown).

Comparison of the plating data revealed that while wildtype G27 showed no decrease in plating efficiency on BB13.5, the resulting colonies were significantly smaller than on non-supplemented media (Fig. 1). This fact suggests that the concentration of NaCl found in the BB13.5 plates was sufficient to create stressful conditions under which the bacteria can grow but at a slower rate. Disruption of *cagA*, *flgR*, *luxS*, *ureB* or *vacA* all resulted in plating efficiencies that were similar to those of the wildtype (data not shown). However, disruption of the ferric uptake regulator, *fur*, or *spoT* resulted in a significant reduction in the plating efficiency of these mutant strains on high salt media (Fig. 1 panel A and data not shown). Given that the *fur* mutant strain showed the most drastic growth defect at elevated salt concentrations and because Fur is known to regulate additional stress responses besides the response to iron ion concentration (Bijlsma *et al.*, 2002; Carpenter *et al.*, 2010), has unique features in *H. pylori* gene regulation (Delany *et al.*, 2001; Ernst *et al.*, 2005; Miles *et al.*, 2010a) and is the scope of on-going research in our laboratory (Gancz *et al.*, 2006; Carpenter *et al.*, 2007, 2009, 2010; Miles *et al.*, 2010a, 2010b), we chose to further investigate the role of *fur* in the response to salt stress.

First, to conclusively demonstrate that the *fur* mutation was responsible for the demonstrated salt sensitive phenotype, we attempted to complement the sensitivity phenotype using pDSM340, which is a plasmid that carries a wildtype copy of *fur* (Carpenter *et al.*, 2007). Transformation of this plasmid into the *Δfur* background resulted in complete restoration of growth to the wildtype phenotype (Fig. 1, panel A). Therefore, we conclude that *fur* is essential for growth under these conditions.

Next, given that *H. pylori* is considered a panmictic species (Salaun *et al.*, 1998) and since increasing evidence indicates the occurrence of strain specific phenomenon (Gancz *et al.*, 2008; Carpenter *et al.*, 2009), we next investigated the necessity of *fur* for growth under high salt conditions in two additional

strain backgrounds. To this end, we compared the plating efficiency of strains HP26695 and HP43504 to that of their isogenic *Δfur* mutants, which were constructed as previously described (Gancz *et al.*, 2006; Carpenter *et al.*, 2009). We noticed that although the various wildtype strains differed in their overall salt sensitivity, each of the *Δfur* strains showed a growth defect in comparison to their respective parental strain (Fig. 1, panel B). This indicates that the requirement of Fur for growth under salt stress conditions is not strain dependent. Similar to G27, the other *H. pylori* strains also exhibited smaller colonies on BB13.5, indicating that they were experiencing stressful conditions (Fig. 2, panel B).

Fur is known to be an important regulatory factor in *H. pylori* that has been shown to control the expression of a large number of genes (Lee *et al.*, 2004; Ernst *et al.*, 2005a; Danielli *et al.*, 2006; Gancz *et al.*, 2006). Included among these genes is *amiE*, which encodes an aliphatic amidase that is regulated by the iron-bound form of Fur (van Vliet *et al.*, 2003), and *pfr*, which encodes a nonheme iron-containing ferritin that is regulated by the apo form of Fur (Bereswill *et al.*, 2000). Thus, to determine if Fur regulated processes were taking place during salt stress in *H. pylori*, we monitored the relative transcription of these two known Fur dependent genes by RNAse protection assay (RPA) as previously described (Carpenter *et al.*, 2007). Additionally, we monitored the Fur-dependent expression of *vacA* and *cagA*, two virulence genes whose salt-dependent transcription we previously examined (Gancz *et al.*, 2008). RPAs were conducted with RNA harvested from cultures of the wildtype strain and the isogenic *fur* mutant strain that were grown in BB5 and then exposed to a salt shock for one hour. To accomplish this, overnight cultures of each strain were harvested by centrifugation and then exposed to salt shock conditions by resuspension of the cells in BB containing 25 g/L of NaCl (BB25). RNA was extracted from bacterial cells, the *amiE*, *pfr*, *cagA*, and *vacA* riboprobe templates were generated by PCR, radiolabeled riboprobes were produced, and the RPAs were performed all as previously described (Gancz *et al.*, 2008; Carpenter *et al.*,




Fig. 1. Growth characteristics of *H. pylori* and isogenic Δfur mutants on normal (BB5) and elevated salt (BB13.5) solidified media. (A) Relative growth of G27 wildtype, the isogenic Δfur mutant (DSM300), and the *fur* complemented strain (DSM343) on BB5 and BB13.5 plates is shown. Ten-fold dilutions of the indicated strain were spotted on solidified Brucella broth based medium containing 5 g/L NaCl (BB5) (upper panel) or 13.5 g/L NaCl (BB13.5) (lower panel). (B) Relative growth of strains 43054 and 26695 and their isogenic Δfur mutant strains on BB5 and BB13.5 plates is shown. Ten-fold dilutions of the indicated strain were spotted on solidified Brucella broth based medium containing 5 g/L NaCl (BB5) (upper panel) or 13.5 g/L NaCl (BB13.5) (lower panel). The results shown are representative of multiple biological repeats.

2010). The gels were subsequently exposed to phosphor screens, which were scanned using a FLA-5100 scanner (FujiFilm). Data were then analyzed/quantitated using the Multi-Gauge software (version 3.0, FujiFilm).

Examination of the response of the virulence factors *vacA*




Fig. 2. Transcription of *amiE*, *pfr*, *vacA* and *cagA* following salt shock as assessed by RNAse protection assay (RPA). Logarithmically growing liquid cultures of G27 and the isogenic Δfur mutant strain were salt shocked by replacing the BB5 growth medium with BB25 medium (containing 25 g/L NaCl). Samples were taken immediately after the cells were resuspended in BB25 (t=0) and 1 h after the change of the medium (t=60). RNA was harvested, and RPAs were conducted as described in the text. Protected bands for the indicated transcript are shown for each strain. The fold change in expression from the t=0 time point is indicated. These data are representative of three independent biological experiments.

and *cagA* to salt shock by RPAs demonstrated that, as expected for strain G27 (Gancz *et al.*, 2008), upon exposure to increased salt, *vacA* expression was increased by 10.55 fold, while *cagA* expression was virtually unchanged (Fig. 2). These responses were almost identical in the *fur* mutant strain (Fig. 2), indicating that Fur does not function as a critical regulatory factor for *cagA* and *vacA* in these conditions. Conversely, Fur does function as a critical regulator of *amiE* and *pfr* in these conditions; in the wildtype strain, *amiE* expression was increased upon salt shock, while *pfr* expression was decreased, but these changes were completely lost in the *fur* mutant strain (Fig. 2). Interestingly, these changes in expression closely resemble the previously characterised transcriptional response of these genes to iron depletion (Carpenter *et al.*, 2007, 2009). For example, *amiE* expression in response to salt stress increased 2.03 fold in comparison to 5.8 fold under iron depletion (Carpenter *et al.*, 2007), while the *pfr* transcript showed a 5.6 decrease under salt shock conditions and a 3.3 fold decrease under iron depletion (Carpenter *et al.*, 2007). Taken together, these data suggest that for *H. pylori*, salt shock mimics iron depletion and that Fur plays a crucial role in mediating changes in gene expression that allow growth under these conditions.

The involvement of Fur in the response to high concentrations of salt was previously also suggested for other micro-

organisms: *Desulfovibrio vulgaris* (Mukhopadhyay *et al.*, 2006; Bender *et al.*, 2007; He *et al.*, 2010), the halophilic bacterium *Chromohalobacter salexigens* (Argandona *et al.*, 2010), *Bacillus subtilis* (Hoffmann *et al.*, 2002; Steil *et al.*, 2003; Hahne *et al.*, 2010), and *Bacillus cereus* (den Besten *et al.*, 2009). For *D. vulgaris*, microarray analysis demonstrated that many of the genes showing marked changes in transcription following salt exposure belonged to the Fur regulon. Despite this, the wildtype *D. vulgaris* and the *Afur* strain display virtually identical suppressed growth curves under elevated salt conditions. However, the *fur* mutant strain differs from the wildtype in that while growth of the wildtype in salt can be restored by the addition of glycine betaine as an osmoprotectant, the *fur* mutant strain is unable to be rescued from salt induced poor growth (Bender *et al.*, 2007). In *B. subtilis* the connection between Fur and salt stress was inferred from proteomic examination of the response to high salt conditions (Hoffmann *et al.*, 2002) and was further studied by comparative transcriptional profiling in response to iron depletion and salt shock (Steil *et al.*, 2003; Hahne *et al.*, 2010). Interestingly, in this organism addition of iron to the growth media partially reversed the *B. subtilis* growth defect induced by salt-stress (Hoffmann *et al.*, 2002). Similarly, in the closely related bacterium, *B. cereus*, salt exposure causes induction of genes involved in iron homeostasis (den Besten *et al.* 2009). Thus, while it is clear that in *H. pylori* Fur acts as a global transcriptional regulator that responds to and/or mediates the response to an assortment of environmental signals (Bijlsma *et al.*, 2002; Carpenter *et al.*, 2010), we suggest that the link between salt, iron and mediation of the osmotic stress response by Fur crosses bacterial taxa and may prove to be a general one.

We thank members of the Merrell lab for useful discussions and Jay Solnick for the G27 *luxS* mutant. Research in the laboratory of D. Scott Merrell is made possible by grants R073PW from USUHS and AI065529 from the NIAID. Contents of this manuscript are the sole responsibility of the authors and do not necessarily represent the official views of the NIH or the DOD.

References

1994. Schistosomes, liver flukes and *Helicobacter pylori*. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Lyon, 7-14 June 1994. *IARC Monogr Eval. Carcinog. Risks Hum.* 61, 1-241.
- Alamuri, P., N. Mehta, A. Burk, and R.J. Maier. 2006. Regulation of the *Helicobacter pylori* Fe-S cluster synthesis protein NifS by iron, oxidative stress conditions, and fur. *J. Bacteriol.* 188, 5325-5330.
- Amieva, M.R., N.R. Salama, L.S. Tompkins, and S. Falkow. 2002. *Helicobacter pylori* enter and survive within multivesicular vacuoles of epithelial cells. *Cell Microbiol.* 4, 677-690.
- Argandona, M., J.J. Nieto, F. Iglesias-Guerra, M.I. Calderon, R. Garcia-Esteva, and C. Vargas. 2010. Interplay between iron homeostasis and the osmotic stress response in the halophilic bacterium *Chromohalobacter salexigens*. *Appl. Environ. Microbiol.* 76, 3575-3589.
- Bender, K.S., H.C. Yen, C.L. Hemme, Z. Yang, Z. He, Q. He, J. Zhou, and *et al.* 2007. Analysis of a ferric uptake regulator (Fur) mutant of *Desulfovibrio vulgaris* Hildenborough. *Appl. Environ. Microbiol.* 73, 5389-5400.
- Bereswill, S., S. Greiner, A.H. van Vliet, B. Waidner, F. Fassbinder, E. Schiltz, J.G. Kusters, and M. Kist. 2000. Regulation of ferritin-mediated cytoplasmic iron storage by the ferric uptake regulator homolog (Fur) of *Helicobacter pylori*. *J. Bacteriol.* 182, 5948-5953.
- Bijlsma, J.J., B. Waidner, A.H. Vliet, N.J. Hughes, S. Hag, S. Bereswill, D.J. Kelly, C.M. Vandenbroucke-Grauls, M. Kist, and J.G. Kusters. 2002. The *Helicobacter pylori* homologue of the ferric uptake regulator is involved in acid resistance. *Infect. Immun.* 70, 606-611.
- Carpenter, B.M., H. Gancz, S.L. Benoit, S. Evans, C.H. Olsen, S.L. Michel, R.J. Maier, and D.S. Merrell. 2010. Mutagenesis of conserved amino acids of *Helicobacter pylori* Fur reveals residues important for function. *J. Bacteriol.* 192, 5037-5052.
- Carpenter, B.M., H. Gancz, R.P. Gonzalez-Nieves, A.L. West, J.M. Whitmire, S.L. Michel, and D.S. Merrell. 2009. A single nucleotide change affects fur-dependent regulation of *sodB* in *H. pylori*. *PLoS One.* 4, e5369.
- Carpenter, B.M., T.K. McDaniel, J.M. Whitmire, H. Gancz, S. Guidotti, S. Censini, and D.S. Merrell. 2007. Expanding the *Helicobacter pylori* genetic toolbox, modification of an endogenous plasmid for use as a transcriptional reporter and complementation vector. *Appl. Environ. Microbiol.* 73, 7506-7514.
- Covacci, A., S. Censini, M. Bugnoli, R. Petracca, D. Burrone, G. Macchia, A. Massone, and *et al.* 1993. Molecular characterization of the 128-kDa immunodominant antigen of *Helicobacter pylori* associated with cytotoxicity and duodenal ulcer. *Proc. Natl. Acad. Sci. USA* 90, 5791-5795.
- Danielli, A., D. Roncarati, I. Delany, V. Chiarini, R. Rappuoli, and V. Scarlato. 2006. *In vivo* dissection of the *Helicobacter pylori* Fur regulatory circuit by genome-wide location analysis. *J. Bacteriol.* 188, 4654-4662.
- Delany, I., G. Spohn, R. Rappuoli, and V. Scarlato. 2001. The Fur repressor controls transcription of iron-activated and -repressed genes in *Helicobacter pylori*. *Mol. Microbiol.* 42, 1297-1309.
- den Besten, H.M., M. Mols, R. Moezelaar, M.H. Zwietering, and T. Abee. 2009. Phenotypic and transcriptomic analyses of mildly and severely salt-stressed *Bacillus cereus* ATCC 14579 cells. *Appl. Environ. Microbiol.* 75, 4111-4119.
- Ernst, F.D., S. Bereswill, B. Waidner, J. Stoof, U. Mader, J.G. Kusters, E.J. Kuipers, M. Kist, A.H. van Vliet, and G. Homuth. 2005a. Transcriptional profiling of *Helicobacter pylori* Fur- and iron-regulated gene expression. *Microbiology* 151, 533-546.
- Ernst, F.D., G. Homuth, J. Stoof, U. Mader, B. Waidner, E.J. Kuipers, M. Kist, J.G. Kusters, S. Bereswill, and A.H. van Vliet. 2005b. Iron-responsive regulation of the *Helicobacter pylori* iron-cofactored superoxide dismutase SodB is mediated by Fur. *J. Bacteriol.* 187, 3687-3692.
- Gancz, H., S. Censini, and D.S. Merrell. 2006. Iron and pH homeostasis intersect at the level of Fur regulation in the gastric pathogen *Helicobacter pylori*. *Infect. Immun.* 74, 602-614.
- Gancz, H., K.R. Jones, and D.S. Merrell. 2008. Sodium chloride affects *Helicobacter pylori* growth and gene expression. *J. Bacteriol.* 190, 4100-4105.
- Hahne, H., U. Mader, A. Otto, F. Bonn, L. Steil, E. Bremer, M. Hecker, and D. Becher. 2010. A comprehensive proteomics and transcriptomics analysis of *Bacillus subtilis* salt stress adaptation. *J. Bacteriol.* 192, 870-882.
- He, Z., A. Zhou, E. Baidoo, Q. He, M.P. Joachimiak, P. Benke, R. Phan, and *et al.* 2010. Global transcriptional, physiological, and metabolite analyses of the responses of *Desulfovibrio vulgaris* hildenborough to salt adaptation. *Appl. Environ. Microbiol.* 76, 1574-1586.
- Hoffmann, T., A. Schutz, M. Brosius, A. Volker, U. Volker, and E. Bremer. 2002. High-salinity-induced iron limitation in *Bacillus subtilis*. *J. Bacteriol.* 184, 718-727.
- Joyce, E.A., J.V. Gilbert, K.A. Eaton, A. Plaut, and A. Wright. 2001. Differential gene expression from two transcriptional units in the *cag* pathogenicity island of *Helicobacter pylori*. *Infect. Immun.* 69,

- 4202-4209.
- Kato, S., T. Tsukamoto, T. Mizoshita, H. Tanaka, T. Kumagai, H. Ota, T. Katsuyama, M. Asaka, and M. Tatematsu. 2006. High salt diets dose-dependently promote gastric chemical carcinogenesis in *Helicobacter pylori*-infected Mongolian gerbils associated with a shift in mucin production from glandular to surface mucous cells. *Int. J. Cancer* 119, 1558-1566.
- Lee, H.W., Y.H. Choe, D.K. Kim, S.Y. Jung, and N.G. Lee. 2004. Proteomic analysis of a ferric uptake regulator mutant of *Helicobacter pylori*, regulation of *Helicobacter pylori* gene expression by ferric uptake regulator and iron. *Proteomics* 4, 2014-2027.
- Loh, J.T., V.J. Torres, and T.L. Cover. 2007. Regulation of *Helicobacter pylori* *cagA* expression in response to salt. *Cancer Res.* 67, 4709-4715.
- Malaty, H.M. 2007. Epidemiology of *Helicobacter pylori* infection. *Best. Pract. Res. Clin. Gastroenterol.* 21, 205-214.
- Matysiak-Budnik, T. and F. Megraud. 2006. *Helicobacter pylori* infection and gastric cancer. *Eur. J. Cancer* 42, 708-716.
- Mbulaiteye, S.M., M. Hisada, and E.M. El-Omar. 2009. *Helicobacter pylori* associated global gastric cancer burden. *Front Biosci.* 14, 1490-1504.
- Miles, S., B.M. Carpenter, H. Gancz, and D.S. Merrell. 2010a. *Helicobacter pylori* *apo-Fur* regulation appears unconserved across species. *J. Microbiol.* 48, 378-386.
- Miles, S., M.B. Piazuelo, C. Semino-Mora, M.K. Washington, A. Dubois, R.M. Peek, Jr., P. Correa, and D.S. Merrell. 2010b. Detailed *in vivo* analysis of the role of *Helicobacter pylori* *Fur* in colonization and disease. *Infect. Immun.* 78, 3073-3082.
- Mukhopadhyay, A., Z. He, E.J. Alm, A.P. Arkin, E.E. Baidoo, S.C. Borglin, W. Chen, and *et al.* 2006. Salt stress in *Desulfovibrio vulgaris* Hildenborough, an integrated genomics approach. *J. Bacteriol.* 188, 4068-4078.
- Pounder, R.E. and D. Ng. 1995. The prevalence of *Helicobacter pylori* infection in different countries. *Aliment. Pharmacol. Ther.* 9, 33-39.
- Rogers, A.B., N.S. Taylor, M.T. Whary, E.D. Stefanich, T.C. Wang, and J.G. Fox. 2005. *Helicobacter pylori* but not high salt induces gastric intraepithelial neoplasia in B6129 mice. *Cancer Res.* 65, 10709-10715.
- Salaun, L., C. Audibert, L.G. Le, C. Burucoa, J.L. Fauchere, and B. Picard. 1998. Panmictic structure of *Helicobacter pylori* demonstrated by the comparative study of six genetic markers. *FEMS Microbiol. Lett.* 161, 231-239.
- Steil, L., T. Hoffmann, I. Budde, U. Volker, and E. Bremer. 2003. Genome-wide transcriptional profiling analysis of adaptation of *Bacillus subtilis* to high salinity. *J. Bacteriol.* 185, 6358-6370.
- Tomb, J.F., O. White, A.R. Kerlavage, R.A. Clayton, G.G. Sutton, R.D. Fleischmann, K.A. Ketchum, and *et al.* 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* 388, 539-547.
- Tsugane, S. and S. Sasazuki. 2007. Diet and the risk of gastric cancer, review of epidemiological evidence. *Gastric. Cancer* 10, 75-83.
- van Vliet, A.H., J. Stoof, S.W. Poppelaars, S. Bereswill, G. Homuth, M. Kist, E.J. Kuipers, and J.G. Kusters. 2003. Differential regulation of amidase- and formamidase-mediated ammonia production by the *Helicobacter pylori* *fur* repressor. *J. Biol. Chem.* 278, 9052-9057.
- Wang, X.Q., P.D. Terry, and H. Yan. 2009. Review of salt consumption and stomach cancer risk, epidemiological and biological evidence. *World J. Gastroenterol.* 15, 2204-2213.
- Zhou, Y.N., W.G. Coleman, Jr., Z. Yang, Y. Yang, N. Hodgson, F. Chen, and D.J. Jin. 2008. Regulation of cell growth during serum starvation and bacterial survival in macrophages by the bifunctional enzyme SpoT in *Helicobacter pylori*. *J. Bacteriol.* 190, 8025-8032.