

REVIEW

Transcriptional Regulatory Elements in Fungal Secondary Metabolism

Wenbing Yin¹ and Nancy P. Keller^{1,2*}

¹Department of Medical Microbiology and Immunology, ²Department of Bacteriology, University of Wisconsin-Madison, WI 53706, USA

(Received January 7, 2011 / Accepted March 15, 2011)

Filamentous fungi produce a variety of secondary metabolites of diverse beneficial and detrimental activities to humankind. The genes required for a given secondary metabolite are typically arranged in a gene cluster. There is considerable evidence that secondary metabolite gene regulation is, in part, by transcriptional control through hierarchical levels of transcriptional regulatory elements involved in secondary metabolite cluster regulation. Identification of elements regulating secondary metabolism could potentially provide a means of increasing production of beneficial metabolites, decreasing production of detrimental metabolites, aid in the identification of 'silent' natural products and also contribute to a broader understanding of molecular mechanisms by which secondary metabolites are produced. This review summarizes regulation of secondary metabolism associated with transcriptional regulatory elements from a broad view as well as the tremendous advances in discovery of cryptic or novel secondary metabolites by genomic mining.

Keywords: transcriptional regulatory elements, velvet complex, Zn(II)₂Cys₆, bZIP, StuA, genome mining

Secondary metabolites (SMs) are a vast array of small molecules produced by microbes and plants. These molecules are not essential for normal growth of an organism but play important roles, for example by functioning as defense compounds (Rohlfis *et al.*, 2007; Rohlfis and Churchill, 2011) or signaling molecules (Böhnert *et al.*, 2004; Dufour and Rao, 2011; Rodrigues *et al.*, 2011) in ecological interactions. Many SM often possess pharmacological activities and have been exploited in medicine as antibiotics, anticancer and anti-infective agents and for a broad range of other applications (Fox and Howlett, 2008a; Osbourn, 2010). Considering this importance, much effort has focused on studying the genetic regulation of SM gene expression. Fungi are among the most prolific SM producers. One significant hallmark of fungal SM biosynthetic genes is that genes for any particular metabolite (e.g. penicillin) are clustered and act as a single genetic locus (Keller *et al.*, 2005; Yu and Keller, 2005). Furthermore, the SM clusters are not randomly distributed in the genome but most often subtelomerically located (Perrin *et al.*, 2007; Palmer and Keller, 2010). These findings have provided a convenient platform to elucidate the regulatory mechanisms of SM biosynthetic gene expression.

The co-regulation of SM biosynthetic genes can be in part explained by transcriptional control through hierarchical levels of transcriptional regulatory elements. One group consists of bonafide transcription factors ranging from pathway specific transcription factors (e.g. AflR regulating the sterigmatocystin (ST)/aflatoxin (AF) SM clusters, (Brown *et al.*, 1996; Fernandes *et al.*, 1998) to broad domain transcription factors (e.g. PacC mediating fungal responses to changes in pH, (Tilburn *et al.*,

1995) and bZip proteins involved in cluster activation [e.g. MeaB (Polley and Caddick, 1996) and RsmA (Shaaban *et al.*, 2010)]. Another class consists of heteromeric protein complexes participating in global regulation of multiple SM clusters coupled with other developmental processes (e.g. the Velvet complex, (Bayram *et al.*, 2008a) and the CCAAT-binding complex AnCF /PENR1 (Brakhage *et al.*, 1999).

In this short review, we focus primarily on these and associated transcriptional regulatory elements of secondary metabolism and how this knowledge, in turn, has led to tremendous advances in genomic mining of fungal secondary metabolites.

Cluster specific regulatory elements

Fungal secondary metabolic gene clusters often contain one or more transcription factors that are required for expression of the genes for the biosynthetic enzymes (Hoffmeister and Keller, 2007; Osbourn, 2010). AF is one of the best-characterized fungal secondary metabolites and, along with the related metabolite ST, among the first compounds found to have all of their biosynthetic genes organized within a gene cluster. The biosynthetic gene clusters were found to include ca. 25 genes arranged in about 70 kb clusters in *A. parasiticus*, *A. flavus*, and *A. nidulans* (Yu *et al.*, 1995; Brown *et al.*, 1996). Two pathway specific regulatory genes, *aflR* and *aflJ*, located divergently adjacent to each other within the AF/ST cluster are involved in the regulation of AF/ST gene expression. These two genes are now referred to as *aflR* and *aflS* in both *A. flavus* and *A. parasiticus*, respectively (Yu *et al.*, 2004).

a) Transcription factors

AflR, encoding a sequence-specific DNA-binding binuclear zinc cluster (Zn(II)₂Cys₆) protein, is required for AF and ST activation by binding to a palindromic sequence, called AflR

* For correspondence. E-mail: npkeller@wisc.edu; Tel: +1-608-262-9795

Table 1. Examples of identified Zn(II)₂Cys₆ transcription factors (TFs) involvement in secondary metabolism in fungi

Protein	Regulated gene cluster	Regulatory mode	Species	Reference
AflR	AF/ST	Positive	<i>A. nidulans</i> , <i>A. parasiticus</i> , <i>A. flavus</i>	Chang <i>et al.</i> (1995b), Brown <i>et al.</i> (1996), Yu <i>et al.</i> (1996), Fernandes <i>et al.</i> (1998), Ehrlich <i>et al.</i> (1999)
GliZ	gliotoxin	Positive	<i>A. fumigatus</i>	Bok <i>et al.</i> (2006a)
SirZ	sirodesmin PL	Positive	<i>Leptosphaeria maculans</i>	Fox <i>et al.</i> (2008)
MlcR	compactin	Positive	<i>Penicillium citrinum</i>	Abe <i>et al.</i> (2002b)
Bik5	bikaverin	Positive	<i>Fusarium fujikuroi</i>	Wiemann <i>et al.</i> (2009)
DEP6	depudecin	Positive	<i>Alternaria brassicicola</i>	Wight <i>et al.</i> (2009)
ZFR1	fumonisin	Positive	<i>Fusarium verticillioides</i>	Flaherty and Woloshuk (2004), Brown <i>et al.</i> (2007)
FUM21				
CTB8	cercosporin	Positive	<i>Cercospora nicotianae</i>	Chen <i>et al.</i> (2007)
GIP2	aurofusarin	Positive	<i>Gibberella zeae</i>	Kim <i>et al.</i> (2006)
CtnA	citrinin	Positive	<i>Monascus purpureus</i>	Shimizu <i>et al.</i> (2007)
LovE	lovastatin	Positive	<i>A. terreus</i>	Kennedy <i>et al.</i> (1999), Huang and Li (2009)
ApdR	aspyridone	Positive	<i>A. nidulans</i>	Bergmann <i>et al.</i> (2007)
CtnR	asperfuranone	Positive	<i>A. nidulans</i>	Chiang <i>et al.</i> (2009b)
MdpE	monodictyphenone/ emodin analogs	Positive	<i>A. nidulans</i>	Chiang <i>et al.</i> (2010)
Cmr1p	melanin	Positive	<i>Colletotrichum lagenarium</i>	Tsuji <i>et al.</i> (2000)
Pig1p	melanin	Positive	<i>Magnaporthe grisea</i>	Tsuji <i>et al.</i> (2000)
MokH	monacolin K	Positive	<i>Monascus pilosus</i>	Chen <i>et al.</i> (2010)

binding motif, 5'-TCG(N₅)GCA. It is found in most promoters of the AF/ST biosynthetic genes in *A. flavus*, *A. parasiticus*, and *A. nidulans* (Brown *et al.*, 1996; Fernandes *et al.*, 1998; Chang, 2003; Georgianna and Payne, 2009). A second binding site, 5'-TTAGGCCTAA, is reported as important in auto-regulation of *aflR* transcript in *A. flavus* and *A. parasiticus* (Chang *et al.*, 1995a). Disruption or mutation of *aflR* abolishes expression of AF/ST cluster genes (Woloshuk *et al.*, 1994; Yu *et al.*, 1996; Cary *et al.*, 2000). Conversely, overexpression of *aflR* in the genome increases biosynthetic gene expression and SM production (Chang *et al.*, 1995b; Bok *et al.*, 2006c). Despite clear differences in the sequence of AflR between *A. nidulans* and *A. flavus*, function is conserved. AflR from *A. flavus* is able to drive expression of the ST cluster in an *A. nidulans aflR* deletion strain (Yu *et al.*, 1996).

To more carefully understand the role of AflR in gene regulation, Price *et al.* (2006) assessed gene expression in wild-type and *aflR* deletion strains of *A. parasiticus* using microarrays. The results indicated that AflR regulates genes not only in the biosynthetic cluster but also outside it. Ehrlich *et al.* (1999) thoroughly examined promoters for AflR binding in eleven different genes from AF cluster by using electrophoretic mobility shift assays (EMSA). All of these genes demonstrate some degree of AflR binding in EMSA assays except three genes: *aflR*, *aflJ*, and *avnA* having sites that deviate from the predicted AflR binding motif. Moreover, all of these genes were differentially expressed between WT and the $\Delta aflR$ mutant, suggesting AflR is required to activate their expression (Price *et al.*, 2006). However, the differences in AF gene expression do not always correspond with the differences in AF production as seen in the comparison (Price *et al.*, 2006). This suggests that there are other regulatory factors modulating AF gene

cluster expression and/or post-transcriptional and translational modifications important in regulation. For example, it is known that AflR is regulated by protein kinase A by phosphorylation (Shimizu *et al.*, 2003).

AflJ (also called AflS) is another regulator which resides next to AflR. The mechanism of AflJ regulation of AF biosynthesis remains elusive although data suggest AflJ may act as an enhancer and it is required for full AflR activity and modulates AF expression through its interaction with AflR (Chang, 2003). Over-expression of *aflJ* in the aflatoxin-producing strains resulted in increased AF (Du *et al.*, 2007). Inactivation or mutation of *aflJ* gives a phenotype similar to an *aflR* deletion, i.e., a great reduction in AF production (Meyers *et al.*, 1998). The dual presence of AflR and AflJ-like proteins is present in several SM clusters such as the *A. nidulans* monodictyphenone cluster where both *mdpA* (a putative *aflJ* homolog) and *mdpE* (a putative *aflR* homolog) are required for activation of *mdp* biosynthetic genes (Chiang *et al.*, 2010). In fact, evolutionary evaluations of the AF biosynthetic cluster suggest *aflR/aflJ(aflS)* exists as a conserved regulatory module in SM clusters (Carbone *et al.*, 2007).

Zinc binuclear (Zn(II)₂Cys₆) proteins like AflR are the most common type of in-cluster pathway regulators and are unique to fungi (Table 1). *A. fumigatus* GliZ is required for expression of the gliotoxin cluster (Bok *et al.*, 2006a). Gliotoxin is an epipolythiodioxopiperazine (ETP) metabolite and has been shown to play a significant role in enabling the virulence of *A. fumigatus* (Fox and Howlett, 2008b). Due to its role in invasive aspergillosis, there has been much focus on the biosynthesis of gliotoxin and its regulation mechanism (Sugui *et al.*, 2007; Schrettl *et al.*, 2010b). The putative gliotoxin (*gli*) cluster spans about 28 kb in the genome of *A. fumigatus*

(Gardiner and Howlett, 2005). Deletion of *gliZ* resulted in the complete inhibition of all gliotoxin cluster gene expression and effective diminution of gliotoxin production (Bok *et al.*, 2006a). Placement of multiple copies of *gliZ* in the genome increased gliotoxin production. SirZ, bearing 57% identity to GliZ, is a Zn(II)₂Cys₆ DNA binding protein present in the ETP sirodesmin cluster in *Leptosphaeria maculans* and is required for sirodesmin gene expression (Fox *et al.*, 2008a). Other characterized SM Zn(II)₂Cys₆ proteins include MclR, which is required for compactin biosynthesis by *Penicillium citrinum* (Abe *et al.*, 2002a), the fumonisin regulators ZFR1 and FUM21 found in *Fusarium verticillioides* (Flaherty and Woloshuk, 2004; Brown *et al.*, 2007). ZFR1, reported as a positive regulator of fumonisin production requiring a cyclin-like (C-Type) gene FCC1 for function, is an example of a regulatory gene found outside of the SM cluster it regulates (Flaherty and Woloshuk, 2004). Little or no fumonisin production in FUM21 deletion mutants also indicates its positive role in the biosynthesis of fumonisin (Brown *et al.*, 2007). CTB8 was identified to regulate the transcription of seven clustered genes responsible for the biosynthesis of cercosporin in *Cercospora nicotianae* (Chen *et al.*, 2007). CtnA is a major activator of citrinin biosynthesis in *Monascus purpureus*. A strain of *M. purpureus* in which *ctnA* was disrupted exhibited a decrease in citrinin production (Shimizu *et al.*, 2007), whereas heterologous expression of an additional copy of *ctnA* in *A. oryzae* results in 400-fold higher citrinin production than that of the parental strain (Sakai *et al.*, 2008). LovE regulates biosynthesis of lovastatin in *A. terreus* (Kennedy *et al.*, 1999; Huang and Li, 2009). GIP2 regulates aurofusarin biosynthetic gene cluster in *Gibberella zeae* (Kim *et al.*, 2006). Zn(II)₂Cys₆ protein MokH is required for monacolin K synthesis (Chen *et al.*, 2010). ApdR activates the biosynthesis of the PKS-NRPS metabolites aspyridones (Bergmann *et al.*, 2007) and CntR is the transcriptional regulator of asperfuranone. As mentioned earlier, *mdpE* (a putative *afIR* homolog) activates monodictyphenone biosynthesis (Chiang *et al.*, 2010). Cmr1p and Pig1p, which regulate melanin synthesis in *Colletotrichum lagenarium* and *Magnaporthe grisea*, respectively, contain a Cys₂His₂ zinc finger motif as well as a Zn(II)₂Cys₆ binuclear cluster motif (Tsuji *et al.*, 2000). Aside from the few mentioned in this review, there are numerous examples of Zn(II)₂Cys₆ proteins in SM clusters which have not yet been genetically or biochemically characterized (e.g. the Zn(II)₂Cys₆ protein in the solanapyrone *Alternaria solani* (Kasahara *et al.*, 2010). Commonly, Zn(II)₂Cys₆ DNA binding domains interact with DNA binding sites consisting of conserved terminal trinucleotides, which are usually in a symmetrical configuration and are spaced by an internal variable sequence of defined length, for example, CCG(N)_xCGG (MacPherson *et al.*, 2006).

Other types of pathway specific transcription factors have been found to be important in cluster transcriptional regulation. These include several Cys₂His₂ zinc finger proteins such as Tri6 and MRTRI6 for trichothecene production in *Fusarium sporotrichioides* and *Myrothecium roridum*, respectively (Trapp *et al.*, 1998; Hohn *et al.*, 1999), and an ankyrin repeat protein ToxE for HC-toxin production in *Cochliobolus carbonum* (Pedley and Walton, 2001). Hohn and coworker determined the Tri6 binding sequence motif to be YNAGGCC in the promoter regions of nine other *tri* genes. Loss of this protein leads to

decreased expression of *tri* cluster genes and accumulation of low levels of the trichothecene precursor trichodiene (Hohn *et al.*, 1999). Tri6 in turn is regulated by the master regulatory gene Tri10 (Tag *et al.*, 2001). Recently, two more putative Cys₂His₂ transcription factors ScpR and AfoA from *A. nidulans* were found to interact with each other where ScpR not only regulated expression of two adjacent NRPS genes *inpA* and *inpB* but also AfoA, the specific regulator of the asperfuranone gene cluster (Bergmann *et al.*, 2010). The proposed product of *inp* cluster is a tripeptide with a reduced C-terminus. The final product of this cluster is unknown. Furthermore, two transcription factors CPCR1 (cephalosporin C regulator 1) and AcFKH1 (forkhead transcription factor 1) required for cephalosporin C production were characterized from *Acremonium chrysogenum* (Schmitt and Kuck, 2000). They are members of subfamilies of winged helix transcription factors (Schmitt *et al.*, 2004; Hoff *et al.*, 2005). The CPCR1 protein belongs to the conserved family of eukaryotic regulatory factor X (RFX) transcription factors. Transcription factor AcFKH1 belongs to the family of forkhead proteins and is characterized by two conserved domains, the forkhead-associated domain (FHA), which might be involved in phospho-protein interactions, and the C-terminal forkhead DNA-binding domain (FKH). CPCR1 and AcFKH1 regulate cephalosporin C biosynthesis by binding to regulatory sequences in the promoter region of the cephalosporin C biosynthesis genes *pcbAB-pcbC* (Schmitt and Kuck, 2000; Schmitt *et al.*, 2004).

b) Alternative in-cluster regulation

In addition to pathway specific transcription factors, other members in SM clusters can affect expression of the remaining cluster genes. For example, the disruption of *gliP*, encoding a multimodular nonribosomal peptide synthetase (NRPS) in the gliotoxin cluster, resulted in the inhibition of expression of other *gli* genes (Sugui *et al.*, 2007). On the other hand, *gliT* encoding a gliotoxin reductase required for self-protection against gliotoxin, is independently regulated from other *gli* genes; that is, it is still expressed in a *A. fumigatus* Δ *gliZ* background (Schrettl *et al.*, 2010b). Another interesting case of in-cluster regulation has been reported in the fungus *Leptosphaeria maculans* (Gardiner *et al.*, 2005). Here, the disruption of a putative ABC transporter, *sirA*, in the sirodesmin cluster resulted in increased expression of *sirP*, the sirodesmin NRPS (Gardiner *et al.*, 2005).

Transcriptional linkage to environmental programs

In addition to pathway-specific regulators, fungal secondary metabolite production is also responsive to general environmental factors called broad domain transcription factors. Carbon and nitrogen sources, temperature, light, and pH are well known parameters which greatly affect metabolite output. Fungi are able to turn the energetically costly process of SM production off and on under certain environmental conditions presumably as an advantage to the producing fungus. Many studies have indicated that these environmental signals influencing SM production are mediated through Cys₂His₂ zinc finger global transcription factors such as CreA for carbon signaling (Dowzer and Kelly, 1989, 1991), AreA for nitrogen signaling (Hynes, 1975), and PacC for pH signaling (Tilburn *et al.*, 1995). There are several reports of all three proteins affecting both negative and positive regulation of gene




Fig. 1. Regulation of fungal secondary metabolism by transcriptional regulatory elements. SM gene clusters are often silent in laboratory conditions. Activation of SM clusters is complex and involves several layers of transcriptional regulators from pathway specific regulators to more global regulators such as bZip proteins, StuA and Velvet (LaeA, VeA, and VelB) and Hap-like complexes. This activation is associated with conversion of heterochromatin marks (silent cluster) to euchromatin marks (active cluster). In addition, environmental stimuli are translated by signal transduction cascades, including MAPK and PkaA, to activate SM synthesis.

clusters.

For example, carbon catabolite repression by the CreA-transcriptional repressor is widespread in filamentous fungi (Dowzer and Kelly, 1989). The best known example is ethanol metabolism in *A. nidulans* (Felenbok *et al.*, 2001). In the ethanol utilization pathway, CreA directly represses the expression of *alcA* and *aldA* encoding alcohol and aldehyde dehydrogenases, respectively, and also directly represses *alcR*, encoding a transcriptional activator required for *alcA* and *aldA* expression (Felenbok *et al.*, 2001). Although penicillin production is carbon catabolite repressed (Brakhage *et al.*, 1992), extensive investigations in *A. nidulans* support only a minimal role for negative regulation by CreA in this species (Espeso and Penalva, 1992; Espeso *et al.*, 1993). AreA is a GATA transcription factor that mediates nitrogen metabolite repression in response to NH_4^+ or glutamine (Wilson and Arst, 1998). Its binding sites are found in promoter regions throughout the AF biosynthetic cluster in *A. parasiticus*. This indicates that AreA may play a role in the regulation of AF biosynthesis by nitrogen (Chang *et al.*, 2000). More concrete data comes from studies in *Gibberella fujikuroi* where AreA is directly involved in positively regulating gibberellin (GA) gene expression (Tudzynski *et al.*, 1998; Mihlan *et al.*, 2003). PacC is a zinc finger transcription factor that regulates gene expression depending on ambient pH (Tilburn *et al.*, 1995). Penicillin production is positively regulated by PacC but it appears to negatively regulate ST gene expression (Espeso *et al.*, 1993; Keller *et al.*, 1997; Martin, 2000). Additionally, transcriptional studies have shown that ochratoxin A biosynthetic genes are down regulated in alkali pH, possibly an indication of PacC regulation (O'Callaghan *et al.*, 2006). Some evidence also exists for possible negative regulation of fumonisin by PacC in *F. verticillioides* (Flaherty *et al.*, 2003). Sometimes SM clusters are regulated by two or more of these broad domain factors such as the case of bikaverin in *F. fujikuroi* where the *bik* gene cluster is repressed by both AreA and PacC (Wiemann *et al.*, 2009).

Transcriptional complexes

a) Velvet complex

In fungi, SM production is regulated not only by pathway specific transcription factors or broad domain transcription factors but also heteromeric complexes encoded by genes unlinked to the SM biosynthetic gene clusters. A significant advance in our understanding of SM cluster regulation was the finding of a conserved fungal complex known as the Velvet complex linking light regulation of both secondary metabolism and sexual development (Bayram *et al.*, 2008a) (Fig. 1). This heterotrimeric complex is composed of at least three proteins, LaeA, VeA, and VelB. Of these, VeA has been known for several decades but it was LaeA that first captured the attention of the SM community.

LaeA (loss of *aflR* expression-A) was first identified in *A. nidulans* through a genetic mutagenesis strategy to find regulators of AflR (Butchko *et al.*, 1999; Bok and Keller, 2004). The conserved LaeA sequence led quickly to identification of orthologs in other Aspergilli including *A. terreus*, *A. fumigatus* (Bok and Keller, 2004) and subsequently *A. flavus* (Kale *et al.*, 2008). Overexpression of *laeA* greatly increases penicillin and lovastatin production in *A. nidulans* and *A. terreus* respectively (Bok and Keller, 2004). Loss of LaeA results in inactivation of numerous secondary metabolite gene clusters, including ST, penicillin, and terrequinone A production in *A. nidulans*, lovastatin production in *A. terreus*, AF and aflatoxin production in *A. flavus* and gliotoxin production in *A. fumigatus* (Keller *et al.*, 2005; Bok *et al.*, 2006c; Kale *et al.*, 2008). Deletion of *laeA* in *A. fumigatus* decreases the virulence of this human pathogen (Bok *et al.*, 2005; Dagenais and Keller, 2009). A similar effect was observed in *A. flavus laeA* deletants which are reduced in virulence on host seed (Amaike and Keller, 2009). LaeA is also used as a genomic mining tool providing a novel method for identifying new SMs (Bok *et al.*, 2006b) including terrequinone A (Bouhired *et al.*, 2007). Microarray analysis of wild-type and $\Delta laeA$ strains of *A. fumigatus* indicated that LaeA controls up to 9.5% of the transcriptome and up to 13 of its 22 secondary metabolite

gene clusters (Perrin *et al.*, 2007). LaeA orthologs have since been identified in other filamentous fungi. For example, PclaeA was found from *P. chrysogenum* where it controls penicillin biosynthesis, pigmentation and sporulation (Kosalkova *et al.*, 2009), *P. citrinum* where it is proposed to regulate mevastatin synthesis (Xing *et al.*, 2010), *M. pilosus* where it is associated with monicolin J production (Zhang and Miyake, 2009) and *Fusarium fujikuroi* FflaeA1 where it both positively (GAs, fumonisins and fusarin C) and negatively (bikaverin) regulates secondary metabolism (Wiemann *et al.*, 2010). It has now been shown that all *laeA* mutants are also aberrant in sporulation as well as secondary metabolism (Kosalkova *et al.*, 2009; Dagenais *et al.*, 2010; Shaaban *et al.*, 2010; Wiemann *et al.*, 2010).

LaeA was found to interact with VeA and VelB in 2008 (Bayram *et al.*, 2008a) and, as a complex, this heterotrimeric unit, called the Velvet complex, jointly links SM production to sporulation processes (Fig. 1). Although little is known about VelB function, VeA, also called *velvet*, was first described in 1965 (Kafer, 1965) and early on associated with differentiation of asexual and sexual development. Recently, VeA has been found to function as a key global metabolic regulator in the biosynthesis of many SMs concomitant with sexual development (Calvo, 2008). Kato *et al.* (2003) found that *A. nidulans veA* regulates the expression of ST and penicillin genes. *veA* is necessary for activation of the ST-specific transcription factor *afIR* expression leading to the production of this mycotoxin (Calvo *et al.*, 2004). The deletion of *veA* decreased production of numerous SMs (Kato *et al.*, 2003; Dreyer *et al.*, 2007; Myung *et al.*, 2009). Similarly, *A. parasiticus* and *A. flavus veA* genes were necessary for the transcription of both *afIR* and *afIJ* (Calvo, 2008). Furthermore, VeA can activate production of other secondary metabolites including cyclopiazonic acid and aflatoxin in *A. flavus* (Duran *et al.*, 2007; Amaike and Keller, 2009). Interestingly, although VeA is required for penicillin biosynthesis, over-expression of VeA represses expression of penicillin synthesis genes (Brakhage *et al.*, 2004). Chanda *et al.* (2009) reported a novel role of VeA in *A. parasiticus* where it mediates not only AF gene expression (shown earlier in Kato *et al.*, 2003) but also accumulation of vesicles known as aflatoxisomes which contain AF and AF precursors. Orthologs of VeA have been found in other fungal genera, such as FvVe1 in the plant pathogen *F. verticillioides* (Li *et al.*, 2006), FfVe1 from *F. fujikuroi* (Wiemann *et al.*, 2010), AcveA in *Acremonium chrysogenum* (Dreyer *et al.*, 2007), *velA* in *Trichoderma virens* (Mukherjee and Kenerley, 2010) and *veA-1* in *N. crassa* (Bayram *et al.*, 2008b) and MVE1 in *Mycosphaerella graminicola* (Choi and Goodwin, 2011).

It is now clear that the overlapping properties of *laeA* and *veA* mutations are due to the formation of the Velvet complex (Bayram *et al.*, 2008a). Both proteins are members of this heterotrimeric velvet complex (also composed of VelB) which connects light-responding development with secondary metabolism. In light conditions, nuclear levels of VeA are low, asexual development (asexual sporulation) is induced and genes involved in ST biosynthesis minimally expressed. By contrast, in dark conditions VeA nuclear levels are high and this protein acts as a bridge between LaeA and VelB to form a heterotrimeric complex that triggers induction of genes in-

involved in SM and sexual development (Bayram *et al.*, 2008a) (Fig. 1). Although it is unknown how the Velvet complex identifies, binds to and/or activates SM gene clusters, biochemical analysis of *laeA* and heterochromatin mutants (e.g. histone deacetylase and histone methyltransferase mutants) in *A. nidulans* reveal SM gene expression is associated with removal of heterochromatin marks, which in part require LaeA activity (Bok *et al.*, 2009; Palmer and Keller, 2010; Reyes-Dominguez *et al.*, 2010). The Velvet complex has recently been characterized in other fungal species such as *F. fujikuroi* (Wiemann *et al.*, 2010) and *P. chrysogenum* (Hoff *et al.*, 2010). In all genera examined so far (*Aspergillus*, *Penicillium*, and *Fusarium*), the Velvet complex couples sporulation processes with SM. Examination of the *F. fujikuroi* Velvet-like complex revealed that it can simultaneously act as a positive (GAs, fumonisins and fusarin C) or negative (bikaverin) regulator of secondary metabolism.

b) HAP-like CCAAT-binding complex

Another important regulatory complex is the HAP-like CCAAT-binding complex which is evolutionary conserved in eukaryotic organisms (Kato, 2005). Different names have been designated for the homologous complexes in various eukaryotic organisms: Hap in *Saccharomyces cerevisiae* (Pinkham and Guarente, 1985; McNabb *et al.*, 1995), *Kluyveromyces lactis* (Mulder *et al.*, 1994), and *Arabidopsis thaliana* (Edwards *et al.*, 1998), AnCF/PENR1 in *Aspergillus* species (Brakhage *et al.*, 1999), CBF in *Xenopus laevis*, and NF-Y in mammals (Vanhuijsduijnen *et al.*, 1990; Li *et al.*, 1998), respectively. The *A. nidulans* CCAAT-binding factor (AnCF, formerly called PENR1) consists of the subunits HapB, HapC, and HapE. All of the three subunits are necessary for DNA binding. AnCF can activate the expression of numerous genes including the anabolic penicillin biosynthesis genes *ipnA* and *aatA* and is necessary for penicillin production in a number of fungal species (Bergh *et al.*, 1996; Litzka *et al.*, 1996, 1998). Like the Velvet complex, AnCF-mediated repression of gene expression has also been described, e.g. the homoaconitase-encoding *lysF* and the *hapB* gene are repressed by AnCF (Steidl *et al.*, 2001; Weidner *et al.*, 2001). In addition, there is growing evidence that the expression of certain CCAAT-regulated genes requires additional proteins to complex with AnCF. For example, in *A. nidulans*, AnCF represses the transcription of genes that encode iron-containing proteins during iron-replete growth; and during iron-depleted conditions it interacts with the iron-sensing bZIP protein HapX (Hortschansky *et al.*, 2007). The HapX ortholog in *A. fumigatus* is required for full virulence, in part as it regulates siderophore and ribotoxin AspF1 production (Schrettl *et al.*, 2010a).

c) bZIPs

The finding that the bZIP factor HapX is necessary for siderophore regulation by AnCF is intriguing, especially considering two recent findings of two other bZIP proteins involved in SM. Basic leucine zipper (bZIP) proteins are transcription factors that contain a basic region mediating sequence specific DNA-binding which is followed by a leucine zipper region required for bZIP dimerization. In fungi, bZIP transcription factors regulate stress responses, development and morphology as well as multiple metabolic processes.

MeaB, a bZIP transcription factor, was first characterized in *A. nidulans* as affecting nitrogen metabolite repression

(Polley and Caddick, 1996b). Later MeaB was found to tie into AreA function by transcriptionally activating NmrA, an AreA repressor (Wong *et al.*, 2007). As noted above, AreA is associated with regulation of several SM. Wong *et al.* (2007) showed that MeaB binds the promoter region of *nmrA*, TTGCACCAT, and activates its expression in both nitrogen-sufficient and starved conditions. A direct role for MeaB in regulating a SM cluster comes from work in *F. fujikuroi* where it is negatively associated with the production of bikaverin and GAs in *F. fujikuroi* (Wagner *et al.*, 2010). Deletion of MeaB increases the expression of nitrogen-regulated genes under nitrogen-limiting conditions, such as the GA and bikaverin biosynthesis genes (Wagner *et al.*, 2010). MeaB expression also appears to impact SM expression in *A. nidulans* (Amaike, Yin and Keller, unpublished data).

Recently, a putative bZIP protein, RsmA (remediation of secondary metabolism), was identified by using a multicopy-suppressor genetics approach in *A. nidulans* (Shaaban *et al.*, 2010). Overexpression of *rsmA* partially remediates secondary metabolism in $\Delta laeA$ and ΔveA backgrounds and greatly increases SM in strains of *A. nidulans* with an intact Velvet complex. The finding that both RsmA and MeaB affect expression of multiple metabolites suggests that like Velvet and AnCF, bZIP proteins may be involved in global SM regulation (Shaaban *et al.*, 2010).

d) StuA

The complexes described above also impact sporulation processes of the fungi; it is not surprising therefore to find that transcriptional activators first identified as sporulation regulators affect SM. In particular, the transcriptional activator StuA, first identified in *A. nidulans* as a spatial modifier of conidiphore morphology (Miller *et al.*, 1991), has now been associated with concomitant regulation of SM in several fungal genera. StuA orthologs associated with both developmental processes and SM activation include those in *A. fumigatus* (Sheppard *et al.*, 2005; Twumasi-Boateng *et al.*, 2009), *P. chrysogenum* (Sigl *et al.*, 2010), *Stagonospora nodorum* (Ipcho *et al.*, 2010), and *Fusarium graminearum* (Lysoe *et al.*, 2011). StuA activation is closely allied with BrlA activity, another transcription factor first identified in the regulation of *A. nidulans* conidiphore formation (Adams *et al.*, 1988). In the above studies, BrlA has been found to be important in a subset of StuA regulated SM clusters in *A. fumigatus* but not *P. chrysogenum* (Twumasi-Boateng *et al.*, 2009; Sigl *et al.*, 2011).

Genome mining

With the identification of pathway-specific transcription factors, global transcription regulators as well as transcriptional complexes coupled with available genomic sequences, several studies have recently focused on application of this knowledge to mine genomes for SM. Many promising approaches based on chimeric regulation have been developed for the discovery of SMs, principally using variations of microarray analysis as the basis of genomic mining and/or engineering silent SM cluster expression through chimeric promoters, inter-organismal confrontations or chromatin remodeling (Cichewicz, 2010; Brakhage and Schroeckh, 2011).

a) Microarray approach

More and more available fungal genome sequences make it possible to analyze SM-regulating gene clusters at a genomic

level using microarray approach. LaeA-based genomic mining of microarrays analysis from *A. nidulans* $\Delta laeA$ mutants showed that many SM gene clusters are down-regulated containing the best-characterized fungal SM gene clusters ST and PN clusters as well as some indole alkaloid biosynthesis (Bok *et al.*, 2006b). Conversely, microarray analysis of an overexpression *laeA* strain identified many up-regulated SM clusters. This led to the identification of the anti-tumor compound terrequinone A (Bok *et al.*, 2006b). Consistently, genomic profiling of *laeA* mutants in *A. fumigatus* showed similar regulation effect on SM gene clusters (Perrin *et al.*, 2007). Microarray data revealed that 13 SM gene clusters was influenced by LaeA. Ten of them with a majority of genes within these clusters are significantly down-regulated in $\Delta laeA$ background. In addition, three additional clusters have at least one gene encoding a critical enzyme such as a nonribosomal peptide synthetase (NRPS) or a polyketide synthase showing decreased expression in $\Delta laeA$ strain (Perrin *et al.*, 2007). Microarray analysis of *A. flavus* $\Delta laeA$ and *OE::laeA* strains revealed LaeA to regulate up to 50% of the 55 putative SM clusters in this species (Georgianna *et al.*, 2010). Furthermore, microarray technology was employed for the first time to monitor the selective induction of silent fungal SM genes through bacterial-fungal interactions (Schroeckh *et al.*, 2009). Very recently, a functional genomic profiling of *A. fumigatus* biofilm was developed. From proteome and transcriptome studies, genes and proteins involved in the biosynthesis of the mycotoxin gliotoxin were upregulated in biofilm-grown cultures (Bruns *et al.*, 2010).

b) Promoter activation

As an inducible promoter, the *alcA* promoter (*alcA(p)*) has been broadly applied to activate cryptic gene clusters (Scherlach and Hertweck, 2009). Bergmann *et al.* (2007) found a silent hybrid PKS-NRPS gene cluster in the genome of *A. nidulans* and activated the cluster by driving the cluster transcription factor *apdR* with the *alcA* promoter. This resulted in the production of aspyrones. Northern blot analyses revealed that five other predicted pathway genes (*apdB*, *apdC*, *apdD*, *apdE*, and *apdG*) are only expressed upon the *alcA* induction condition (Bergmann *et al.*, 2007). By replacing the original promoter of the regulatory gene *ctnR* with the *alcA* promoter, Chiang *et al.* (2009b) activated a cryptic gene cluster containing two polyketide synthases (PKS) for asperfuranone biosynthesis. Another silent gene cluster containing two NRPS genes *inpA* and *inpB* was induced by the chimeric *alcA(p)-scpR* construct (Bergmann *et al.*, 2010). The proposed product of *inp* cluster is a tripeptide with a reduced C-terminus. As mentioned earlier, the *alcA(p)-scpR* construct also activated the asperfuranone cluster (Bergmann *et al.*, 2010). Similarly, the *mdp* gene cluster for monodictyphenone biosynthesis was activated by expression of regulatory gene *mdpE* with *alcA(p)* (Chiang *et al.*, 2010).

gdpA(p), a strong constitutive promoter, firstly characterized in *A. nidulans* (Punt *et al.*, 1990) has also been used to increase or activate SM genes. Flaherty and Payne (1997) examined the effects of *aflR* overexpression with *gpdA* promoter on AF production, and expression of AF biosynthetic genes in *A. flavus*. The results showed that AF production greatly increased in *gpdA(p)::aflR* strain and transcriptional activities of AF pathway genes *nor-1* and *pksA* also increased dramatically. Chen *et al.* (2010) expressed the monacolin K transcrip-

tion factor *mokH* with the *gpdA(p)* promoter in *M. pilosus*. The transcripts of *mokH* in the transformants were expressed at a much higher level than those of the wild-type strain. Consistently, the production of monacolin K in the transformant showed a 1.7-fold higher than the wild-type strain (Chen *et al.*, 2010). Similarly, expression of the bZip protein RsmA under *gpdA(p)* control conditions greatly increases ST production in *A. nidulans* (Shaaban *et al.*, 2010).

c) Fungus-microbe confrontation

Fungi and bacteria co-inhabit a wide variety of environments and their interactions constitute a driving force for SM production (Daniel, 2004). Simulating these habitats by culturing two or more strains together should be a rational way to obtain new SM through their cross-talk. Co-cultivation of two marine microorganisms (Oh *et al.*, 2007), an *Emericella* sp. tentatively identified as *Emericella parvathecica* and the actinomycete *Salinispora arenicola* greatly enhanced the expression of the emericellamides biosynthesis gene cluster (recently identified in *A. nidulans* by Chiang *et al.* (2008)). In addition, Schroeckh *et al.* (2009) co-cultivated *A. nidulans* with a collection of 58 actinomycetes. Using microarray analyses of both *Aspergillus* secondary metabolism and full-genome arrays, in combination with Northern blot and quantitative RT-PCR analyses, they identified a bacterium, *Streptomyces rapamycinicus* that selectively triggered the expression of silent gene clusters under standard laboratory conditions. Another interesting example is identification of biosynthetic gene cluster of mutanobactin A from the oral pathogen *Streptococcus mutans*. Mutanobactin A is capable of suppressing the morphological transition of *Candida albicans* from yeast to mycelium under co-culture conditions (Joyner *et al.*, 2010). Losada *et al.* (2009) demonstrated that co-culture of different *Aspergillus* spp. together resulted in the up-regulation of different SM than when each species was grown alone. Consequently, the concept of interspecies cross-talk leading to regulatory complex and chemical diversity provides a conceptual framework for the elucidation of secondary metabolites and the understanding their cross-talk regulation.

Conclusions

Filamentous fungi are well known prolific producers of low-molecular weight molecules with various biological activities (Hoffmeister and Keller, 2007). However, bioinformatic analyses of the sequenced fungal genomes indicate that secondary metabolism gene clusters are silent under standard laboratory conditions and it appears that many "cryptic" SM await discovery (Hertweck, 2009). In order to customize the biosynthesis of natural products or activate silent gene clusters we must understand how SMs are regulated. Undoubtedly, the research of transcriptional regulatory elements and their functional mechanism plays a major role in secondary metabolism. This review provides a broad view of our understanding of SM regulation mechanism from transcriptional regulatory elements and cluster specific regulatory elements to global regulators and transcription complexes. This would be very helpful to the development of new approaches for discovering new natural products. In combination with genomic mining and microarray approaches, genome sequence availability has led to rapid progress in understanding fungal secondary metabo-

lism regulation. Many successful approaches in the basis of understanding SM gene regulation have been developed to activate silent or cryptic genes (Chiang *et al.*, 2009a; Brakhage and Schroeckh, 2011). These include overexpression of transcription factors or other pleiotropic regulators, promoter exchange and simulation of the natural habitat by co-cultivation of microorganisms. A major future goal is to investigate the molecular mechanisms that interconvert the transcriptional regulatory elements into active and inactive forms and yield unexpected discoveries and significant new insights.

Acknowledgements

The project was supported by PO1GM084077 from the National Institute of General Medical Sciences to N.P.K.

References

- Abe, Y., C. Ono, M. Hosobuchi, and H. Yoshikawa. 2002a. Functional analysis of *mlcR*, a regulatory gene for ML-236B (compactin) biosynthesis in *Penicillium citrinum*. *Mol. Genet. Genomics* 268, 352-361.
- Abe, Y., T. Suzuki, C. Ono, K. Iwamoto, M. Hosobuchi, and H. Yoshikawa. 2002b. Molecular cloning and characterization of an ML-236B (compactin) biosynthetic gene cluster in *Penicillium citrinum*. *Mol. Genet. Genomics* 267, 636-646.
- Adams, T.H., M.T. Boylan, and W.E. Timberlake. 1988. BrIA is necessary and sufficient to direct conidiophore development in *Aspergillus nidulans*. *Cell* 54, 353-362.
- Amaike, S. and N.P. Keller. 2009. Distinct roles for VeA and LaeA in development and pathogenesis of *Aspergillus flavus*. *Eukaryot. Cell* 8, 1051-1060.
- Bayram, O., S. Krappmann, M. Ni, J.W. Bok, K. Helmstaedt, O. Valerius, S. Braus-Stromeier, and *et al.* 2008a. VelB/VeA/LaeA complex coordinates light signal with fungal development and secondary metabolism. *Science* 320, 1504-1506.
- Bayram, O., S. Krappmann, S. Seiler, N. Vogt, and G.H. Braus. 2008b. *Neurospora crassa* ve-1 affects asexual conidiation. *Fungal Genet. Biol.* 45, 127-138.
- Bergh, K.T., O. Litzka, and A.A. Brakhage. 1996. Identification of a major cis-acting DNA element controlling the bidirectionally transcribed penicillin biosynthesis genes *acvA* (*pcbAB*) and *ipnA* (*pcbC*) of *Aspergillus nidulans*. *J. Bacteriol.* 178, 3908-3916.
- Bergmann, S., A.N. Funk, K. Scherlach, V. Schroeckh, E. Shelest, U. Horn, C. Hertweck, and A.A. Brakhage. 2010. Activation of a silent fungal polyketide biosynthesis pathway through regulatory cross talk with a cryptic nonribosomal peptide synthetase gene cluster. *Appl. Environ. Microbiol.* 76, 8143-8149.
- Bergmann, S., J. Schumann, K. Scherlach, C. Lange, A.A. Brakhage, and C. Hertweck. 2007. Genomics-driven discovery of PKS-NRPS hybrid metabolites from *Aspergillus nidulans*. *Nat. Chem. Biol.* 3, 213-217.
- Böhnert, H.U., I. Fudal, W. Diod, D. Tharreau, J.L. Notteghem, and M.H. Lebrun. 2004. A putative polyketide synthase peptide synthetase from *Magnaporthe grisea* signals pathogen attack to resistant rice. *Plant Cell* 16, 2499-2513.
- Bok, J.W., S.A. Balajee, K.A. Marr, D. Andes, K.F. Nielsen, J.C. Frisvad, and N.P. Keller. 2005. LaeA, a regulator of morphogenetic fungal virulence factors. *Eukaryot. Cell* 4, 1574-1582.
- Bok, J.W., Y.M. Chiang, E. Szweczyk, Y. Reyes-Dominguez, A.D. Davidson, J.F. Sanchez, H.C. Lo, and *et al.* 2009. Chromatin-level regulation of biosynthetic gene clusters. *Nat. Chem. Biol.* 5, 462-464.
- Bok, J.W., D. Chung, S.A. Balajee, K.A. Marr, D. Andes, K.F. Nielsen, J.C. Frisvad, K.A. Kirby, and N.P. Keller. 2006a. GliZ, a transcriptional regulator of gliotoxin biosynthesis, contributes to

- Aspergillus fumigatus* virulence. *Infect. Immun.* 74, 6761-6768.
- Bok, J.W., D. Hoffmeister, L.A. Maggio-Hall, R. Murillo, J.D. Glasner, and N.P. Keller. 2006b. Genomic mining for *Aspergillus* natural products. *Chem. Biol.* 13, 31-37.
- Bok, J.W. and N.P. Keller. 2004. LaeA, a regulator of secondary metabolism in *Aspergillus* spp. *Eukaryot. Cell.* 3, 527-535.
- Bok, J.W., D. Noordermeer, S.P. Kale, and N.P. Keller. 2006c. Secondary metabolic gene cluster silencing in *Aspergillus nidulans*. *Mol. Microbiol.* 61, 1636-1645.
- Bouhired, S., M. Weber, A. Kempf-Sontag, N.P. Keller, and D. Hoffmeister. 2007. Accurate prediction of the *Aspergillus nidulans* terrequinone gene cluster boundaries using the transcriptional regulator LaeA. *Fungal Genet. Biol.* 44, 1134-1145.
- Brakhage, A.A., A. Andrianopoulos, M. Kato, S. Steidl, M.A. Davis, N. Tsukagoshi, and M.J. Hynes. 1999. HAP-Like CCAAT-binding complexes in filamentous fungi: implications for biotechnology. *Fungal Genet. Biol.* 27, 243-252.
- Brakhage, A.A., P. Browne, and G. Turner. 1992. Regulation of *Aspergillus nidulans* penicillin biosynthesis and penicillin biosynthesis genes *acvA* and *ipnA* by glucose. *J. Bacteriol.* 174, 3789-3799.
- Brakhage, A.A. and V. Schroeckh. 2011. Fungal secondary metabolites - Strategies to activate silent gene clusters. *Fungal Genet. Biol.* 48, 15-22.
- Brakhage, A.A., P. Sprote, Q. Al-Abdallah, A. Gehrke, H. Plattner, and A. Tuncher. 2004. Regulation of penicillin biosynthesis in filamentous fungi. *Adv. Biochem. Eng. Biotechnol.* 88, 45-90.
- Brown, D.W., R.A.E. Butchko, M. Busman, and R.H. Proctor. 2007. The *Fusarium verticillioides* FUM gene cluster encodes a Zn(II)₂-Cys₆ protein that affects FUM gene expression and fumonisin production. *Eukaryot. Cell.* 6, 1210-1218.
- Brown, D.W., J.H. Yu, H.S. Kelkar, M. Fernandes, T.C. Nesbitt, N.P. Keller, T.H. Adams, and T.J. Leonard. 1996. Twenty-five coregulated transcripts define a sterigmatocystin gene cluster in *Aspergillus nidulans*. *Proc. Natl. Acad. Sci. USA* 93, 1418-1422.
- Bruns, S., M. Seidler, D. Albrecht, S. Salvenmoser, N. Remme, C. Hertweck, A.A. Brakhage, O. Kniemeyer, and F.M.C. Muller. 2010. Functional genomic profiling of *Aspergillus fumigatus* biofilm reveals enhanced production of the mycotoxin gliotoxin. *Proteomics* 10, 3097-3107.
- Butchko, R.A.E., T.H. Adams, and N.P. Keller. 1999. *Aspergillus nidulans* mutants defective in *stc* gene cluster regulation. *Genetics* 153, 715-720.
- Calvo, A.M. 2008. The VeA regulatory system and its role in morphological and chemical development in fungi. *Fungal Genet. Biol.* 45, 1053-1061.
- Calvo, A.M., J. Bok, W. Brooks, and N.P. Keller. 2004. VeA is required for toxin and sclerotial production in *Aspergillus parasiticus*. *Appl. Environ. Microbiol.* 70, 4733-4739.
- Carbone, I., J.H. Ramirez-Prado, J.L. Jakobek, and B.W. Horn. 2007. Gene duplication, modularity and adaptation in the evolution of the aflatoxin gene cluster. *BMC Evol. Biol.* 9, 111.
- Cary, J.W., K.C. Ehrlich, M. Wright, P.K. Chang, and D. Bhatnagar. 2000. Generation of *afIR* disruption mutants of *Aspergillus parasiticus*. *Appl. Microbiol. Biot.* 53, 680-684.
- Chanda, A., L.V. Roze, S. Kang, K.A. Artymovich, G.R. Hicks, N.V. Raikhel, A.M. Calvo, and J.E. Linz. 2009. A key role for vesicles in fungal secondary metabolism. *Proc. Natl. Acad. Sci. USA* 106, 19533-19538.
- Chang, P.K. 2003. The *Aspergillus parasiticus* protein AFLJ interacts with the aflatoxin pathway-specific regulator AFLR. *Mol. Genet. Genomics* 268, 711-719.
- Chang, P.K., D. Bhatnagar, T.E. Cleveland, and J.W. Bennett. 1995a. Sequence variability in homologs of the aflatoxin pathway gene *afIR* distinguishes species in *Aspergillus* section flavi. *Appl. Environ. Microbiol.* 61, 40-43.
- Chang, P.K., K.C. Ehrlich, J.J. Yu, D. Bhatnagar, and T.E. Cleveland. 1995b. Increased expression of *Aspergillus parasiticus* AfIR, encoding a sequence-specific DNA-binding protein, relieves nitrate inhibition of aflatoxin biosynthesis. *Appl. Environ. Microbiol.* 61, 2372-2377.
- Chang, P.K., J. Yu, D. Bhatnagar, and T.E. Cleveland. 2000. Characterization of the *Aspergillus parasiticus* major nitrogen regulatory gene, *areA*. *Biochim. Biophys. Acta.* 1491, 263-266.
- Chen, H.Q., M.H. Lee, M.E. Daub, and K.R. Chung. 2007. Molecular analysis of the cercosporin biosynthetic gene cluster in *Cercospora nicotianae*. *Mol. Microbiol.* 64, 755-770.
- Chen, Y.P., G.F. Yuan, S.Y. Hsieh, Y.S. Lin, W.Y. Wang, L.L. Liaw, and C.P. Tseng. 2010. Identification of the *mokH* gene encoding transcription factor for the upregulation of monacolin K biosynthesis in *Monascus pilosus*. *J. Agr. Food Chem.* 58, 287-293.
- Chiang, Y.M., K.H. Lee, J.F. Sanchez, N.P. Keller, and C.C. Wang. 2009a. Unlocking fungal cryptic natural products. *Nat. Prod. Commun.* 4, 1505-1510.
- Chiang, Y.M., E. Szewczyk, A.D. Davidson, R. Entwistle, N.P. Keller, C.C. Wang, and B.R. Oakley. 2010. Characterization of the *Aspergillus nidulans* monodictyphenone gene cluster. *Appl. Environ. Microbiol.* 76, 2067-2074.
- Chiang, Y.M., E. Szewczyk, A.D. Davidson, N. Keller, B.R. Oakley, and C.C.C. Wang. 2009b. A gene cluster containing two fungal polyketide synthases encodes the biosynthetic pathway for a polyketide, asperuranone, in *Aspergillus nidulans*. *J. Am. Chem. Soc.* 131, 2965-2970.
- Chiang, Y.M., E. Szewczyk, T. Nayak, A.D. Davidson, J.F. Sanchez, H.C. Lo, W.Y. Ho, and *et al.* 2008. Molecular genetic mining of the *Aspergillus* secondary metabolome: Discovery of the emericellamide biosynthetic pathway. *Chem. Biol.* 15, 527-532.
- Choi, Y.E. and S.B. Goodwin. 2011. MVE1 encoding the velvet gene in *Mycosphaerella graminicola* is associated with aerial mycelium formation, melanin biosynthesis, hyphal swelling, and light signaling. *Appl. Environ. Microbiol.* 77, 942-953.
- Cichewicz, R.H. 2010. Epigenome manipulation as a pathway to new natural product scaffolds and their congeners. *Nat. Prod. Rep.* 27, 11-22.
- Dagenais, T.R., S.S. Giles, V. Aimaniana, J.P. Latge, C.M. Hull, and N.P. Keller. 2010. *Aspergillus fumigatus* LaeA-mediated phagocytosis is associated with a decreased hydrophobin layer. *Infect. Immun.* 78, 823-829.
- Dagenais, T.R. and N.P. Keller. 2009. Pathogenesis of *Aspergillus fumigatus* in invasive aspergillosis. *Clin. Microbiol. Rev.* 22, 447-465.
- Daniel, R. 2004. The soil metagenome - a rich resource for the discovery of novel natural products. *Curr. Opin. Biot.* 15, 199-204.
- Dowzer, C.E. and J.M. Kelly. 1989. Cloning of the *creA* gene from *Aspergillus nidulans*: a gene involved in carbon catabolite repression. *Curr. Genet.* 15, 457-459.
- Dowzer, C.E. and J.M. Kelly. 1991. Analysis of the *creA* gene, a regulator of carbon catabolite repression in *Aspergillus nidulans*. *Mol. Cell. Biol.* 11, 5701-5709.
- Dreyer, J., H. Eichhorn, E. Friedlin, H. Kurnsteiner, and U. Kuck. 2007. A homologue of the *Aspergillus* velvet gene regulates both cephalosporin C biosynthesis and hyphal fragmentation in *Acremonium chrysogenum*. *Appl. Environ. Microbiol.* 73, 3412-3422.
- Du, W., G.R. O'Brien, and G.A. Payne. 2007. Function and regulation of *afII* in the accumulation of aflatoxin early pathway intermediate in *Aspergillus flavus*. *Food Addit. Contam.* 24, 1043-1050.
- Dufour, N. and R.P. Rao. 2011. Secondary metabolites and other small molecules as intercellular pathogenic signals. *FEMS Microbiol. Lett.* 314, 10-17.
- Duran, R.M., J.W. Cary, and A.M. Calvo. 2007. Production of cyclopiazonic acid, aflatrem, and aflatoxin by *Aspergillus flavus* is regulated by *veA*, a gene necessary for sclerotial formation. *Appl. Microbiol. Biot.* 73, 1158-1168.
- Edwards, D., J.A.H. Murray, and A.G. Smith. 1998. Multiple genes encoding the conserved CCAAT-box transcription factor complex are expressed in *Arabidopsis*. *Plant Physiol.* 117, 1015-1022.
- Ehrlich, K.C., B.G. Montalbano, and J.W. Cary. 1999. Binding of the C6-zinc cluster protein, AFLR, to the promoters of aflatoxin

- pathway biosynthesis genes in *Aspergillus parasiticus*. *Gene* 230, 249-257.
- Espeso, E.A. and M.A. Penalva. 1992. Carbon catabolite repression can account for the temporal pattern of expression of a penicillin biosynthetic gene in *Aspergillus nidulans*. *Mol. Microbiol.* 6, 1457-1465.
- Espeso, E.A., J. Tilburn, H.N. Arst, and M.A. Penalva. 1993. PH regulation is a major determinant in expression of a fungal penicillin biosynthetic gene. *EMBO J.* 12, 3947-3956.
- Felenbok, B., M. Flipphi, and I. Nikolaev. 2001. Ethanol catabolism in *Aspergillus nidulans*: A model system for studying gene regulation. *Prog. Nucleic Acid Res. Mol. Biol.* 69, 149-204.
- Fernandes, M., N.P. Keller, and T.H. Adams. 1998. Sequence-specific binding by *Aspergillus nidulans* AflR, a C6 zinc cluster protein regulating mycotoxin biosynthesis. *Mol. Microbiol.* 28, 1355-1365.
- Flaherty, J.E. and G.A. Payne. 1997. Overexpression of *aflR* leads to upregulation of pathway gene transcription and increased aflatoxin production in *Aspergillus flavus*. *Appl. Environ. Microbiol.* 63, 3995-4000.
- Flaherty, J.E., A.M. Pirttila, B.H. Bluhm, and C.P. Woloshuk. 2003. PAC1, a pH-regulatory gene from *Fusarium verticillioides*. *Appl. Environ. Microbiol.* 69, 5222-5227.
- Flaherty, J.E. and C.P. Woloshuk. 2004. Regulation of fumonisin biosynthesis in *Fusarium verticillioides* by a zinc binuclear cluster-type gene, ZFR1. *Appl. Environ. Microbiol.* 70, 2653-2659.
- Fox, E.M., D.M. Gardiner, N.P. Keller, and B.J. Howlett. 2008. A Zn(II)₂Cys₆ DNA binding protein regulates the sirodesmin PL biosynthetic gene cluster in *Leptosphaeria maculans*. *Fungal Genet. Biol.* 45, 671-682.
- Fox, E.M. and B.J. Howlett. 2008a. Biosynthetic gene clusters for epipolythiodioxopiperazines in filamentous fungi. *Mycol. Res.* 112, 162-169.
- Fox, E.M. and B.J. Howlett. 2008b. Secondary metabolism: regulation and role in fungal biology. *Curr. Opin. Microbiol.* 11, 481-487.
- Gardiner, D.M. and B.J. Howlett. 2005. Bioinformatic and expression analysis of the putative gliotoxin biosynthetic gene cluster of *Aspergillus fumigatus*. *FEMS Microbiol. Lett.* 248, 241-248.
- Gardiner, D.M., R.S. Jarvis, and B.J. Howlett. 2005. The ABC transporter gene in the sirodesmin biosynthetic gene cluster of *Leptosphaeria maculans* is not essential for sirodesmin production but facilitates self-protection. *Fungal Genet. Biol.* 42, 257-263.
- Georgianna, D.R., N.D. Fedorova, J.L. Burroughs, A.L. Dolezal, J.W. Bok, S. Horowitz-Brown, C.P. Woloshuk, J. Yu, N.P. Keller, and G.A. Payne. 2010. Beyond aflatoxin: four distinct expression patterns and functional roles associated with *Aspergillus flavus* secondary metabolism gene clusters. *Mol. Plant Pathol.* 11, 213-226.
- Georgianna, D.R. and G.A. Payne. 2009. Genetic regulation of aflatoxin biosynthesis: from gene to genome. *Fungal Genet. Biol.* 46, 113-125.
- Hertweck, C. 2009. Hidden biosynthetic treasures brought to light. *Nat. Chem. Biol.* 5, 450-452.
- Hoff, B., J. Kamerewerd, C. Sigl, R. Mitterbauer, I. Zadra, H. Kurnsteiner, and U. Kuck. 2010. Two components of a velvet-like complex control hyphal morphogenesis, conidiophore development, and penicillin biosynthesis in *Penicillium chrysogenum*. *Eukaryot. Cell.* 9, 1236-1250.
- Hoff, B., E.K. Schmitt, and U. Kuck. 2005. CPR1, but not its interacting transcription factor AcFKH1, controls fungal arthrospore formation in *Acremonium chrysogenum*. *Mol. Microbiol.* 56, 1220-1233.
- Hoffmeister, D. and N.P. Keller. 2007. Natural products of filamentous fungi: enzymes, genes, and their regulation. *Nat. Prod. Rep.* 24, 393-416.
- Hohn, T.M., R. Krishna, and R.H. Proctor. 1999. Characterization of a transcriptional activator controlling trichothecene toxin biosynthesis. *Fungal Genet. Biol.* 26, 224-235.
- Hortschansky, P., M. Eisendle, Q. Al-Abdallah, A.D. Schmidt, S. Bergmann, M. Thon, O. Kniemeyer, and *et al.* 2007. Interaction of HapX with the CCAAT-binding complex - a novel mechanism of gene regulation by iron. *EMBO J.* 26, 3157-3168.
- Huang, X. and H.M. Li. 2009. Cloning and bioinformatic analysis of lovastatin biosynthesis regulatory gene *lovE*. *Chinese Med. J.* 122, 1800-1805.
- Hynes, M.J. 1975. Studies on role of *area* gene in regulation of nitrogen catabolism in *Aspergillus nidulans*. *Aust. J. Biol. Sci.* 28, 301-313.
- Ipcho, S.V.S., K.C. Tan, G. Koh, J. Gummer, R.P. Oliver, R.D. Trengove, and P.S. Solomon. 2010. The transcription factor StuA regulates central carbon metabolism, mycotoxin production, and effector gene expression in the wheat pathogen *Stagonospora nodorum*. *Eukaryot. Cell.* 9, 1100-1108.
- Joyner, P.M., J. Liu, Z. Zhang, J. Merritt, F. Qi, and R.H. Cichewicz. 2010. Mutanobactin A from the human oral pathogen *Streptococcus mutans* is a cross-kingdom regulator of the yeast-mycelium transition. *Org. Biomol. Chem.* 8, 5486-5489.
- Kafer, E. 1965. Origins of translocations in *Aspergillus nidulans*. *Genetics* 52, 217-232.
- Kale, S.P., L. Milde, M.K. Trapp, J.C. Frisvad, N.P. Keller, and J.W. Bok. 2008. Requirement of LaeA for secondary metabolism and sclerotial production in *Aspergillus flavus*. *Fungal Genet. Biol.* 45, 1422-1429.
- Kasahara, K., T. Miyamoto, T. Fujimoto, H. Oguri, T. Tokiwano, H. Oikawa, Y. Ebizuka, and I. Fujii. 2010. Solanapyrone synthase, a possible diels-alderase and iterative type I polyketide synthase encoded in a biosynthetic gene cluster from *Alternaria solani*. *Chembiochem.* 11, 1245-1252.
- Kato, M. 2005. An overview of the CCAAT-box binding factor in filamentous fungi: Assembly, nuclear translocation, and transcriptional enhancement. *Biosci. Biotechnol. Biochem.* 69, 663-672.
- Kato, N., W. Brooks, and A.M. Calvo. 2003. The expression of sterigmatocystin and penicillin genes in *Aspergillus nidulans* is controlled by *veA*, a gene required for sexual development. *Eukaryot. Cell.* 2, 1178-1186.
- Keller, N.P., C. Nesbitt, B. Sarr, T.D. Phillips, and G.B. Burow. 1997. pH Regulation of sterigmatocystin and aflatoxin biosynthesis in *Aspergillus* spp. *Phytopathology* 87, 643-648.
- Keller, N.P., G. Turner, and J.W. Bennett. 2005. Fungal secondary metabolism - from biochemistry to genomics. *Nat. Rev. Microbiol.* 3, 937-947.
- Kennedy, J., K. Auclair, S.G. Kendrew, C. Park, J.C. Vederas, and C.R. Hutchinson. 1999. Modulation of polyketide synthase activity by accessory proteins during lovastatin biosynthesis. *Science* 284, 1368-1372.
- Kim, J.E., J.M. Jin, H. Kim, J.C. Kim, S.H. Yun, and Y.W. Lee. 2006. GIP2, a putative transcription factor that regulates the aurofusarin biosynthetic gene cluster in *Gibberella zeae*. *Appl. Environ. Microbiol.* 72, 1645-1652.
- Kosalkova, K., C. Garcia-Estrada, R.V. Ullan, R.P. Godio, R. Feltrer, F. Teijeira, E. Mauriz, and J.F. Martin. 2009. The global regulator LaeA controls penicillin biosynthesis, pigmentation and sporulation, but not roquefortine C synthesis in *Penicillium chrysogenum*. *Biochimie* 91, 214-225.
- Li, Q., M. Herrler, N. Landsberger, N. Kaludov, V.V. Ogryzko, Y. Nakatani, and A.P. Wolffe. 1998. Xenopus NF-Y pre-sets chromatin to potentiate p300 and acetylation-responsive transcription from the Xenopus hsp70 promoter *in vivo*. *EMBO J.* 17, 6300-6315.
- Li, S.J., K. Myung, D. Guse, B. Donkin, R.H. Proctor, W.S. Grayburn, and A.M. Calvo. 2006. FvVE1 regulates filamentous growth, the ratio of microconidia to macroconidia and cell wall formation in *Fusarium verticillioides*. *Mol. Microbiol.* 62, 1418-1432.
- Litzka, O., K.T. Bergh, and A.A. Brakhage. 1996. The *Aspergillus nidulans* penicillin-biosynthesis gene *aat* (*penDE*) is controlled by a CCAAT-containing DNA element. *Eur. J. Biochem.* 238, 675-682.
- Litzka, O., P. Papagiannopoulos, M.A. Davis, M.J. Hynes, and A.A. Brakhage. 1998. The penicillin regulator PENR1 of *Aspergillus nidulans* is a HAP-like transcriptional complex. *Eur. J. Biochem.*

- 251, 758-767.
- Losada, L., O. Ajayi, J.C. Frisvad, J.J. Yu, and W.C. Nierman. 2009. Effect of competition on the production and activity of secondary metabolites in *Aspergillus* species. *Med. Mycol.* 47, S88-S96.
- Lysøe, E., M. Pasquali, A. Breakspear, and H.C. Kistler. 2011. The transcription factor FgStuA influences spore development, pathogenicity, and secondary metabolism in *Fusarium graminearum*. *Mol. Plant Microbe Interact.* 24, 54-67.
- MacPherson, S., M. Larochelle, and B. Turcotte. 2006. A fungal family of transcriptional regulators: The zinc cluster proteins. *Microbiol. Mol. Biol. Rev.* 70, 583-604.
- Martin, J.F. 2000. Molecular control of expression of penicillin biosynthesis genes in fungi: Regulatory proteins interact with a bidirectional promoter region. *J. Bacteriol.* 182, 2355-2362.
- McNabb, D.S., Y.Y. Xing, and L. Guarente. 1995. Cloning of yeast Hap5 - a novel subunit of a heterotrimeric complex required for ccaat binding. *Gene. Dev.* 9, 47-58.
- Meyers, D.M., G. Obrian, W.L. Du, D. Bhatnagar, and G.A. Payne. 1998. Characterization of *aflJ*, a gene required for conversion of pathway intermediates to aflatoxin. *Appl. Environ. Microbiol.* 64, 3713-3717.
- Mihlan, M., V. Homann, T.W.D. Liu, and B. Tudzynski. 2003. AREA directly mediates nitrogen regulation of gibberellin biosynthesis in *Gibberella fujikuroi*, but its activity is not affected by NMR. *Mol. Microbiol.* 47, 975-991.
- Miller, K.Y., T.M. Toennis, T.H. Adams, and B.L. Miller. 1991. Isolation and transcriptional characterization of a morphological modifier - the *Aspergillus nidulans* stunted (*stua*) gene. *Mol. General Genet.* 227, 285-292.
- Mukherjee, P.K. and C.M. Kenerley. 2010. Regulation of morphogenesis and biocontrol properties in *Trichoderma virens* by a VELVET protein, Vel1. *Appl. Environ. Microbiol.* 76, 2345-2352.
- Mulder, W., I.H.J.M. Scholten, R.W. Deboer, and L.A. Grivell. 1994. Sequence of the Hap3 transcription factor of *Kluyveromyces-Lactis* predicts the presence of a novel 4-cysteine zinc-finger motif. *Mol. General Genet.* 245, 96-106.
- Myung, K., S.J. Li, R.A.E. Butchko, M. Busman, R.H. Proctor, H.K. Abbas, and A.M. Calvo. 2009. FvVE1 regulates biosynthesis of the mycotoxins fumonisins and fusarins in *Fusarium verticillioides*. *J. Agr. Food Chem.* 57, 5089-5094.
- O'Callaghan, J., P.C. Stapleton, and A.D.W. Dobson. 2006. Ochratoxin A biosynthetic genes in *Aspergillus ochraceus* are differentially regulated by pH and nutritional stimuli. *Fungal Genet. Biol.* 43, 213-221.
- Oh, D.C., C.A. Kauffman, P.R. Jensen, and W. Fenical. 2007. Induced production of emericellamides A and B from the marine-derived fungus *Emericella* sp. in competing co-culture. *J. Nat. Prod.* 70, 515-520.
- Osborn, A. 2010. Secondary metabolic gene clusters: evolutionary toolkits for chemical innovation. *Trends Genet.* 26, 449-457.
- Palmer, J.M. and N.P. Keller. 2010. Secondary metabolism in fungi: does chromosomal location matter? *Curr. Opin. Microbiol.* 13, 431-436.
- Pedley, K.F. and J.D. Walton. 2001. Regulation of cyclic peptide biosynthesis in a plant pathogenic fungus by a novel transcription factor. *Proc. Natl. Acad. Sci. USA* 98, 14174-14179.
- Perrin, R.M., N.D. Fedorova, J.W. Bok, R.A. Cramer, J.R. Wortman, H.S. Kim, W.C. Nierman, and N.P. Keller. 2007. Transcriptional regulation of chemical diversity in *Aspergillus fumigatus* by LaeA. *PLoS Pathog.* 3, e50.
- Pinkham, J.L. and L. Guarente. 1985. Cloning and molecular analysis of the Hap2 locus - a global regulator of respiratory genes in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 5, 3410-3416.
- Polley, S.D. and M.X. Caddick. 1996. Molecular characterisation of *meaB*, a novel gene affecting nitrogen metabolite repression in *Aspergillus nidulans*. *FEBS Lett.* 388, 200-205.
- Price, M.S., J.J. Yu, W.C. Nierman, H.S. Kim, B. Pritchard, C.A. Jacobus, D. Bhatnagar, T.E. Cleveland, and G.A. Payne. 2006. The aflatoxin pathway regulator AfIR induces gene transcription inside and outside of the aflatoxin biosynthetic cluster. *FEMS Microbiol. Lett.* 255, 275-279.
- Punt, P.J., M.A. Dingemans, A. Kuyvenhoven, R.D.M. Soede, P.H. Pouwels, and C.A.M.J. Vandenbondel. 1990. Functional elements in the promoter region of the *Aspergillus nidulans* *gpdA* gene encoding glyceraldehyde-3-phosphate dehydrogenase. *Gene.* 93, 101-109.
- Reyes-Dominguez, Y., J.W. Bok, H. Berger, E.K. Shwab, A. Basheer, A. Gallmetzer, C. Scazzocchio, N. Keller, and J. Strauss. 2010. Heterochromatic marks are associated with the repression of secondary metabolism clusters in *Aspergillus nidulans*. *Mol. Microbiol.* 76, 1376-1386.
- Rodrigues, A.P., A.S. Carvalho, A.S. Santos, C.N. Alves, J.L. do Nascimento Snr, and E.O. Silva. 2011. Kojic acid, a secondary metabolite from *Aspergillus* sp., acts as an inducer of macrophage activation. *Cell Biol. Int.* 35, 335-343.
- Rohlf, M., M. Albert, N.P. Keller, and F. Kempken. 2007. Secondary chemicals protect mould from fungivory. *Biol. Lett.* 3, 523-525.
- Rohlf, M. and A.C. Churchill. 2011. Fungal secondary metabolites as modulators of interactions with insects and other arthropods. *Fungal Genet. Biol.* 48, 23-34.
- Sakai, K., H. Kinoshita, T. Shimizu, and T. Nihira. 2008. Construction of a citrinin gene cluster expression system in heterologous *Aspergillus oryzae*. *J. Biosci. Bioeng.* 106, 466-472.
- Scherlach, K. and C. Hertweck. 2009. Triggering cryptic natural product biosynthesis in microorganisms. *Org. Biomol. Chem.* 7, 1753-1760.
- Schmitt, E.K., B. Hoff, and U. Kuck. 2004. AcFKH1, a novel member of the forkhead family, associates with the RFX transcription factor CPC1 in the cephalosporin C-producing fungus *Acremonium chrysogenum*. *Gene.* 342, 269-281.
- Schmitt, E.K. and U. Kuck. 2000. The fungal CPC1 protein, which binds specifically to beta-lactam biosynthesis genes, is related to human regulatory factor X transcription factors. *J. Biol. Chem.* 275, 9348-9357.
- Schrettl, M., N. Beckmann, J. Varga, T. Heinekamp, I.D. Jacobsen, C. Jochl, T.A. Moussa, and *et al.* 2010a. HapX-mediated adaptation to iron starvation is crucial for virulence of *Aspergillus fumigatus*. *PLoS Pathog.* 6, e1001124.
- Schrettl, M., S. Carberry, K. Kavanagh, H. Haas, G.W. Jones, J. O'Brien, A. Nolan, J. Stephens, O. Fenelon, and S. Doyle. 2010b. Self-protection against gliotoxin--a component of the gliotoxin biosynthetic cluster, GliT, completely protects *Aspergillus fumigatus* against exogenous gliotoxin. *PLoS Pathog.* 6, e1000952.
- Schroeckh, V., K. Scherlach, H.W. Nutzmann, E. Shelest, W. Schmidt-Heck, J. Schuermann, K. Martin, C. Hertweck, and A.A. Brakhage. 2009. Intimate bacterial-fungal interaction triggers biosynthesis of archetypal polyketides in *Aspergillus nidulans*. *Proc. Natl. Acad. Sci. USA* 106, 14558-14563.
- Shaaban, M.I., J.W. Bok, C. Lauer, and N.P. Keller. 2010. Suppressor mutagenesis identifies a velvet complex mediator of *Aspergillus nidulans* secondary metabolism. *Eukaryot. Cell.* 9, 1816-1824.
- Sheppard, D.C., T. Doedt, L.Y. Chiang, H.S. Kim, D. Chen, W.C. Nierman, and S.G. Filler. 2005. The *Aspergillus fumigatus* StuA protein governs the up-regulation of a discrete transcriptional program during the acquisition of developmental competence. *Mol. Biol. Cell.* 16, 5866-5879.
- Shimizu, K., J. Hicks, T.P. Huang, and N.P. Keller. 2003. Pka, Ras and RGS protein interactions regulate activity of AfIR, a Zn(II)₂Cys₆ transcription factor in *Aspergillus nidulans*. *Genetics* 165, 1095-1104.
- Shimizu, T., H. Kinoshita, and T. Nihira. 2007. Identification and *in vivo* functional analysis by gene disruption of *ctnA*, an activator gene involved in citrinin biosynthesis in *Monascus purpureus*. *Appl. Environ. Microbiol.* 73, 5097-5103.
- Sigl, C., H. Haas, T. Specht, K. Pfaller, H. Kurnsteiner, and I. Zadra. 2011. Among developmental regulators StuA but not BrlA is essential for penicillin V production in *Penicillium chrysogenum*.

- Appl. Environ. Microbiol.* 77, 972-982.
- Steidl, S., M.J. Hynes, and A.A. Brakhage. 2001. The *Aspergillus nidulans* multimeric CCAAT binding complex AnCF is negatively autoregulated via its *hapB* subunit gene. *J. Mol. Biol.* 306, 643-653.
- Sugui, J.A., J. Pardo, Y.C. Chang, A. Muellbacher, K.A. Zarembek, E.M. Galvez, L. Brinster, and *et al.* 2007. Role of *laeA* in the regulation of *alb1*, *gliP*, conidial morphology, and virulence in *Aspergillus fumigatus*. *Eukaryot. Cell.* 6, 1552-1561.
- Tag, A.G., G.F. Garifullina, A.W. Peplow, C. Ake, Jr., T.D. Phillips, T.M. Hohn, and M.N. Beremand. 2001. A novel regulatory gene, *Tri10*, controls trichothecene toxin production and gene expression. *Appl. Environ. Microbiol.* 67, 5294-5302.
- Tilburn, J., S. Sarkar, D.A. Widdick, E.A. Espeso, M. Orejas, J. Mungroo, M.A. Penalva, and H.N. Arst. 1995. The *Aspergillus* Pacc zinc-finger transcription factor mediates regulation of both acid-expressed and alkaline-expressed genes by ambient pH. *EMBO J.* 14, 779-790.
- Trapp, S.C., T.M. Hohn, S. McCormick, and B.B. Jarvis. 1998. Characterization of the gene cluster for biosynthesis of macrocyclic trichothecenes in *Myrothecium roridum*. *Mol. General Genet.* 257, 421-432.
- Tsuji, G., Y. Kenmochi, Y. Takano, J. Sweigard, L. Farrall, I. Furusawa, O. Horino, and Y. Kubo. 2000. Novel fungal transcriptional activators, *Cmr1p* of *Colletotrichum lagenarium* and *pig1p* of *Magnaporthe grisea*, contain Cys₂His₂ zinc finger and Zn(II)₂Cys₆ binuclear cluster DNA-binding motifs and regulate transcription of melanin biosynthesis genes in a developmentally specific manner. *Mol. Microbiol.* 38, 940-954.
- Tudzynski, B., H. Kawaide, and Y. Kamiya. 1998. Gibberellin biosynthesis in *Gibberella fujikuroi*: Cloning and characterization of the copalyl diphosphate synthase gene. *Curr. Genet.* 34, 234-240.
- Twumasi-Boateng, K., Y. Yu, D. Chen, F.N. Gravelat, W.C. Nierman, and D.C. Sheppard. 2009. Transcriptional profiling identifies a role for *brlA* in the response to nitrogen depletion and for *StuA* in the regulation of secondary metabolite clusters in *Aspergillus fumigatus*. *Eukaryot. Cell.* 8, 104-115.
- Vanhuijsduijnen, R.H., X.Y. Li, D. Black, H. Matthes, C. Benoist, and D. Mathis. 1990. Coevolution from yeast to mouse - cDNA cloning of the 2 Nf-y (Cp-1/Cbf) subunits. *EMBO J.* 9, 3119-3127.
- Wagner, D., A. Schmeinck, M. Mos, I.Y. Morozov, M.X. Caddick, and B. Tudzynski. 2010. The bZIP transcription factor *MeaB* mediates nitrogen metabolite repression at specific loci. *Eukaryot. Cell.* 9, 1588-1601.
- Weidner, G., S. Steidl, and A.A. Brakhage. 2001. The *Aspergillus nidulans* homoaconitase gene *lysF* is negatively regulated by the multimeric CCAAT-binding complex AnCF and positively regulated by GATA sites. *Arch. Microbiol.* 175, 122-132.
- Wiemann, P., D.W. Brown, K. Kleigrewe, J.W. Bok, N.P. Keller, H.U. Humpf, and B. Tudzynski. 2010. FfVe1 and FfLae1, components of a velvet-like complex in *Fusarium fujikuroi*, affect differentiation, secondary metabolism and virulence. *Mol. Microbiol.* 77, 972-994.
- Wiemann, P., A. Willmann, M. Straeten, K. Kleigrewe, M. Beyer, H.U. Humpf, and B. Tudzynski. 2009. Biosynthesis of the red pigment bikaverin in *Fusarium fujikuroi*: genes, their function and regulation. *Mol. Microbiol.* 72, 931-946.
- Wight, W.D., K.H. Kim, C.B. Lawrence, and J.D. Walton. 2009. Biosynthesis and role in virulence of the histone deacetylase inhibitor depudecin from *Alternaria brassicicola*. *Mol. Plant Microbe Interact.* 22, 1258-1267.
- Wilson, R.A. and H.N. Arst. 1998. Mutational analysis of *AREA*, a transcriptional activator mediating nitrogen metabolite repression in *Aspergillus nidulans* and a member of the "streetwise" GATA family of transcription factors. *Microbiol. Mol. Biol. Rev.* 62, 586-596.
- Woloshuk, C.P., K.R. Foutz, J.F. Brewer, D. Bhatnagar, T.E. Cleveland, and G.A. Payne. 1994. Molecular characterization of *AflR*, a regulatory locus for aflatoxin biosynthesis. *Appl. Environ. Microbiol.* 60, 2408-2414.
- Wong, K.H., M.J. Hynes, R.B. Todd, and M.A. Davis. 2007. Transcriptional control of *nmrA* by the bZIP transcription factor *MeaB* reveals a new level of nitrogen regulation in *Aspergillus nidulans*. *Mol. Microbiol.* 66, 534-551.
- Xing, W., C. Deng, and C.H. Hu. 2010. Molecular cloning and characterization of the global regulator *LaeA* in *Penicillium citrinum*. *Biotechnol. Lett.* 32, 1733-1737.
- Yu, J.H., R.A. Butchko, M. Fernandes, N.P. Keller, T.J. Leonard, and T.H. Adams. 1996. Conservation of structure and function of the aflatoxin regulatory gene *aflR* from *Aspergillus nidulans* and *A. flavus*. *Curr. Genet.* 29, 549-555.
- Yu, J.J., P.K. Chang, J.W. Cary, M. Wright, D. Bhatnagar, T.E. Cleveland, G.A. Payne, and J.E. Linz. 1995. Comparative mapping of aflatoxin pathway gene clusters in *Aspergillus parasiticus* and *Aspergillus flavus*. *Appl. Environ. Microbiol.* 61, 2365-2371.
- Yu, J.J., P.K. Chang, K.C. Ehrlich, J.W. Cary, D. Bhatnagar, T.E. Cleveland, G.A. Payne, J.E. Linz, C.P. Woloshuk, and J.W. Bennett. 2004. Clustered pathway genes in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* 70, 1253-1262.
- Yu, J.H. and N. Keller. 2005. Regulation of secondary metabolism in filamentous fungi. *Annu. Rev. Phytopathol.* 43, 437-458.
- Zhang, M.Y. and T. Miyake. 2009. Development and media regulate alternative splicing of a methyltransferase pre-mRNA in *Monascus pilosus*. *J. Agr. Food Chem.* 57, 4162-4167.