

Prasanna Datar · Prashant Desai · Evans Coutinho ·
Krishna Iyer

CoMFA and CoMSIA studies of angiotensin (AT1) receptor antagonists

Published online: 8 February 2003
© Springer-Verlag 2003

J Mol Model (2002)
DOI 10.1007/s00894-002-0097-6

Due to an unfortunate error Table 1a was incomplete. The correct table is printed below.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00894-002-0097-6>


P. Datar · P. Desai · E. Coutinho (✉) · K. Iyer
Department of Pharmaceutical Chemistry,
Bombay College of Pharmacy, Kalina, Santacruz (E),
Mumbai 400 098, India
e-mail: evans-im@eth.net
Tel.: +91-22-6126284
Fax: +91-22-6100935

Present address:

P. Desai, Department of Medicinal Chemistry, School of Pharmacy,
University of Mississippi, MI 38677-1848, USA

Table 1a Structure of molecules that form the training and test sets for CoMFA and CoMSIA studies

a. Imidazole derivatives:



ID	R ₁	R ₂	R ₃
1	Cl	CH ₂ OH	
36	H	COOCH ₃	
37	H	COOH	
38	H	COOH	
40	C ₂ H ₅	COOCH ₃	
41	H	COOCH ₃	
51	Cl	COOCH ₃	
52	H	COOH	
53	H	COOCH ₃	