

Peter Politzer · Jane S. Murray · Pat Lane ·
Monica C. Concha · Ping Jin · Zenaida Peralta-Inga

An unusual feature of end-substituted model carbon (6,0) nanotubes

Published online: 27 January 2006
© Springer-Verlag 2006

J Mol Model (2005) 11: 258–264

In the original article, the color portions of Figs. 6 and 7 were interchanged, so that each was with the wrong caption. The correct figures and captions are shown.




Fig. 6 HF/STO-5G electrostatic potential (a) and average local ionization energy (b) on outer surface of open (6,1) $C_{68}H_{13}OH$. The hydroxyl group is at the left end of the tube. Color ranges are the same as in Fig. 5




Fig. 7 HF/STO-5G electrostatic potential (a) and average local ionization energy (b) on outer surface of open (6,0) $C_{72}H_{11}OH$. The hydroxyl group is at the left end of the tube. Color ranges are the same as in Fig. 5

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00894-005-0265-6>

P. Politzer (✉) · J. S. Murray · P. Lane · M. C. Concha · P. Jin ·
Z. Peralta-Inga
Department of Chemistry, University of New Orleans,
New Orleans, LA 70148, USA
e-mail: ppolitze@uno.edu