

## Erratum to: Computational structural analysis of proteins of *Mycobacterium tuberculosis* and a resource for identifying off-targets

Sameer Hassan · Abhimita Debnath ·  
Vasanth Mahalingam · Luke Elizabeth Hanna

Published online: 5 July 2012  
© Springer-Verlag 2012

**Erratum to: J Mol Model**  
**DOI 10.1007/s00894-012-1412-5**

The original version of this article unfortunately contained a mistake. On page 3, line 15 - the sentence “The  $R^2$  statistic can range from  $-1$  to  $1$ , with  $1$  representing perfect positive correlation and  $-1$  representing perfect negative correlation” should be corrected to “The values of  $R$  squared range from  $0$  to  $1$  and the  $R$  correlation can range from  $-1$  to  $1$ , with  $1$  representing perfect positive correlation and  $-1$  representing perfect negative correlation”.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00894-012-1412-5>.

---

S. Hassan · A. Debnath · L. E. Hanna (✉)  
Department of Biomedical Informatics,  
National Institute for Research in Tuberculosis,  
Chetpet, Chennai, India  
e-mail: [hanna@trcchennai.in](mailto:hanna@trcchennai.in)

V. Mahalingam  
Department of Statistics,  
National Institute for Research in Tuberculosis,  
Chetpet, Chennai, India

L. E. Hanna  
Department of Clinical Research,  
National Institute for Research in Tuberculosis,  
Chetpet, Chennai 600031, India