

# A theoretical study on the halogen bonding interactions of C<sub>6</sub>F<sub>5</sub>I with a series of group 10 metal monohalides

Na Cheng · Yongjun Liu · Changqiao Zhang · Chengbu Liu

Received: 6 March 2013 / Accepted: 3 June 2013 / Published online: 25 June 2013  
© Springer-Verlag Berlin Heidelberg 2013

**Abstract** The halogen bonding interactions between C<sub>6</sub>F<sub>5</sub>I and a series of transition metal monohalides *trans*-[M(X)(2-C<sub>5</sub>NF<sub>4</sub>)-(PR<sub>3</sub>)<sub>2</sub>] (M = Ni, Pd, Pt; X = F, Cl, Br; R = Me, Cy) have been studied with quantum chemical calculations. Optimized geometries of the halogen bonding complexes indicate that angles C<sub>1</sub>-I⋯X are basically linear (178–180°) and angles I⋯X-M mainly range from 90 to 150°. The strength of these metal-influenced halogen bonds alters with different metal centers, metal-bound halogen atoms and the substituents on phosphine ligands. Electrostatic potential and natural bond orbital analysis show that both of the electrostatic and orbital interactions make a contribution to the formation of halogen bonds, while the electrostatic term plays a dominant role. AIM analysis suggests that, for *trans*-[M(F)(2-C<sub>5</sub>NF<sub>4</sub>)-(PR<sub>3</sub>)<sub>2</sub>] (M = Ni, Pd, Pt) monomers, the formed halogen bonding complexes are stabilized by local concentration of the charge of intermediate character, while for the metal monomers containing chlorine and bromine, a typical closed-shell interaction exist. These results prove that the structures and geometries of these halogen bonding complexes can be tuned by changing the halogen atoms and metal centers, which may provide useful information for the design and synthesis of new functional materials.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00894-013-1910-0) contains supplementary material, which is available to authorized users.

N. Cheng · Y. Liu (✉) · C. Zhang · C. Liu  
Key Laboratory of Theoretical and Computational Chemistry in Universities of Shandong, School of Chemistry and Chemical Engineering, Shandong University, Jinan, Shandong 250100, China  
e-mail: yongjunliu\_1@sdu.edu.cn

Y. Liu  
Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining, Qinghai 810001, China

**Keywords** DFT · Halogen bonds · Iodoperfluoroarenes · Metal halide · NBO

## Introduction

Halogen bonds are specific noncovalent interactions between a covalently bonded halogen atom in compounds of R-X (X = F, Cl, Br and I) and a Lewis base [1–7]. It has attracted widespread interest of chemists in recent years due to its potential applications in crystal engineering [1, 8, 9], materials science [10], molecular recognition [3, 11] and biological systems [12–14]. Politzer et al. [2] have explained the formation of halogen bonds with the electrostatic potential of covalent halogen atoms. They found that the charge density on the covalent halogen atom is anisotropic distribution, i.e., the electrostatic potential along the extension the C-X bond is positive and perpendicular to covalent bond is negative. The region of positive electrostatic potential is called  $\sigma$ -hole, which can form halogen bond with Lewis bases which are usually N, O and F atoms with lone-pair electrons, or molecules with  $\pi$  electrons such as ethylene, ethyne and benzene [15–17]. Meanwhile, the covalent halogen atoms may also form hydrogen bond as Lewis bases with hydrogen atom [18, 19].

In the past decades, numerous experimental and computational studies have been devoted to better understand the structure, directionality and nature of halogen bonds [20–25], and a fair amount is known about this noncovalent interaction. For example, its strength is similar to that of hydrogen bond, and increases in the order of chlorine < bromine < iodine. In general, fluorine does not form halogen bond because it lacks a positive  $\sigma$ -hole. However, when fluorine attaches to a strong electron withdrawing groups, it can display a  $\sigma$ -hole and form halogen bonds with Lewis bases [11, 26]. The structural characterization of halogen

bonds has been established in a wide range of situations encompassing organic compounds and main group elements.

In recent years, two types of metal-influenced halogen bonds attracted much attention. One type is the halogen bonds of  $M-C-X\cdots X'$  ( $M$  represents metal), which exist widely in the supramolecular networks [27, 28]. Many kinds of halogen bonds  $M-C-X\cdots X'$  have been studied from both theoretical and experimental points of view. For example, Xu et al. [29] studied two model complexes  $PyCl\cdots X$  ( $PyCl = NC_5H_4Cl-4$ ;  $X = F^-, Cl^-,$  or  $Br^-$ ) and  $MPyCl\cdots X$  ( $M = Cu^+, Zn^+$ ) by using DFT calculations. They predicted that the anisotropy of electrostatic potential on the outer surface of covalent chlorine is enhanced due to the metal coordination, and the strength and directionality of halogen bond are thus strengthened.

Another type of halogen bonds exists in the metal complexes  $C-X\cdots X'-M$  (usually,  $M$  is transition metal atom) [30–33]. Unlike the traditional halogen bond, this kind of halogen bond is formed between an organic halide ( $C-X$ ) and an inorganic metal halide ( $X'-M$ ), which also plays an important role in supramolecular self-assembly process. In these halogen bond complexes, the group which the covalent halogen is bound to and the halometallate also have significant influence on the strength and directionality of such halogen bonds [28, 29, 34, 35]. Thus we can design or suitably tune the structures and geometries of the metal-influenced halogen bonds by changing the covalent halogen atom  $X$ , the metal atom  $M$ , and the Lewis base  $X'$ . For example, Zordan et al. [36] synthesized a series of  $trans-[MCl_2(3-X'-py)_2]$  compounds ( $M = Pd, Pt$ ;  $X' = Cl, Br,$  and  $I$ ). They found that two different networks, propagated through  $M-Cl\cdots X'-C$  halogen bonds, exist in the crystals structures of these compounds. The  $trans-[PdCl_2(3-I-py)_2]$  and  $trans-[PtCl_2(3-X'-py)_2]$  ( $X' = Cl, Br$ ) are the pattern of one dimensional tapes, while the  $trans-[PdCl_2(3-X'-py)_2]$  ( $X' = Cl, Br$ ) and  $trans-[PtCl_2(3-X'-py)_2]$  ( $X' = Br, I$ ) form the two dimensional layers. In a recent review, Bertani et al. [37] demonstrated that  $C-X\cdots X'-M$  synthon is effective in controlling the crystals structures and chemical behavior of supramolecules.

Due to the fluorination of benzene ring, perfluorinated iodoalkanes and iodoarenes have also been identified as halogen bond donors. Substituted iodoperfluoroarenes and 1,4-diiodo-tetrafluorobenzenes have been widely used in designing liquid crystal and constructing halogen bond-based supramolecular architectures [38, 39]. Libri et al. studied the strong hydrogen bonds and halogen bonds formed by iodopentafluorobenzene and  $trans$ -(tetrafluoropyrid-2-yl)-bis(triethylphosphine) fluoronickel ( $NiF$ ) in solution [40]. Recently, Perutz and co-workers [41] synthesized a series of structurally similar group 10 metal monofluoride compounds (Chart 1) and studied the thermodynamics characteristics of the halogen bonding interactions between  $C_6F_5I$  and these monofluoride complexes by using  $^{19}F$ -NMR titration


experiments and computational approaches. They showed that the data for  $-\Delta H^\circ$  for the halogen bonding follow a trend  $Ni < Pd < Pt$ , and the fluoropyridyl ligand has a negligible influence on the thermodynamic data, but that of the phosphine ligand is significant. Perutz et al. [41] suggested that these group 10 metal fluorides are excellent halogen bond acceptors. But how about the group 10 metal chlorides and bromides? Why the dates of halogen bonding energies follow a trend of  $Ni > Pd > Pt$ ? To explore the bonding nature of  $C-X\cdots X'-M$  halogen bond, the interactions between substituted iodoperfluoroarenes and a series of group 10 metal monohalide compounds  $trans-[M(X)(2-C_5NF_4)-(PR_3)_2]$  ( $M = Ni, Pd, Pt$ ;  $X = F, Cl, Br$ ;  $R = Me, Cy$ ) (Chart 1) have been systematically studied by using quantum chemistry method. We are interested in the influences of different metal atoms and halogen atoms on the strength and bonding characteristics of halogen bonds. These studies may enrich our understanding on metal-bound halogen bond, and provide useful information for developing new supramolecular synthons  $C-X\cdots X'-M$  in supramolecular chemistry and crystal engineering.

## Computational methods

For all monomers and halogen bonding complexes, geometrical optimizations and vibrational frequency calculations were performed by using density functional theory as implemented in the Gaussian 09 program package [42]. The density functional theory (DFT) method has been used in the study of large systems containing metal atoms such as  $MPyCl\cdots X$  and  $PyCl\cdots X$  ( $M = Cu, Zn$ ) systems [11], and BHandHLYP functional can provide accurate energetics for noncovalently bonded systems [2, 43, 44]. Therefore, in this study, the BHandHLYP functional was used in conjunction with the SDD effective core potential and associated basis set on  $I, P, Ni, Pd$  and  $Pt$  and a 6-31G\* basis set on the main group atoms  $C, H, N, F, Cl,$  and  $Br$ . Simultaneously, for all fluorine, chlorine and bromine atoms, diffuse functions were also added. To exam the efficiency of DFT method, a single-point energy calculation at the MP2/def2-TZVP level was also performed. Due to the size of the complexes and the computing cost of MP2, the single-point energy calculation was only applied for the complexes formed by  $C_6F_5I$  and  $trans-[Pd(X)(C_5F_4)(PMe_3)_2]$  ( $X = F$  and  $Cl$ ).

The interaction energies of all given complexes (Chart 2) were calculated as the difference between the energy of complexes and the sum of energies of isolated iodoperfluoroarenes and metal halides. The basis set superposition error (BSSE) was eliminated using the Boys and Bernardi [45] counterpoise method. Libri et al. [40] have revealed that the influence of solvation effects on halogen bonding can not be negligible. Thus, after all the stationary points were located, single point electronic energy calculations with the polarized continuum

**Chart 1** Substituted iodoperfluoroarenes and a series of group 10 metal monohalogenides



model (PCM) were performed at the same level of theory [46, 47]. All interaction energies reported here also were corrected by the effect of zero-point energies. Natural bond orbital (NBO) analysis was performed for our selected systems utilizing the Gaussian NBO module [48] at the same level. To further study the bonding characteristic of the halogen bond, the atoms in molecules (AIM) analysis by means of Bader's "atoms in molecules" theory was carried out with the AIMAll program package [49].

## Results and discussion


### Geometries and binding energies

Figure 1 depicts the optimized structures of the halogen-bonded complexes. The iodoperfluoroarenes is the halogen bond donor, and the group 10 metal monohalide compounds function as halogen bond acceptors. For a convenient and clear illustration, we number the important atoms, i.e., the carbon atoms bonding with iodine and transition metals are denoted as  $C_1$  and  $C_2$ , while the two phosphorus atoms


connecting to metals are named as  $P_1$  and  $P_2$ , as shown in 1-2a. The interaction energies ( $\Delta E$ ), enthalpies ( $\Delta H$ ), and some key geometrical parameters of the complexes are collected in Table 1. While some important bond distances of monomers (iodoperfluoroarenes and metal halides) are listed in Table S1 of Supporting information. All the optimized structures of metal halide monomers are shown in Fig. S1.

Table 1 shows that the binding distances ( $R_{X\cdots I}$ ) between I and X (X = F, Cl, and Br) are in the range of 2.217 to 3.536 Å, which are clearly shorter than the sum of van der Waals radii of I and X atoms (the sum of van der Waals radii of I $\cdots$ F, I $\cdots$ Cl and I $\cdots$ Br are 3.11, 3.73 and 3.83 Å, respectively), implying a typical halogen bonding interaction between the monomers of iodoperfluoroarenes and metal halides.  $R_{X\cdots I}$  of complex 1-5 is the shortest, which is consistent with the strongest halogen-bonding interaction of 1-5 with the  $\Delta E$  value of  $-66.30$  kJ mol $^{-1}$ . For the complexes formed by  $C_6F_5I$  and *trans*-[M(X)(C<sub>5</sub>F<sub>4</sub>)(PMe<sub>3</sub>)<sub>2</sub>] (M = Ni, Pd, Pt), the binding distances  $R_{X\cdots I}$  follow the order F < Cl < Br, corresponding to their tendency of  $\Delta E$  value. We also note that the binding distances  $R_{X\cdots I}$  of F complexes (1-2a, 1-3a and 1-4a) are much shorter than that of other complexes, indicating that the fluorides are

**Chart 2** A series of halogen-bonding complexes



**Fig. 1** The optimized structures of halogen-bonding complexes at BhandHLYP/6-31G\*-SDD level



the strongest halogen bonding acceptors. As for the changes of bond lengths of  $R_{C_1-I}$ ,  $R_{M-X}$ , and  $R_{M-C_2}$ , Table 1 shows that, after the formation of halogen bonded complexes, all bond lengths  $R_{C_1-I}$  and  $R_{M-X}$  are elongated, and  $R_{M-C_2}$  are contracted.

Politzer et al. have reported that halogen bonding is an electrostatic driven directional molecular interaction [2]. From Table 1, one can see that angles  $\theta_{C_1-I \cdots X}$  of all complexes are basically  $180^\circ$ , which is consistent with the linear characteristic of halogen bond. However, the  $I \cdots X-M$  moieties of all

**Table 1** Corrected interaction energies ( $\Delta E$ ,  $\text{kJ mol}^{-1}$ ), enthalpies ( $\Delta H$ ,  $\text{kJ mol}^{-1}$ ), binding distance ( $R$ ,  $\text{\AA}$ ), changes of key bond lengths ( $\Delta R$ ,  $\text{\AA}$ ) and bond angles ( $\theta$ , degree) in selected halogen-bonded complexes

Complexes	$\Delta E$	$\Delta H$	$R_{X-I}$	$\Delta R_{C_1-I}$	$\Delta R_{M-X}$	$\Delta R_{M-C_2}$	$\theta_{C_1-I \cdots X}$	$\theta_{I \cdots X-M}$
1-2a	-19.87	-15.93	2.725	0.009	0.016	-0.006	180.0	179.9
1-2b	-9.29	-5.22	3.398	0.007	0.010	-0.002	179.4	118.8
1-2c	-6.63	-2.73	3.536	0.008	0.010	-0.002	178.2	92.8
1-2d	-16.59	-11.05	2.753	0.008	0.021	-0.006	179.9	179.7
1-3a	-23.68	-20.00	2.372	0.015	0.021	-0.004	179.9	147.3
1-3b	-10.49	-6.43	3.370	0.009	0.011	-0.003	179.5	117.0
1-3c	-8.98	-4.89	3.499	0.010	0.010	-0.003	179.1	103.8
1-4a	-22.24	-18.26	2.688	0.013	0.019	-0.004	179.5	142.7
1-4b	-10.27	-6.02	3.384	0.008	0.008	-0.002	179.3	115.0
1-4c	-9.01	-4.71	3.503	0.010	0.007	-0.001	178.8	106.4
1-5	-66.30	-61.28	2.217	0.151			180.0	

metal complexes except nickel fluoride complexes are distinctly bent, agreeing well with the calculation results of Perutz et al. and the observation in a survey of  $H\cdots X-M$  angles for hydrogen bonding involving metal halides [41, 50]. The reason that the  $I\cdots X-M$  angles are bent is that the interactions are with the sides of the halogen X. Such angles also agree with some experimental results of  $M-X\cdots X$  angles [33, 41].

Table 1 shows that the interaction enthalpy ( $\Delta H$ ) of complex **1-2a** is  $15.93 \text{ kJ mol}^{-1}$ , agreeing well with the experimental value of  $16 \text{ kJ mol}^{-1}$  [40]. Although there is a difference of  $4 \text{ kJ mol}^{-1}$  between interaction enthalpy ( $\Delta H$ ) and interaction energy ( $\Delta E$ ), both of them give the similar tendency of halogen bonding. Thus, we just analyze these halogen bonded complexes based on their interaction energies ( $\Delta E$ ) in the following sections. For all these metal-influenced complexes **1-2**, **1-3**, and **1-4**, the interaction energies follow the order of  $F < Cl < Br$ . The fluorides correspond to the strongest halogen bonds, and bromides correspond to the weakest halogen bonds. As for the influence of metal atoms, one can see that there are only several  $\text{kJ mol}^{-1}$  differences of interaction energies between these complexes. The interaction energies of fluoride complex follow the order of  $Pd < Pt < Ni$ . The influence of phosphine ligand is also not very significant. By changing the substitute on phosphine ligand from methyl (Me) to cyclohexyl (Cy), the interaction energy only changes from  $-19.87$  to  $-16.59 \text{ kJ mol}^{-1}$ .

To assess the efficiency of DFT method, the interaction energies of **1-3a** and **1-3b** were also calculated at the MP2/def2-TZVP level on the BHandHLYP optimized geometries. After the corrections of zero-point vibrational energies (ZPE) and BSSE, the interaction energies  $\Delta E_{MP2}$  of complexes **1-3a** and **1-3b** including solvation effect are  $-33.49$  and  $-22.78 \text{ kJ mol}^{-1}$ , respectively. With respect to the MP2 results, BhandHLYP method underestimates the interaction energies of


halogen bonds, but both of the MP2 and BhandHLYP methods give the same trend of interaction energies. Thus, we used the BhandHLYP method for the calculation of interaction energies.

For further understanding the formation of halogen bonds, electrostatic potential analysis was applied to all monomers. The electrostatic potentials of  $C_6F_5I$  (**1**) and some selected metal halide monomers are shown in Fig. 2. It can be seen that there is a positive electrostatic potential cap ( $\sigma$ -hole) on the molecular surface of monomer **1** along the  $C_1-I$  bond, but for the metal monohalides, there are negative electrostatic regions on the outermost portions of halogen atoms X. Table 2 lists the most positive electrostatic potentials ( $V_{max}$ ) for iodine atom, and the most negative electrostatic potential ( $V_{min}$ ) for X atoms in the metal halide monomers. The  $V_{max}$  value for iodine atom is  $37.3 \text{ kcal mol}^{-1}$ . The  $V_{min}$  values for X atoms follow the order of  $F < Cl < Br$ , which is consistent with their binding energies of the halogen bond complexes, which indicates that the contribution of electrostatic interaction is significant.

#### NBO analysis

To have a deeper understanding of the nature and properties of noncovalent interactions, we perform a NBO analysis at the BhandHLYP/6-31G(d)-SDD level of theory. The important donor-acceptor orbital interactions and corresponding second-order perturbation stabilization energies  $E(2)$  are given in Table 3. There are four types of orbital interactions:  $LP(1)X \rightarrow \sigma^*(C-I)$ ,  $LP(2)X \rightarrow \sigma^*(C-I)$ ,  $LP(3)X \rightarrow \sigma^*(C-I)$ , and  $LP(4)X \rightarrow \sigma^*(C-I)$ . They indicate that the electron transfer always occurs from the lone electron pairs of X to C-I antibonding  $\sigma^*$  orbital. The stabilization energy  $E(2)$  of the  $LP(3)X \rightarrow \sigma^*(C-I)$  orbital interaction is larger than that of the  $LP(1)X$  or  $LP(2) \rightarrow \sigma^*(C-I)$  in all complexes except

**Fig. 2** Calculated electrostatic potential on the 0.001 a.u. molecular surface of selected monomers



**Table 2** The most positive electrostatic potentials ( $V_{\max}$ , kcal mol<sup>-1</sup>) on 0.001 a.u. surface of I atom in C<sub>6</sub>F<sub>5</sub>I, and the most negative electrostatic potentials ( $V_{\min}$ , kcal mol<sup>-1</sup>) on 0.001 a.u. outer surface of halogen atom in metal monohalogenides

Complexes	$V_{\max}$	$V_{\min}$
1	37.3	
2a		-43.7
2b		-30.7
2c		-27.1
2d		-44.4
3a		-50.7
2b		-34.3
3c		-30.1
4a		-48.8
4b		-32.6
4c		-28.9
5		-173.7

complex 1–5. Nevertheless, the stabilization energy  $E(2)$  of LP(4)X  $\rightarrow$   $\sigma^*(\text{C-I})$  orbital interaction is extremely large in complexes 1–5. Table 3 shows that the largest stabilization energy  $E(2)$  of each complex correlates at some extent with the corresponding interaction energy, which implies that orbital interaction plays a role in the formation of halogen bond.

As we know, the charge transfer is the concomitant of orbital interaction. Table 3 lists the charge transfer (CT) between monomer C<sub>6</sub>F<sub>5</sub>I and metal monohalides upon forming complexes. With the increase of atomic number of X, the charge transfer gradually increases for complexes

**Table 3** Stabilization energies ( $E(2)$ , kJ mol<sup>-1</sup>), charge transfer (CT, e), and differences between NBO electron density in the complexes and isolated C<sub>6</sub>F<sub>5</sub>I in C-I sigma bonding ( $\sigma$ ) and sigma antibonding ( $\sigma^*$ ) orbitals in complexes

Complexes	$E(2)$	CT	$\Delta\sigma(\text{C-I})$	$\Delta\sigma^*(\text{C-I})$
1–2a	25.22 <sup>c</sup>	0.0157	-0.0055	0.0212
1–2b	3.18 <sup>a</sup> , 15.93 <sup>c</sup>	0.0226	-0.0030	0.0240
1–2c	1.76 <sup>a</sup> , 19.45 <sup>c</sup>	0.0237	-0.0029	0.0317
1–2d	23.67 <sup>c</sup>	0.0059	-0.0061	0.0211
1–3a	1.25 <sup>b</sup> , 30.57 <sup>c</sup>	0.0241	-0.0067	0.0283
1–3b	1.80 <sup>b</sup> , 17.82 <sup>c</sup>	0.0271	-0.0035	0.0273
1–3c	1.76 <sup>a</sup> , 20.41 <sup>c</sup>	0.0348	-0.0033	0.0349
1–4a	2.93 <sup>b</sup> , 31.03 <sup>c</sup>	0.0214	-0.0064	0.0265
1–4b	2.47 <sup>a</sup> , 17.90 <sup>c</sup>	0.0258	-0.0031	0.0262
1–4c	2.38 <sup>a</sup> , 20.49 <sup>c</sup>	0.0343	-0.0030	0.0340
1–5	46.75 <sup>a</sup> , 223.40 <sup>d</sup>	0.1962	-0.0324	0.1725


<sup>a</sup> the LP(1)X  $\rightarrow$   $\sigma^*(\text{C-I})$  orbital interaction. <sup>b</sup> the LP(2)X  $\rightarrow$   $\sigma^*(\text{C-I})$  orbital interaction

<sup>c</sup> the LP(3)X  $\rightarrow$   $\sigma^*(\text{C-I})$  orbital interaction. <sup>d</sup> the LP(4)X  $\rightarrow$   $\sigma^*(\text{C-I})$  orbital interaction

X = F, Cl, Br

**Table 4** Bond critical point properties (in a.u.) for halogen bond in studied complexes

Complexes	$\rho_b$	$\nabla^2\rho_b$	$V_b$	$G_b$	$H_b$	$ V_b /G_b$
1–2a	0.0224	0.0900	-0.0206	0.0204	-0.0002	1.0098
1–2b	0.0109	0.0395	-0.0061	0.0077	0.0016	0.7922
1–2c	0.0107	0.0332	-0.0053	0.0066	0.0013	0.8030
1–2d	0.0212	0.0848	-0.0193	0.0191	-0.0002	1.0105
1–3a	0.0256	0.1004	-0.0235	0.0227	-0.0008	1.0352
1–3b	0.0116	0.0417	-0.0066	0.0082	0.0016	0.8049
1–3c	0.0114	0.0354	-0.0058	0.0071	0.0013	0.8169
1–4a	0.0248	0.0972	-0.0227	0.0220	-0.0007	1.0318
1–4b	0.0113	0.0407	-0.0063	0.0080	0.0017	0.7875
1–4c	0.0113	0.0352	-0.0057	0.0070	0.0013	0.8143
1–5	0.0726	0.3389	-0.0643	0.0560	-0.0083	1.1482



**Fig. 3** **a** The relationship between the interaction energies and the electron density at bond critical points; **b** The relationship between the interaction energies and the Laplacian of electron density at bond critical points

$C_6F_5I$  *trans*-[M(X)(C<sub>5</sub>F<sub>4</sub>)(PMe<sub>3</sub>)<sub>2</sub>]. For example, the CT value of **1-2a** is 0.0157, while that of **1-2c** is 0.0237. However, the CT values do not directly correct with interaction energies, i.e., the largest CT value does not correspond to the smallest interaction energy, which means the charge transfer plays a minor role in the formation of halogen bond. In addition, these orbital interactions in all complexes also cause an increase of electron density for  $\sigma^*(C-I)$  and a decrease of electron density for  $\sigma(C-I)$ . From Table 3, one can see the overall electron-density changes of  $\sigma(C-I)$  are very small. The electron-density increase of  $\sigma^*(C-I)$  contributes mainly to the elongation of C-I bond.

### AIM analysis

The AIM topological analysis has been widely used in the study of noncovalent interactions, such as hydrogen and halogen bonds. It is another effective method for understanding the nature of halogen bond. Therefore, we analyze the electron density ( $\rho_b$ ) and corresponding Laplacian ( $\nabla^2\rho_b$ ) at the bond critical point (BCP) of halogen bonds for all complexes. The calculation results are listed in Table 4. Koch and Popelier [51] have proposed that for closed-shell interaction, both the parameters of  $\rho_b$  and  $\nabla^2\rho_b$  are positive and should be within the range of 0.002–0.035 a.u. and 0.024–0.139 a.u., respectively. Table 4 shows that all the values of  $\rho_b$  and  $\nabla^2\rho_b$ , except that of complex **1-5**, are within the proposed range. However,  $\rho_b$  at BCP of the complex **1-5** is beyond the above range. We found that the halogen bond in **1-5** has the small interaction distance R, large interaction energy  $\Delta E$ , large  $\rho_b$ , positive  $\nabla^2\rho_b$  and negative  $H_b$  (total energy density). These characteristics indicate that a strong intermolecular interaction is formed in complex **1-5**. All these topological parameters can well describe the interaction strength of the halogen bond, because they relate directly with the interaction energy. The linear relationship between  $\rho_b$  and  $\Delta E$  is shown in Fig. 3a, and that of  $\nabla^2\rho_b$  and  $\Delta E$  is shown in Fig. 3b.

Moreover, it may be insufficient to characterize an intermolecular interaction only based on the values of  $\rho_b$  and the sign of  $\nabla^2\rho_b$ , therefore, the local potential energy density ( $V_b$ ), local kinetic energy ( $G_b$ ) and total energy density ( $H_b=V_b+G_b$ ) at BCPs are also listed in Table 4.  $H_b$  is a more precise index to characterize weak interaction. Later, Jenkins and Morrison [52] used the ratio  $|V_b|/G_b$  to describe the noncovalent interaction. For complexes involving metal monofluorides (**1-2a**, **1-2d**, **1-3a**, **1-4a**, and **1-5**), Table 4 shows that the values of  $|V_b|/G_b$  is between 1 and 2 ( $1 < |V_b|/G_b < 2$ ), and  $H_b$  is negative, which indicates that these monofluorides complexes are stabilized by local concentration of the charge of intermediate character. For the other complexes, the values of  $|V_b|/G_b$  are less than 1 ( $|V_b|/G_b < 1$ ) and  $\Delta H_b > 0$ , implying that these complexes are closed-shell interactions.

### Conclusions

In this paper, a series of metal-influenced halogen bonding complexes formed by  $C_6F_5I$  and *trans*-[M(X)(2-C<sub>5</sub>NF<sub>4</sub>)-(PR<sub>3</sub>)<sub>2</sub>] (M = Ni, Pd, Pt; X = F, Cl, Br; R = Me, Cy) has been studied with quantum chemical calculation. The interaction energies of the metal-influenced halogen bonds in these complexes do not monotonically depend on the atom number of group 10 metals. The data of  $-\Delta E$  for the halogen bonds follow the trend Ni < Pt < Pd when X is F and Cl. Electrostatic potential and NBO analysis show that both of the electrostatic and orbital interactions contribute to the formation of halogen bonds, but the electrostatic interaction plays a dominant role. AIM calculation indicates that for *trans*-[M(F)(2-C<sub>5</sub>NF<sub>4</sub>)-(PR<sub>3</sub>)<sub>2</sub>] (M = Ni, Pd, Pt) monomers, the formed halogen bonding complexes are stabilized by local concentration of the charge of intermediate character, while for the metal monomers containing chlorine and bromine, a typical closed-shell interaction exist. All these results show that the metal-bound halogen atoms can be halogen bond acceptors, and the interaction strength and structures of these complexes can be controlled by changing the metal and halogen atoms. The present study may help to understand the nature of C-I...X-M halogen bond and provide useful information for the design and synthesis of new functional materials.

**Acknowledgments** This work was supported by the Natural Science Foundation of China (21173129, 91127014).

### References

1. Metrangolo P, Meyer F, Pilati T, Resnati G, Terraneo G (2008) Halogen bonding in supramolecular chemistry. *Angew Chem Int Ed* 47:6114–6127
2. Politzer P, Murray JS, Clark T (2010) Halogen bonding: an electrostatically-driven highly directional noncovalent interaction. *Phys Chem Chem Phys* 12:7748–7757
3. Metrangolo P, Neukirch H, Pilati T, Resnati G (2005) Halogen bonding based recognition processes: a world parallel to hydrogen bonding. *Acc Chem Res* 38:386–395
4. Mohajeri A, Alipour M, Mousaee M (2011) Halogen-hydride interaction between Z–X (Z = CN, NC; X = F, Cl, Br) and H–Mg–Y (Y = H, F, Cl, Br, CH<sub>3</sub>). *J Phys Chem A* 115:4457–4466
5. Politzer P, Murray JS (2013) Halogen bonding: an interim discussion. *ChemPhysChem* 14:278–294
6. Feng G, Evangelisti L, Gasparini N, Caminati W (2012) On the Cl...N halogen bond: a rotational study of CF<sub>3</sub>Cl...NH<sub>3</sub>. *Chem Eur J* 18:1364–1368
7. Politzer P, Murray JS, Clark T (2013) Halogen bonding and other  $\sigma$ -hole interactions: a perspective. *Phys Chem Chem Phys*. doi:10.1039/c3cp00054k
8. Metrangolo P, Carcenac Y, Lahtinen M, Pilati T, Rissanen K, Vij A, Resnati G (2009) Nonporous organic solids capable of dynamically resolving mixtures of diiodoperfluoroalkanes. *Science* 323:1461–1464
9. Martí-Rujas J, Colombo L, Lü J, Dey A, Terraneo G, Metrangolo P, Pilati T, Resnati G (2012) Hydrogen and halogen bonding drive the

- orthogonal self-assembly of an organic framework possessing 2D channels. *Chem Commun* 48:8207–8209
10. Metrangolo P, Resanti G, Pilati T, Liantonio R, Meyer F (2007) Engineering functional materials by halogen bonding. *J Polym Sci Part A: Polym Chem* 45:1–15
  11. Metrangolo P, Murray JS, Pilati T, Politzer P, Resnati G, Terraneo G (2011) Fluorine-centered halogen bonding: a factor in recognition phenomena and reactivity. *Cryst Growth Des* 11:4238–4246
  12. Johnson CM, Linsky TW, Yoon DW, Person MD, Fast W (2011) Discovery of halopyridines as quiescent affinity labels: inactivation of dimethylarginine dimethylaminohydrolase. *J Am Chem Soc* 133:1553–1562
  13. Hardegger LA, Kuhn B, Spinnler B, Anselm L, Ecabert R, Stihle M, Gsell B, Thoma R, Diez J, Benz J, Plancher JM, Hartmann G, Banner DW, Haap W, Diederich F (2011) Systematic investigation of halogen bonding in protein-ligand interactions. *Angew Chem Int Ed* 50:314–318
  14. Ibrahim MAA (2011) Molecular mechanical study of halogen bonding in drug discovery. *J Comput Chem* 32:2564–2574
  15. Li QZ, Zhao JL, Li R, Li WZ, Cheng JB, Gong BA (2011) Rare gas atomic number dependence of the halogen bond in HRgF-XCCF (Rg = He, Ar, Kr; X = Cl, Br) complex. *Comput Theor Chem* 970:61–65
  16. Li RY, Li ZR, Wu D, Li Y, Chen W, Sun CC (2005) Study of  $\pi$  halogen bonds in complexes  $C_2H_{4-n}F_n-ClF$  ( $n = 0-2$ ). *J Phys Chem A* 109:2608–2013
  17. Duarte DJR, de las Vallejos MM, Peruchena NM (2010) Topological analysis of aromatic halogen/hydrogen bonds by electron charge density and electrostatic potentials. *J Mol Model* 16:737–748
  18. Liu XF, Cheng JB, Li QZ, Li WZ (2013) Competition of hydrogen, halogen, and pnictogen bonds in the complexes of HARF with XH<sub>2</sub>P (X = F, Cl, and Br). *Spectrochim Acta A (Molecular and Biomolecular Spectroscopy)* 101:172–177
  19. Wang WZ, Tian AM, Wong NB (2005) Theoretical study on the bromomethane-water 1:2 complexes. *J Phys Chem A* 109:8035–8040
  20. Riley KE, Murray JS, Fanfrlík J, Řezáč J, Solá RJ, Concha MC, Ramos FM, Politzer P (2011) Halogen bond tunability I: the effects of aromatic fluorine substitution on the strengths of halogen-bonding interactions involving chlorine, bromine, and iodine. *J Mol Model* 17:3309–3318
  21. Riley KE, Murray JS, Fanfrlík J, Řezáč J, Solá RJ, Concha MC, Ramos FM, Politzer P (2012) *J Mol Model*. doi:10.1007/s00894-012-1428-x
  22. Lu JM, Zhang B, Deng QM, Wang JA, Lu YX, Zhu WL (2011) The nature and magnitude of specific halogen bonds between iodo-perfluorobenzene and heterocyclic systems. *Int J Quantum Chem* 111:2352–2358
  23. Lu YX, Li HY, Zhu X, Zhu WL, Liu HL (2011) How does halogen bonding behave in solution? A theoretical study using implicit solvation model. *J Phys Chem A* 115:4467–4475
  24. Grabowski SJ, Bilewicz E (2006) Cooperativity halogen bonding effect – Ab initio calculations on H<sub>2</sub>CO (ClF)<sub>n</sub> complexes. *Chem Phys Lett* 427:51–55
  25. Forni A, Rendineb S, Pieraccini S, Sironi M (2012) Solvent effect on halogen bonding: the case of the I ⋯ O interaction. *J Mol Graph Model* 38:31–39
  26. Metrangolo P, Murray JS, Pilati T, Politzer P, Resnati G, Terraneo G (2011) The fluorine atom as a halogen bond donor, viz. a positive site. *CrystEngComm* 13:6593–6596
  27. Zhao Q, Feng DC (2012) Influence of transition metal M (M = Cu, Ag, Au) on the strength of halogen bonding interaction X⋯Cl (X = F, Cl, Br). *Acta Physico-Chimica Sinica* 28:1361–1367
  28. Xu L, Lv J, Sang P, Zou JW, Yu QS (2010) DFT model investigations on copper(II) chloride complexes: halogen bonding of pyridyl bromine of CuCl<sub>2</sub>(NH<sub>3</sub>)(NC<sub>5</sub>H<sub>4</sub>Br-3) with several metal ligands. *J Mol Struct (THEOCHEM)* 953:170–174
  29. Xu L, Lv J, Sang P, Zou JW, Yu QS, Xu MB (2011) Comparative insight into the halogen bonding of 4-chloropyridine and its metal [Cu<sup>I</sup>, Zn<sup>II</sup>] coordinations with halide ions: A theoretical study on M–C–X X'. *Chem Phys* 379:66–72
  30. Bhosekar G, Jeß I, Lehnert N, Näther C (2008) Synthesis, crystal structure and thermal reactivity of [ZnX<sub>2</sub>(2-chloropyrazine)] (X = Cl, Br, I) coordination compounds. *Eur J Inorg Chem* 4:605–611
  31. Wriedt M, Jeß I, Näther C (2009) Synthesis, crystal structures, and thermal properties of new [ZnX<sub>2</sub>(2,5-dimethylpyrazine)] (X = Cl, Br, I) coordination compounds. *Eur J Inorg Chem* 3:363–372
  32. Espallargas GM, Brammer L, van de Streek J, Shankland K, Florence AJ, Adams H (2006) Reversible extrusion and uptake of HCl molecules by crystalline solids involving coordination bond cleavage and formation. *J Am Chem Soc* 128:9584–9585
  33. Zordan F, Brammer L, Sherwood P (2005) Supramolecular chemistry of halogens: complementary features of inorganic (M–X) and organic (C–X') halogens applied to M–X X'–C halogen bond formation. *J Am Chem Soc* 127:5979–5989
  34. Awwadi FF, Willett RD, Haddad SF, Twamley B (2006) The electrostatic nature of aryl-bromine halide synthons: The role of aryl-bromine-halide synthons in the crystal structures of the trans-Bis(2-bromopyridine)dihalocopper(II) and trans-Bis(3-bromopyridine)dihalocopper(II) complexes. *Cryst Growth Des* 6:1833–1838
  35. Metrangolo P, Meyer F, Pilati T, Resnati G, Terraneo G (2008) Mutual induced coordination in halogen-bonded anionic assemblies with (6,3) cation-templated topologies. *Chem Commun* 14:1635–1637
  36. Zordan F, Brammer L (2006) M–X X'–C halogen-bonded network formation in MX<sub>2</sub>(4-halopyridine)<sub>2</sub> complexes (M = Pd, Pt; X = Cl, I; X' = Cl, Br, I). *Cryst Growth Des* 6:1374–1379
  37. Bertani R, Sgarbossa P, Venzo A, Lelj F, Amati M, Resnati G, Pilati T, Metrangolo P, Terraneo G (2010) Halogen bonding in metal-organic-supramolecular networks. *Coord Chem Rev* 254:677–695
  38. Roper LC, Präsang C, Kozhevnikov VN, Whitwood AC, Karadakov PB, Bruce DW (2010) Experimental and theoretical study of halogen-bonded complexes of DMAP with di- and triiodofluorobenzenes. A complex with a very short N ⋯ I halogen bond. *Cryst Growth Des* 10:3710–3720
  39. Nguyen HL, Horton PN, Hursthouse MB, Legon AC, Bruce DW (2004) Halogen bonding: a new interaction for liquid crystal formation. *J Am Chem Soc* 126:16–17
  40. Libri S, Jasim NA, Perutz RN, Brammer L (2008) Metal fluorides form strong hydrogen bonds and halogen bonds: measuring interaction enthalpies and entropies in solution. *J Am Chem Soc* 130:7842–7844
  41. Beweries T, Brammer L, Jasim NA, McGrady JE, Perutz RN, Whitwood AC (2011) Energetics of halogen bonding of group 10 metal fluoride complexes. *J Am Chem Soc* 133:14338–14348
  42. Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Mennucci B, Petersson GA, Nakatsuji H, Caricato M, Li X, Hratchian HP, Izmaylov AF, Bloino J, Zheng G, Sonnenberg JL, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Vreven T, Montgomery JA Jr, Peralta JE, Ogliaro F, Bearpark M, Heyd JJ, Brothers E, Kudin KN, Staroverov VN, Kobayashi R, Normand J, Raghavachari K, Rendell A, Burant JC, Iyengar SS, Tomasi J, Cossi M, Rega N, Millam JM, Klene M, Knox JE, Cross JB, Bakken V, Adamo C, Jaramillo J, Gomperts R, Stratmann RE, Yazyev O, Austin AJ, Cammi R, Pomelli C, Ochterski JW, Martin RL, Morokuma K, Zakrzewski VG, Voth GA, Salvador P, Dannenberg JJ, Dapprich S, Daniels AD, Farkas O, Foresman JB, Ortiz JV, Cioslowski J, Fox DJ (2009) Gaussian 09, Revision A.1. Gaussian Inc, Wallingford
  43. Csontos J, Palermo NY, Murphy RF, Lovas S (2008) Calculation of weakly polar interaction energies in polypeptides using density



- functional and local Møller-Plesset perturbation theory. *J Comput Chem* 29:1344–1352
44. Waller MP, Robertazzi A, Platts JA, Hibbs DE, Williams PA (2006) Hybrid density functional theory for  $\pi$ -stacking interactions: application to benzenes, pyridines, and DNA bases. *J Comput Chem* 27:491–504
  45. Boys SF, Bernardi F (1970) The calculation of small molecular interaction by the differences of separate total energies. Some procedures with reduced errors. *Mol Phys* 19:553–566
  46. Barone V, Cossi M (1997) A new definition of cavities for the computation of solvation free energies by the polarizable continuum model. *J Chem Phys* 107:3210–3221
  47. Tomasi J, Mennucci B, Cammi R (2005) Quantum mechanical continuum solvation models. *Chem Rev* 105:2999–3093
  48. Reed AF, Curtiss LA, Weinhold F (1988) Intermolecular interactions from a natural bond orbital, donor-acceptor viewpoint. *Chem Rev* 88:899–926
  49. Keith TA (2012) AIMAll, Version 12.06.03, Professional
  50. Brammer L, Bruton EA, Sherwood P (1999) Fluoride ligands exhibit marked departures from the hydrogen bond acceptor behavior of their heavier halogen congeners. *New J Chem* 23:965–968
  51. Koch U, Popeliter PLA (1995) Characterization of C-H-O hydrogen bonds on the basis of the charge density. *J Phys Chem* 99:9747–9754
  52. Jenkins S, Morrison I (2000) The chemical character of the intermolecular bonds of seven phases of ice as revealed by ab initio calculation of electron densities. *Chem Phys Lett* 317:97–102