

CYCLOPEPTIDE ALKALOIDS. PHENCYCLOPEPTINES¹ FROM THE POLYMORPHIC SPECIES *CEANOOTHUS INTEGERRIMUS*

J. CLARK LAGARIAS, DANE GOFF, FREDERICK K. KLEIN AND HENRY RAPOPORT

*Department of Chemistry and Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720*

ABSTRACT.—Seven cyclopeptide alkaloids, phencyclopeptines 1–7, have been found distributed among three forms of the shrub *Ceanothus integerrimus*. Chemical degradation, mass spectroscopy, and ¹H nmr spectroscopy have established structures for these seven compounds, three of which have been previously reported. The use of cyclopeptide alkaloid structure and distribution for chemotaxonomic assignments is discussed.

Ceanothus integerrimus (“Deer Brush”) is a polymorphic species of the family Rhamnaceae occurring from southern Washington through California into western New Mexico. Although as many as eight varieties of this semi-deciduous shrub have been characterized, only two of the seven found in California are present in significant population. *C. integerrimus* H. and A. var. *integerrimus* inhabits the inner South Coast Range and *C. integerrimus* var. *californicus* (Kell.) G. T. Benson is found in the Sierra Nevada northward through the Cascade and Klamath Ranges (1).

As part of more comprehensive alkaloid structure studies of Pacific North American Rhamnaceae, we have begun a phytochemical investigation of *Ceanothus integerrimus*. Our investigation of three specimens of this shrub, one of *C. integerrimus* var. *californicus*, and two from different populations of *C. integerrimus* var. *integerrimus* has led to the identification of four new cyclopeptide alkaloids, phencyclopeptines 1–4, in addition to three previously reported alkaloids 5–7 (table 1). The distribution of phencyclopeptines among the three plants was determined by reversed phase high performance liquid chromatography and mass and ¹H nmr spectroscopy (figures 1, 2, and 3; tables 1 and 2).


EXPERIMENTAL²

PLANT MATERIAL.³—Root bark of *C. integerrimus* var. *integerrimus* was obtained from its

¹We propose the name phencyclopeptine to represent the fundamental para-bridged 14-membered ring nucleus most common in this large class of widely occurring alkaloids. This basic nucleus and the numbering system shown in table 1 allow individual alkaloids to be simply and unambiguously designated. Thus we can avoid the multitude of trivial names based on botanical anagrams that have no structural significance.

²Hplc was performed with a Spectra Physics Model SP 3500B Chromatograph and a model 748 oven, Santa Clara, CA. UV absorbance was monitored with an Altex Model 151 Dual Wavelength Detector, Altex Scientific Inc., Berkeley, CA. Hplc grade solvents from Burdick and Jackson Laboratories, Muskegon, MI, and water purified with a Milli-Q system, Millipore Corp., Bedford, MA, were used for hplc. Uncorrected melting points were determined on a Kofler Micro Hot Stage (μ mp). A model AEI-MS12 mass spectrometer, AEI Scientific Apparatus Ltd, Manchester, England, with INCOS data system was used for determining low resolution mass spectra. High resolution mass spectra were obtained with a Consolidated ElectroDynamics CEC-110B instrument. Amino acid analyses were performed on a Beckman 120C Chromatograph, Fullerton, CA; unless otherwise indicated ¹H nmr spectra were taken in CDCl₃ solution (CHCl₃ at 7.21 ppm) at 22°C on a home-made spectrometer based on a Bruker 63kG magnet operating at 270 MHz with a Nicolet 1180 data system. Evaporations were done *in vacuo* with a Buchi rotary evaporator.

³Identification of plant materials was performed by Dr. L. R. Heckard, University of California, Berkeley, CA, and Dr. M. A. Nobs, Carnegie Institution of Washington, Stanford, CA. All three plants were collected in the months of May, June, and September.

TABLE 1. *Phencyclopeptide constituents of Ceanothus integrerrimus.*


	R ₉	R ₅	R ₃	MW
1, 5-β-Indolylmethyl-8-N-methylvalyl-9-phenylphencyclopeptide	C ₆ H ₅	β-indolyl-CH ₂	NMeVal	579
2, 5-β-Indolylmethyl-8-N,N-dimethylvalyl-9-isopropylphencyclopeptide	(CH ₃) ₂ CH	β-indolyl-CH ₂	NMe ₂ Val	559
3, 5-Benzyl-8-N,N-dimethylisoleucyl-9-phenylphencyclopeptide	C ₆ H ₅	C ₆ H ₅ CH ₂	NMe ₂ Ile	568
4, 5-Isobutyl-8-N-methylisoleucyl-9-phenylphencyclopeptide	C ₆ H ₅	(CH ₃) ₂ CHCH ₂	NMeIle	520
5, 5-β-Indolylmethyl-8-N,N-dimethylisoleucyl-9-isopropylphencyclopeptide (Discarine B) ^a	(CH ₃) ₂ CH	β-indolyl-CH ₂	NMe ₂ Ile	573
6, 5-β-Indolylmethyl-8-N,N-dimethylvalyl-9-phenylphencyclopeptide (Integerrine) ^b	C ₆ H ₅	β-indolyl-CH ₂	NMe ₂ Val	593
7, 5-Isobutyl-8-N,N-dimethylisoleucyl-9-phenylphencyclopeptide (Integerrenine) ^c	C ₆ H ₅	(CH ₃) ₂ CHCH ₂	NMe ₂ Ile	534
8, 5-Benzyl-8-N-methylvalyl-9-phenylphencyclopeptide (Integerressine) ^d	C ₆ C ₅	C ₆ H ₅ CH ₂	NMe ₂ Val	554

^aFirst identified in *Discaria longespina* H. and A. (2).

^bFirst identified in *C. integrerrimus* H. and A. (4).

^cFirst identified in *C. integrerrimus* H. and A. (5).

^dIdentified constituent of *C. integrerrimus* H. and A. (3), not observed in any of the three plants in the present investigation.

type locality in the Santa Cruz Mountains of California and from a population in the North Coast Ranges of Mendocino Co., California, while root bark of *C. integrerrimus* var. *californicus* came from its type locality in the Sierra Nevada Mountains of Calaveras Co., California. Counting annuli revealed the plant from Santa Cruz Co. was 11 years old, the one from Mendocino Co. was 16 years old, and the var. *californicus* was considerably older. Herbarium voucher specimens were submitted to the University Herbarium, University of California, Berkeley, California.

EXTRACTION PROCEDURE.—Plant material (500 g), frozen in liquid nitrogen, was ground to a fine powder in a Waring blender and extracted with 0.1N HCl (2 x 2 liters) over a period of 8–12 hours at room temperature. After filtration, the extracts were combined, adjusted to pH 10 with saturated NaOH, and extracted with CH₂Cl₂ (2 x 1 liter). The combined CH₂Cl₂ layers were concentrated to 100 ml and extracted with 0.1N HCl (5 x 20 ml or until further acid extracts were alkaloid free). The combined acid extracts were made alkaline with saturated Na₂CO₃ to pH 10, extracted with CH₂Cl₂ (5 x 50 ml), and evaporated, affording the following alkaloidal yields: *C. integrerrimus* var. *integrerrimus* (Santa Cruz Co.), 0.09%; *C. integrerrimus* var. *californicus*, 0.33%; *C. integrerrimus* var. *integrerrimus* (Mendocino Co.), 0.14% (root bark).

HPLC ISOLATION OF PHENCYCLOPEPTINES.—Semi-preparative hplc was performed on a LiChrosorb C2 column (10 μ, 10 x 150 mm or 10 x 250 mm, E. M. Merck). The crude alkaloidal mixtures were dissolved in methanol-acetonitrile (1:1) at a concentration of 3 mg/ml, and injection volumes ranged from 10–250 μl. The mobile phase was a mixture of acetonitrile and 0.0015% (v/v) aqueous ammonia with the aqueous ammonia comprising 10–30%; the flow rate was usually 2 ml/min; and the temperature was maintained at 40°C. Alkaloid components were detected at 254 nm. Figure 1 shows a typical hplc tracing for the alkaloid mixtures from each plant variety; 10–20 injections provided sufficient material of each component for structural analysis. Fractions were evaporated *in vacuo* and dried under high vacuum immediately after collection.




FIG. 1. Hplc of crude alkaloidal extracts of the polymorphic species *C. integerrimus*. Hplc system employed: LiChrosorb C2 (10 μ , 10 x 150 mm); mobile phase CH₃CN/10% aq. NH₃ (9/1, v/v); flow rate, 2 ml/min; 35°C; A 254 nm; injection volume, 100 μ l; c~3 mg/ml.

YIELDS OF PHENCYCLOPEPTINE COMPONENTS FROM *C. integerrimus*.—*C. integerrimus* var. *integerrimus* (Santa Cruz Co.). Of the eight components separated by hplc shown in figure 1, five showed mass spectral patterns characteristic of the phencyclopeptide nucleus. These components were obtained in the following relative yield: 7 (70%), 2 (16%), 4 (10%), 3 (4%), 5 trace.

C. integerrimus var. *integerrimus* (Mendocino Co.). Three of seven components contained the phencyclopeptide nucleus, 6 (70%), 5 (15%), and 1 (15%).

C. integerrimus var. *californicus*. Four phencyclopeptides were identified by mass spectroscopy in relative amounts as follows: 7 (45%), 5 (45%), 4 (5%) and 3 (5%).

STRUCTURES⁴ OF PHENCYCLOPEPTINE COMPOUNDS FROM *C. integerrimus*.—5- β -Indolylmethyl-8-*N*-(*N*¹-methylvalyl)-9-phenylphencyclopeptide 1. C₃₄H₃₇N₅O₄; μ mp >350°; ms: M⁺ C₃₄H₃₇N₅O₄ requires 579.2845, found 579.2788, M-43 C₃₁H₃₀N₅O₄ requires 536.2298, found 536.2302, bp C₅H₁₂N requires 86.0970, found 86.0970 (see figure 2 for complete mass spectra); amino acid analysis after acid hydrolysis: no amino acids were observed; ¹H NMR, high field region:⁵ δ 0.27 (d, 3H, *J*=6.9 Hz val- γ -CH₂), 0.54 (d, 3H, *J*=6.9 Hz val- γ -CH₃).

5- β -Indolylmethyl-8-*N*-(*N*¹,*N*¹-dimethylvalyl)-9-isopropylphencyclopeptide 2. C₃₂H₄₁N₅O₄; μ mp 233°; ms: M⁺ C₃₂H₄₁N₅O₄ requires 559.3158, found 559.3146, M-43 *m/e* 516, bp C₆H₁₄N requires 100.1126, found 100.1130 (see figure 2); ¹H nmr, high field region; δ 0.84 (d, 3H, *J*=6.8 Hz, (CH₃)₂CH), 0.93 (d, 3H, *J*=6.8 Hz, (CH₃)₂CH), 0.96 (d, 3H, *J*=6.9 Hz, val- γ -CH₃), 1.18 (d, 3H, *J*=6.9 Hz val- γ -CH₃).

5-Benzyl-8-*N*-(*N*¹,*N*¹-dimethylisoleucyl)-9-phenylphencyclopeptide 3. C₃₄H₄₀N₄O₄; μ mp >350°; ms: M⁺ *m/e* 568, M-57, C₃₀H₃₁N₄O₄ requires 511.2345, found 511.2332, bp C₇H₁₆N requires 114.1282, found 114.1279 (see figure 2); ¹H nmr, high field region: δ 0.18 (d, 3H, *J*=6.9 Hz, ileu- γ -CH₃), 0.80 (t, 3H, *J*=6.9 Hz, ileu- δ -CH₃).

5-Isobutyl-8-*N*-(*N*¹-methylisoleucyl)-9-phenylphencyclopeptide 4. C₃₀H₄₀N₄O₄; μ mp 213°; ms: M⁺ C₃₀H₄₀N₄O₄ requires 520.3049, found 520.3053, M-57 C₂₆H₃₁N₄O₄ requires 463.2345, found 463.2356, bp C₆H₁₄N requires 100.1126 found 100.1131 (see figure 2); amino acid analysis after acid hydrolysis: 1.0 leucine; ¹H nmr, high field region: δ 0.57 (d, 3H, *J*=6.9 Hz, ileu- γ -CH₃), δ 0.66 (m, 6H, ileu- δ -CH₃ and leu(C δ)- δ -CH₃), 0.76 (d, 3H, *J*=6.5 Hz, leu-(C δ)- δ -CH₃).

⁴Compounds of the same structure isolated from different plants had the same mp's and ¹H nmr spectra.

⁵A complete analysis of the ¹H nmr spectra of the phencyclopeptide system will be dealt with in a future report.

TABLE 2. Mass spectra of the hplc purified phenylcyclopeptide components of *C. integerrimus*.

Fragment ^a	Compound						
	1	2	3	4	5	6	7
M ⁺	579 ^b	559 ^b	568	520 ^b	573 ^b	593 ^b	534 ^b
BP a.....	86 ^b	100 ^b	114 ^b	100 ^b	114 ^b	100 ^b	114 ^b
b.....	536 ^b	516	511 ^b	463 ^b	516	550	477 ^b
c.....	215	195	229	215	195	229	229
d.....	187	167	201	187	167	201	201
e.....	410 ^c	376 ^c	371	337	376		337
f.....	224 ^c	190	224	224	190	224	224
g.....	494			421			
h.....	347 ^c	347 ^c	308	274	347	347 ^c	
i.....	135	135	135	135	135	135	135
j.....	451	417 ^c	412 ^c	378			
k.....	317 ^c	283	278 ^c	244	283	317	244
l.....	289	255 ^c	250 ^c	216	255		216
m.....	131	97 ^c	131	131	97	131	131
n.....	170 ^c	170	131	97	170	170	97
o.....	159	159	120	86	159	159	86
p.....	130	130	91	57	130	130	57
other.....	117	117	98	505 ^d	117	117	519 ^d
				491 ^e	85 ^f	85 ^f	505 ^e
				477 ^f			491 ^f

^aFragment ions refer to structures in Figure 2.

^bHigh resolution mass spectral data obtained.

^cWeak ion intensity in some spectra.

^dM⁺-15.

^eM⁺-29.

^fM⁺-43.

^gFragments from rearrangement of BP, diagnostic of *N*-alkylated amino acid *N*-terminal moiety: *m/e* 58, MeLeu; 72, Me₂Leu; 85, Me₂Val and Me₂Ileu. Taken from ref. (5).

5-β-Indolylmethyl-8-*N*-(*N*',*N*'-dimethylisoleucyl)-9-isopropylphenylcyclopeptide (Discarine B) 5. C₃₃H₄₅N₅O₄; μmp 233°, lit (2) mp 235-236°; ms: M⁺ C₃₃H₄₅N₅O₄ requires 573.3315, found 573.3297, M-57 *m/e* 516, bp C₇H₁₅N requires 114.1282 found 114.1284 (see figure 2); ¹H nmr (identical to literature (2,3)), high field region: δ 0.82 (d, 3H, *J*=6.7 Hz, ileu-γ-CH₃), 0.90 (t, 3H, *J*=7.5 Hz, ileu-δ-CH₃), 0.91 (d, 3H, *J*=6.8 Hz, (CH₃)₂CH), 1.18 (d, 3H, *J*=6.8 Hz, (CH₃)₂CH).

5-β-Indolylmethyl-8-*N*-(*N*',*N*'-dimethylvalyl)-9-phenylphenylcyclopeptide (Integerrine) 6. C₃₅H₃₉N₅O₄; μmp 246°, lit (4) mp 258°; ms: M⁺ C₃₅H₃₉N₅O₄ requires 593.3002, found 593.2924, M-43 *m/e* 550, bp C₆H₁₄N requires 100.1126, found 100.1127 (see figure 2); ¹H nmr, high field region: δ 0.16 (d, 3H, *J*=6.8 Hz, val-γ-CH₃), 0.70 (d, 3H, *J*=6.8 Hz, val-γ-CH₃).

5-Isobutyl-8-*N*-(*N*',*N*'-dimethylisoleucyl)-9-phenylphenylcyclopentene (Integerrenine) 7. C₃₁H₄₂N₄O₄; μmp 259°, lit (5) mp 278°; ms: M⁺ C₃₁H₄₂N₄O₄ requires 534.3205, found 534.3200, M-57 C₂₇H₃₂N₄O₄ requires 477.2502, found 477.2515, bp C₇H₁₅N requires 114.1282, found 114.1283 (see figure 2); ¹H nmr (identical to literature (5)), high field region: δ 0.36 (d, 3H, *J*=6.7 Hz, ileu-γ-CH₃), 0.78 (d, 3H, *J*=6.5 Hz, leu-δ-CH₃), 0.85 (d, 3H, *J*=6.5 Hz, leu-δ-CH₃), 0.86 (t, 3H, *J*=7.3 Hz, ileu-δ-CH₃).

DISCUSSION

The identification of the hplc-purified constituents of *C. integerrimus* is based mainly on their characteristic electron impact mass spectra. According to the fragmentation schemes previously proposed (6, 7) (figure 2), the mass spectra of the seven alkaloids from the three plants (table 2) confirm the structural assignments made in table 1. In addition, the total alkaloid acid hydrolytic products from each plant (table 3) are consistent with the distribution of phenylcyclopeptides




FIG. 2. Electron impact mass spectral fragmentation of phencyclopeptides.

1-7 among the three plants shown in figure 3. Since tryptophan is destroyed by acid hydrolysis and *N*-alkylated amino acids are not detected by the usual automatic amino acid analysis due to their low color yield, the failure to detect any other amino acids in the acidic hydrolysate of *C. integerrimus* var. *integerrimus* (Mendocino Co.) corroborates the observation of only indolic phenylcyclopeptide components in this plant.

Leucine, isoleucine, and valine and their methylated derivatives were distinguished from one another by mass spectroscopy and ^1H nmr spectroscopy, and amino acid analysis in some cases. Amino acid analysis of the acid hydrolysates of each hplc-purified phenylcyclopeptide confirmed the identity of the ring amino acid (R_5) suggested by mass spectroscopy. Fragments produced from the rearrangement of the base peak (BP, a) in the mass spectrum of the phenylcyclopeptide provided diagnostic evidence for the structure of the *N*-alkylated amino acid residue, R_8 (table 2).

^1H nmr spectroscopy furnished the most definitive information regarding the nature of the *N*-terminal amino acid moiety (R_8), since the two methyl groups of isoleucine manifest different multiplicity in their nmr signals, the γ -methyl being a doublet and the δ -methyl a triplet. Both the δ -methyls of leucine and the γ -methyls of valine are two sets of doublets. Furthermore, the chemical shifts of the methyl groups on the *N*-terminal amino acid (R_8) are also diagnostic. In the cases of phenylcyclopeptides where R_5 is phenyl, a pronounced upfield shift (as much as 0.6 ppm) has been observed in the *N*-methyl and γ -methyl resonances of the *N*-terminal amino acids (5). Such high field resonances do not occur in the spectra of alkaloids which have *N*-terminal leucine residues since there are no γ -methyl groups. Thus the chemical shift of the leucine δ -methyls in creatine A 9 occurs within the expected range (8), two doublets at δ 0.78 and 0.83 ppm in CDCl_3 ; whereas the doublet occurring at 0.24 ppm in the spectrum of integerrine 6 is attributable to the γ -methyl of the *N*-terminal isoleucine residue.

Our observation of unusually high field doublets in the spectra of phenylcyclopeptides 1, 3, 4, and 7, as well, establishes that the *N*-terminal amino acids are either derivatives of valine or isoleucine. Such high field resonances were not observed in the ^1H nmr spectrum of discarine B 5 and phenylcyclopeptide 2 in agreement with the literature (2, 3).

It is unusual that only one of the phenylcyclopeptides, discarine B 5, found in *C. integerrimus* var. *integerrimus* from Santa Cruz Co. was observed in the extract of the plant of the same species from Mendocino Co. (figure 3). In contrast, the

TABLE 3. Amino acid identification and yields from acid hydrolysis of *C. integerrimus* root bark total alkaloid mixtures.^a

Product	<i>C. integerrimus</i> var. <i>integerrimus</i> Santa Cruz Co.	<i>C. integerrimus</i> var. <i>integerrimus</i> Medocino Co.	<i>C. integerrimus</i> var. <i>californicus</i>
NH ₃	769	516	590
Ile.....	—	—	6
Leu.....	352	—	190
Phe.....	29	—	26

^aCrude alkaloid mixtures (250 mg) were hydrolyzed with 1-2 ml 6N HCl containing 1 drop glacial acetic acid for solubilization in sealed ampules for 24 hr at 135°C. Yields are reported in nanomoles/250 mg mixture hydrolyzed.





FIG. 3. Phencyclopeptine distribution in *C. integrerrimus* var. *integrerrimus* H. and A. from Santa Cruz Co., *C. integrerrimus* var. *integrerrimus* H. and A. from Mendocino Co., and *C. integrerrimus* var. *californicus* (Kell) and G. T. Benson.

total alkaloidal mixture from var. *integrerrimus* of Santa Cruz Co. and that from var. *californicus* continued four common phencyclopeptines 3, 4, 5, and 7.

Integerressine 8 has been reported as the major alkaloid of *C. integrerrimus* var. *integrerrimus* roots, integerrenine 7 as a minor alkaloid, and integerrine 6 as a trace component (4, 5). Our results are different from this reported estimation. In the extract of *C. integrerrimus* var. *integrerrimus* from Santa Cruz Co., integerrenine 7 was the major alkaloid, whereas integerrine 6 and integerressine 8 were absent. On the other hand, integerrine 6 was the major constituent of *C. integrerrimus* var. *integrerrimus* obtained from Mendocino Co.

Conservative botanical opinion has been that the polymorphic forms of *C.*



integerrimus may represent responses to varying amounts of moisture and, therefore, should be included in a single species, *C. integerrimus* H. and A. (1). It is possible that qualitative differences in alkaloid composition between plants from different populations of *C. integerrimus* may similarly reflect the response of the plants to local environmental conditions.

The phytochemical investigation of *C. integerrimus* also poses a difficult challenge to both the botanist and the chemist because interspecific hybridization within the genus *Ceanothus* is widespread. Thus the variation in the alkaloidal characters could be representative of the degree of interspecific hybridization in *Ceanothus*. This concept might explain the disparities among the alkaloid contents of the three examples of *C. integerrimus* var. *integerrimus* examined in this investigation and those observed by others (4, 5). Furthermore, the reported association of nitrogen-fixing actinomycetes with the roots of *Ceanothus* (9), as well as with other plants which produce phenylcyclopeptines, may implicate the symbionts in the production of cyclopeptide alkaloids. These intriguing possibilities further complicate the phytochemical investigation of *C. integerrimus* and will be addressed in future studies of *Ceanothus*.

The chemotaxonomic use of the phenylcyclopeptines must rely upon the examination of many plants from each different population of *C. integerrimus*. The procedure outlined here, involving standardized isolation, hplc purification, and mass spectral identification, provides a quick and objective means upon which to base plant taxonomic and evolutionary relationships.

ACKNOWLEDGMENTS

The authors thank Dr. L. R. Heckard, University of California, Berkeley, California, and Dr. M. A. Nobs, Carnegie Institution of Washington, Stanford, California, for plant identification and enlightening discussions of basic plant taxonomy. Willy C. Shih was of great help in the acquisition of nmr spectra.

Received 30 November 1978.

LITERATURE CITED

1. M. VAN RENSSLAER and H. E. McMINN, "*Ceanothus*", Santa Barbara Botanic Garden, Santa Barbara, Calif., 1942, p. 179-83.
2. O. A. MASCARETTI, V. M. MERKUZA, G. E. FERRARO, E. A. RUVEDA, C.-J. CHANG, and E. WENKERT, *Phytochemistry*, **11**, 1133 (1972).
3. C.-J. CHANG, E. W. HAGAMAN, E. WENKERT, M. G. SIERRA, O. A. MASCARETTI, V. M. MERKUZA, and E. A. RUVEDA, *Phytochemistry* **13**, 1273 (1974).
4. R. TSCHESCHE, E. FROBERG and H.-W. FEHLHABER, *Tetrahedron Lett.*, 1311 (1968).
5. R. TSCHESCHE, J. RHEINGANS, H.-W. FEHLHABER and G. LEGLER, *Chem. Ber.*, **100**, 3924 (1967).
6. R. TSCHESCHE and E. U. KAUSSMAN, "The Alkaloids", Ed. R. H. F. Manske, Vol. XV, Academic Press, New York, N.Y., 1975, p. 165.
7. F. K. KLEIN and H. RAPOPORT, *J. Am. Chem. Soc.*, **90**, 2398 (1968).
8. M. SILVA, D. S. BHAKUNI, P. G. SAMMES, M. PAIS and F. X. JARREAU, *Phytochemistry* **13**, 861 (1974).
9. G. BOND, "The Biology of Nitrogen Fixation", Ed. A. Quispel, North Holland Publishing Co., Amsterdam, 1974, p. 342.