- 8. J.G. Diaz, B.M. Fraga, A.G. González, M.G. Hernández, and A. Perales, *Phytochemistry*, 25, 1161 (1986).
- 9. Y. Hashidoko, S. Tahara, and J. Mizutani, Phytochemistry, 28, 425 (1989).
- T. Ishida, H. Ohishi, M. Inoue, M. Kamigauchi, M. Sugiura, N. Takao, S. Kato, Y. Hamada, and T. Shioiri, J. Org. Chem., 54, 5337 (1989).

Received 27 November 1989

ERRATA

For the paper by Guerriero *et al.* entitled "Hydroxyicosatetraenoic, Hydroxyicosapentaenoic, Hydroxydocosapentaenoic, and Hydroxydocosahexaenoic Acids from the Sponge *Echinochalina mollis* of the Coral Sea," *J. Nat. Prod.*, **53**, 57 (1990), the authors request the following addition and deletions:

On pages 57 and 61, the $12R^*$ stereochemical designation for compounds (+)-1, (+)-3, and (+)-4, as well as reference 6 on page 57, line 6, should be deleted.

On page 57 add "(12R)-HETE has also been reported to have endogenous origins (1)."

The absolute configuration of compound (+)-1, isolated as a semisynthetic derivative from the red seaweed Murrayella periclados, has been extrapolated to be 12S from optical rotation data in comparison with (12S)-HETE methyl ester [M. Bernart and W.H. Gerwick, Tetrahedron Lett., 29, 2015 (1988)].

For the paper by Vasanth et al., entitled "Isolation and Characterization of Vicodiol, a New Monoterpenediol from Vicoa indica," J. Nat. Prod., 53, 354 (1990), the following figure was omitted from the paper:




FIGURE 1. Stereoviews of molecules A, B, and C in the crystal structure of vicodiol.