

Errata. Studies on Peroxides of *Artemisia lancea*

Luo Shide, Ning Ning Bingmei, Hu Wangyu, and Xie Jinlun

J. Nat. Prod., **1991**, 54 (5), 1415-1415• DOI: 10.1021/np50077a600

• Publication Date (Web): 08 September 2004

Downloaded from <http://pubs.acs.org> on April 4, 2009

More About This Article

The permalink <http://dx.doi.org/10.1021/np50077a600> provides access to:

- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article



ACS Publications

High quality. High impact.

Journal of Natural Products is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

17-Oxosparteine (1 mg), α -isolupanine (3 mg), and *N*-methylcytisine (2 mg) were identified by ms and tlc comparisons with authentic samples (6,8).

Catalytic reduction (2 mg) of 5,6-dehydrolupanine (4 mg), oil, ms (9) with 5% Pd/C in glacial HOAc afforded lupanine, compared by ms and tlc.

13-Hydroxysparteine (65 mg), mp 151–153°, $[\alpha]_D +25^\circ$ ($c = 0.01$, MeOH), compared by ir, ms (6) and R_f values with 13-hydroxysparteine [Si gel, petroleum ether-diethylamine (2:0.5)] obtained after reduction of 13-hydroxylupanine with LiAlH₄ in THF.

LITERATURE CITED

1. B.K. Shishkin, *Flora USSR*, **11**, 37 (1945).
2. F. Bohlmann, E. Winterfeldt, and H. Bräckel, *Chem. Ber.*, **91**, 2194 (1958).
3. W. Wysocka, *Bull. Acad. Polon. Sci. Ser. Sci. Chim.*, **27**, 343 (1979).
4. H.M. Fales, H.A. Lloyd, and G.W.A. Milne, *J. Am. Chem. Soc.*, **92**, 1590 (1970).
5. A.S. Sadykov, H.A. Aslanov, and Y.K. Kushmuradov, "Alkaloidi Quinolizidinovogo Ryada," Nauka, Moskow, 1975, p. 75.
6. N. Neuner-Jehle, H. Nesvadba, G. Spittel, *Monatsh. Chem.*, **95**, 687 (1964).
7. F. Bohlmann and E. Winterfeldt, *Chem. Ber.*, **93**, 1956 (1960).
8. D. Schumann, N. Neuner-Jehle, and G. Spittel, *Monatsh. Chem.*, **99**, 390 (1968).
9. Y.D. Cho and R.O. Martin, *Can. J. Chem.*, **49**, 265 (1971).

Received 29 November 1990

ERRATA

For the paper by Were *et al.*, entitled "Pyrrolizidine Alkaloids of *Senecio hadiensis*," *J. Nat. Prod.*, **54**, 491 (1991), the physical properties reported for hadienecine [**10**] were those of a salt. The free base has the following properties: mp 140–141° (Me₂CO), ν max (KBr) 3343 (br s), 2945–2885 (br s), 1350 (w), 1341 (w), 1319 (w), 1281 (m), 1260 (s), 1192 (s), 1125 (s), 1076 (w), 1061 (w), 1045 (s), 1020 (s), 974 (w), 882 (w), 835 (s), 746 (s), 669 (m), 602 (m), 534 (m), 488 (m), 475 (m) cm⁻¹; ¹H-nmr (pyridine-*d*₅, ref. δ 123.5) δ 82.4 (C-1), 36.9 (C-2), 55.5 (C-3), 54.7 (C-5), 37.8 (C-6), 70.9 (C-7), 80.5 (C-8), 66.4 (C-9) ppm; ¹H nmr (pyridine-*d*₅, ref. 7.19) δ 2.65 (1H, ddd, $J = 9.6, 9.6$, and 12 Hz, H_a-2), 1.99 (1H, ddd, $J = 2.3, 6.8$, and 12 Hz, H_b-2), 3.77 (1H, ddd, $J = 9.8, 9.8$, and 6.8 Hz, H_a-3), 2.95 (1H, ddd, $J = 9.6, 9.6$, and 2.3 Hz, H_b-3), δ 3.36 (1H, dd, $J = 8.0$ and 8.3 Hz, H_a-5), 3.12 (1H, ddd, $J = 6.3, 8.3$, and 11.4 Hz, H_b-5), 2.04 (1H, dd, $J = \text{ca. } 6$ and ca. 12 Hz, H_a-6), 1.90 (1H, dddd, $J = 3.4, 8.2, 11.6$, and 11.9 Hz, H_b-6), 4.59 (1H, dd, $J = \text{ca. } 3$ and ca. 3 Hz, H-7), 3.76 (1H, d, $J = 3$ Hz, H-8), 4.49 (1H, d, $J = 10.9$ Hz, H_a-9), 4.46 (1H, d, $J = 10.9$ Hz, H_b-9); eims *m/z* (%) [M]⁺ 173.1050 (5) (calcd for C₈H₁₅NO₃, 173.1052), 156 (12), 155 (40), 129 (28), 112 (20), 99 (92), 98 (95), 82 (100), 56 (25), 55 (25), 51 (8), 41 (43); hydrochloride salt mp 160–161° (EtOH/Et₂O).

Also, in Table 2, the missing resonance for the acetyl carbonyl is δ 170.5 ppm.

For the paper by Shide *et al.*, *J. Nat. Prod.*, **54**, 573 (1991) the title should read "Studies on Peroxides of *Artemisia lancea*."