

## Isolation and Synthesis of a New 9,11-Secosterol from the Sponge *Spongia officinalis*

Riccardo Adinolfi, Anna Migliuolo, Vincenzo Piccialli, and Donato Sica

*J. Nat. Prod.*, **1994**, 57 (9), 1220-1226 • DOI:

10.1021/np50111a005 • Publication Date (Web): 01 July 2004

Downloaded from <http://pubs.acs.org> on April 4, 2009

### More About This Article

---

The permalink <http://dx.doi.org/10.1021/np50111a005> provides access to:

- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article



**ACS Publications**  
High quality. High impact.

Journal of Natural Products is published by the American  
Chemical Society, 1155 Sixteenth Street N.W., Washington,  
DC 20036

ISOLATION AND SYNTHESIS OF A NEW 9,11-SECOSTEROL  
FROM THE SPONGE *SPONGIA OFFICINALIS*

RICCARDO ADINOLFI, ANNA MIGLIUOLO, VINCENZO PICCIALI, and DONATO SICA\*

*Dipartimento di Chimica Organica e Biologica, Università degli Studi di Napoli Federico II,  
via Mezzocannone 16, 80134 Napoli, Italy*

ABSTRACT.—A new 9,11-secosterol [**1**] has been isolated from the sponge *Spongia officinalis*. The structure of the new metabolite has been assigned by interpretation of spectral data and confirmed by synthesis starting from 7-dehydrocholesterol.

The sponge *Spongia officinalis* L. (order Dictyoceratida, family Spongiidae) is a particularly rich source of steroidal metabolites. Previous studies on this organism have led to the isolation of a number of new polyoxygenated steroids, namely,  $\Delta^7$ -3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -trihydroxysterols (1),  $\Delta^7$ -3 $\beta$ ,5 $\alpha$ ,6 $\beta$ ,9 $\alpha$ -tetrahydroxysterols (2),  $\Delta^8$ - and  $\Delta^{8(14)}$ -5 $\alpha$ ,6 $\alpha$ -epoxysterols (3), and 9,11-secosterols (4,5). The sponge also produces  $\Delta^{5,7}$ -sterols (6) which can be seen hypothetically as biogenetic precursors of all of the above-mentioned polyoxysteroids.

A reinvestigation of the polar fractions of the extracts of this sponge has now led to the isolation of a further 9,11-secosterol [**1**], related to the previously isolated 9,11-secosterols from the same source, whose structure elucidation and synthesis are discussed in the present paper.

## RESULTS AND DISCUSSION

Compound **1** was obtained as a crystalline, optically active substance that gave no molecular ion peak in the eims spectrum. Nevertheless, a careful examination of the ms and  $^{13}\text{C}$ - and  $^1\text{H}$ -nmr spectra of **1** (the latter two recorded in pyridine-*d*<sub>5</sub>, Table 1) unequivocally established a molecular formula of  $\text{C}_{29}\text{H}_{46}\text{O}_6$  for the new metabolite. The high-field region of the  $^1\text{H}$ -nmr spectrum displayed four methyl resonances (two singlets at  $\delta$  1.73 and 0.90, and two doublets at  $\delta$  0.97 and 0.85, the latter integrating for six protons), which suggested that the new metabolite was a steroid belonging to the cholestane series. Ir absorptions at 3452, 1725, 1713, 1679, and 1265  $\text{cm}^{-1}$  indicated the presence of ester,  $\alpha$ , $\beta$ -unsaturated ketone, and hydroxyl functions in the molecule. The uv absorption at 231 nm ( $\epsilon$  2977) and  $^{13}\text{C}$ -nmr ( $\text{CDCl}_3$ ) resonances at  $\delta$  203.95 (s, C-9), 139.11 (d, C-7) and 136.80 (s, C-8) confirmed the presence of the enone moiety. In the  $^{13}\text{C}$ -nmr spectrum recorded in pyridine-*d*<sub>5</sub>, the signal for C-8 is submerged by the solvent signal at  $\delta$  135.5. The ester function was part of an acetate group because the  $^1\text{H}$ -nmr spectrum displayed a three-proton singlet signal at  $\delta$  2.00 (acetoxymethyl), while the mass spectrum included an  $\text{M}^+$ - $\text{CH}_3\text{COOH}$  ion peak at  $m/z$  430. The eims also revealed the presence of two hydroxyl groups in **1** exhibiting fragment ions at  $m/z$  472 ( $\text{M}^+$ - $\text{H}_2\text{O}$ , highest mass ion observed in the spectrum), 412 ( $\text{M}^+$ - $\text{H}_2\text{O}$ - $\text{CH}_3\text{COOH}$ )




TABLE 1.  $^1\text{H}$ - and  $^{13}\text{C}$ -nmr Data for Compound 1.<sup>a</sup>

Position	$\delta_c^b$	$\delta_H^c$ (m, J)
1 .....	28.29	H <sub>ax</sub> 2.67 (ddd, 13.9, 13.9, 4.0)
2 .....	27.29	
3 .....	72.31 <sup>d</sup>	5.73 (m)
4 .....	36.89	H <sub>ax</sub> 2.75 (dd, 12.6, 12.6) H <sub>eq</sub> 2.38 (br dd, 12.6, 4.3)
5 .....	76.38	
6 .....	71.49 <sup>d</sup>	4.71 (br d, 5.1)
7 .....	142.35	6.94 (d, 5.1)
8 .....	*	
9 .....	205.15	
10 .....	46.54	
11 .....	204.09	10.30 (d, 3.5)
12 .....	50.92	Ha 2.44 (dd, 16.3, 3.5) Hb 2.31 (d, 16.3)
13 .....	48.87	
14 .....	43.78	3.94 (dd, 11.8, 8.0)
15 .....	26.58 <sup>e</sup>	
16 .....	26.45 <sup>e</sup>	
17 .....	51.95	
18 .....	16.57	0.90 (s)
19 .....	21.70 <sup>f</sup>	1.73 (s)
20 .....	35.20	
21 .....	19.40	0.97 (d, 6.7)
22 .....	35.73	
23 .....	24.53	
24 .....	39.63	
25 .....	28.15	
26 .....	22.66 <sup>g</sup>	0.85 (d, 6.7)
27 .....	22.90 <sup>g</sup>	0.85 (d, 6.7)
OH .....		7.49 (s), 6.69 (s)
CH <sub>3</sub> CO- .....	21.85 <sup>f</sup>	2.00 (s)
CH <sub>3</sub> CO- .....	170.40	

<sup>a</sup> $^1\text{H}$ - and  $^{13}\text{C}$ -nmr spectra were recorded in pyridine-*d*<sub>5</sub> at 400 and 100.1 MHz, respectively. The signal indicated with an asterisk overlapped with the solvent signal at  $\delta$  135.5. This carbon resonated at  $\delta$  136.80 in the spectrum recorded in CDCl<sub>3</sub>.

<sup>b</sup>Assignment based on DEPT experiments and comparison with literature data (2,3) and 3 $\beta$ ,6 $\alpha$ -dihydroxy-9-oxo-9,11-seco-5 $\alpha$ -cholest-7-en-11-al [2] (4) taken as a model compound. Pyridine-*d*<sub>5</sub> as internal reference (149.9 ppm).

<sup>c</sup>Assignments based on decoupling experiments. Residual pyridine as internal reference (8.71 ppm). Coupling constants are given in Hz.

<sup>d-e</sup>Values with identical superscripts may be interchanged.

and 394 ( $\text{M}^+ - 2\text{H}_2\text{O} - \text{CH}_3\text{COOH}$ ), for two successive  $\text{H}_2\text{O}$  losses. The presence of two exchangeable protons in the high-field region of the pyridine-*d*<sub>5</sub>  $^1\text{H}$ -nmr spectrum of **1** ( $\delta$  7.49 and 6.69) confirmed this deduction. On the other hand,  $^{13}\text{C}$ - and  $^1\text{H}$ -nmr resonances at  $\delta$  204.09 (d), and 10.30 (d,  $J=3.5$  Hz), respectively, indicated the presence of an aldehyde group in the molecule as well.

The placement of the above-mentioned functionalities (2 $\times$ OH, 1 $\times$ CHO, 1 $\times$ OCOCH<sub>3</sub>, 1 $\times$ C=C-C=O) within the steroidal nucleus was established on the basis of the following evidence. A typical seven-line multiplet attributable to the 3 $\alpha$ -H  $^1\text{H}$ -nmr resonance was observed at  $\delta$  5.73. Past experience suggested that the unusually high chemical shift value for this proton could not be solely ascribed to acetylation of the C-3 OH group, but also required the presence in the molecule of an  $\alpha$ -oriented hydroxyl

group at C-5 (7). That C-5 was indeed a non-protonated carbon was confirmed by the multiplicity of the H<sub>2</sub>-4 protons that resonated as two mutually coupled double doublets at  $\delta$  2.75 ( $J=12.6$  and  $12.6$  Hz, H<sub>ax</sub>-4) and  $\delta$  2.38 ( $J=12.6$  and  $4.3$  Hz, H<sub>eq</sub>-4). The second splitting observed for each of these signals was due to coupling with the adjacent 3 $\alpha$ -H proton, as indicated by decoupling experiments. Evidence corroborating the presence of a C-5, axially disposed OH group was given by the strong pyridine-induced shifts experienced by both the H<sub>ax</sub>-1 and H<sub>ax</sub>-3 protons, which resonated downfield at  $\delta$  2.67 (ddd,  $J=13.9$ ,  $13.9$ , and  $4.0$  Hz) and  $5.73$ , respectively [ $\Delta\delta$  (H<sub>ax</sub>-1)= $0.67$ ;  $\Delta\delta$  (H<sub>ax</sub>-3)= $0.64$  in comparison with the spectrum recorded in CDCl<sub>3</sub>], due to their 1,3-diaxial relationship, and hence accounted for the vicinity in space of the hydroxyl group in question (5-OH) (7).


The unsaturation count and consideration of the functionalities present in the molecule indicated a tricyclic skeleton, and therefore a secosterol structure for the metabolite under investigation [**1**]. The similarity of the proton spectrum of **1** with that of compound **2**, a recently isolated 9,11-secosterol from the same sponge (4), strongly suggested that the two steroids were closely related metabolites and that compound **1** was itself a 9,11-secosterol. This suggested the placement of the enone moiety between C-6 and C-10 and the aldehyde group at C-11, as in compound **2**, leaving C-6 as the probable carbon atom carrying the second OH group. These considerations were corroborated by the following spectral data. Decoupling experiments demonstrated that the H-6 proton, resonating at  $\delta$  4.71 (br d,  $J=5.1$  Hz), was coupled with the H-7 olefinic proton centered at  $\delta$  6.94 (d,  $J=5.1$  Hz) and homoallylically coupled with the H-14 proton at  $\delta$  3.94 (dd,  $J=11.8$  and  $8.0$  Hz). The  $\beta$  orientation of the C-6 OH group was indicated by the strong pyridine-induced shifts experienced by both the Me-19 and H<sub>ax</sub>-4 protons ( $\Delta\delta=0.38$  and  $0.55$  ppm, respectively, in comparison with the spectrum recorded in CDCl<sub>3</sub>). On the contrary, in sterol **2**, these protons experienced only limited downfield shifts due to the inversion in the configuration at C-6 ( $\alpha$ -OH), while H<sub>eq</sub>-4 underwent a more consistent pyridine-induced deshielding for the same reason (4).

The presence in **1** of a -CH<sub>2</sub>(12)-CHO(11) moiety was proposed from the following evidence. The H<sub>2</sub>-12 protons resonated as a pair of mutually coupled ( $J=16.3$  Hz) signals centered at  $\delta$  2.44 (dd) and 2.31 (d) with only the proton resonating at lower field coupled with the aldehydic proton at  $\delta$  10.30 (d,  $J=3.5$  Hz), as seen for **2**, while the mass spectrum exhibited intense peaks at  $m/z$  447 ( $M^+ - \text{CH}_2\text{CHO}$ ) and 429 ( $M^+ - \text{CH}_2\text{CHO} - \text{H}_2\text{O}$ ). That this grouping was linked to C-13 followed from the nOe enhancements exhibited by both the H<sub>2</sub>-12 protons when H<sub>3</sub>-18 was irradiated. This nOe experiment also revealed the vicinity in space of Me-18 and H-7 (irradiation on H<sub>3</sub>-18 also resulted in a strong enhancement of the H-7 signal), a fact that further supported a ring-C seco-structure for **1**. In fact, examination of a Dreiding model of the molecule indicated that the protons in question (H-7 and H<sub>3</sub>-18) can be brought near in space only if the right-hand (ring-D-containing) part of the molecule is free to rotate around the 8,14 bond; that is, only if ring C is fragmented at a point of the C-9-C-13 segment (at the C-9-C-11 bond, in our case).

The presence of a cholesterol-type side-chain was indicated by <sup>1</sup>H-nmr and ms evidence. As previously noted, the proton spectrum of **1** included resonances for an isopropyl group (H<sub>3</sub>-26 and H<sub>3</sub>-27) at  $\delta$  0.85 and for a methyl linked to a methine group at  $\delta$  0.97 (H<sub>3</sub>-21), while the ms spectrum exhibited fragment ions at  $m/z$  377 ( $M^+ - \text{C}_8\text{H}_{17}$ ), 341 ( $M^+ - \text{C}_8\text{H}_{17} - 2\text{H}_2\text{O}$ ), 299 ( $M^+ - \text{C}_8\text{H}_{17} - \text{CH}_3\text{COOH} - \text{H}_2\text{O}$ ) and 281 ( $M^+ - \text{C}_8\text{H}_{17} - \text{CH}_3\text{COOH} - 2\text{H}_2\text{O}$ ), that indicated the presence of a C<sub>8</sub>H<sub>17</sub> saturated side-chain.

Final confirmation of the stereostructure of **1** was obtained by synthesis starting from 7-dehydrocholesteryl acetate [**3**] following a synthetic protocol similar to that

previously used for the synthesis of compound **2** (Scheme 1). Thus, 7-dehydrocholesteryl acetate [**3**] was oxidized with  $\text{Na}_2\text{Cr}_2\text{O}_7$  in  $\text{CH}_3\text{COOH}/\text{C}_6\text{H}_6$ , as described by Fieser *et al.* for its  $\Delta^{22}$  C-24 methyl homolog (ergosterol) (**8**), to give  $\alpha$ -ketol **4**, which was reduced with  $\text{NaBH}_4$  in EtOH to afford  $5\alpha$ -cholest-7-ene- $3\beta,5,6\beta$ -triol 3-acetate [**5**]. Reaction of **5** with  $\text{Hg}(\text{OAc})_2/\text{CH}_3\text{COOH}$  in  $\text{CHCl}_3$  (**9**) yielded the 7,9(11)-diene **6**, a key intermediate when the scission of the 9(11)-double bond is to be accomplished. Dihydroxylation at the C-9-C-11 positions was obtained by reacting **6** with  $\text{OsO}_4$  in 1,4-dioxane (**10**) for 1 h followed by treatment with  $\text{NaHSO}_3$ , which gave the tetrahydroxysterol **7**, along with a small amount of the  $\alpha,\beta$ -unsaturated ketone **8** derived from further oxidation at the C-6 carbon atom. Compound **8** was easily converted into **7** by  $\text{NaBH}_4$  reduction in EtOH for 30 min. Finally, compound **7** was reacted with crystalline  $\text{Pb}(\text{OAc})_4$  in  $\text{CH}_3\text{COOH}$  (**11**) for 5 min. This reaction led to the



SCHEME 1

exclusive scission of the C-9-C-11 bond leaving the C-5-C-6 trans-diol system unaffected, to give a product which had spectral ( $^1\text{H}$ -nmr,  $^{13}\text{C}$ -nmr, ir, uv, ms) and chromatographic properties identical to those exhibited by natural **1**. In addition, the optical rotations of the synthetic and natural materials were identical, thus establishing that the absolute configuration of the new compound is the one indicated.

From a biogenetic point of view it seems reasonable that both the 9,11-secosterols **1** and **2** could be derived from a common 5,7,9(11)-triene sterol through oxidation at the C-5 and C-6 carbons, in the case of **1**, or only at C-6, in the case of **2**, with concomitant oxidative cleavage of the 9,11 double bond.

## EXPERIMENTAL

**GENERAL EXPERIMENTAL PROCEDURES.**— $^1\text{H}$ - and  $^{13}\text{C}$ -nmr spectra were recorded on Bruker WM 270 and 400 spectrometers in pyridine-*d*<sub>5</sub> or  $\text{CDCl}_3$ .  $^1\text{H}$ -Nmr chemical shifts were referenced to the residual  $\text{CHCl}_3$  and pyridine signals (7.26 and 8.71 ppm, respectively).  $^{13}\text{C}$ -Nmr chemical shifts were referenced to the solvent signals ( $\text{CDCl}_3$ : 77.0 ppm;  $\text{C}_6\text{D}_6$ : 149.9 ppm). The multiplicity of the  $^{13}\text{C}$ -nmr resonances was determined by DEPT experiments (12). NOe nmr spectra were obtained at 400 MHz in a degassed pyridine solution. Hreims were recorded on a Kratos AEI-MS mass spectrometer. Lreims were recorded on a TRIO 2000 mass spectrometer. Ft-ir spectra were obtained with a Perkin-Elmer 1760-X Ft-ir spectrophotometer. Uv spectra were recorded with a Perkin-Elmer model 550-S spectrophotometer. Hplc separations were performed using a Varian 2510 pump equipped with a Waters R403 differential refractometer. Mps were determined on a Reichert Termovar type 300429 Kofler hot-stage melting point apparatus and are uncorrected. Optical rotations were measured on a Perkin-Elmer model 141 polarimeter. Cc was performed on Merck Si gel 40 (70–230 mesh). Tlc analyses were performed on Merck precoated Si gel F<sub>254</sub> plates (0.25 mm thick).

**EXTRACTION OF THE SPONGE AND ISOLATION OF 1.**—*Spongia officinalis* was collected in the Bay of Naples in March 1990. Reference specimens are on file at the Dipartimento di Chimica Organica e Biologica dell'Università di Napoli. Freshly collected animals (433 g dry wt after extraction) were extracted twice with  $\text{Me}_2\text{CO}$  and twice with  $\text{CHCl}_3$ - $\text{MeOH}$  (1:1). The combined extracts were concentrated under reduced pressure and the aqueous residue was extracted with  $\text{Et}_2\text{O}$ . The combined  $\text{Et}_2\text{O}$  extracts were evaporated and the oily residue (29.64 g) was chromatographed on a Si gel column (600 g, 4 cm diameter) eluted with  $\text{CHCl}_3$  and increasing amounts of  $\text{CH}_3\text{OH}$  in  $\text{CHCl}_3$ , with 200-ml fractions collected. Fractions 23–26 (345 mg), eluted with  $\text{CHCl}_3$ - $\text{CH}_3\text{OH}$  (99:1), were combined and subjected to hplc separation on a Hibar LiChrosorb Si-60 column (250×10 mm; flow 2.5 ml/min) using hexane- $\text{EtOAc}$  (3:7) as the mobile phase. The first eluted fraction from this separation (*R*<sub>f</sub> 8.8 min; 9.7 mg) was further purified by normal-phase hplc on a Hibar LiChrosorb Si-60 column (250×4 mm) eluted with hexane- $\text{EtOAc}$  (6:4) to give pure **1** (3.3 mg).

**3 $\beta$ -Acetoxy-5,6 $\beta$ -dihydroxy-9-oxo-9,11-seco-5 $\alpha$ -cholest-7-ene-11-ol [1].**—Mp 164–165° [petroleum ether (80–100°)- $\text{CHCl}_3$ , 8:2]; [ $\alpha$ ]<sub>D</sub> -52.5° (*c*=0.3,  $\text{CHCl}_3$ ); ir (neat)  $\nu$  max 3452, 1725, 1713, 1679, 1265  $\text{cm}^{-1}$ ; uv (MeOH)  $\lambda$  max 231 nm ( $\epsilon$  2977);  $^1\text{H}$  nmr (pyridine-*d*<sub>5</sub>) and  $^{13}\text{C}$  nmr (pyridine-*d*<sub>5</sub>), see Table 1;  $^1\text{H}$  nmr ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  9.89 (1H, d, *J*=3.7 Hz, H-12), 6.48 (1H, d, *J*=4.9 Hz, H-6), 5.09 (1H, m, H <sub>$\alpha$</sub> -3), 4.06 (1H, d, *J*=4.9 Hz, H <sub>$\alpha$</sub> -6), 3.58 (1H, dd, *J*=11.0 and 11.0 Hz, H-14), 2.27 (1H, dd, *J*=15.9 and 3.7 Hz, H <sub>$\alpha$</sub> -12), 2.20 (1H, dd, *J*=12.8 and 12.8 Hz, H <sub>$\alpha$</sub> -4), 2.06 (m, H <sub>$\alpha$</sub> -2, overlapped with other signals), 2.05 (3H, s, acetate), 2.01 (1H, d, *J*=15.9 Hz, H <sub>$\beta$</sub> -12), 2.00 (m, H <sub>$\alpha$</sub> -1, overlapped with other signals), 1.89 (1H, dd, *J*=12.8 and 4.9 Hz, H <sub>$\alpha$</sub> -4), 1.63 (1H, dddd, *J*=12.8, 12.8, 12.8, and 4.3 Hz, H <sub>$\alpha$</sub> -2), 1.35 (3H, s, H <sub>$\beta$</sub> -19), 0.92 (3H, d, *J*=6.7 Hz, H <sub>$\beta$</sub> -21), 0.86 (6H, d, *J*=6.7 Hz, H <sub>$\beta$</sub> -26 and H <sub>$\beta$</sub> -27), 0.79 (3H, s, H <sub>$\beta$</sub> -18); eims *m/z* 472 ( $\text{M}^+$ - $\text{H}_2\text{O}$ , 10), 448 ( $\text{M}^+$ - $\text{CH}_3\text{CO}$ , 5), 447 ( $\text{M}^+$ - $\text{CH}_2\text{CHO}$ , 12), 430 ( $\text{M}^+$ - $\text{CH}_3\text{COOH}$ , 8), 429 ( $\text{M}^+$ - $\text{CH}_2\text{CHO}$ - $\text{H}_2\text{O}$ , 15), 412 ( $\text{M}^+$ - $\text{H}_2\text{O}$ - $\text{CH}_3\text{COOH}$ , 18), 394 ( $\text{M}^+$ - $2\text{H}_2\text{O}$ - $\text{CH}_3\text{COOH}$ , 20), 377 ( $\text{M}^+$ -side chain, 12), 369 ( $\text{M}^+$ - $\text{CH}_2\text{CHO}$ - $\text{H}_2\text{O}$ - $\text{CH}_3\text{COOH}$ , 35), 351 ( $\text{M}^+$ - $\text{CH}_2\text{CHO}$ - $2\text{H}_2\text{O}$ - $\text{CH}_3\text{COOH}$ , 20), 341 ( $\text{M}^+$ -side chain- $2\text{H}_2\text{O}$ , 10), 299 ( $\text{M}^+$ -side chain- $\text{CH}_3\text{COOH}$ - $\text{H}_2\text{O}$ , 15), 281 ( $\text{M}^+$ -side chain- $\text{CH}_3\text{COOH}$ - $2\text{H}_2\text{O}$ , 20); hreims *m/z* 472.3175 ( $\text{M}^+$ - $\text{H}_2\text{O}$ ),  $\text{C}_{29}\text{H}_{44}\text{O}_5$ , requires 472.3189.

**SYNTHESIS OF 1 (SCHEME 1).**—5 $\alpha$ -Cholest-7-ene-3 $\beta$ ,5,6 $\beta$ -triol 3-acetate [**5**].—5 $\alpha$ -Cholesta-5,7-diene-3 $\beta$ -yl acetate (5.0 g, 12 mmol) was oxidized with  $\text{Na}_2\text{Cr}_2\text{O}_7$  dihydrate (1.5 g, 5 mmol) according to Fieser's procedure (8). The reaction mixture was chromatographed on a Si gel column (350 g, 4.5 cm i.d.) eluted with increasing amounts of  $\text{Et}_2\text{O}$  in petroleum ether. The fractions eluted with petroleum ether- $\text{Et}_2\text{O}$  (72:28) contained 1.54 g of crude ketone **4** which was subjected to  $\text{NaBH}_4$  reduction without further purification as follows. To a solution of crude **4** (950 mg) in  $\text{EtOH}$  (30 ml), excess  $\text{NaBH}_4$  was added and the suspension stirred at room temperature for 2 h, then excess reagent was destroyed by dropwise addition of  $\text{CH}_3\text{COOH}$ . The reaction mixture was treated with brine (20 ml) and extracted with  $\text{Et}_2\text{O}$  (3×20 ml).

The combined Et<sub>2</sub>O extracts were washed with a saturated NaHCO<sub>3</sub> solution, dried (MgSO<sub>4</sub>), and concentrated. The residue was chromatographed on a Si gel column eluting with petroleum ether/Et<sub>2</sub>O mixtures of increasing polarity. Petroleum ether-Et<sub>2</sub>O (1:1) was used to elute 620 mg of a material which was further separated by hplc on a Hibar LiChrosorb Si-60 (250×10 mm) column using hexane-EtOAc (65:35) as eluent to give 410 mg of pure 5 $\alpha$ -cholest-7-ene-3 $\beta$ ,5,6 $\beta$ -triol 3-acetate [**5**].

Compound **5** exhibited: Mp 236–238° [petroleum ether (80–100°)-CHCl<sub>3</sub>, 8:2]; [ $\alpha$ ]<sub>D</sub> -31.6° ( $c$ =1.5, CHCl<sub>3</sub>); ir (film)  $\nu$  max 3414, 1728, 1714, 1255 cm<sup>-1</sup>; <sup>1</sup>H nmr (CDCl<sub>3</sub>, 270 MHz)  $\delta$  5.33 (1H, br ddd,  $J$ =5.4, 2.4, and 2.4 Hz, H-7), 5.13 (1H, m, H<sub>a</sub>-3), 3.60 (1H, br dd,  $J$ =5.4 and 5.4 Hz, H<sub>a</sub>-6), 2.23 (1H, dd,  $J$ =12.2 and 12.2 Hz, H<sub>a</sub>-4), 2.02 (3H, s, acetate), 1.08 (3H, s, H<sub>3</sub>-19), 0.92 (3H, d,  $J$ =6.7 Hz, H<sub>3</sub>-21), 0.86 (6H, d,  $J$ =6.1 Hz, H<sub>3</sub>-26 and H<sub>3</sub>-27), 0.57 (3H, s, H<sub>3</sub>-18); <sup>13</sup>C nmr (CDCl<sub>3</sub>, 100.1 MHz)  $\delta$  170.67 (s), 143.94 (s), 117.46 (d), 75.67 (s), 73.71 (d), 71.15 (d), 56.26 (d), 54.64 (d), 43.80 (d), 43.23 (s), 39.48 (t), 39.34 (t), 37.16 (s), 36.15 (d), 36.06 (t), 35.81 (t), 32.64 (t), 27.99 (d), 27.76 (t), 26.95 (t), 23.95 (t), 22.94 (t), 22.79 (q), 22.54 (q), 21.95 (t), 21.41 (q), 18.83 (q), 18.60 (q), 12.08 (q); eims  $m/z$  442 (M<sup>+</sup>-H<sub>2</sub>O, 12), 382 (M<sup>+</sup>-H<sub>2</sub>O-CH<sub>3</sub>COOH, 96), 367 (M<sup>+</sup>-H<sub>2</sub>O-CH<sub>3</sub>COOH-CH<sub>3</sub>, 77), 353 (100), 329 (M<sup>+</sup>-H<sub>2</sub>O-side-chain, 7), 311 (M<sup>+</sup>-side-chain-2H<sub>2</sub>O, 5), 269 (M<sup>+</sup>-side-chain-H<sub>2</sub>O-CH<sub>3</sub>COOH, 35), 251 (M<sup>+</sup>-side-chain-2H<sub>2</sub>O-CH<sub>3</sub>COOH, 27).

5 $\alpha$ -Cholesta-7,9(11)-diene-3 $\beta$ ,5,6 $\beta$ -triol 3-acetate [**6**].—To a solution of 232 mg (0.5 mmol) of 5 $\alpha$ -cholest-7-ene-3 $\beta$ ,5,6 $\beta$ -triol 3-acetate [**5**] in 6.5 ml of CHCl<sub>3</sub>, 368 mg (1.2 mmol) of Hg(OAc)<sub>2</sub> in 12.7 ml of CH<sub>3</sub>COOH was added and the suspension was stirred at room temperature (approximately 25°) for 20 h and then filtered. The filtrate was treated with saturated aqueous NaHCO<sub>3</sub> solution and extracted with CHCl<sub>3</sub>. The extract was washed with H<sub>2</sub>O, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. The residue (215 mg) was chromatographed over a Si gel column eluted with petroleum ether-Et<sub>2</sub>O (4:6) to give 163 mg of a mixture which, on tlc analysis (CHCl<sub>3</sub>-MeOH, 95:5), was shown to be mainly composed of two uv-visible spots. Hplc separation by reversed-phase hplc on a Hibar LiChrosorb RP-18 (250×10 mm) column (eluent MeOH-H<sub>2</sub>O, 96:4) gave 40 mg of 5 $\alpha$ -cholesta-7,9(11)-diene-3 $\beta$ ,5,6 $\beta$ -triol 3-acetate [**6**] still contaminated by another product. Final purification of **6** was achieved by analytical tlc (CHCl<sub>3</sub>-EtOAc, 85:15) thus affording 25 mg of pure **6**: mp 184–185° [petroleum ether (80–100°)-CHCl<sub>3</sub>, 8:2]; [ $\alpha$ ]<sub>D</sub> +11.1° ( $c$ =0.8, CHCl<sub>3</sub>); ir (neat)  $\nu$  max 3597, 3446, 1713, 1278 cm<sup>-1</sup>; uv (CH<sub>3</sub>OH)  $\lambda$  max 245 nm ( $\epsilon$  10086); <sup>1</sup>H nmr (CDCl<sub>3</sub>, 270 MHz)  $\delta$  5.72 (1H, br d,  $J$ =6.8 Hz, H-11), 5.42 (1H, br d,  $J$ =5.9 Hz, H-7), 5.17 (1H, m, H<sub>a</sub>-3), 3.81 (1H, br d,  $J$ =5.9 Hz, H<sub>a</sub>-6), 2.38 (1H, dd,  $J$ =17.6 and 6.8 Hz, Ha-12), 2.14 (1H, br d,  $J$ =17.6 Hz, Hb-12), 2.03 (3H, s, acetate), 1.28 (3H, s, H<sub>3</sub>-19), 0.91 (3H, d,  $J$ =6.3 Hz, H<sub>3</sub>-21), 0.86 (6H, d,  $J$ =6.8 Hz, H<sub>3</sub>-26 and H<sub>3</sub>-27), 0.57 (3H, s, H<sub>3</sub>-18); <sup>13</sup>C nmr (CDCl<sub>3</sub>, 67.9 MHz)  $\delta$  170.57 (s), 139.91 (s), 138.92 (s), 126.37 (d), 118.10 (d), 75.04 (s), 73.89 (d), 71.08 (d), 56.32 (d), 51.29 (d), 42.65 (s), 42.37 (t), 40.60 (s), 39.47 (t), 35.98 (t), 34.17 (t), 31.06 (t), 28.24 (t), 27.99 (d), 26.91 (t), 26.18 (q), 23.91 (t), 23.10 (t), 22.79 (q), 22.53 (q), 21.40 (q), 18.42 (q), 11.35 (q); eims  $m/z$  458 (M<sup>+</sup>, 4), 440 (M<sup>+</sup>-H<sub>2</sub>O, 5), 425 (M<sup>+</sup>-H<sub>2</sub>O-CH<sub>3</sub>, 4), 398 (M<sup>+</sup>-CH<sub>3</sub>COOH, 20), 380 (M<sup>+</sup>-CH<sub>3</sub>COOH-H<sub>2</sub>O, 65), 365 (M<sup>+</sup>-CH<sub>3</sub>COOH-H<sub>2</sub>O-CH<sub>3</sub>, 28), 327 (M<sup>+</sup>-side-chain-H<sub>2</sub>O, 8), 285 (M<sup>+</sup>-side-chain-CH<sub>3</sub>COOH, 25), 267 (M<sup>+</sup>-side-chain-CH<sub>3</sub>COOH-H<sub>2</sub>O, 38), 95 (100).

5 $\alpha$ -Cholest-7-ene-3 $\beta$ ,5,6 $\beta$ ,9 $\alpha$ ,11 $\alpha$ -pentol 3-acetate [**7**] and 3 $\beta$ -acetoxy-5,9 $\alpha$ ,11 $\alpha$ -trihydroxy-5 $\alpha$ -cholest-7-en-6-one [**8**].—To 14.6 mg (0.03 mmol) of **6** dissolved in 4 ml of freshly distilled 1,4-dioxane, excess osmium tetroxide was added, and the mixture stirred at room temperature for 90 min. Removal of the solvent under reduced pressure gave the crude osmate ester as a dark brown material which was hydrolyzed as follows. The osmate ester was dissolved in 2 ml of 1,4-dioxane and 2 ml of a saturated NaHSO<sub>3</sub> solution were added under stirring. After 15 min the mixture was filtered and the filtrate was washed two times with a 2 N HCl solution, dried (MgSO<sub>4</sub>), and taken to dryness. The residue (15.0 mg) was separated by hplc on a Hibar LiChrosorb Si-60 (250×4 mm) column using hexane-EtOAc (85:15) as eluent to afford 10.6 mg of pure tetrol **7** and 2.5 mg of ketone **8**. NaBH<sub>4</sub> reduction of compound **8** in the same conditions used for the reduction of **4** gave, after hplc purification in the above conditions (hexane-EtOAc, 85:15), 2.2 mg of **7**. Compound **7** exhibited mp 210–211° [petroleum ether (80–100°)-CHCl<sub>3</sub>, 8:2]; [ $\alpha$ ]<sub>D</sub> -4.1° ( $c$ =0.5, CHCl<sub>3</sub>); ir (neat)  $\nu$  max 3401, 1714, 1265 cm<sup>-1</sup>; <sup>1</sup>H nmr (pyridine-*d*<sub>5</sub>, 400 MHz)  $\delta$  5.87 (1H, dd,  $J$ =5.4 and 1.6 Hz, H-7), 5.77 (1H, m, H<sub>a</sub>-3), 4.50 (1H, dd,  $J$ =11.4 and 5.1 Hz, H<sub>a</sub>-11), 4.37 (1H, dd,  $J$ =5.4 and 2.2 Hz, H<sub>a</sub>-6), 2.88 (1H, dd,  $J$ =12.4 and 12.4 Hz, H<sub>a</sub>-4), 2.38 (1H, dd,  $J$ =12.1 and 5.1 Hz, H<sub>eq</sub>-12), 2.33 (1H, dd,  $J$ =12.4 and 5.1 Hz, H<sub>ax</sub>-4), 2.00 (3H, s, acetate), 1.95 (1H, dd,  $J$ =12.1 and 12.1 Hz, H<sub>ax</sub>-12), 1.73 (3H, s, H<sub>3</sub>-19), 0.94 (3H, d,  $J$ =5.7 Hz, H<sub>3</sub>-21), 0.87 (6H, d,  $J$ =6.4 Hz, H<sub>3</sub>-26 and H<sub>3</sub>-27), 0.68 (3H, s, H<sub>3</sub>-18); <sup>13</sup>C nmr (pyridine-*d*<sub>5</sub>, 67.9 MHz)  $\delta$  170.47 (s), 141.58 (s), 123.38 (d), 78.65 (s), 76.96 (s), 73.36 (d), 71.95 (d), 70.04 (d), 56.32 (d), 51.02 (d), 47.10 (t), 42.87 (s), 42.42 (s), 39.71 (t), 37.82 (t), 36.44 (d), 36.29 (t), 29.38 (t), 28.23 (t), 28.23 (d), 28.02 (t), 24.20 (t), 23.51 (t), 22.96 (q), 22.69 (q), 21.40 (q), 21.40 (q), 18.92 (q), 12.78 (q); eims  $m/z$  492 (M<sup>+</sup>, 2), 474 (M<sup>+</sup>-H<sub>2</sub>O, 20), 456 (M<sup>+</sup>-2H<sub>2</sub>O, 18), 414 (M<sup>+</sup>-H<sub>2</sub>O-CH<sub>3</sub>COOH, 7), 396 (M<sup>+</sup>-2H<sub>2</sub>O-CH<sub>3</sub>COOH, 33), 283 (M<sup>+</sup>-side-chain-2H<sub>2</sub>O-CH<sub>3</sub>COOH, 37), 265 (M<sup>+</sup>-side-chain-3H<sub>2</sub>O-CH<sub>3</sub>COOH, 24), 93 (100), 81 (100), 69 (100).

Compound **8** exhibited:  $[\alpha]_D -6.0^\circ$  ( $c=0.2$ ,  $\text{CHCl}_3$ ); ir (neat)  $\nu$  max 3385, 1735, 1683, 1242  $\text{cm}^{-1}$ ; uv ( $\text{CH}_3\text{OH}$ )  $\lambda$  max 232 nm ( $\epsilon$  5820);  $^1\text{H}$  nmr (pyridine-*d*<sub>5</sub>, 400 MHz)  $\delta$  5.97 (1H, br s, H-7), 5.62 (1H, m, H $_{\alpha}$ -3), 4.40 (1H, dd,  $J=12.1$  and 4.4 Hz, H $_{\beta}$ -11), 2.95 (1H, br dd,  $J=8.3$  and 8.3 Hz, H-14), 2.70 (1H, br dd,  $J=12.1$  and 4.4 Hz, H $_{\alpha}$ -4), 2.40 (1H, dd,  $J=12.1$  and 5.1 Hz, H $_{\alpha}$ -12), 2.17 (1H, dd,  $J=12.1$  and 12.1 Hz, H $_{\alpha}$ -4), 2.06 (1H, d,  $J=12.1$  and 12.1 Hz, H $_{\alpha}$ -12), 1.98 (3H, s, acetate), 1.31 (3H, s, H $_{\beta}$ -19), 0.92 (3H, d,  $J=5.7$  Hz, H $_{\beta}$ -21), 0.87 (6H, d,  $J=7.0$  Hz, H $_{\beta}$ -26 and H $_{\beta}$ -27), 0.64 (3H, s, H $_{\beta}$ -18); eims  $m/z$  490 ( $\text{M}^+$ , 1), 472 ( $\text{M}^+ - \text{H}_2\text{O}$ , 12), 430 ( $\text{M}^+ - \text{CH}_3\text{COOH}$ , 5), 412 ( $\text{M}^+ - \text{CH}_3\text{COOH} - \text{H}_2\text{O}$ , 48), 394 ( $\text{M}^+ - \text{CH}_3\text{COOH} - 2\text{H}_2\text{O}$ , 16), 359 ( $\text{M}^+ - \text{side-chain} - \text{H}_2\text{O}$ , 27), 299 ( $\text{M}^+ - \text{side-chain} - \text{H}_2\text{O} - \text{CH}_3\text{COOH}$ , 10), 281 ( $\text{M}^+ - \text{side-chain} - 2\text{H}_2\text{O} - \text{CH}_3\text{COOH}$ , 20), 181 (100), 121 (100), 93 (100).

*Reaction of 5 $\alpha$ -cholest-7-ene-3 $\beta$ ,5,6 $\beta$ ,9 $\alpha$ ,11 $\alpha$ -pentol 3-acetate [7] with lead tetraacetate to produce synthetic 1.*—To a solution of 5 $\alpha$ -cholest-7-ene-3 $\beta$ ,5,6 $\beta$ ,9 $\alpha$ ,11 $\alpha$ -pentol 3-acetate [7] (6.0 mg, 0.012 mmol) in  $\text{CH}_3\text{COOH}$  (1 ml), crystalline lead tetraacetate (6.0 mg) was added portionwise over a 5 min period at room temperature. When the reaction was complete [10 min, tlc analysis ( $\text{CHCl}_3$ - $\text{CH}_3\text{OH}$ , 9:1)], two drops of ethylene glycol were added and the mixture diluted with ice- $\text{H}_2\text{O}$  and extracted with  $\text{CHCl}_3$ . The organic layer was washed with aqueous  $\text{NaHCO}_3$ , dried ( $\text{MgSO}_4$ ), and evaporated. The residue was chromatographed on a Hibar LiChrosorb Si-60 (250 $\times$ 4 mm) column using  $\text{CHCl}_3$ - $\text{CH}_3\text{OH}$  (96:4) as eluent to give 4.0 mg of ketoaldehyde **1** which had spectral ( $^1\text{H}$ -nmr,  $^{13}\text{C}$ -nmr, ir, uv, ms) and chromatographic properties identical to those exhibited by the natural product, **1**. Furthermore, synthetic **1** had  $[\alpha]_D -50.8^\circ$  ( $c=0.4$ ,  $\text{CHCl}_3$ ).

#### ACKNOWLEDGMENTS

We thank Stazione Zoologica di Napoli for supplying the sponge. This work was supported by the Ministero dell' Università e della Ricerca Scientifica e Tecnologica (MURST), Italy. Mass spectral data were provided by the Servizio di Spettrometria di Massa del CNR e dell' Università di Napoli. The assistance of the staff is gratefully acknowledged.

#### LITERATURE CITED

1. A. Madaio, V. Piccialli, D. Sica, and G. Corriero, *J. Nat. Prod.*, **52**, 952 (1989).
2. A. Migliuolo, G. Notaro, V. Piccialli, and D. Sica, *J. Nat. Prod.*, **53**, 1414 (1990).
3. A. Migliuolo, V. Piccialli, D. Sica, and F. Giordano, *Steroids*, **58**, 134 (1993).
4. A. Migliuolo, V. Piccialli, and D. Sica, *Tetrahedron*, **47**, 7937 (1991).
5. A. Migliuolo, V. Piccialli, and D. Sica, *Steroids*, **57**, 344 (1992).
6. M. De Rosa, L. Minale, and G. Sodano, *Comp. Biochem. Physiol. [B]*, **46**, 823 (1973).
7. Y. Fujimoto, T. Yamada, and N. Ikekawa, *Chem. Pharm. Bull.*, **33**, 3129 (1985).
8. M. Fieser, A. Quilico, A. Nickon, W.E. Rosen, E.J. Tarlton, and L.F. Fieser, *J. Am. Chem. Soc.*, **75**, 4066 (1953).
9. W.V. Ruyle, T.A. Jacob, J.A. Chernerda, E.M. Chamberlin, D.W. Rosemburg, G.E. Sita, R.L. Erickson, L.M. Aliminosa, and M. Tishler, *J. Am. Chem. Soc.*, **75**, 2604 (1953).
10. M. Schroeder, *Chem. Rev.*, **80**, 187 (1980).
11. A.H. Haines, "Methods for the Oxidation of Organic Compounds." Ed. by A.R. Katritzky, O. Meth-Cohn, and C.W. Rees, Academic Press, London, 1988, pp. 287-292.
12. M.R. Bendall, D.M. Doddrell, D.T. Pegg, and W.E. Hull, "DEPT Bruker Information Bulletin," Bruker Analytische Masstechnik, Karlsruhe, Germany, 1982.

Received 17 February 1994