

Bromotyrosine Alkaloids from the Sponge *Pseudoceratina verrucosa*

A. Benharref¹ and M. Païs*²

¹Institut de Chimie des Substances Naturelles, CNRS, 91198 Gif-sur-Yvette, France

C. Debitus

²Centre ORSTOM, BP A5, Nouméa, New Caledonia


Received August 29, 1995³

Three new bromotyrosine alkaloids, pseudoceratinines A (**1**), B (**2**), and C (**3**), have been isolated from two specimens of the sponge *Pseudoceratina verrucosa*. In addition, the known alkaloids aplysamine-1 (**4**), aplysamine-2 (**5**), purealin (**6**), and purealidins A (**7**) and B (**8**) were found in both specimens. The structures and absolute configurations of the new products were elucidated by spectral methods.

Bromotyrosine-derived alkaloids are well known in marine Verongida sponges. We report here the isolation and structural elucidation of three new bromotyrosines, pseudoceratinines A (**1**), B (**2**), and C (**3**), from two specimens of the sponge *Pseudoceratina verrucosa* Bergquist (order Verongida, family Aplysiniellidae) collected off the coast of two different New Caledonian islands. Compounds **1** and **2** were isolated from a specimen collected at Ile Longue (Chesterfield Archipelago), whereas compounds **2** and **3** were found in a specimen collected at Ile Walpole. In addition, the known alkaloids aplysamine-1 (**4**),² aplysamine-2 (**5**),² purealin (**6**),³ and purealidins A (**7**)⁴ and B (**8**)⁵ have been isolated from both specimens.

The MeOH extract of each specimen was partitioned between CH₂Cl₂ and H₂O, and the aqueous layer was subsequently extracted with *n*-BuOH. Repeated Si gel column chromatography of the CH₂Cl₂ extracts with CH₂Cl₂/MeOH mixtures yielded aplysamine-2 (**5**). The butanol extracts were first chromatographed on a Sephadex LH-20 column with CH₂Cl₂/MeOH (1:1) and further on a Si gel column with EtOAc/butanone containing increasing amounts of HCO₂H/H₂O (5:3:0.5:0.5 to 5:3:1.5:1.5) to give compounds **1**, **2**, and **4–8** from the specimen from Ile Longue and compounds **2–9** from the specimen from Ile Walpole. The new alkaloids **1–3** were converted into their hydrochloride salts on an Amberlite IR-45 anion exchange column.

Pseudoceratinine A (**1**) showed an isotopic cluster of MH⁺ ions in the ratio 1:2:1, consistent with two bromine atoms, in the FABMS at *m/z* 490, 492, and 494. The molecular formula C₁₅H₁₈Br₂N₅O₄ was determined by HRFABMS (*m/z* 489.9725, MH⁺, Δ 0.2). The ¹H-NMR spectrum exhibited the characteristic signals of a spirocyclohexadiene–isoxazole ring at δ 4.08 (1H, s, H-1), 6.40 (1H, s, H-5) and 3.09, 3.75 (two d, 1H each, *J* = 18 Hz). This partial structure was further supported by the ¹³C resonances for carbons 1–9 (Table 1). The remaining part of **1** consisted of a 2-aminohistamine residue as suggested, in conjunction with the elemental composition of the molecule, by the two triplets at δ 3.51 and 2.76 (2H each), by the singlet at δ 6.55 observed in the ¹H-NMR spectrum, and by analysis of the ¹³C-NMR spectrum (Table 1). Thus, pseudoceratinine A was assigned structure **1**, which differs from the previously



reported aerophobin-2⁶ only by the lack of one methylene in the central chain.

Pseudoceratinine B (**2**) displayed an isotopic cluster of MH⁺ ions at *m/z* 502, 504, and 506 (ratio 1:3:1) in the FABMS. Accurate mass measurement confirmed a formula C₁₈H₂₆Br₂N₅O₂ (*m/z* 502.0432, MH⁺, Δ -2.1). The ¹H-NMR spectrum showed two singlets at δ 3.82 (3H) and 3.32 (9H) corresponding to an MeO and to a Me₃N⁺ group, respectively. Two additional singlets were observed at δ 7.55 (2H) and 6.42 (1H). The signal at δ 7.55 was assigned to a 1,2,4,6-tetrasubstituted aryl ring, while the signal at δ 6.42 was assigned to the heterocyclic proton of a 2-aminoimidazole. This assignment was confirmed by characteristic ¹³C signals (Table 1), which corresponded to the aromatic carbons of an *O*-methyl-3,5-dibromotyrosine moiety and to a 2-aminoimidazole ring, respectively (Table 1). In the NMR spectra, the signals of the CH₂CHNMe₃CO group of the bromotyrosine residue were observed at δ_H 3.30 (2H, m, H-7) and 4.25 (1H, dd, H-8) and δ_C 32.0 (C-7), 75.8 (C-8), and 166.2 (C-9). The 2-aminoimidazole was part of an aminohistamine moiety showing the usual resonances for the CH₂CH₂CO group (Table 1). Thus, pseudoceratinine B was assigned structure **2**, which was further supported by HMBC cross peaks H-7/C-1, C-6, C-8, C-9 and H-8/C-7, C-9, and Me₃.

³ Abstract published in *Advance ACS Abstracts*, January 15, 1996.

Table 1. ^{13}C - (75 MHz) and ^1H -NMR (400 MHz) Data^a for Pseudoceratines A (**1**), B (**2**), and C (**3**)

| 1 | | | 2 | | 3 | | | |
|---------------------|------------------|----------------------------|------------------|------------------------|---------------------|------------------|------------------------------|--------------------------|
| position | δC | δH (JHz) | δC | δH (JHz) | position | δC | δH (JHz) | HMBC |
| 1 | 75.6 | 4.08 s | 134.0 | | 1 | 75.4 | 4.10 s | 3, 4, 5, 6 |
| 2 | 123.1 | | 135.2 | 7.55 s | 2 | 122.7 | | |
| 3 | 149.0 | | 118.9 | | 3 | 149.3 | | |
| 4 | 114.5 | | 154.8 | | 4 | 114.1 | | 1, 2, 3, 4, 7 |
| 5 | 132.6 | 6.40 s | 118.9 | | 5 | 132.3 | 6.40 s | |
| 6 | 92.9 | | 132.5 | 7.55 s | 6 | 92.5 | | |
| 7 | 39.5 | 3.75 d (18) 3.09 d (18) | 32.0 | 3.30 m | 7 | 40.2 | 3.78, d (18) 3.10, d (18) | 1, 5, 6, 8 1, 5, 6, 8 |
| 8 | 155.6 | | 75.8 | 4.25 dd (12, 4) | 8 | 155.3 | | |
| 9 | 162.1 | | 166.2 | | 9 | 161.1 | | |
| | | | | | 10 | 38.3 | 3.60, t (7) | 10,12 |
| | | | | | 11 | 30.6 | 2.12 m | 10,11 |
| | | | | | 12 | 72.3 | 4.02 t (7) | 13 |
| | | | | | 13 | 154.1 | | |
| | | | | | 14,14' | 119.2 | 7.52 s | 13, 14, 16, 17 |
| | | | | | 15,15' | 135.0 | | |
| | | | | | 16 | 136.0 | | |
| | | | | | 17 | 38.0 | 3.07 m | 16, 18, 19 |
| | | | | | 18 | 56.0 | 4.07 m | 16, 17, 19 |
| | | | | | 19 | 170.1 | | |
| 10 | 40.5 | 3.51 t (7) | 38.7 | 3.30 m | 20 | 39.3 | 3.45 m | 19, 21, 22 |
| 11 | 26.1 | 2.76 t (7) | 25.2 | 2.45 m | 21 | 25.7 | 2.65 m | 20, 21, 22 |
| 12 | 126.3 | | 125.3 | | 22 | 125.9 | | |
| 13 | 111.3 | 6.55 s | 110.5 | 6.42 s | 23 | 110.6 | 6.55 s | 22, 24 |
| 14 | 149.7 | | 148.5 | | 24 | 149.9 | | |
| MeO | 60.8 | 3.72 s | 61.0 | 3.82 s | MeO | 60.4 | 3.72 s | |
| Me ₃ N | | | 53.2 | 3.32 s | | | | |
| NH-9 | | 8.75 br s | | 9.20 t (6) | NH-9 | | 8.60 t (6) | |
| | | | | | NH-19 | | 8.70 t (6) | |
| NH-12 ^b | | 11.70 s | | 11.70 s | NH-22 | | 11.80 s | |
| NH-13 ^b | | 12.30 s | | 12.25 s | NH-23 | | 12.35 s | |
| NH ₂ -14 | | 7.45 s | | 7.25 br s | NH ₂ -24 | | 7.35 br s | |
| OH-1 | | 6.55 br s | | | OH-1 | | 6.50 br s | |
| | | | | | NH ₂ -18 | | 4.18 br | |

^a Measurements were taken in CD₄O, except for OH and NH, which were taken in DMSO-*d*₆. Assignments in CD₄O were based on 2D experiments for compounds **2** and **3**. ^b Respective values for NH-12 and NH-13 are from Nakamura *et al.*³

Pseudoceratinine C (**3**) exhibited an isotopic cluster of five MH⁺ ions separated by 2 amu (ratio 1:4:6:4:1) centered at MH⁺ 870 in the FABMS, indicating the presence of four bromine atoms. The molecular formula C₂₇H₃₂Br₄N₇O₆ was established by HRFABMS (*m/z* 869.9088, MH⁺, Δ -1.9). Comparison of the NMR data of **3** (Table 1) with those of the known purealin **6**³ showed similar signals for the C₁-C₁₆ moiety and the aminohistamine residue (C₁₉-C₂₄), respectively. The only difference between the two alkaloids was an 18-CHNH₂ group in **3** instead of the 18-C=NOH group in **6**. In the ^1H -NMR spectrum, the NH₂ signal was observed as a broad singlet at δ_{H} 4.18. The CH-18 group appeared at δ_{C} 56.0 and δ_{H} 4.07 and was coupled in the COSY with CH₂-17 (δ_{C} 38.0 and δ_{H} 3.07). The HBMBC experiment (Table 1), which displayed correlations from H-17 and H-18 to C-16 and C-19, further supported these assignments.


The absolute configuration at C-6 of compounds **1**-**3** was established as illustrated on the basis of the CD spectra, which showed negative Cotton effects near 250 and 290 nm.³ To our knowledge, there is only one report of a bromotyrosine alkaloid containing a nonoxidized amino group. However, the absolute configuration of the α carbon has not been determined.⁷ The stereochemistry of an aromatic amino acid such as tyrosine can be deduced from the Cotton effect observed in the region of 220 nm.⁸ Both pseudoceratinines B (**2**) and C (**3**) exhibited positive Cotton effects at 212 and 215 nm, respectively, corresponding to an *S*-configuration of the tyrosine residue. The Me₃ group did not affect the CD

results, inasmuch as **10**, which was obtained by treatment of 3,5-dibromo-L-tyrosine (**9**)⁹ with MeI,¹⁰ also exhibited a positive Cotton effect in the same region. Acid hydrolysis (HCl 6 N, 24 h, reflux) of pseudoceratinine B (**2**) did not yield *O*-methyl-3,5-dibromo-NMe₃-L-tyrosine (**10**), probably because of steric hindrance.

It is worthwhile to note that the alkaloids **4**-**8**, which have been found previously in two different species [i.e., aplysamine-1 (**4**) and aplysamine-2 (**5**) in *Aplysina* sp. and purealin (**6**) and purealidins A (**7**) and B (**8**) in *Psammaphysilla* (= *Pseudoceratina*) *purea*] were in this case all isolated from a sponge belonging to another species. The specimens of *Pseudoceratina verrucosa* studied here were shown to have a somewhat different metabolic composition, for pseudoceratinine A (**1**) was isolated only from one specimen and pseudoceratinine C (**3**) only from the other. However, this often occurs for specimens coming from different areas and has been observed previously for a species belonging to the same genus, *Pseudoceratina crassa*.¹¹

Experimental Section

General Experimental Procedures. Optical rotations at 20 °C were determined on a Perkin-Elmer 241 polarimeter. UV spectra were recorded on a Shimadzu UV-1616 UV-vis spectrophotometer. CD spectra were recorded on a Jobin-Yvon Mark 5; IR on Nicolet 205 FT-IR spectrometer; FABMS on a Kratos MS 80; HRFABMS on a VG-Zab-Seq spectrometer; and NMR on Bruker AC 300 (^1H and ^{13}C spectra) and AM 400 (2D spectra) spectrometers. UV spectra were recorded in



MeOH. Column chromatography was performed using Si gel Merck H60.

Animal Material. The samples of *Pseudoceratina verrucosa* Bergquist (order Verongida, family Aplysinellidae) were collected at Ile Longue (Chesterfield Archipelago, 15 m depth, sponge I) and at Ile Walpole (8–10 m depth, sponge II) as part of the CNRS-ORSTOM program "Substances Marines d'Intérêt Biologique" (SMIB). Samples of the sponges I (ref R1576) and II (ref R 1593) were identified by P. R. Bergquist and C. Lévi and are preserved at ORSTOM, Nouméa, New Caledonia.

Extraction and Purification. The freeze-dried animal material (225 g of sponge I and 200 g of sponge II) was extracted with 80% EtOH (3 × 2 L) at room temperature. After filtration, each solution was concentrated *in vacuo* to an aqueous suspension, which was extracted with CH₂Cl₂. The organic extract (4.8 g for I, 4.2 g for II) was subjected to Si gel column chromatography with CH₂Cl₂ containing increasing concentrations of MeOH (+ minor amounts of H₂O for the fraction eluted with CH₂Cl₂ containing more than 10% of MeOH). The fractions eluted with CH₂Cl₂/MeOH/H₂O (80:20:1) were further purified by column chromatography on Si gel, yielding aplysamine-2 (5) [eluent CH₂Cl₂/MeOH (95:5)] in yields of 30 mg from sponge I and 14 mg from sponge II. The aqueous layer (from the CH₂Cl₂ extract of each sponge) was extracted with *n*-BuOH. The organic layers were evaporated to give a crude

residue (4.8 g for I; 4.2 g for II). The *n*-BuOH extracts (0.5 g for I, 0.75 g for II) were chromatographed on Sephadex LH 20 with CH₂Cl₂/MeOH (1:1), discarding the first and the last deep-colored fractions. The middle fractions were further chromatographed on Si gel using EtOAc/butanone containing increasing amounts of HCO₂H/H₂O (5:3:0.5:0.5 to 5:3:1.5:1.5). Sponge I yielded purealin A (6) (10 mg, eluent 5:3:0.5:0.5), purealidin B (8) (9 mg, 5:3:0.5:0.5), purealidin A (7) (11 mg, 5:3:1:1), pseudoceratinine B (2) (120 mg, 5:3:1:1), pseudoceratinine A (1) (26 mg, 5:3:1:1), and aplysamine-1 (4) (11 mg, 5:3:1.5:1.5). Sponge II yielded purealin A (6) (22 mg, eluent 5:3:0.5:0.5), purealidin B (8) (14 mg, 5:3:0.5:0.5), pseudoceratinine C (3) (45 mg, 5:3:1:1), purealidin A (7) (11 mg, 5:3:1:1), pseudoceratinine B (2) (120 mg, 5:3:1.5:1.5), and aplysamine-1 (4) (11 mg, 5:3:1.5:1.5). The MeOH solutions of the three alkaloids pseudoceratinins A (1), B (2), and C (3) were passed through a small column of Amberlite IR-45 (Cl⁻) to yield the hydrochloride salts. Pseudoceratinine C HCl (3) was further purified on a C-18 Sep-Pak column (Waters) using MeOH/H₂O as eluent.


Pseudoceratinine A (1): [α]_D -158° (MeOH, *c* 1); UV λ max nm 220 (log ϵ 4.47) and 284 (log ϵ 4.08); IR ν max (KBr) 3400, 1680, 1543, 1437, 1381, 1250 cm⁻¹; CD (MeOH) λ ext 248 ($\Delta\epsilon$ - 9.66) and 290 ($\Delta\epsilon$ - 8.48); FABMS *m/z* 490, 492, 494 (MH⁺); NMR see Table 1.

Pseudoceratinine B (2): [α]_D +17° (MeOH, *c* 1); UV 285 (log ϵ 3.17) and 330 (log ϵ 2.69); IR ν max (KBr) 3406, 1680, 1556, 1475, 1262 cm⁻¹; CD (H₂O) λ ext 212 ($\Delta\epsilon$ + 12.6); FABMS *m/z* 502, 504, 506 (MH⁺); NMR see Table 1.

Pseudoceratinine C (3): [α]_D -10° (MeOH, *c* 1); UV 220 (log ϵ 4.54) and 228 (log ϵ 4.11); IR ν max (KBr) 3375, 1680, 1543, 1456, 1387, 1250 cm⁻¹; FABMS *m/z* 866, 868, 870, 872, 874 (MH⁺); CD (H₂O) λ ext 215 ($\Delta\epsilon$ + 17.4), 255 ($\Delta\epsilon$ - 1.91) and 293 ($\Delta\epsilon$ - 2.44); NMR see Table 1.

3,5-Dibromo-L-tyrosine HBr (9):⁹ CD (H₂O) λ ext 210 ($\Delta\epsilon$ + 15.3); ¹H NMR (MeOH) δ 7.45 (2H, s, H-4 and H-6), δ 4.25 (1H, dd, *J* = 5.5, 7; H-8), δ 3.15 (2H, ddd, *J* = 15, 5.5, 7; H-8).

O-Methyl-3,5-dibromo-NMe₃-L-tyrosine (10). A stirred solution of 3,5-dibromotyrosine (0.17 g) in MeOH (10 mL) was treated with HCO₃K (0.50 g) and MeI (0.5 mL), and stirring was continued overnight. The solution was neutralized with concentrated HCl and evaporated to dryness. MeOH was added, and the MeOH-soluble material was passed through an Amberlite IR-120 column (2 mL). The column was eluted with 5% NH₄OH. The alkaline solution was evaporated and the residue (35 mg) was chromatographed on Si gel yielding **10** (15 mg) (eluent CH₂Cl₂/MeOH/H₂O 80:20:1); [α]_D +30° (MeOH, *c* 0.8) (lit.¹⁰ [α]_D -8.3°), probably the



9 R₁ = H, R₂ = NH₂
10 R₁ = Me, R₂ = NMe₃

R-isomer]; FABMS m/z 394, 396, 398 (MH^+); CD (H_2O) λ ext 215 ($\Delta\epsilon + 7.10$); 1H NMR, $^{11}\delta$ 3.75 (1H, dd, $J = 12, 3.5$; H-8, splitting pattern similar to that of H-8 of **2**, see Table 1).

Acknowledgment. We thank Prof. P. R. Bergquist, Department of Zoology, University of Auckland, Auckland, New-Zealand, and Prof. C. Levi, Muséum d'Histoire Naturelle, Paris, France, for their kind identification of the sponge specimens.

Note Added in Proof. During the preparation of this paper, the isolation from the sponge *Psammaphysilla purea* and the structure elucidation of compound **1**, though of reverse absolute configuration, named purealidin J, have been published: Kobayashi, J.; Honma, K.; Sasaki, T.; Tsuda, M. *Chem. Pharm. Bull.* **1995**, *50*, 403–407.

References and Notes

- (1) Present address: Laboratoire de Chimie des Substances Naturelles, Faculté des Sciences, Université Cadi Ayyad, BPS 15, Marrakech, Morocco.
- (2) Xynas, R.; Capon, R. *J. J. Aust. J. Chem.* **1989**, *42*, 1427–1433.
- (3) Nakamura, H.; Wu, H.; Kobayashi, J.; Nakamura, Y.; Ohizumi, Y.; Hirata, Y. *Tetrahedron Lett.* **1985**, *37*, 4517–4520.
- (4) Ishibashi, M.; Tsuda, M.; Ohizumi, Y.; Sasaki, T.; Kobayashi, Y. *Experientia* **1991**, *47*, 299–300.
- (5) Kobayashi, J.; Tsuda, M.; Agemi, K.; Shigemori, H.; Ishibashi, M.; Sasaki, T.; Mikami, Y. *Tetrahedron* **1991**, *47*, 6617–6622.
- (6) Cimino, C.; De Rosa, S.; De Stefano, S.; Self, R.; Sodano, G. *Tetrahedron Lett.* **1983**, *29*, 3029–3032.
- (7) Kernan, M. R.; Cambie, R. C.; Bergquist, P. R. *J. Nat. Prod.* **1990**, *53*, 720–723.
- (8) Sakota, N.; Okita, K.; Matsui, Y. *Bull. Soc. Japan* **1970**, *43*, 1138–1141.
- (9) Nikolics, K.; Teplan, I.; Medzihradsky, K. *J. Labelled Compd. Radiopharm.* **1976**, *12*, 163–171.
- (10) Chen, F. C. M.; Benoiton, N. L. *Can. J. Chem.* **1976**, *54*, 3310–3311.
- (11) Ciminello, P.; Fattorusso, E.; Magno, S.; Pansini, M. *J. Nat. Prod.* **1995**, *58*, 689–696.
- (12) Albrizio, S.; Ciminello, P.; Fattorusso, E.; Magno, S.; Pansini, M. *Tetrahedron* **1994**, *50*, 783–788.

NP9600457