

Isolation, Identification, and Antioxidant Activity of Three Stilbene Glucosides Newly Extracted from *Vitis vinifera* Cell Cultures

Pierre Waffo Teguo,[†] Bernard Fauconneau,[‡] Gérard Deffieux,[†] François Huguet,[‡] Joseph Vercauteren,[†] and Jean-Michel Mérillon^{*,†}

Groupe d'Etude des Substances Naturelles à Intérêt Thérapeutique, EA 491, Faculté des Sciences Pharmaceutiques, Université de Bordeaux 2, 3 place de la Victoire, 33000 Bordeaux, France, and Centre d'Etudes et de Recherche sur les Xénobiotiques, EA1223, Faculté de Médecine et de Pharmacie, Université de Poitiers, 34 rue du Jardin des Plantes, 86005 Poitiers, France

Received October 29, 1997

Suspension cultures of *Vitis vinifera* L. (Vitaceae) produce many hydroxylated stilbene glucosides found in red wine. From these cells, we isolated and characterized glycosylated stilbenes, (*Z*)-piceatannol (3,5,3',4'-tetrahydroxystilbene)-3-*O*- β -D-glucopyranoside (**6**) and (*E*)- and (*Z*)-resveratrol (3,5,4'-trihydroxystilbene)-4'-*O*- β -D-glucopyranoside (**2** and **7**, respectively), which have not previously reported to be constituents of *Vitis vinifera* or wine. The ability of these compounds to act as radical scavengers was investigated using 1,1 diphenyl-2-picrylhydrazyl, a stable free radical. Antioxidant activities were assessed by their capacity to prevent Cu²⁺-induced lipid peroxidation in human low-density lipoprotein.

Numerous epidemiological studies in France have shown a negative correlation between moderate red wine consumption and the incidence of cardiovascular diseases.^{1–3} This is the so-called French paradox. Wine contains natural plant phenolic compounds that may protect circulating lipoproteins from oxidative damage.⁴ Stilbene has attracted a great deal of interest because relatively high quantities are found in grapes and wine, which are considered the most important dietary sources of these substances.^{2,5} On the other hand, stilbene derivatives seem to have a variety of biological activities.^{6,7} We reported previously that (*E*)- and (*Z*)-piceid and (*E*)-astringin, stilbene glucosides isolated from cell cultures of *Vitis vinifera* L. (Vitaceae),^{8,9} inhibit the lipid peroxidation induced by Cu²⁺.¹⁰


In this study we isolated and characterized stilbene glucosides, (*E*)- and (*Z*)-resveratrolside and (*Z*)-astringin, not previously reported to be constituents of *Vitis vinifera* or of wine. We assessed the relative activities of a range of stilbene glucosides as scavengers of radicals and their properties to prevent Cu²⁺-induced lipid peroxidation in low-density lipoprotein (LDL). We further analyzed the structure–antioxidant activity relationship of these compounds *in vitro*.

Stilbene compounds **1**, **2**, **3**, and **6–8** were purified from the EtOAc extract of cell suspension of *Vitis vinifera* by a combination of chromatographic techniques. Compound **1** was identified as (*E*)-astringin.⁸ Compounds **3** and **8** were identified as (*E*)- and (*Z*)-piceid.^{8,9}

The structures of **2**, **6**, and **7**, three stilbene glucosides newly isolated from cell suspension of *Vitis vinifera*, were deduced by spectrometric methods. Assignments of proton and carbon resonances were deduced from

analysis of ¹H–¹H COSY,¹¹ heteronuclear HMQC,¹² and HMBC¹³ 2D chemical shift correlations. Compounds **2** and **7** were characterized as (*E*)- and (*Z*)-3,5,4'-trihydroxystilbene-4'-*O*- β -D-glucopyranoside, respectively, by comparison with literature data.¹⁴ These compounds have been found in the roots of *Polygonum cuspidatum*¹⁴ and are also called (*E*)- and (*Z*)-resveratrolside. For **6**, our results are similar to those previously reported by Strack et al.¹⁵ and indicate that this compound is (*Z*)-piceatannol-3-*O*- β -D-glucopyranoside or (*Z*)-astringin.

The antioxidant activities of compounds **1–9** were studied (Table 1). Coexistence of an antioxidant A and a free radical R° (such as reactive oxygen species generated by an oxidative stress, or 1,1 diphenyl-2-picrylhydrazyl (DPPH) leads to the disappearance of this free radical and to the appearance of the free radical A° according to the reaction: A + R° → A° + R. On the stilbenes studied, the conjugation between rings A and B via a planar C2 unsaturated structure allows an electron delocalization across the molecules for stabilization of the radical, which explains the relative antioxidant properties of all these compounds.



	R ₁	R ₂	R ₃	R ₄
1	GlcO	OH	OH	OH
2	OH	OH	H	GlcO
3	GlcO	OH	H	OH
4	OH	OH	OH	OH
5	OH	OH	H	OH

	R ₁	R ₂	R ₃	R ₄
6	GlcO	OH	OH	OH
7	OH	OH	H	GlcO
8	GlcO	OH	H	OH
9	OH	OH	H	OH

On Cu²⁺-induced lipid peroxidation on the LDL, the results (Table 1) show no important difference between

* To whom correspondence should be addressed. Tel.: (33) 5 57 57 18 22. Fax: (33) 5 56 91 79 88. E-mail: jean-michel.merillon@phyto.u-bordeaux2.fr.

[†] Groupe d'Etude des Substances Naturelles à Intérêt Thérapeutique.

[‡] Centre d'Etudes et de Recherche sur les Xénobiotiques.

Table 1. IC₅₀ Values^a for the Antioxidant Activities of Stilbenes Extracted from *Vitis Vinifera* Cells and of Trolox

compound	IC ₅₀ values (μM)	
	DPPH	LDL
1	30.2 ± 1.8	3.0 ± 0.2
2	1000 ± 95	100 ± 11.7
3	198 ± 16.8	19.1 ± 3.0
4	28 ± 1.2	1.8 ± 0.2
5	72 ± 4.5	2.4 ± 0.2
6	29 ± 1.9	2.5 ± 0.3
7	>1000	150 ± 16.1
8	142 ± 7.9	16.8 ± 1.6
9	95 ± 3.8	18 ± 2.0
trolox	10.1 ± 0.5	4.7 ± 0.4

^a Each value is the mean of at least three independent experiments ± SD. Statistical analysis was performed by using Student's *t*-test.

(*E*) and (*Z*) structures of each molecule, except for (*E*- and (*Z*-resveratrol, with a better activity for (*E*-resveratrol.

The glycosylation of (*E*-stilbenes reduces their activity when compared to the corresponding aglycons (respectively, seven times for (*E*-piceid vs. (*E*-resveratrol, $p < 0.001$; 35 times for (*E*-resveratrol vs. (*E*-resveratrol, $p < 0.001$; 1.6 times for astringin vs. piceatannol, $p < 0.05$). This difference is less important in the (*Z*) structure. Blocking the 4'-hydroxyl group in the B ring by a glycosyl moiety decreases dramatically the antioxidant activity [(*E*-resveratrol vs. (*E*-resveratrol)] compared to glycosylation in the 3-position on A ring [(*E*-piceid vs. (*E*-resveratrol)].

Considering the antioxidant activities of these molecules, it is worth noting the importance of the two hydroxyl groups in the *ortho*-diphenolic arrangement in the B ring. Actually, astringin, which possesses this catechol structure and consequently a supplementary OH on the B ring as compared to piceid, has an activity six times higher for the (*E*) and (*Z*) structure ($p < 0.001$). Astringin, despite the presence of glycoside in the 3-position on ring A, has an activity close to that observed with (*E*-resveratrol.

Among these molecules, the most potent antioxidant is piceatannol, which possesses four hydroxyl groups, including the catechol structure in the B ring. Furthermore, piceatannol is two times more efficient than trolox, the water-soluble vitamin E analogue ($p < 0.01$).

On DPPH, Table 1 shows no important difference between (*E*) and (*Z*) structures of each molecule. The glycosylation of (*E*- and (*Z*-stilbenes reduces their activity when compared to the corresponding aglycons [(*E*- and (*Z*-piceid vs. (*E*- and (*Z*-resveratrol; (*E*- and (*Z*-resveratrol vs. (*E*- and (*Z*-resveratrol)], but the difference of activity between (*E*-astringin and piceatannol is less than that observed on the LDL test.

The glycosylation of resveratrol in the 3-position in the A ring leads to piceid, which has an activity about two times lower for (*E*) and (*Z*) structures ($p < 0.01$). When glycosylation is performed in the 4'-position in B ring, the antioxidant activities of the molecules obtained, that is, (*E*- and (*Z*-resveratrol, decrease dramatically, as compared to those of (*E*- and (*Z*-resveratrol.

These results show that the catechol structure is essential for the antioxidant activities of stilbenes, as reported by Rice-Evans et al.¹⁶ for flavonoids. But (*E*-

and (*Z*-resveratrols may be hydrolyzed by glycosidases in the human gastrointestinal tract.² Further work is being undertaken to characterize these substances (**1**, **2**, **4**, **6**, **7**) in wine.

Experimental Section

General Experimental Procedures. UV spectra were measured in MeOH using a Hitachi U-2000 spectrophotometer. IR spectra were obtained on a KBr disk using a Shimadzu IR-470 spectrophotometer. NMR spectra were performed with a Bruker AMX-500 spectrometer. FABMS were recorded using glycerol as matrix, in positive-ion mode. (*E*-Piceatannol (**4**, 6 mg) and (*Z*-resveratrol (**9**, 5 mg) were obtained by enzymatic hydrolysis of (*Z*-piceid (10 mg) and (*E*-astringin (10 mg), respectively.

Cell Culture. Cell suspension cultures of *Vitis vinifera* L. cv Gamay Fréaux var. Teinturier were maintained as previously described.¹⁷ The maintenance medium (MM) contained B5 macroelements,¹⁸ microelements,¹⁹ and vitamins,²⁰ and was supplemented with 58 mM sucrose, 250 mg/L casein hydrolysate, 0.54 μM 1-naphthaleneacetic acid, and 0.93 μM kinetin. Experiments were carried out by inoculating a 7-day-old cell suspension into an induction medium (IM₁) at a 1:8 (v/v) ratio, for one transfer.¹⁷ IM₁ was the same as MM, but contained 2 mM (NH₄)₂SO₄, 2.2 mM NaH₂PO₄, 2 mM MgSO₄, and 175 mM sucrose. Harvesting was made on day 12 corresponding to the optimal period for the production of stilbenes by grape cells in this medium.⁸ Cells were collected through filtration under partial vacuum (nylon cloth, 30 μm), rapidly washed with cold distilled H₂O, and then extracted.

Test on Low-Density Lipoproteins (LDL). This assay was previously described.¹⁰ Briefly, human LDL were oxidized by cupric ions. Lipid peroxidation was assessed by TBARS (thiobarbituric acid reactive substances) measurement, using thiobarbituric acid (TBA) colorimetric method.²¹ The absorbance is read at 532 nm. The stilbenes added in this mixture inhibited this peroxidation, and the efficient concentration (IC₅₀) was the concentration that inhibited 50% of coloration.

Test on DPPH. DPPH is a dyed free radical. The trapping effect of the molecules tested was assessed by measuring the absorbance change at 515 nm of a DPPH solution²² (100 μM) in the presence of different concentrations of the stilbenes. Measurements were performed at least in triplicate. The efficient concentration (IC₅₀) is the concentration that inhibited 50% of coloration.

Extraction, Isolation, and Identification of the Stilbene Compounds. Frozen cells (900 g) were homogenized with Me₂CO–H₂O as previously described.^{8,9} The extract was concentrated *in vacuo*, and the aqueous mixture was extracted with EtOAc. The EtOAc extract was chromatographed over a cation-exchange resin column (1.5 × 60 cm) and eluted by H₂O–MeOH gradient. The stilbenes were eluted by 50% MeOH. For the further fractionation, the crude stilbenes were divided into fractions on a Sephadex LH-20 column (1.5 × 60 cm). Two main fractions were obtained. The mixture of (*Z*-stilbenes was eluted by 20% MeOH and the mixture of (*E*-stilbenes by 30% MeOH. Extracts were constantly protected from light to avoid (*E*–*Z*) isomerization.

Compounds **1** (*E*)-astringin (12 mg), **2** (*E*)-resveratrolside (8 mg), and **3** (*E*)-piceid (67 mg), **6** (*Z*)-astringin (7 mg), **7** (*Z*)-resveratrolside (7 mg), and **8** (*Z*)-piceid (52 mg), were obtained as pure compounds by semi-prep. HPLC on an Ultrasep RP18 (6 μ m) reversed-phase C18 column (8 mm i.d. \times 250 mm) with column guard eluted by gradient system solvent: A, H₂O adjusted to pH 2.4 with TFA; B, 20% A with 80% MeCN. The elution program at 3 mL min⁻¹ was as follows: 18% B (0–10 min); 18–23% B (10–17 min); 23–24.5% B (17–21 min); 24.5–31.5% B (21–27 min); 31.5–50% B (27–30 min); 50–60% B (30–35 min); 60–100% B (35–40 min). The chromatogram was monitored at dual mode 286–306 nm using an UV detector.

Compound **2**: UV (MeOH) λ_{\max} (log ϵ) 241 (4.52), 261 (4.52), 304 (4.89) nm; IR (KBr) ν_{\max} 3400, 1600 cm⁻¹.

Compound **6**: UV (MeOH) λ_{\max} (log ϵ) 302 (4.08) nm; IR (KBr) ν_{\max} 3400, 1600 cm⁻¹.

Compound **7**: UV (MeOH) λ_{\max} (log ϵ) 283 (4.02) nm; IR (KBr) ν_{\max} 3450, 1600 cm⁻¹.

Acknowledgment. This work was supported by grants from the Ministère de la Recherche and the O.N.I. VINS. We thank N. Pinaud, R. Pontcharraud, and A. Soriano for technical assistance, and P. Villalba for the typing.

References and Notes

- (1) Renaud, S.; De Lorgeril, M. *Lancet* **1992**, *339*, 1523–1526.
- (2) Goldberg, D. M. *Clinic. Chem.* **1995**, *41*, 14–16.

- (3) Waterhouse, A. L. *Chem. Ind.* **1995**, 338–341.
- (4) Fuhrman, B.; Lavy, A.; Aviram, M. *Am. J. Clin. Nutr.* **1996**, *42*, 1591–1593.
- (5) Mattivi, F.; Reniero, F.; Korhammer, S. *J. Agric. Food Chem.* **1995**, *43*, 1820–1823.
- (6) Frankel, E. N.; Waterhouse, A. L.; Kinsella, J. E. *Lancet* **1993**, *341*, 1103–1104.
- (7) Jang, M.; Cai, L.; Udeani, G. O.; Slowing, K. V.; Thomas, C. F.; Beecher, C. W. W.; Fong, H. H. S.; Farnsworth, N. R.; Kinghorn, A. D.; Mehta, R. G.; Moon, R. C.; Pezzuto, J. M. *Science* **1997**, *275*, 218–220.
- (8) Waffo Tegu, P.; Decendit, A.; Krisa, S.; Vercauteren, J.; Deffieux, G.; Mérillon, J.-M. *J. Nat. Prod.* **1996**, *59*, 1189–1191.
- (9) Waffo Tegu, P.; Decendit, A.; Vercauteren, J.; Deffieux, G.; Mérillon, J.-M. *Phytochemistry* **1996**, *42*, 1591–1593.
- (10) Mérillon, J.-M.; Fauconneau, B.; Waffo Tegu, P.; Barrier, L.; Vercauteren, J.; Hugué, F. *Clinic. Chem.* **1997**, *43*, 1092–1093.
- (11) Aue, W. P.; Bartholdi, E.; Ernst, R. R. *J. Chem. Phys.* **1976**, *64*, 2229–2246.
- (12) Bax, A.; Subramanian, S. *J. Magn. Res.* **1986**, *67*, 565–569.
- (13) Bax, A.; Summer, M. F. *J. Am. Chem. Soc.* **1986**, *108*, 2093–2094.
- (14) Jayatilake, G. S.; Jayasuriya, H.; Lee, E. S.; Koonchanok, N. M.; Geahlen, R. L.; Ashendel, C. L.; McLaughlin, J. L.; Chang C. J. *J. Nat. Prod.* **1993**, *56*, 1805–1810.
- (15) Strack, D.; Heilemann, J.; Wray, V.; Dirks, H. *Phytochemistry* **1989**, *28*, 2071–2078.
- (16) Rice-Evans, C. A.; Miller, N. J.; Bolwell, P. G.; Bramley, P. M.; Pridham, J. B. *Free Rad. Res.* **1995**, *22*, 375–383.
- (17) Decendit, A.; Mérillon, J.-M. *Plant Cell Rep.* **1996**, *15*, 762–765.
- (18) Gamborg, O. L.; Miller, R. A.; Ojima, K. *Exp. Cell. Res.* **1968**, *50*, 151–156.
- (19) Murashige, T.; Skoog, F. *Physiol. Plant.* **1962**, *15*, 473–497.
- (20) Morel, G. *Physiol. Vég.* **1970**, *8*, 189–191.
- (21) Buege, J. A.; Aust, S. D. *Methods Enzymol.* **1978**, *52*, 302–310.
- (22) Tamura, A.; Sato, T.; Fuhu, T. *Chem. Pharm. Bull.* **1990**, *38*, 255–257.

NP9704819