

Oligosaccharide Esters from the Roots of *Polygala arillata*

Wataru Kobayashi,[†] Toshio Miyase,^{*,†} Sayako Suzuki,[†] Hiroshi Noguchi,[†] and Xin-Min Chen[‡]

School of Pharmaceutical Sciences, University of Shizuoka, 52-1, Yada, Shizuoka 422-8526, Japan, and Division of Phytochemistry, Chengdu Institute of Biology, Chinese Academy of Sciences, 4-9 Ren Ming Nan Lu, Chengdu, Sichuan, People's Republic of China

Received February 4, 2000



Two new sucrose esters, arillatoses A (**1**) and B (**2**), and four new trisaccharide esters, arillatoses C–F (**3**–**6**), were isolated from the roots of *Polygala arillata*, together with four known sucrose esters, glomeratose E (**7**) and sibiricoses A₁ (**8**), A₅ (**9**), and A₆ (**10**). The structures of the new compounds were elucidated on the basis of chemical and spectroscopic evidence.

In the course of conducting a research program on the oligosaccharide esters from *Polygala* species,² we have investigated *P. arillata* Buch.–Ham. (Polygalaceae). This species is widely distributed in the People's Republic of China, and its roots are used as a traditional medicine in a manner similar to "Yuan zhi" (the roots of *P. tenuifolia* Willd.) to tranquilize, as a tonic, and to prevent loss of memory.³ No previous investigation has been reported on the oligosaccharide esters of *P. arillata*. We now report the isolation and structure elucidation of two sucrose esters, arillatoses A (**1**) and B (**2**), and four new trisaccharide esters, arillatoses C–F (**3**–**6**). Four known compounds isolated from this plant were identified by comparison of the spectral data with reported data as glomeratose E (**7**)¹ sibiricoses A₁ (**8**), A₅ (**9**), and A₆ (**10**).²


with mixtures of water and methanol. The 50% MeOH eluate was chromatographed further to afford six sucrose esters (**1**, **2**, and **7**–**10**) and four trisaccharide esters (**3**–**6**).

The FABMS of arillatose A (**1**) showed quasimolecular ion peaks at m/z 753 $[M + H]^+$ and 775 $[M + Na]^+$, consistent with a molecular formula of C₃₄H₄₀O₁₉. On alkaline hydrolysis, **1** afforded sucrose, while on acid hydrolysis, it gave D-glucose and D-fructose.⁴ The ¹H NMR spectrum of **1** exhibited two methine protons [δ 4.81 (1H, br s) and δ 3.98 (1H, d, $J = 2$ Hz)], a vinyl proton at a highly deshielded position [δ 7.77 (1H, s)], and two aromatic protons [δ 6.92 (1H, s) and δ 6.34 (2H, s)], in addition to the signals due to sucrose. The aromatic proton (δ 6.34) indicated the presence of a 1,3,4,5-tetrasubstituted benzene unit in **1**. All proton and carbon signals were assigned by COSY, HOHAHA, HMBC, and HMQC NMR experiments. On irradiation of the aromatic proton at δ 6.34 due to H-2', ROEs were observed at a methoxyl at δ 3.70 (6H, s), a methine proton at δ 4.81 due to H-1, a methine proton at δ 3.98 due to H-2, and an olefinic proton at δ 7.77 due to H-4 (Figure 1). From this ROE correlation between H-2' and H-4, the bond C-1–C-1' was concluded to be quasi-axial. The coupling constant between H-1 and H-2 (4 Hz) indicated that the bond C-2–H-2 was quasi-equatorial, and if the bond C-2–H-2 had been quasi-axial, the H-4 signal would have been a doublet induced by the allylic coupling with H-2.⁵ The CD spectrum of **1**, which is identical with (1*S*,2*R*)-1,2-dihydro-6,7-dihydroxy-1-(3',4'-dihydroxyphenyl)-naphthalene 2,3-dicarboxylic acid dimethyl ester,⁶ showed a positive first Cotton effect at 346 nm. Therefore, **1** has a 1*S*,2*R* configuration. From these data, the structure of arillatose A was deduced as cyclic 3'→3:6→2-[(1*S*,2*R*)-1-(4-hydroxy-3,5-dimethoxyphenyl)-1,2-dihydro-7-hydroxy-6,8-dimethoxy-2,3-naphthalenedicarboxyl]- β -D-fructofuranosyl α -D-glucopyranoside.

Arillatose B (**2**) was isolated as an amorphous powder. The positive mode FABMS revealed a quasimolecular ion peak at m/z 541 $[M + Na]^+$, consistent with a molecular formula of C₂₂H₃₀O₁₄. On alkaline hydrolysis, **2** afforded sucrose and ferulic acid, while on acid hydrolysis, it gave D-glucose and D-fructose. In the ¹H NMR spectrum of **2**, one feruloyl signal was observed, in addition to the signals due to sucrose. All proton and carbon signals in the NMR spectra (Tables 2 and 3) of **2** were assigned from its ¹H–¹H COSY, HOHAHA, HMBC, and HMQC spectra. The position of the feruloyl group in the sucrose moiety of **2** was deduced from the HMBC experiment. In this spectrum, a long-range correlation (³J_{COCH}) was observed between the feruloyl carbonyl carbon signal at δ 169.2 and the proton



	R ₁	R ₂	R ₃
2	H	A	H
3	Glc	A	H
4	Glc	B	H
5	Glc	H	A
6	Glc	H	B



Results and Discussion

The air-dried roots of *P. arillata* were extracted with MeOH under reflux. The MeOH extract was suspended in H₂O and extracted with ether. The H₂O layer was adsorbed on a porous polymer gel (Diaion HP-20) column and eluted

* To whom correspondence should be addressed: Tel.: 054-264-5661. Fax: 054-264-5661. E-mail: miyase@ys7.u-shizuoka-ken.ac.jp.

[†] University of Shizuoka.

[‡] Chinese Academy of Sciences.




Figure 1. Correlations observed in the ROE difference NMR spectrum of arillatose A (1).

Table 1. ^1H NMR and ^{13}C NMR Data of 1 in CD_3OD at 35°C

	^1H NMR	^{13}C NMR	HMBC (C-H)
sugar moiety			
Glc-1	5.53 d (4)	92.8	Fru-2
2	3.37 dd (4, 10)	73.7	
3	3.46 dd (10, 9)	75.6	Glc-2
4	3.10 dd (10, 9)	72.1	Glc-3, -5
5	3.70 m	73.2	Glc-6
6	4.06 dd (10, 2)	65.7	α
	4.57 dd (10, 9)		Glc-5, α
Fru-1	3.66 d (12)	66.7	
	3.77 d (12)		
2		105.3	
3	5.28 d (9)	80.3	Fru-4, α'
4	4.33 dd (10, 9)	72.2	Fru-3, -5, -6
5	3.66 ^a	81.9	Fru-2, -3
6	3.76 ^a	61.0	
	3.86 ^a		
lignan moiety			
1	4.81 br s	42.0	2, 8, 8a, 9, 1', 2', 3', 6'
2	3.98 d (2)	49.6	1, 4, 9, 10, 1'
3		123.1	
4	7.77 s	141.5	2, 3, 4a, 5, 10
4a		124.3	
5	6.92 s	109.7	4, 4a, 6, 7
6		149.5	
7		143.8	
8		146.9	
8a		124.8	
9		174.7	
10		168.6	
1'		135.4	
2'	6.34 s	106.0	1, 1', 3', 6'
3'		149.1	
4'		135.6	
5'		149.1	
6'	6.34 s	106.0	1, 1', 2', 5'
MeO-6	3.91 s	56.9	6
MeO-8	3.53 s	60.8	8
MeO-3'	3.70 s	56.8	3'
MeO-5'	3.70 s	56.8	5'

^a Overlapped.

signal at δ 4.27 due to H-6 of glucose. From these data, the structure of **2** was elucidated as β -D-fructofuranosyl 6-O-feruloyl- α -D-glucopyranoside.

Arillatose C (**3**) was obtained as an amorphous powder. The FABMS of **3** showed a quasimolecular ion peak at m/z 703 $[\text{M} + \text{Na}]^+$. Compound **3** gave D-glucose and D-fructose in the ratio 2:1 on acid hydrolysis, while on alkaline hydrolysis it afforded ferulic acid. In the ^1H NMR spectrum of **3**, one feruloyl signal was observed. Full assignments of the proton and carbon signals were secured by a HOHAHA difference spectrum, on irradiating at the glucosyl anomeric proton signal and H-3 of the fructosyl moiety, and from ^1H - ^1H COSY, HMQC, and HMBC experiments. In the HMBC spectrum, H-6 of Glc 1 was correlated to an ester

carbonyl carbon at δ 169.1. In a ROE difference spectrum, ROE was observed at δ 3.92 (1H, dd, $J = 9, 9$ Hz) due to H-3 of Glc 1 on irradiation at the anomeric proton signal of Glc 2 at δ 4.53 (1H, d, $J = 8$ Hz). Accordingly, **3** was deduced as a O - β -D-glucopyranosyl-(1 \rightarrow 3)-6- O -feruloyl- α -D-glucopyranosyl β -D-fructofuranoside.

The FABMS of arillatose D (**4**) showed a quasimolecular ion peak at m/z 733 $[\text{M} + \text{Na}]^+$. The ^1H NMR spectrum was similar to that of arillatose C (**3**), but it showed the presence of a sinapoyl residue. Compound **4** gave D-glucose and D-fructose in the ratio 2:1 on acid hydrolysis, while alkaline hydrolysis gave sinapic acid. The HMBC experiment showed a correlation between H-6 of Glc 1 and an ester carbonyl carbon at δ 169.0. Thus, **4** was determined as a O - β -D-glucopyranosyl-(1 \rightarrow 3)-6- O -sinapoyl- α -D-glucopyranosyl β -D-fructofuranoside.

The ^1H NMR spectra of arillatoses E (**5**) and F (**6**) displayed patterns similar to those of arillatoses C (**3**) and D (**4**), respectively, except for downfield-shifted oxymethine protons due to H-3 of a fructosyl moiety at δ 5.44 (1H, d, $J = 7.5$ Hz). On alkaline hydrolysis, **5** afforded ferulic acid and **6** afforded sinapic acid. Compounds **5** and **6** gave D-glucose and D-fructose in the ratio 2:1 on acid hydrolysis. In the HMBC spectra of **5** and **6**, H-3 of a fructosyl moiety was correlated to an ester carbonyl carbon. Therefore, **5** was deduced as O - β -D-glucopyranosyl-(1 \rightarrow 3)- α -D-glucopyranosyl 3'- O -feruloyl- β -D-fructofuranoside and **6** as O - β -D-glucopyranosyl-(1 \rightarrow 3)- α -D-glucopyranosyl 3'- O -sinapoyl- β -D-fructofuranoside.

Experimental Section

General Experimental Procedures. Optical rotations were measured on a JASCO DIP-1000 digital polarimeter. UV spectra were recorded on Hitachi U-3410 spectrometer and CD spectra on a JASCO J-20A spectropolarimeter. ^1H (400 MHz) and ^{13}C (100 MHz) NMR spectra were recorded on a JEOL α -400 FT-NMR spectrometer with TMS as an internal standard. Inverse-detected heteronuclear correlations were measured using HMQC (optimized for $^1J_{\text{C-H}} = 145$ Hz) and HMBC (optimized for $^nJ_{\text{C-H}} = 8$ Hz) pulse sequences with a pulse-field gradient. Positive-mode FABMS were recorded on a JEOL JMS-SX102 spectrometer, using a *m*-nitrobenzyl alcohol matrix. GC was carried out with Hitachi G-3000 gas chromatograph. HPLC was performed using a JASCO System 800.

Plant Material. *P. arillata* Buch.-Ham. was collected in June 1996, in Sichuan, People's Republic of China. The plant was identified by Prof. Zhaoguang Liu, Chengdu Institute of Biology, Academia Sinica, People's Republic of China, and a voucher specimen (no. 960715) has been deposited in the Herbarium, School of Pharmaceutical Sciences, University of Shizuoka.

Extraction and Isolation. The dried and powdered roots of *P. arillata* (1.98 kg) were extracted twice with MeOH under reflux. After evaporation of the solvent under reduced pressure, the MeOH extract was suspended in H_2O and extracted with diethyl ether. The H_2O layer was subjected to passage over a porous polymer gel Mitsubishi Diaion HP-20 column (15 \times 31.5 cm). The adsorbed material was eluted with 50% aqueous MeOH, 70% aqueous MeOH, and MeOH, successively, after washing with H_2O . The 50% aqueous MeOH eluate (11.7 g) was chromatographed on a Si gel (330 g) column using CHCl_3 -MeOH- H_2O (80:18:2) as an eluent to afford fractions A-R. Fractions H + I (686 mg) were subjected to preparative HPLC [ODS 5 \times 100 cm; CH_3CN - H_2O (9:91) \rightarrow (17:83) linear gradient] to afford **2** (6 mg). Fractions J + K (565 mg) were subjected to preparative HPLC [ODS 5 \times 100 cm; CH_3CN - H_2O (9:91) \rightarrow (17:83) linear gradient] to afford **10** (86 mg). Fraction L (764 mg) was subjected to preparative HPLC [ODS 5 \times 100 cm; CH_3CN - H_2O (8:92) \rightarrow (16:84) linear gradient] to afford **1** (15 mg), **7** (31 mg), and **8** (16 mg). Fraction M (807 mg) was subjected to preparative HPLC [ODS 5 \times 100 cm;

Table 2. ¹H NMR Data of **2–6** in CD₃OD at 35 °C

	2	3	4	5	6
Sugar moiety					
Glc1–1	5.42 d (4)	5.47 d (3.5)	5.47 d (3.5)	5.48 d (4)	5.48 d (3.5)
2	3.46 dd (10, 4)	3.68 dd (9, 3.5)	3.68 dd (9, 3.5)	3.62 dd (10, 4)	3.62 dd (10, 3.5)
3	3.75 dd (10, 9)	3.92 dd (9, 9)	3.92 dd (9, 9)	3.65 dd (10, 10)	3.63 dd (10, 10)
4	3.33 dd (9, 9)	3.44 dd (9, 10)	3.44 dd (9, 10)	3.49 dd (10, 10)	3.49 dd (10, 10)
5	4.11 m	4.17 m	4.18 m	3.94 m	3.95 m
6	4.27 dd (12.5, 6)	4.28 dd (12, 6)	4.28 dd (12, 6)	3.78 dd (12, 6)	3.78 dd (12, 6)
	4.51 dd (12.5, 2)	4.53 dd (12, 1.5)	4.53 dd (12, 1.5)	3.87 dd (12, 1.5)	3.87 dd (12, 3)
Glc2–1		4.53 d (8)	4.53 d (8)	4.28 d (7.5)	4.25 d (7)
2		3.30 ^a	3.30 ^a	3.23 ^a	3.20 ^a
3		3.40 dd (9, 9)	3.40 dd (9, 9)	3.26 ^a	3.26 ^a
4		3.30 ^a	3.30 ^a	3.27 ^a	3.22 ^a
5		3.34 m	3.34 m	3.00 m	2.92 m
6		3.65 dd (12, 6)	3.64 dd (12, 6)	3.54 dd (12, 6)	3.52 dd (12, 6)
		3.89 ^a	3.89 ^a	3.64 ^a	3.60 ^a
Fru-1	3.60 d (12.5)	3.62 d (12)	3.61 d (12)	3.61 d (12)	3.61 d (12)
	3.60 d (12.5)	3.64 d (12)	3.63 d (12)	3.68 d (12)	3.69 d (12)
3	4.09 d (8)	4.08 d (8)	4.08 d (8)	5.44 d (7.5)	5.44 d (7.5)
4	4.06 dd (8, 8)	4.08 ^a	4.08 ^a	4.38 dd (9, 7.5)	4.39 dd (9, 7.5)
5	3.80 ^a	3.80 ^a	3.80 ^a	3.96 ^a	3.97 ^a
6	3.78 ^a	3.80 ^a	3.80 ^a	3.81 ^a	3.81 ^a
	3.82 ^a	3.80 ^a	3.80 ^a	3.81 ^a	3.81 ^a
Acid					
β	6.42 d (16)	6.43 d (16)	6.45 d (16)	6.44 d (16)	6.47 d (16)
γ	7.63 d (16)	7.63 d (16)	7.62 d (16)	7.72 d (16)	7.71 d (16)
2	7.22 d (2)	7.23 d (2)	6.94 s	7.24 d (2)	6.96 s
5	6.81 d (7.5)	6.81 d (9)		6.84 d (9)	
6	7.09 dd (7.5, 2)	7.09 dd (9, 2)	6.94 s	7.14 dd (9, 2)	6.96 s
OMe	3.89 s	3.89 s	3.88 s	3.91 s	3.90 s

^a Overlapped.**Table 3.** ¹³C NMR Data of **2–6** in CD₃OD at 35 °C

	2	3	4	5	6
sugar moiety					
Glc1–1	93.3	93.0	92.9	93.1	93.1
2	73.2	72.3	72.3	72.1	72.1
3	74.7	85.4	85.4	86.9	86.8
4	71.9	70.3	70.4	69.8	69.9
5	72.1	71.9	71.9	74.4	74.4
6	65.1	65.0	65.2	62.4	62.4
Glc2–1		105.3	105.2	105.3	105.4
2		75.5	75.5	75.5	75.5
3		77.9	77.9	77.6	77.6
4		71.5	71.6	71.1	71.0
5		78.2	78.2	77.9	77.8
6		62.6	62.6	62.3	62.1
Fru-1	64.3	64.4	64.5	65.5	65.5
2	105.2	105.2	105.3	105.5	105.5
3	79.4	79.5	79.5	80.0	80.0
4	76.2	76.2	76.2	74.2	74.2
5	83.9	84.0	84.0	84.6	84.7
6	64.1	64.2	64.3	62.9	62.9
acid					
α	169.2	169.1	169.0	168.3	168.2
β	115.4	115.4	115.9	115.2	115.6
γ	147.0	147.0	147.2	147.7	147.9
1	127.8	127.8	126.7	127.7	126.6
2	111.8	111.8	107.1	112.1	107.4
3	149.4	149.4	149.5	149.5	149.5
4	150.6	150.6	139.7	150.9	139.9
5	116.5	116.5	149.5	116.7	149.5
6	124.2	124.2	107.1	124.5	107.4
OMe	56.6	56.6	57.0	56.6	57.1

CH₃CN–H₂O (8:92) → (16:84) linear gradient] to afford **2** (23 mg), **8** (16 mg), **9** (61 mg), and **10** (18 mg). Fraction N (748 mg) was subjected to a preparative HPLC [ODS 5 × 100 cm; CH₃CN–H₂O (8:92) → (16:84) linear gradient] to afford **3** (9 mg), **7** (53 mg), and **9** (15 mg). Fraction O (812 mg) was subjected to a preparative HPLC [ODS 5 × 100 cm; CH₃CN–H₂O (8:92) → (16:84) linear gradient] to afford **3** (19 mg), **4** (13 mg), **5** (33 mg), and **6** (23 mg).

Arillatose A (1): amorphous powder, [α]²⁷_D +25.1° (*c* 0.12, MeOH); UV (MeOH) λ_{max} (log ε) 205 (4.69), 250 (4.34), 340

(4.17) nm; CD (MeOH) λ_{max} (Δε) 346 (+7.4), 253 (–6.4), 229 (–12.9) nm; ¹H and ¹³C NMR, see Table 1; FABMS *m/z* 775 [M + Na]⁺, 753 [M + H]⁺.

Arillatose B (2): amorphous powder, [α]²⁷_D +15.8° (*c* 0.13, MeOH); UV (MeOH) λ_{max} (log ε) 235 (3.73), 295 (3.66), 326 (3.76) nm; ¹H and ¹³C NMR, see Tables 2 and 3; FABMS *m/z* 703 [M + Na]⁺.

Arillatose C (3): amorphous powder, [α]²⁷_D +15.8° (*c* 0.13, MeOH); UV (MeOH) λ_{max} (log ε) 245 (3.81), 295 (3.75), 326 (3.83) nm; ¹H and ¹³C NMR, see Tables 2 and 3; FABMS *m/z* 703 [M + Na]⁺.

Arillatose D (4): amorphous powder, [α]²⁷_D +2.0° (*c* 0.10, MeOH); UV (MeOH) λ_{max} (log ε) 239 (4.21), 330 (4.13) nm; ¹H and ¹³C NMR, see Tables 2 and 3; FABMS *m/z* 733 [M + Na]⁺.

Arillatose E (5): amorphous powder, [α]²⁷_D –20.6° (*c* 0.09, MeOH); UV (MeOH) λ_{max} (log ε) 203 (4.44), 240 (4.05), 294 (3.94), 327 (4.01), 380 (3.35) nm; ¹H and ¹³C NMR, see Tables 2 and 3; FABMS *m/z* 703 [M + Na]⁺.

Arillatose F (6): amorphous powder, [α]²⁷_D –4.5° (*c* 0.10, MeOH); UV (MeOH) λ_{max} (log ε) 202 (4.50), 240 (4.17), 330 (4.12) nm; ¹H and ¹³C NMR, see Tables 2 and 3; FABMS *m/z* 733 [M + Na]⁺.

Alkaline Hydrolysis of 1–6. Each compound (2 mg) was treated with 1 N NaOH aqueous (50 μL) for 4 h at room temperature in N₂ atmosphere, and the reaction mixture was extracted three times with EtOAc after acidification with 1 N HCl. From the H₂O layer, a sugar was detected by HPLC [Asahipak NH2P-50, 4.6 mm × 25 cm, CH₃CN–H₂O (65:35), 1.0 mL/min, UV 195 nm,⁷] as follows: sucrose (*t*_R 5.2 min) from **1** and **2**. From the EtOAc layer, ferulic acid (*t*_R 9.1 min) was detected from **2**, **3**, and **5**; sinapic acid (*t*_R 8.6 min) was detected from **4** and **6** by HPLC [YMC R–ODS-5, 4.6 mm × 25 cm, CH₃CN–H₂O (22.5:77.5) + 0.05% CF₃COOH, 1.0 mL/min, UV 270 nm].

Acid Hydrolysis of 1–6. Each compound (1 mg) was heated on a boiling water bath with 1 N HCl (50 μL) for 15 min. The reaction mixture was passed through an Amberlite IRA-60E column and the eluate was concentrated. The residue was warmed at 60 °C with a solution of D-cysteine methyl ester in pyridine (3 mg/25 μL) for 90 min and to the reaction mixture hexamethyldisilazane (10 μL) and trimethylsilyl chloride (10 μL) were added, and the reaction mixture was stirred at 60

°C for 30 min. The supernatant was subjected to GC. Conditions: column Supelco SPB-1, 0.25 mm × 27 m; temperature 220 °C; carrier gas, N₂. From **1–6**, D-glucose (*t*_R 18.6 min) and D-fructose (*t*_R 14.3 min) were detected.⁸ In **3–6**, the ratio of these sugars was 2:1.

Acknowledgment. We thank the staff of the Analytical Center, University of Shizuoka, for measurement of FABMS.

References and Notes

- (1) Zhang, D.-M.; Miyase, T.; Kuroyanagi, M.; Umehara, K.; Noguchi, H. *Phytochemistry* **1998**, *47*, 45–52, and references therein.
- (2) Miyase, T.; Noguchi, H.; Chen X.-M. *J. Nat. Prod.* **1999**, *62*, 993–996.

- (3) Jian, S. *New Medicinal College Dictionary of Chinese Drugs*; Shanghai Scientific Technologic Publisher: Shanghai, 1997; p 2174.
- (4) Hara, S.; Okabe, H.; Mihashi, K. *Chem. Pharm. Bull.* **1986**, *34*, 1843–1845.
- (5) Maeda, S.; Masuda, H.; Tokoroyama, T. *Chem. Pharm. Bull.* **1995**, *43*, 84–90.
- (6) Nishizawa, M.; Tuda, M.; Hayashi, K. *Phytochemistry* **1990**, *29*, 2645–2649.
- (7) Binder, H. *J. Chromatogr.* **1980**, *189*, 414–420.
- (8) The *t*_R for L-glucose (17.1 min) and the *t*_R for L-fructose (15.0 min) were obtained from that of the enantiomer (D-fructose + L-cysteine methyl ester).

NP0000567