

## Withanolides from *Vassobia lorentzii*

Rosana I. Misico,<sup>†</sup> Roberto R. Gil,<sup>†</sup> Juan C. Oberti,<sup>†</sup> Adriana S. Veleiro,<sup>‡</sup> and Gerardo Burton<sup>\*‡</sup>

Departamento de Química Orgánica and IMBIV, Facultad de Ciencias Químicas, Universidad Nacional de Córdoba, 5000 Córdoba, Argentina, and Departamento de Química Orgánica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Pabellón 2, Ciudad Universitaria, 1428 Buenos Aires, Argentina

Received January 14, 2000

Eight new withanolides were isolated from the aerial parts of *Vassobia lorentzii* and characterized by spectroscopic methods and with the aid of molecular modeling. The compounds were identified as (1*S*,20*R*,22*R*)-5*β*,6*β*:18,20-diepoxy-18-hydroxy-1-oxowitha-2,5,24-trienolide (**1**); (1*S*,20*R*,22*R*)-18,20-epoxy-4*β*,18-dihydroxy-1-oxowitha-2,5,24-trienolide (**2**); (1*S*,18*R*,20*R*,22*R*)-4*β*-hydroxy-18,20-epoxy-18-methoxy-1-oxowitha-2,5,24-trienolide (**3**); (1*S*,18*S*,20*R*,22*R*)-4*β*-hydroxy-18,20-epoxy-18-methoxy-1-oxowitha-2,5,24-trienolide (**4**); (1*S*,20*R*,22*R*)-4*β*-hydroxy-18,20-epoxy-1,18-dioxowitha-2,5,24-trienolide (**5**); (1*S*,18*R*,20*R*,22*R*)-18,20-epoxy-18-methoxy-1,4-dioxowitha-2,5,24-trienolide (**6**); (1*S*,18*S*,20*R*,22*R*)-18,20-epoxy-18-methoxy-1,4-dioxowitha-2,5,24-trienolide (**7**); and (1*S*,20*R*,22*R*)-5*β*,6*β*-epoxy-4*β*,18,20-trihydroxy-1-oxowitha-2,24-dienolide (**8**). Compounds **1** and **2** were obtained as epimeric mixtures at C-18.

*Vassobia* Rusby is a genus in the Solanaceae represented by five species that grow in South America. It has been related to *Acnistus*, *Dunalia*, and *Iochroma* species,<sup>1</sup> and, considering the different opinions concerning the systematic position of this plant at the trivial level, identification of the withanolides may be important from a chemotaxonomic point of view.

We now report the isolation of eight new withanolides (**1–8**) from *Vassobia lorentzii* (Dammer) A. T. Hunziker. All of these compounds have a functionalized C-18 at various oxidation levels (alcohol, aldehyde, and lactone) and they are closely related to those isolated from *Dunalia brachyacantha*<sup>2</sup> and to the withaphysalins from *Physalis minima*.<sup>3</sup> It is noteworthy that all plants originally classified as *Dunalia* (i.e., *Acnistus*, *Dunalia*, *Iochroma*, and *Vassobia*) have yielded 18-oxygenated withanolides.<sup>4</sup> In contrast to a previous report on *V. lorentzii* (at that time *Acnistus lorentzii*),<sup>5</sup> withaferin A was not detected in the present study. Structures of the new compounds were determined using a combination of techniques including 2D NMR, molecular modeling, and chemical transformations.

### Results and Discussion

Withaphysalin F (**1**), C<sub>28</sub>H<sub>36</sub>O<sub>7</sub>, did not show a molecular ion in its EIMS but a peak at *m/z* 466 (0.3%) corresponding to the [M – H<sub>2</sub>O]<sup>+</sup> ion was observed; an intense fragment at *m/z* 125 (34%) was indicative of an unsaturated  $\delta$ -lactone side chain, typical of many withanolides. The HRFABMS (*m*-nitrobenzyl alcohol, NaCl) showed a [M + Na]<sup>+</sup> ion at *m/z* 507.2375 that was consistent with the proposed formula. The <sup>1</sup>H NMR spectrum of **1** indicated that it was an approximately 1:1 mixture of two stereoisomers, which we were unable to separate (Table 1). Despite this, the low-field part of the spectrum, in conjunction with the H-4 doublet at  $\delta$  3.77 (*J* = 5.8 Hz), indicated a 4*β*-hydroxy-2-en-1-one system in ring A, identical to that described for withaferin A. The H-6 signal at  $\delta$  3.24 confirmed the presence of a 5*β*,6*β*-epoxide.<sup>6</sup> The rest of the spectrum was

identical to that of lactol **9**,<sup>2</sup> the only difference being absence of the acetate at C-4. The <sup>13</sup>C NMR spectrum of **1** (Table 2) was also almost identical to that of **9**, except for the resonances of carbons C-1 to C-6 due to the absence of the C-4 acetate. Lack of a C-18 methyl signal and the presence of two methine carbon resonances at  $\delta$  101.0 and 103.6 indicated the presence of a mixture of epimeric hemiacetals at C-18. This agreed with two signals at  $\delta$  5.15 (s) and 5.26 (s) in the <sup>1</sup>H NMR spectrum, which were assigned to H-18.


Withaphysalin G (**2**) C<sub>28</sub>H<sub>36</sub>O<sub>6</sub>, showed a [M + Na]<sup>+</sup> ion at *m/z* 491.2449 in the HRFABMS (*m*-nitrobenzyl alcohol, NaCl). The EIMS did not show a molecular ion, but a peak at *m/z* 450 (8%) corresponding to the [M – H<sub>2</sub>O]<sup>+</sup> ion; an intense fragment at *m/z* 125 was indicative of an unsaturated  $\delta$ -lactone side chain. The <sup>1</sup>H and <sup>13</sup>C NMR spectra of **2** were very similar to those of **1** and indicated that withaphysalin G was also an approximately 1:1 mixture of hemiacetals, epimeric at C-18 (Tables 1 and 2). As in the previous case, these stereoisomers could not be separated, and their spectral data were obtained from the mixture. The <sup>1</sup>H NMR spectrum of compound **2** (Table 1) exhibited signals typical of a 2,5-dien-1-one system with a substituent at C-4.<sup>7</sup> A 4*β*-hydroxy group was inferred from the presence of a doublet at  $\delta$  4.64 (*J* = 4.4 Hz) corresponding to H-4; coupling of the latter to H-3 confirmed the  $\beta$  stereochemistry of this substituent.<sup>8</sup> The <sup>13</sup>C NMR spectrum (Table 2) showed only four methyl signals and two methine carbons ( $\delta$  101.1 and 103.8) assigned to C-18 of the epimeric hemiacetals. The <sup>1</sup>H and <sup>13</sup>C NMR data for rings C–D, the side chain and the  $\gamma$ -hemiacetal ring were almost identical to that of **1** and of the 5,6-epoxy analogue recently isolated from *D. brachyacantha* in which the isomer with the higher chemical shift for H-18 had the 18*S* configuration.<sup>2</sup> Acetylation of the epimeric mixture **2** (Ac<sub>2</sub>O/pyridine, 25 °C) gave a single diacetyl derivative (**10**). The close similarity of <sup>13</sup>C chemical shifts for carbons 13–22 when compared to (18*R*)-**2** supported the 18*S* stereochemistry for diacetate **10**.

The <sup>1</sup>H and <sup>13</sup>C NMR spectra of compounds **3** and **4** (Tables 1 and 2) were very similar to those of **2**, indicating the same substitution pattern for rings A–C and the side chain. Both compounds showed a single C-18 resonance

\* To whom correspondence should be addressed. Tel./Fax: (54-11)4576-3385. E-mail: burton@qo.fcen.uba.ar.

<sup>†</sup> Universidad de Córdoba.

<sup>‡</sup> Universidad de Buenos Aires.



shifted downfield about 7 ppm with respect to **2**, and an additional methyl signal at  $\delta$  55–56. These data were consistent with compounds **3** and **4** being the corresponding 18-methylacetals of each epimer of withaphysalin G (**2**). Withaphysalin H (**3**) showed a singlet in its <sup>1</sup>H NMR spectrum at  $\delta$  4.62 (H-18) and a sharp three-proton singlet at  $\delta$  3.40 corresponding to the 18-methoxy group (assignments confirmed from HETCOR spectrum). As compared with **3**, withaphysalin I (**4**) showed a downfield shift of the H-18 signal ( $\delta$  4.75) and upfield shifts of the H-21 (to  $\delta$  1.35) and methoxyl (to  $\delta$  3.31) resonances. The FABMS (glycerol) of both compounds showed an intense quasimolecular ion [M+1] at  $m/z$  483, indicating an additional methyl group, and HRFABMS data was consistent with the molecular formula C<sub>29</sub>H<sub>38</sub>O<sub>6</sub> for both compounds. Intense fragments at  $m/z$  125 (100%) in the EIMS in both cases indicated a  $\delta$ -lactone unsaturated side chain.

The stereochemistry at C-18 was determined by the combined analysis of the H-21 shifts, the NOESY spectrum, and molecular modeling of both (18*R* and 18*S*) epimers. Thus, in the 18*R* epimer (**3**), the 1,3-pseudodiaxial relationship between methyl-21 and the 18-methoxy group would give rise to a deshielding effect on the former. In the NOESY spectrum of **4** the correlations observed for the pairs H-12 $\beta$ /H-18 and H-12 $\beta$ /21-CH<sub>3</sub> are only possible for the 18*S* stereoisomer in which H-18 is positioned on the same side as H-12 $\beta$  and H-21. On the other hand, in the NOESY spectrum of **3** a correlation was observed for the pair H-18/H-15 $\beta$  as expected from the geometries obtained by semiempirical AM1 calculations. No correlation was observed for the pair H-18/21-CH<sub>3</sub> in either epimer in accordance with the AM1 calculated structures, which predicted this distance to be larger than 3.5 Å. Thus, the structure of **3** was determined to be (17*S*,18*R*,20*R*,22*R*)-4 $\beta$ -hydroxy-18,20-epoxy-18-methoxy-1-oxowitha-2,5,24-trienolide, and that of **4** as its 18*S* epimer.

The HRFABMS (*m*-nitrobenzyl alcohol, NaCl) of withaphysalin J (**5**) showed a [M + Na]<sup>+</sup> ion at  $m/z$  489.2274 consistent with the formula C<sub>28</sub>H<sub>34</sub>O<sub>6</sub>. The EIMS of **5** showed the molecular ion at  $m/z$  466 (10%) and a fragment

at  $m/z$  125 (35%), indicative of an  $\alpha,\beta$ -unsaturated  $\delta$ -lactone ring. The <sup>1</sup>H and <sup>13</sup>C NMR chemical shifts of compound **5** (Tables 1 and 2) were closely related to those of **2**. The singlet for the 21-methyl at  $\delta$  1.53 indicated the presence of an oxygen function at C-20. The lack of an 18-methyl signal in conjunction with the presence of an IR band at 1748 cm<sup>-1</sup> ( $\gamma$ -lactone carbonyl) and the presence of a carbonyl carbon signal at  $\delta$  177.6 in the <sup>13</sup>C NMR spectrum (Table 2) led to the conclusion that withaphysalin J (**5**) contained an 18,20- $\gamma$ -lactone as in other withaphysalins.<sup>9</sup> The <sup>1</sup>H and <sup>13</sup>C NMR data for rings C–D, the side chain, and  $\gamma$ -lactone ring were coincident with that found for the synthetic lactone obtained by oxidation of an 18,20-hemiacetal of *D. brachyacantha*.<sup>2</sup>

The <sup>1</sup>H NMR spectrum (Table 1) of compounds **6** and **7** were almost identical to those of compounds **3** and **4**, with differences in the substitution pattern of rings A and B. In the low-field region, an AB system at  $\delta$  6.74 and 6.67 ( $J$  = 10.5 Hz) and the proton at  $\delta$  6.84 (dd,  $J$  = 5.5; 2.3 Hz) suggested the presence of a 2,5-diene-1,4-diketone system in rings A and B.<sup>7</sup> The substitution pattern in ring A was further corroborated by the signals at  $\delta$  202.0 and 187.7 in the <sup>13</sup>C NMR spectrum (Table 2) that were assigned to C-1 and C-4, respectively. The <sup>13</sup>C NMR spectrum of the 18*R* epimer (withaphysalin K, **6**) showed five methyl groups, four of which were coincident with C-21, C-27, C-28, and the methoxyl group of compound **3**. The methine signal at  $\delta$  108.1 (C-18) confirmed the presence of a 18-methylacetal group. Its <sup>1</sup>H NMR spectrum showed signals for H-18 and the methoxyl group also coincident with those in **3**. The 18*S* epimer (withaphysalin L, **7**) had H-18 shifted downfield to  $\delta$  4.75 and showed the expected singlet at  $\delta$  3.30 (as in **4**) assigned to the 18-methoxy group. The <sup>13</sup>C NMR spectrum also showed a methoxyl group at the same chemical shift as in **4**. The HRFABMS of **6** and **7** gave quasimolecular ions [M + Na] consistent with the molecular formula C<sub>29</sub>H<sub>36</sub>O<sub>6</sub>.

The HRFABMS (*m*-nitrobenzyl alcohol, NaCl) of 18-hydroxy-withanolide D (**8**) showed a [M + Na]<sup>+</sup> ion at  $m/z$  509.2538 consistent with the formula C<sub>28</sub>H<sub>38</sub>O<sub>7</sub>. Compound **8** did not have an 18,20-hemiacetal group but showed in its <sup>1</sup>H NMR spectrum (Table 1) only four methyl singlets assigned to C-19, C-21, C-27, and C-28. The missing methyl-18 signal and the appearance of an AB system at  $\delta$  3.59 and 3.60 ( $J$  = 10.9 Hz) indicated that C-18 was present as a hydroxymethyl group. The <sup>1</sup>H resonances were coincident with those of its 18-*O*-acetyl derivative isolated from *Ichroma fuschsioides*,<sup>10</sup> except for H-18 which was shifted upfield, supporting the presence of a free hydroxyl at C-18. The <sup>13</sup>C NMR (Table 2) was also almost identical to that of the known 18-acetate, confirming the proposed structure for **8**. The spectral data of **8** were also similar to the 20-deoxy analogue recently isolated from *D. brachyacantha*.<sup>2</sup>

## Experimental Section

**General Experimental Procedures.** <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker AC-200 NMR spectrometer at 200.13 and 50.32 MHz, respectively, using CDCl<sub>3</sub> as solvent. Multiplicity determinations (DEPT) and 2D spectra (COSY-45, HETCOR, NOESY) were obtained using standard Bruker software. Chemical shifts are given in parts per million (ppm) downfield from TMS as internal standard. EIMS were collected on a VG Trio-2 mass spectrometer at 70 eV by direct inlet; FABMS were measured on a VG ZAB-BEQ mass spectrometer, and HRFABMS were measured on a VG 7070-HF mass spectrometer. IR and UV spectra were measured on a Nicolet 5-SXC-FTIR and a Shimadzu UV 260 spectrophotometer,

**Table 1.** <sup>1</sup>H NMR Spectral Data for the Relevant Protons of Compounds **1–8** and **10** (CDCl<sub>3</sub>, 200.13 MHz)<sup>a</sup>

proton	<b>1</b> <sup>b</sup>	<b>2</b> <sup>b</sup>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>10</b>
2	6.18 d (9.9) [6.19 d (9.9)]	5.95 d (10.0)	5.95 d (10.0)	5.95 d (10.0)	5.96 d (10.0)	6.74 d <sup>c</sup> (10.5)	6.74 d <sup>d</sup> (10.5)	6.19 d (9.9)	6.00 dd (10.0)
3	6.94 dd (9.9; 5.8) [6.78 dd (9.9; 5.8)]	6.76 dd (10.0; 4.4) [6.78 dd (10.0; 4.4)]	6.80 dd (10.0; 4.5)	6.78 dd (10.0; 4.5)	6.79 dd (10.0; 4.4)	6.67 d <sup>c</sup> (10.5)	6.67 d <sup>d</sup> (10.5)	6.92 dd (9.9; 5.5)	6.70 dd (10.0; 4.6)
4 $\alpha$	3.77 d (5.8)	4.64 d (4.4)	4.66 d (4.5)	4.66 d (4.5)	4.65 d (4.4)			3.77 d (5.5)	5.79 d (4.6)
6	3.24 br s [3.26 br s]	5.95 br d (3.7)	5.93 br d (5.5)	5.93 br d (5.5)	5.94 br d (3.7)	6.84 dd (5.5; 2.3)	6.84 dd (6.0; 2.1)	3.24 br s	6.07 br d (4.4)
7 $\alpha$	1.34 m	1.60 m	1.71 m	1.60 m	1.35 m	1.83 m		1.28 m	1.80m
7 $\beta$	2.21 m	2.15 m	2.18 m	2.19 m	2.23 m	2.37 m	2.37 m	2.15 m	2.22 m
18	5.15 s [5.26 s]	5.18 s [5.30 s]	4.62 s	4.75 s		4.61 s	4.75 s	3.63 d (10.9) 3.59 d (10.9)	6.22 s
19	1.39 s [1.41 s]	1.42 s [1.44 s]	1.44 s	1.43 s	1.53 s	1.35 s	1.34 s	1.43 s	1.48 s
21	1.48 s [1.28 s]	1.49 s [1.29 s]	1.40 s	1.35 s	1.53 s	1.43 s	1.37 s	1.43 s	1.27 s
22	4.43 dd (13.2; 3.4) [4.52 dd (13.2; 3.4)]	4.44 dd (13.3; 3.5) [4.54 dd (13.3; 3.5)]	4.48 dd (13.1; 3.2)	4.42 dd (13.0; 3.0)	4.54 dd (13.2; 3.7)	4.45 dd (13.0; 3.0)	4.45 dd (13.0; 3.0)	4.28 dd (13.1; 3.6)	4.50 dd (13.5; 3.5)
23 $\alpha$	2.02 m	2.05 m [2.20 m]	2.04 m	2.13 m	2.03 m	2.10 m	2.09 m	2.10 m	2.09 m
23 $\beta$		2.44 m [2.51 m]	2.42 m	2.40 m	2.42 m	2.43 m		2.37 m	2.49 m
27	1.88 br s	1.87 br s	1.89 br s	1.89 br s	1.89 br s	1.89 br s	1.90 br s	1.89 br s	1.89 br s
28	1.93 br s	1.96 br s	1.96 br s	1.94 br s	1.96 br s	1.94 br s	1.94 br s	1.96 br s	1.94 br s
OMe			3.40 s	3.31 s		3.39 s	3.30 s		
OAc									2.07 s 2.08 s

<sup>a</sup> Chemical shifts are in ppm ( $\delta$ ) downfield from TMS,  $J$  couplings (in parentheses) are in Hz. <sup>b</sup> Chemical shift data correspond to the 18*R* epimer. Distinct resonances for the 18*S* epimer observed in the spectrum of the epimeric mixture are shown in square brackets. <sup>c,d</sup> Assignments may be interchanged.

**Table 2.** <sup>13</sup>C NMR Spectral Data of Compounds **1–8** and **10** (CDCl<sub>3</sub>, 50.32 MHz)

carbon	<b>1</b> <sup>a</sup>		<b>2</b> <sup>a</sup>		<b>3</b> (18 <i>R</i> )	<b>4</b> (18 <i>S</i> )	<b>5</b>	<b>6</b> (18 <i>R</i> )	<b>7</b> (18 <i>S</i> )	<b>8</b>	<b>10</b>
	(18 <i>R</i> )	(18 <i>S</i> )	(18 <i>R</i> )	(18 <i>S</i> )							
1	202.5	202.5	203.6	203.9	203.4	203.6	203.6	202.0	202.3	202.3	203.0
2	131.9	132.0	130.5	131.1	130.4	131.0	130.5	140.2	140.1	132.0	133.2
3	142.1	142.4	143.2	143.0	143.1	143.1	143.0	139.0	139.2	141.9	140.2
4	69.7	69.7	69.2	69.2	69.2	69.3	69.3	187.7	187.7	69.7	69.7
5	63.7	63.9	138.9	138.6	139.0	138.7	139.2	139.6	139.1	63.8	134.5
6	62.6	62.7	128.8	128.8	128.8	128.9	128.9	137.1	137.7	62.8	130.7
7	31.3 <sup>b</sup>	31.3	31.1 <sup>d</sup>	31.1	31.1 <sup>f</sup>	31.3	31.3	31.3	31.4	31.6	31.0 <sup>i</sup>
8	30.9	30.3	33.5	32.9	33.8	33.3	30.6	31.0	31.0	29.6	33.2
9	43.9	44.4	42.4	42.9	42.3	42.8	42.0	41.9	42.5	44.5	42.3
10	47.8	47.8	49.4	49.4	49.4	49.3	49.6	51.4	51.4	47.9	49.1
11	24.6	24.6	25.3	25.3	26.1	25.4	22.8	24.5	25.0	22.1	26.3
12	34.6	37.0	34.7	37.2	34.7	36.9	35.3	34.7	34.7	34.8	34.8
13	58.0	58.8	58.0	58.8	57.9	59.0	55.4	58.0	59.2	47.7	57.9
14	56.4 <sup>c</sup>	54.7	56.4 <sup>e</sup>	54.7	56.4	54.0	55.9	56.0 <sup>g</sup>	56.1	55.1 <sup>h</sup>	56.5
15	25.4	25.4	25.2	25.2	25.5	24.9	25.9	25.5	25.6	23.7	24.9
16	26.2	27.5	26.2	27.5	26.0	27.1	27.4	26.1	26.2	31.4	25.5
17	57.0 <sup>c</sup>	57.0	56.9 <sup>e</sup>	56.9	56.3	56.5	52.6	56.3 <sup>g</sup>	56.5	55.9 <sup>h</sup>	56.5
18	101.0	103.6	101.1	103.8	108.3	110.4	177.6	108.1	110.3	58.9	99.3
19	17.6	17.8	22.9	22.9	22.7	22.8	22.9	23.6	23.7	17.8	21.3
20	85.1	84.8	85.1	84.8	84.8	84.5	83.6	84.8	84.6	75.5	86.6
21	21.8	23.7	21.6	24.5	20.9	23.7	23.4	21.0	23.8	21.6	21.1
22	80.5	80.5	80.6	80.5	80.7	81.5	79.0	80.7	81.4	80.6	80.2
23	31.5 <sup>b</sup>	32.1	31.3 <sup>d</sup>	32.0	31.0 <sup>f</sup>	32.2	31.2	32.7	32.3	31.6	31.3 <sup>i</sup>
24	147.7	148.6	148.1	148.9	147.9	148.1	147.2	147.7	147.9	148.4	147.4
25	122.3	122.0	122.2	122.0	122.2	122.2	122.6	122.4	122.4	122.3	122.6
26	165.7	165.7	165.9	165.9	166.0	166.0	165.1	165.9	165.9	165.5	165.9
27	12.5	12.4	12.5	12.4	12.5	12.4	12.5	12.5	12.5	12.5	12.5
28	20.5	20.4	20.5	20.5	20.3	20.0	20.4	20.3	20.4	20.6	20.4
OMe					55.7	55.1		55.7	55.1		
OAc											170.1, 170.2 21.5

<sup>a</sup> Chemical shifts ( $\delta$ ) determined in the *R/S* mixture (ca. 1:1). <sup>b–i</sup> Assignments may be interchanged.

respectively. Melting points were obtained on a Fisher-Johns apparatus and are uncorrected. Chromatographic separations were performed by vacuum liquid chromatography (VLC), column chromatography on Si gel 60 (40–63  $\mu$ m), and preparative TLC (Si gel 60 G F<sub>254</sub> 2 mm thick).

**Plant Material.** Aerial parts of *V. lorentzii* were collected in November 1994, in Department of Andalgalá, Catamarca Province, Argentina. A voucher specimen is deposited at the Museo Botánico, Universidad Nacional de Córdoba [CORD].

**Extraction and Isolation.** The dried and pulverized aerial part of *V. lorentzii* (1.83 kg) was extracted with EtOH at room temperature, the solvent was evaporated, and the residue (78 g) was partitioned between hexane, MeOH, and H<sub>2</sub>O (30:3:1). The aqueous MeOH layer was washed with hexane, concentrated, and extracted with CHCl<sub>3</sub>. The residue (21.5 g) obtained after evaporation of the solvent was fractionated by VLC eluting with hexanes–EtOAc mixtures of increasing polarity (100:0–0:100) to yield four fractions containing withanolides.



These fractions, eluted with hexanes–EtOAc from 50:50 to 10:90, were further fractionated by flash chromatography to yield a fraction (354 mg) containing the mixture **2** of epimers as the major component. The minor fractions were purified by flash chromatography and preparative TLC to yield withanolides **1** (8.3 mg), **3** (22 mg), **4** (6 mg), **5** (6 mg), **6** (3 mg), **7** (4 mg), and **8** (90 mg).

**(17S,20R,22R)-5 $\beta$ ,6 $\beta$ :18,20-diepoxy-18-hydroxy-1-oxowitha-2,5,24-trienolide (1, 18R and 18S)**: white crystals (hexanes–EtOAc); mp 256–258 °C;  $[\alpha]_D^{25} +46.9^\circ$  (*c* 0.06, CHCl<sub>3</sub>); UV (MeOH)  $\lambda_{\max}$  225 nm; IR (film)  $\nu_{\max}$  3485, 2978, 2825, 1701, 1689 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* [M]<sup>+</sup> absent, 466 (0.3) [M – H<sub>2</sub>O]<sup>+</sup>, 152 (6), 125 (34); FABMS (glycerol, KCl) *m/z* 523 (100) [M + K]<sup>+</sup>, 467 (44) [M – H<sub>2</sub>O + 1]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 507.2375 (calcd for C<sub>28</sub>H<sub>36</sub>O<sub>7</sub>Na 507.2359).

**(17S,20R,22R)-18,20-Epoxy-4 $\beta$ ,18-dihydroxy-1-oxowitha-2,5,24-trienolide (2, 18R and 18S)**: white crystals (hexanes–EtOAc); mp 264–266 °C (dec);  $[\alpha]_D^{25} +97.6^\circ$  (*c* 0.13, MeOH); UV (MeOH)  $\lambda_{\max}$  224 nm; IR (film)  $\nu_{\max}$  3350, 2958, 2849, 1703, 1684 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* [M]<sup>+</sup> absent, 450 (8) [M – H<sub>2</sub>O]<sup>+</sup>, 432 (8), 343 (17), 171 (45), 152 (46), 125 (94), 105 (62), 91 (100) and 55 (81); FABMS (glycerol, NaCl) *m/z* 491 (17) [M + Na]<sup>+</sup>, 451 (17) [M – H<sub>2</sub>O + 1]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 491.2449 (calcd for C<sub>28</sub>H<sub>36</sub>O<sub>6</sub>Na 491.2410).

**Acetylation of Compound 2**. The mixture **2** (10 mg) was dissolved in Ac<sub>2</sub>O/pyridine (1:1, 0.3 mL) and left for 18 h at 25 °C. Dilution with CH<sub>2</sub>Cl<sub>2</sub> and evaporation under a stream of nitrogen afforded acetate **10** as a white powder; <sup>1</sup>H and <sup>13</sup>C NMR data, see Tables 1 and 2.

**(17S,18R,20R,22R)-4 $\beta$ -Hydroxy-18,20-epoxy-18-methoxy-1-oxowitha-2,5,24-trienolide (3)**: white amorphous powder;  $[\alpha]_D^{25} +47.9^\circ$  (*c* 0.20, CHCl<sub>3</sub>); UV (MeOH)  $\lambda_{\max}$  224 nm; IR (film)  $\nu_{\max}$  3427, 2953, 2881, 2828, 1704, 1687, 1666 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* [M]<sup>+</sup> absent, 450 (3) [M – CH<sub>3</sub>OH]<sup>+</sup>, 433 (8), 432 (14), 406 (7), 388 (4), 357 (3), 313 (3), 171 (47), 169 (17), 152 (48), 125 (100); FABMS (glycerol) *m/z* 483 (25) [M + 1]<sup>+</sup>, 467 (10), 451 (100) [M – OCH<sub>3</sub>]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 505.2579 (calcd for C<sub>29</sub>H<sub>38</sub>O<sub>6</sub>Na 505.2566).

**(17S,18S,20R,22R)-4 $\beta$ -Hydroxy-18,20-epoxy-18-methoxy-1-oxowitha-2,5,24-trienolide (4)**: white amorphous powder;  $[\alpha]_D^{25} +54.9^\circ$  (*c* 0.18, CHCl<sub>3</sub>); UV (MeOH)  $\lambda_{\max}$  223 nm; IR (film)  $\nu_{\max}$  3480, 2973, 2897, 1700, 1690 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* [M]<sup>+</sup> absent, 464 (4) [M – H<sub>2</sub>O]<sup>+</sup>, 450 (9) [M – CH<sub>3</sub>OH]<sup>+</sup>, 432 (23), 357 (30), 152 (69), 125 (100); FABMS (glycerol) *m/z* 483 (28) [M + 1]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 505.2566 (calcd for C<sub>29</sub>H<sub>38</sub>O<sub>6</sub>Na 505.2566).

**(17S,20R,22R)-4 $\beta$ -Hydroxy-18,20-epoxy-1,18-dioxowitha-2,5,24-trienolide (5)**: white amorphous powder (hexanes–EtOAc); mp 215–217 °C (dec);  $[\alpha]_D^{25} +60.4^\circ$  (*c* 0.15, CHCl<sub>3</sub>); UV (MeOH)  $\lambda_{\max}$  226, 312 nm; IR (film)  $\nu_{\max}$  3456, 1748, 1688, 1655, 1230, 1035 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* 466 (10) [M]<sup>+</sup>, 448 (3), 152 (10), 151 (10), 125 (35), 109 (34), 97 (18), 43 (100); FABMS (glycerol, KCl) *m/z* 505 (100) [M + K]<sup>+</sup>, 467 (8) [M + 1]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 489.2274 (calcd for C<sub>28</sub>H<sub>34</sub>O<sub>6</sub>Na 489.2253).

**(17S,18R,20R,22R)-18,20-Epoxy-18-methoxy-1,4-dioxowitha-2,5,24-trienolide (6)**:  $[\alpha]_D^{25} +41.1^\circ$  (*c* 0.25, CHCl<sub>3</sub>); UV (MeOH)  $\lambda_{\max}$  224, 284, 362 nm; IR film (AgCl)  $\nu_{\max}$  3545, 1700, 1676, 1668, 1620 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* [M]<sup>+</sup> absent, 465 (1) [M – CH<sub>3</sub>]<sup>+</sup>, 449 (5) [M – OCH<sub>3</sub>]<sup>+</sup>, 355 (52), 307 (15), 187 (7), 171 (7), 125 (36); FABMS (glycerol, KCl) *m/z* 481 (9) [M + 1]<sup>+</sup>, 465 (6), 449 (100) [M – OCH<sub>3</sub>]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 503.2440 (calcd for C<sub>29</sub>H<sub>36</sub>O<sub>6</sub>Na 503.2410).

**(17S,18S,20R,22R)-18,20-Epoxy-18-methoxy-1,4-dioxowitha-2,5,24-trienolide (7)**:  $[\alpha]_D^{25} +38.9^\circ$  (*c* 0.04, CHCl<sub>3</sub>); UV (MeOH)  $\lambda_{\max}$  226, 280, 362 nm; IR film (AgCl)  $\nu_{\max}$  3550, 1700, 1680, 1665, 1620 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* [M]<sup>+</sup> absent, 355 (4), 237 (3), 171 (4), 125 (28); FABMS (glycerol, KCl) *m/z* 481 (6) [M + 1]<sup>+</sup>, 449 (100) [M – OCH<sub>3</sub>]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 503.2447 (calcd for C<sub>29</sub>H<sub>36</sub>O<sub>6</sub>Na 503.2410).

**(17S,20R,22R)-5 $\beta$ ,6 $\beta$ -Epoxy-4 $\beta$ ,18,20-trihydroxy-1-oxowitha-2,24-dienolide (8)**: white crystals (hexanes–EtOAc); mp 175–177 °C (dec);  $[\alpha]_D^{25} +59.8^\circ$  (*c* 0.26, MeOH); UV (MeOH)  $\lambda_{\max}$  223 nm; IR (film)  $\nu_{\max}$  3385, 2930, 2880, 1690 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, (Tables 1 and 2); EIMS *m/z* [M]<sup>+</sup> absent, 361 (1.5), 343 (9), 325 (4), 169 (11), 152 (30), 55 (29); FABMS (*m*-nitrobenzyl alcohol, K<sub>2</sub>CO<sub>3</sub>) *m/z* 525 (100) [M + K]<sup>+</sup>, 467 (44) [M – H<sub>2</sub>O + 1]<sup>+</sup>; HRFABMS (*m*-nitrobenzyl alcohol, NaCl) *m/z* [M + Na]<sup>+</sup> 509.2538 (calcd for C<sub>28</sub>H<sub>38</sub>O<sub>7</sub>Na 509.2515).

**Acknowledgment.** We thank Prof. A. T. Hunziker (IMBIV-CONICET) for the collection and identification of the plant material and Prof. A. D. Kinghorn (University of Illinois at Chicago) for the high resolution mass spectra. This work was supported in part by Fundación Antorchas, CONICOR (Argentina), SeCyT–UNC, Universidad de Buenos Aires, and CONICET (Argentina). R.I.M. thanks CONICOR and Fundación Antorchas for fellowships.

**Supporting Information Available:** AM1 calculated structures and NOE data of **3** and **4**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References and Notes

- Hunziker, A. T. *Kurtziana* **1984**, *17*, 91–118.
- Silva, G. L.; Burton, G.; Oberti, J. C. *J. Nat. Prod.* **1999**, *62*, 949–953.
- Glottter, E.; Kirson, I.; Abraham, A.; Subramanian, S. S.; Sethi, P. D. *J. Chem. Soc., Perkin Trans. 1* **1975**, 1370–1374.
- Ray, A. B.; Gupta, M. *Prog. Chem. Org. Nat. Prod.* **1994**, *63*, 1–106.
- Conta, A. G.; Albónico, S. M.; Juliani, H. R. *An. Asoc. Quím. Argent.* **1971**, *59*, 373–375.
- Kupchan, S. M.; Anderson, W. K.; Bollinger, P.; Doskotch, R. W.; Smith, R. M.; Saenz Renauld, J. A.; Schnols, H. K.; Burlingame, A. L.; Smith, D. H. *J. Org. Chem.* **1969**, *34*, 3858–3866.
- Nittala, S. S.; Lavie, D. *Phytochemistry* **1981**, *20*, 2735–2739.
- Shingu, K.; Miyagawa, M.; Yahara, S.; Nohara, T. *Chem. Pharm. Bull.* **1993**, *41*, 1873–1875.
- Sinha, S. C.; Ray, A. B.; Oshima, Y.; Bagchi, A.; Hikino, H. *Phytochemistry* **1987**, *26*, 2115–2117.
- Raffauf, R. F.; Shemluck, M. J.; LeQuesne, P. W. *J. Nat. Prod.* **1991**, *54*, 1601–1606.

NP000022Z