

An Approach to Furolabdanes and Their Photooxidation Derivatives from *R*-(+)-Sclareolide

María C. de la Torre,^{*,†} Isabel García,[†] and Miguel A. Sierra[‡]

Instituto de Química Orgánica, Consejo Superior de Investigaciones Científicas (CSIC), Juan de la Cierva 3, 28006-Madrid, Spain, and Departamento de Química Orgánica, Facultad de Química, Universidad Complutense, 28040-Madrid, Spain

Received November 21, 2001

A synthetic route to furolabdanes from commercially available *R*-(+)-sclareolide is reported, with the specific aim of preparing (12*R* and 12*S*,15*ξ*)-12,15-dihydroxyabda-7,13-dien-16,15-olides (**3** and **5**) and (12*R* and 12*S*,16*ξ*)-12,16-dihydroxyabda-7,13-dien-15,16-olides (**4** and **6**). The key points of our approach are the use of Weinreb's amide **11** to join the furan ring to the terpenic unit. Photooxidation of the furan moiety of compounds **15** and **16**, and of their acetates **19** and **20**, has been used to build the hydroxybutenolide fragment. In this way the four possible isomers at the butenolide moiety, compounds **3–6**, and their C-12 acetyl derivatives **21–24** have been obtained. On the basis of comparison of the spectral data (¹H NMR) of the synthetic peracetates **25–28** (derived from **21–24**) with the reported data for the peracetate **2** (derived from the natural product **1**), the relative configuration at carbon C-12 of the natural product has been corrected. Furthermore, the absolute configuration of the natural product **1**, considered to belong to the *enantio*-series, has to be changed to the *normal*-series on the basis of the optical rotation obtained for the synthetic derivative.

Labdane diterpenoids are among the most common types of diterpenes isolated from higher plants.¹ The interest of this class of compounds resides in their significant anti-mutagenic,² antibacterial, and antifungal activities.³ However, as in the case of many other natural products, they can be isolated only in minute amounts. This fact inhibits their unambiguous structural determination and often precludes the study of biological and chemical reactivity. Therefore, the hemisynthesis of minor components from other abundant natural products is of longstanding interest. As an example, (12*S*,16*ξ*)-12,16-dihydroxy-*ent*-labda-7,13-dien-15,16-olide, **1**, has been recently isolated from *Alomira myriadenia* (Asteraceae) only in a 0.011% yield with respect to plant material. This compound shows significant cytotoxic activity against human oral epidermoid carcinoma and against colon cancer⁴ and is identical to one of the products previously isolated by Bohlmann from *Acritopappus hagei*,⁵ since both compounds yielded the same diacetylated derivative **2**. On the basis of biogenetic grounds, Bohlmann suggested an *enantio* absolute configuration for these compounds, although neither the relative stereochemistry at carbon C-12 nor the absolute configuration of compound **2** has been rigorously established.

Continuing with our research directed toward the hemisynthesis of terpenoids,⁶ we have been interested in the development of synthetic routes to furolabdane diterpenoids. With this in mind, (12*R*- and 12*S*)-12,15-dihydroxy-labda-7,13-dien-16,15-olides **3** and **5**, and their regioisomers (12*R*- and 12*S*)-12,16-dihydroxy-labda-7,13-dien-15,16-olides **4** and **6**, were attractive targets, since their synthesis would provide these labdanes in amounts sufficient for biological assay. Additional reward would be the structural determination of **1**. The study of compounds **3–6** shows that (+)-sclareolide (**7**) may be a suitable starting material for their synthesis. In fact (+)-sclareolide (**7**) bears the methyl groups at carbons C-4, C-8, and C-10 of the decalin moiety, placed in the appropriate positions. The butenolide fragment present in the synthetic targets **3–6** may be derived





Figure 1.

from the oxidation of a furan ring in an intermediate like **8** (Scheme 1). The furan moiety required for this transfor-


Scheme 1. Retrosynthetic Analysis of Compounds **3–6**



* To whom correspondence should be addressed. Tel: 34-915622900. Fax: 34-915644853. E-mail: iqot310@iqog.csic.es.

[†] Instituto de Química Orgánica.

[‡] Departamento de Química Orgánica.

Scheme 2^a

^a (i) 1 equiv of DIBAL-H, toluene, -78 °C (94%); (ii) Me₃Al-MeONHMe-HCl, CH₂Cl₂, 0 °C to rt (88%); (iii) SOCl₂, py, 0 °C; (iv) 3 equiv of 3-bromofuran, 3 equiv BuLi, THF, -78 °C, 1 h, then 1 equiv of **13**, 2 h (72%); (v) 2 equiv of CeCl₃ × 7H₂O, 8 equiv of NaBH₄, MeOH, 0 °C, 30 min (37% for **15**, 33% for **16**); (vi) Ac₂O/py (1:2) 24 h, rt (99% for **19**, 98% for **20**).

mation would be incorporated into the (+)-sclareolide derivative **9**, by reaction of 3-lithiofuran with the electrophilic function at carbon C-12. It should be noted that the proposed synthesis would be effected in a compound having a *normal* configuration instead of the *enantio* proposed for **1**.

Results and Discussion

The synthesis of the key furoabdanes, of general structure **8**, was initially addressed by sequential reduction of the (+)-sclareolide (**7**) to the lactol **10**, followed by coupling with 3-lithiofuran (Scheme 2). Lactol **10** was obtained quantitatively by treatment of (+)-sclareolide with DIBAL-H. In the next step, reaction of 3-lithiofuran with **10** must be performed below 40 °C, to avoid isomerization of the reagent to 2-lithiofuran.⁷ Unfortunately, lactol **10** was unreactive at this temperature. A change of strategy was required, and we devised Weinreb's amides as suitable candidates to react with 3-lithiofuran at the required temperature. Weinreb's amide **11** was prepared in 88% yield by aminolysis of (+)-sclareolide (**7**) with the dimethylaluminum amide derived from *N*-methoxy-*N*-methylamine.⁸ To prevent waste of organometallic reagent as well as undesired side-reactions, the tertiary alcohol at C-8 was dehydrated prior to the furane addition. Treatment of alcohol **11** with SOCl₂ at 0 °C yielded the undesired Δ⁸⁽¹⁷⁾ unsaturated derivative **12**, as the major reaction product (60%), together with the desired Δ⁷-unsaturated isomer **13** (32% yield). However, the *exo* double bond of compound **12** quantitatively rearranged to the desired *endo*-isomer **13**, under acid treatment. Reaction of alkene **13** with 3-lithiofuran at -78 °C yielded 12-ketolabdane **14** in excellent yield. Reduction of the ketone at C-12 with NaBH₄ and excess CeCl₃ × 7H₂O⁹ yielded the epimeric 12-hydroxyfuroabdanes **15** and **16** in a 1:1 ratio. These alcohols were separated by column chromatography, and their absolute configuration at carbon C-12 was established by using Mosher's methodology.¹⁰

Accordingly, the reaction of the alcohols **15** and **16** with (*R*)-(-)-α-methoxy-α-(trifluoromethyl)phenylacetic acid

Table 1. Chemical Shifts of H-14, H-15, H-16, and OMe for **17** and **18** and Δδ Values (δ₁₇-δ₁₈)^a

H	17	18	Δδ
14	6.43	6.27	+0.16
15	7.47	7.42	+0.05
16	7.37	7.36	+0.01
OMe	3.43	3.51	+0.08

^a At 300 MHz in CDCl₃.




Figure 2. Ideal conformations for the Mosher's esters **17** and **18**.


(MTPA) yielded esters **17** and **18**, respectively. The modified Mosher's¹⁰ method is based on the diamagnetic effect caused by the benzene ring of the MTPA moieties on the β-protons of the alcohol. In the case of α-aromatic secondary carbinols, an anisotropic effect due to the aromatic ring produces negligible Δδ values for these β-protons, which may preclude the use of this method to establish absolute configurations. In addition, the β-protons (H-11) on esters **17** and **18** overlapped with the decalin protons, preventing their analysis. Nevertheless, Isobe¹¹ has established that, for secondary alcohols bearing an α-furyl substituent, the analysis of the protons attached to the furan ring may be applied to determine the absolute configuration. As shown in Table 1, the furan protons H-14, H-15, and H-16 of ester **18** are shielded with respect to its epimer **17**. This result is fully consistent with the ideal conformation for ester **18** represented in Figure 2, in which the furan ring is eclipsed with the benzene ring of the MTPA group. Therefore, ester **18** and subsequently alcohol **16** must have a C-12*S* absolute configuration. In agreement with these results,

the -OMe protons of the MTPA moiety in ester **17** are shielded by the diamagnetic effect caused by the furan [$\Delta\delta$ (**17**-**18**) = -0.08], and therefore ester **17** and hence alcohol **15** must have a *12R* absolute configuration.

With alcohols **15** and **16** in hand, we pursued the oxidation of the furan ring to obtain the desired hydroxybutenolides. Among the number of methods that have been developed for the synthesis of 5-hydroxy-2(5*H*)-furanones,¹² the one involving oxidation of furan by ¹O₂ appears to be the most efficient.¹³ In general, the photooxidation of 3-substituted furans is not regioselective, yielding both the 2-alkyl-4-hydroxy- and the 3-alkyl-4-hydroxybutenolide regioisomers.¹⁴ This fact may be advantageous for us, since we were interested in obtaining both regioisomeric hydroxybutenolides to compare with the natural labdane **1**.^{13,14} Irradiation of alcohol **15** in THF (220 W, tungsten lamp, Pyrex well) in the presence of air and a catalytic amount of Rose Bengal for 5 h yielded the regioisomeric hydroxybutenolides **3** and **4**. Both lactones were separated by column chromatography as a mixture of epimers in a 16% and 11% yield, respectively. The ¹H NMR spectrum of **3**, obtained in CDCl₃, is consistent with an α -alkyl-substituted-15-hydroxybutenolide. Specially relevant are the signals at δ 7.04, which must be assigned to proton H-14, placed at the β -position of an α,β -unsaturated butenolide, and the signal at δ 6.12 attributable to H-15.¹⁵ With respect to the regioisomeric lactone derivative **4**, the most deshielded signal in the ¹H NMR spectrum appeared at δ 6.20, and therefore the structure of α,β -alkyl-substituted-16-hydroxybutenolide has been assigned to this compound.¹⁵ Therefore, we established the structure of (12*R*)-12,15-dihydroxyabda-7,13-dien-16,15-olide for compound **3** and the structure of (12*R*)-12,16-dihydroxyabda-7,13-dien-15,16-olide for compound **4**. Alcohol **16**, having a *12S* configuration, was submitted to analogous photooxidation conditions, yielding compounds **5** and **6** in 20% and 10% yields, respectively. The ¹H NMR of product **5** showed signals at δ 7.08 and 7.04 attributable to H-14, and at δ 6.14 for proton H-15. Compound **6** showed a signal at δ 6.23 that could be assigned to proton H-16 in a 16-hydroxy-15,16-butenolide. In addition, the olefinic proton (H-14) of **6** appears at δ 6.07, 1 ppm shielded with respect to its regioisomer **5**. Consequently, we established a structure of (12*S*)-12,15-dihydroxyabda-7,13-dien-16,15-olide for compound **5** and (12*S*)-12,16-dihydroxyabda-7,13-dien-15,16-olide for compound **6**.


In an attempt to increase the yield of the photooxidation reaction, we checked the oxidation of the 12-acetyl derivatives **19** and **20**, quantitatively obtained from alcohols **15** and **16**, respectively, by treatment with Ac₂O-Pyr. Acetate **19** yielded upon sun light irradiation, under the above conditions, the regioisomeric hydroxybutenolides **21** and **22** (35% and 33% yield, respectively). Both compounds were obtained as mixtures of epimers at carbons C-15 and C-16. Again the 15-hydroxy-16,15-butenolide isomer **21** showed signals for protons H-14 and H-15 at δ 6.94 and 6.09, respectively. On the contrary, the ¹H NMR of derivative **22** showed a signal for δ H-14, at 5.97, shielded with respect to **21**, while H-16 appeared at δ 6.20 and 6.03. Thus, compound **22** must be the 16-hydroxy-15,16-butenolide. In turn, 12*S*-acetyl derivative **20** yielded lactones **23** (42%) and **24** (23%) under analogous reaction conditions. As before, lactone **23** showed signals at δ 7.02-7.01 for H-14 and at 6.13-6.10 for H-15, and therefore it was assigned the structure of (12*S*)-12-acetoxy-15-hydroxyabda-7,13-dien-16,15-olide, derivative **24** being the 16-hydroxy-15,16-butenolide isomer. From these results it is clear that

Scheme 3



although both free alcohols **15** and **16** and their acetates **19** and **20** are suitable substrates for the photooxidation reaction, clearly the presence of the free alcohol group considerably decreases the reaction yields.





Despite the reported data,⁴ 12-hydroxybutenolides **3**-**6** were very unstable and their ¹H NMR spectra could not be obtained in *py-d*₅. This instability thwarted the direct comparison with the literature data for **1**.⁴ Thus, the diacetates **25**-**28** were prepared as mixtures of epimers at the hemiacetalic carbon, by Ac₂O-py treatment of lactones **21**-**24**, to compare their ¹H NMR data with those reported in the literature for **2**.⁴ Comparison of the ¹H NMR spectra for the 15-acetoxy-16,15-butenolides **25** and **27** (C12*R* and C-12*S*, respectively) with the ¹H NMR spectra for the corresponding 15-hydroxy analogues **21** and **23** (Table 2) showed a strong downfield shift of the signal attributed to H-15, confirming the acetylation of its geminal hydroxyl group ($\Delta\delta \approx 0.9$ ppm). Similarly, the ¹H NMR of the 16-acetoxy-15,16-butenolides **26** and **28** showed the signals for H-16 deshielded ($\Delta\delta \approx 0.9$ ppm), with respect to the 16-hydroxy derivatives **22** and **24**, while the signal for the olefinic proton H-14 remained unchanged upon acetylation. Thus, the structures of (12*R* and 12*S*)-12,15-diacetoxyabda-7,13-dien-16,15-olide (**25** and **27**) and of (12*R* and 12*S*)-12,16-diacetoxyabda-7,13-dien-15,16-olide (**26** and **28**) have been unambiguously established.



- 25** R¹ = OAc, R² = H, X = H, OAc, Y = O (from **21** 80%)
26 R¹ = OAc, R² = H, X = O, Y = H, OAc (from **22** 90%)
27 R¹ = H, R² = OAc, X = H, OAc, Y = O (from **23** 90%)
28 R¹ = H, R² = OAc, X = O, Y = H, OAc (from **24** 85%)

Figure 3.

Table 2. Chemical Shifts of H-14, H-15, H-16, and H-12 of Compounds **21–28** and **2**

	Compound	H-14	H-15	H-16	H-12
	21 R=H	6.94 m	6.09 m	-----	5.70 dd (6.7, 3.8)
	25 R=25	6.97 m, 6.87 t (1.6)	6.89, 6.97 m	-----	5.76 td (7.8, 1.6)
	22 R=H	5.97 m	-----	6.20, 6.03 m	5.56 d (10.6)
	26 R=Ac	6.12 dd (1.6, 1.1), 6.04 t (1.1)	-----	7.02 d (1.0), 6.94 d (0.9)	5.81 dt (11.7, 1.1), 5.62 d (13.5)
	23 R=H	7.01, 7.00 m	6.12, 6.09 m	-----	5.57 t (6.5)
	27 R=Ac	7.05 m	6.92, 6.98 d (1.1)	-----	5.46 t (7.1), 5.62 t (7.3)
	24 R=H	6.02 m	-----	6.24, 6.05 m	5.36 t (7.7)
	28 R=Ac	6.16, 6.07 t (1.1)	-----	7.04 m, 6.97 d (1.0)	5.68 td (8.1, 1.2), 5.46 (overlapped)
	2^a	6.99, 7.05 dd (0.8, 1.0)	-----	6.09, 6.18 dd (1.1, 0.8)	5.70, 5.10 dt (7.5, 1.2)

^a Taken from ref 4.

Comparison of the ¹H NMR data reported for **2⁴** with the ¹H NMR of diacetates **25–28** clearly shows that derivatives **2** and **28** have identical ¹H NMR spectra. According to this, diacetate **2** and hence the natural product **1** have in fact a 15,16-butenolide moiety. Therefore, the assignment of H-14 and H-16 protons given in the literature for the diacetate **2** must be changed, while the stereochemistry at C-12 in **2** must be the reverse. Consequently, overlooking the absolute stereochemistry, the structure of the reported natural product should be amended to **6**.

Regarding the absolute configuration of compound **1** (isolated from *A. myriadenia*), the sign of the specific rotation reported for its diacetate **2** is negative, like our synthetic diacetate **28** obtained from **24**. Assuming that the ratio of anomeric acetates obtained from **1** and **24** are the same, we conclude that **1** and **28** possess the same absolute configuration. Therefore, diterpene **1** should be linked to the *normal* labdane series.

Experimental Section

General Experimental Procedures. Melting points were determined on a Kofler block. Optical rotations were measured on a Perkin-Elmer 241 MC polarimeter. IR spectra were obtained on a Perkin-Elmer 681 spectrophotometer. ¹H NMR spectra were recorded using a Bruker AM 200 apparatus at 200 MHz or a Varian INOVA-300 spectrometer at 300 MHz. ¹³C NMR spectra were recorded at 50.3 or 75 MHz. Chemical shifts for ¹H NMR are reported with respect to residual CHCl₃ (δ 7.25) and with respect to CDCl₃ (δ 77.00) for ¹³C NMR spectra. MS were recorded in the positive EI mode on a Hewlett-Packard HP 5989A instrument (70 eV). Elemental analyses were made with a Carlo Erba EA 1108 apparatus. *R*-(+)-Sclareolide was from Aldrich, and it was used as received. All reagents were used as obtained from commercial sources. Methylene chloride (CH₂Cl₂), toluene (C₆H₅-CH₃), and tetrahydrofuran (THF) were distilled under positive pressure of argon from CaH₂ or Na-benzophenone. Other

solvents were HPLC grade and were used without purification. Na₂SO₄ was used to remove water from the organic layer in reaction workups. Silica gel 60 F₂₅₄ plates were used for TLC. Flash column chromatography was performed using silica gel (Merck, no. 9385, 230–400 mesh) and mixtures of AcOEt–*n*-hexane as eluents.

Preparation of 10 from (+)-Sclareolide (7). To a solution of (+)-sclareolide (**7**, 400 mg, 1.6 mmol) in 60 mL of toluene, at –78 °C, was added DIBAL-H (1.7 mL, 1.7 mmol, 1.0 M in toluene) dropwise. After 45 min of stirring 10 mL of H₂SO₄ aqueous solution (10%) was added, the reaction mixture was allowed to reach room temperature, and it was stirred for an additional 15 min. The mixture was extracted with CHCl₃ (3 × 30 mL), and the combined organic phases were dried and solvents removed under reduced pressure. The residue, chromatographed with hexanes–AcOEt (1:4), yielded 380 mg (94%) of pure **10** as a syrup: IR (film) ν_{max} 3314, 2925, 2854, 1460, 1379, 1333, 1205, 1153, 1131, 1074, 965, 888 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 5.42 (1H, dd, *J* = 6.1, 5.3 Hz, H-12), 2.02 (1H, ddd, *J* = 18.5, 12.30, 6.3 Hz, H_B-11), 1.93–1.88 (2H, m, H_A-11 and H-9), 1.75–1.65 (3H, m), 1.22 (3H, s), 1.47–0.96 (8H, m), 0.85 (2 × 3H, s), 0.81 (3H, s); ¹³C NMR (CDCl₃, 50.3 MHz) δ 101.8 (d, C-12), 81.2 (s, C-8), 60.2 (d, C-9), 57.0 (d, C-5), 42.4 (t, C-3), 40.0 39.8 (t, C-1), 39.7 (t, C-7), 36.1 (s, C-10), 33.5 (q, C-18), 33.1 (s, C-4), 30.8 (t, C-11), 23.5 (q, C-17), 21.0 (q, C-19), 20.8 (t, C-6), 18.4 (t, C-2), 15.2 y 15.3 (q, C-20); EIMS *m/z* 252 [M]⁺ (5), 237 [M – 15]⁺ (100), 219 (8), 201 (16), 177 (24), 137 (33), 125 (40), 109 (30), 95 (37), 81 (34), 69 (39), 43 (43); *anal.* C 76.05%, H 11.12%, calcd for C₁₆H₂₈O₂, C 76.14%, H 11.18%.

Preparation of (1S,2S,4aS,8aS)-*N*-Methoxy-*N*-methyl 1-(2-hydroxy-2,5,5,8a-tetramethyldecahydronaphthalenyl)acetamide (11) from (R)-(+)-Sclareolide (7). To a stirred suspension of *N,O*-dimethylhydroxylamine hydrochloride (1.5 g, 15.4 mmol) in CH₂Cl₂ (40 mL) at 0 °C was added Me₃Al (8 mL, 16.0 mmol, 2 M in toluene). The mixture was warmed to room temperature and stirred for 2 h until a clear solution was obtained. Sclareolide (**7**, 2.0 g, 8.0 mmol) was added in CH₂Cl₂ (20 mL) and the mixture stirred for another 2 h. After cooling to 0 °C, 15 mL of 10% aqueous H₂SO₄ was added slowly.

The reaction mixture was allowed to reach room temperature, and the two layers were separated. The aqueous layer was extracted with CH_2Cl_2 (3×30 mL), and the combined organic extracts were dried and concentrated under reduced pressure. The residue was purified over silica gel using hexanes–AcOEt (2:3) as eluent to give **11** (2.2 g, 88%) as an amorphous solid: mp 107–109 °C; $[\alpha]_{\text{D}}^{20} +26.8^\circ$ (c 0.112, CHCl_3); IR (KBr) ν_{max} 3439, 2934, 1643, 1645, 1387, 1167, 1107, 1010, 942 cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz) δ 3.71 (3H, s), 3.18 (3H, s), 2.57 (1H, dd, $J = 6.5, 16.6$ Hz, $\text{H}_{\text{B-11}}$), 1.93 (1H, dt, $J = 14.3, 2.9$ Hz, $\text{H}_{\text{A-11}}$), 1.94 (1H, s, OH), 1.72–0.92 (10H, m), 1.15 (3H, s), 0.87 (s, 3H), 0.79 (s, 3H); ^{13}C NMR (CDCl_3 , 50.3 MHz) δ 176.1 (s, C-12), 72.9 (s, C-8), 61.2 (q, OMe), 56.32 (d, C-9), 55.91 (d, C-5), 44.57 (t, C-3), 41.78 (t, C-7), 39.20 (t, C-1), 38.72 (s, C10), 33.4 (q, NMe and C-18), 33.3 (s, C-4), 26.9 (t, C-11), 23.3 (q, C-17), 21.4 (q, C-19), 20.6 (t, C-6), 18.5 (t, C-2), 15.8 (t, C-20); EIMS m/z 311 $[\text{M}]^+$ (1), 278 $[\text{M} - \text{Me} - \text{H}_2\text{O}]^+$ (2), 251 $[\text{M} - \text{N}(\text{Me})\text{OMe}]^+$ (13), 191 (18), 137 (34), 116 (100), 109 (35), 95 (24), 61 (47), 55 (27), 43 (34), 41 (28); anal. C 69.22%, H 10.77%, N 4.41%, calcd for $\text{C}_{18}\text{H}_{33}\text{NO}_3$, C 69.41%, H 10.68%, N 4.50%.

Preparation of (1S,4aS,8aS)-*N*-Methoxy-*N*-methyl-1-(5,5,8a-trimethyl-2-methylenedecahydronaphthalenyl)-acetamide (12) and (1S,4aS,8aS)-*N*-methoxy-*N*-methyl-1-(2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8-octahydronaphthalenyl)acetamide (13) from 11. To a solution of **11** (0.5 g, 1.6 mmol) in pyridine (10.0 mL) at 0 °C was added dropwise 1.2 mL (16.0 mmol) of SOCl_2 in pyridine (5 mL). The reaction mixture was stirred for 30 min, poured into an ice–water mixture (30 mL), and extracted with CH_2Cl_2 (3×40 mL). The combined organic layers were dried and concentrated under vacuum. The residue was chromatographed with hexanes–AcOEt (95:5) as eluent, yielding **12** (280 mg, 60%) and **13** (150 mg, 32%).

Compound 12: amorphous white solid, mp 90–93 °C; $[\alpha]_{\text{D}}^{20} +45.7^\circ$ (c 0.094, CHCl_3); IR (KBr) ν_{max} 2960, 2926, 1662, 1645, 1459, 1383, 1364, 1174, 1000, 904 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 4.70 (1H, s, $\text{H}_{\text{B-17}}$), 4.40 (1H, s, $\text{H}_{\text{A-17}}$), 3.69 (3H, s, OMe), 3.13 (3H, s, NMe), 2.67 (1H, dd, $J = 15.8, 9.9$ Hz, $\text{H}_{\text{B-11}}$), 2.47 (1H, broad d, $J = 10.3$ Hz, H-9 α), 2.36 (1H, dd, $J = 15.7, 3.7$ Hz, $\text{H}_{\text{A-11}}$), 2.42–2.30 (2H, m), 2.11 (1H, td, $J = 12.8, 5.5$ Hz, H-7a), 1.76–1.08 (9H, m), 0.86 (3H, s), 0.79 (3H, s), 0.71 (3H, s); ^{13}C NMR (50.3 MHz, CDCl_3) δ 174.7 (s, C-12), 149.7 (s, C-8), 105.8 (t, C-17), 61.3 (q, OMe), 55.1 (d, C-9), 51.7 (d, C-5), 42.1 (t, C-3), 39.0 (s, C-10), 38.9 (t, C-1), 37.63 (t, C-7), 33.6 (q, C-18), 33.5 (s, C-4), 32.6 (q, NMe), 27.3 (t, C-11), 24.1 (t, C-6), 21.8 (q, C-19), 19.4 (q, C-19), 19.3 (t, C-2), 14.7 (q, C-20); EIMS m/z 293 $[\text{M}]^+$ (22), 278 $[\text{M} - 15]^+$ (43), 262 $[\text{M} - \text{OMe}]^+$ (13), 233 (25), 215 (18), 205 (18), 190 (20), 175 (28), 158 (56), 149 (330), 137 (46), 123 (45), 121 (59), 109 (79), 95 (66), 81 (69), 69 (91), 61 (100), 55 (58), 41 (71); anal. C 73.79%, H 10.43%, N 4.52%, calcd for $\text{C}_{18}\text{H}_{31}\text{NO}_2$, C 73.67%, H 10.65%, N 4.77%.

Compound 13: syrup, $[\alpha]_{\text{D}}^{24} +15.2^\circ$ (c 0.250, CHCl_3); IR (KBr) ν_{max} 2924, 2847, 2862, 1776, 1668, 1444, 1383, 1173, 1100, 1007 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 5.41 (1H, br s, H-7), 3.70 (3H, s), 3.19 (3H, s), 2.80–1.00 (11H, m), 1.53 (3H, br s), 0.89 (3H, s), 0.87 (3H, s), 0.80 (3H, s); ^{13}C NMR (75.0 MHz, CDCl_3) δ 175.4 (s, C-12), 134.1 (s, C-8), 122.1 (d, C-7), 61.1 (q, OMe), 49.7 (d, C-9), 48.9 (d, C-5), 42.0 (t, C-3), 38.8 (t, C-1), 35.8 (s, C-10), 33.0 (q, C-18), 32.8 (s, C-4), 32.7 (q, NMe), 29.1 (t, C-11), 23.6 (t, C-6), 21.7 (q, C-17), 21.3 (q, C-19), 18.7 (t, C-2), 14.2 (q, C-20); EIMS m/z 293 $[\text{M}]^+$ (56), 278 (25), 262 (15), 233 (20), 215 (38), 190 (53), 175 (27), 137 (33), 119 (88), 109 (94), 103 (100), 95 (47), 81 (55), 69 (44), 61 (53), 41 (30); anal. C 73.89%, H 10.56%, N 4.89%, calcd for $\text{C}_{18}\text{H}_{31}\text{NO}_2$, C 73.67%, H 10.65%, N 4.77%.

Preparation of (1S,4aS,8aS)-*N*-Methoxy-*N*-methyl-1-(2,5,5,8a-tetramethyl-1,4,4a,5,6,7,8-octahydronaphthalenyl)acetamide (13) from 11. A mixture of **12** (250 mg, 0.85 mmol) and *p*-TsOH (25 mg) in toluene (25 mL) was heated at 50 °C for 2 h. The reaction mixture was cooled at room temperature and diluted with AcOEt (30 mL). The mixture was washed with NaHCO_3 (saturated solution, 3×50 mL). The organic layer was dried, and the solvents were evaporated.

Chromatography of the residue using hexanes–AcOEt (95:5) yielded 223 mg (89%) of pure **13**.

Preparation of 15,16-Epoxy-12-oxolabda-7,13(16),14-triene (14) from 13. 3-Bromofuran (0.23 mL, 2.5 mmol) was added dropwise to a solution of *n*-butyllithium (1.2 M in hexanes, 2.5 mL, 3.0 mmol) in dry THF (2.5 mL), at –78 °C under argon. The mixture was stirred for 1 h and then transferred via cannula into a solution of amide **13** (0.3 g, 1.0 mmol) in dry THF (40 mL) cooled at –78 °C. After 2 h the substrate was consumed and the reaction was quenched by adding 50 mL of NH_4Cl (saturated solution). The aqueous phase was extracted with AcOEt (3×50 mL), the combined organic layers were dried, and the solvent was evaporated. The residue was chromatographed using hexanes– CH_2Cl_2 (4:1) as eluent. Thus, 220 mg of pure ketone **14** (72%) was obtained: spontaneous white crystals mp 70–73 °C; $[\alpha]_{\text{D}}^{27} +42.9^\circ$ (c 0.219, CHCl_3); IR (KBr) ν_{max} 2963, 2926, 2847, 1668, 1560, 1512, 1385, 1158, 873 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 8.04 (1H, d, $J = 0.9$ Hz, H-16), 7.43 (1H, t, $J = 1.8$ Hz, H-15), 6.78 (1H, dd, $J = 1.8, 0.9$ Hz, H-14), 5.41 (1H, br s, H-7), 2.80 (1H, m), 2.77 (1H, d $J = 16.5$ Hz, $\text{H}_{\text{B-11}}$), 2.67 (1H, dd, $J = 16.5, 14.3$ Hz, $\text{H}_{\text{A-11}}$), 2.04 (1H, m), 1.86 (1H, m), 1.64 (1H, br d, $J = 12.6$ Hz), 1.45 (3H, br s, Me-17), 1.50–1.00 (7H, m), 0.88 (3H, s), 0.87 (3H, s), 0.81 (3H, s); ^{13}C NMR (50.0 MHz, CDCl_3) δ 194.9 (s, C-12), 145.6 (d, C-15), 144.0 (d, C-16), 133.8 (s, C-8), 127.8 (s, C-13), 122.4 (d, C-7), 108.7 (d, C-14), 45.6 (d, C-5), 48.2 (d, C-9), 42.0 (t, C-3), 39.1 (t, C-1), 38.3 (t, C-11), 35.7 (s, C-10), 33.0 (q, C-18), 32.8 (s, C-4), 23.6 (t, C-6), 21.9 (q, C-17)*, 21.7 (q, C-19)*, 18.7 (t, C-2), 14.3 (q, C-20), assignments marked with an asterisk may be interchanged; EIMS m/z 300 $[\text{M}]^+$ (12), 285 (2), 215 (2), 190 (100), 175 (39), 119 (35), 109 (61), 105 (30), 95 (99); anal. C 79.56%, H 9.52%, calcd for $\text{C}_{20}\text{H}_{28}\text{O}_2$, C 79.96%, H 9.39%.

Preparation of (12R)- and (12S)-15,16-Epoxy-12-hydroxylabda-7,13(16),14-triene (15 and 16) from 14. To a solution of ketone **14** (820 mg, 2.7 mmol) in MeOH (53 mL) at room temperature was added 2.0 g of $\text{CeCl}_3 \times 7\text{H}_2\text{O}$ (5.5 mmol), and the mixture was stirred for 30 min. The resulting slurry was cooled until 0 °C, and NaBH_4 (625 mg, 16.4 mmol) was added in portions. The mixture was allowed to reach room temperature and stirred for 24 h. After this time, H_2O (50 mL) was added and the mixture stirred for an additional 1 h. The resulting solution was concentrated under vacuum, and the aqueous phase was extracted with AcOEt (3×50 mL). The combined organic layers were dried and evaporated under reduced pressure. Chromatography of the residue using hexanes–AcOEt (1:1) yielded in increasing order of polarity unreacted ketone **14** (150 mg), alcohol **15** (250 mg, 37%), and alcohol **16** (220 mg, 33%).

Compound 15: oil; $[\alpha]_{\text{D}}^{21} +25.6^\circ$ (c 0.340, CHCl_3); IR (film) ν_{max} 3400, 2922, 2847, 1502, 1455, 1387, 1159, 1024, 874 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.38 (2H, m, H-16 and H-15), 6.41 (H, dd, $J = 1.5, 0.8$ Hz, H-14), 5.43 (1H, br s, H-7), 4.72 (1H, dd, $J = 8.4, 1.6$ Hz, H-12), 2.20–0.90 (12 H, m), 1.66 (3H, br s), 0.89 (3H, s), 0.87 (3H, s), 0.76 (3H, s); ^{13}C NMR (50.3 MHz, CDCl_3) δ 143.3 (d, C-15), 138.5 (d, C-16), 134.6 (s, C-8), 130.1 (s, C-13), 123.0 (d, C-7), 108.5 (d, C-14), 67.6 (d, C-12), 50.5 (d, C-9)*, 50.4 (d, C-5)*, 42.3 (t, C-3), 39.4 (t, C-1), 36.3 (t, C-11), 36.3 (s, C-10)*, 33.1 (q, C-18), 33.0 (s, C-4)*, 23.9 (t, C-6), 22.3 (q, C-17)*, 21.8 (q, C-19)*, 18.8 (t, C-2), 13.6 (q, C-20), assignments marked with an asterisk may be interchanged; EIMS m/z 302 $[\text{M}]^+$ (absent), 284 $[\text{M} - 18]^+$ (11), 205 (32), 190 (40), 175 (29), 160 (82), 119 (31), 109 (72), 97 (100), 82 (54), 69 (48), 41 (54); anal. C 79.60%, H 9.59%, calcd for $\text{C}_{20}\text{H}_{30}\text{O}_2$, C 79.42%, H 10.00%.

Compound 16: oil; $[\alpha]_{\text{D}}^{21} -18.6^\circ$ (c 0.200, CHCl_3); IR (film) ν_{max} 3367, 2922, 2847, 1503, 1456, 1387, 1155, 1022, 874, 790 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.38 (1H, m, H-16), 7.36 (1H, m, H-15), 6.44 (1H, dd, $J = 1.6, 1.0$ Hz, H-14), 5.42 (1H, br s, H-7), 4.75 (1H, dd, $J = 9.4, 5.4$ Hz, H-12), 2.20–0.40 (12H, m), 1.76 (3H, br s, Me-17), 0.84 (3H, s), 0.81 (3H, s), 0.74 (3H, s); ^{13}C NMR (50.3 MHz, CDCl_3) δ 143.4 (d, C-15), 139.7 (d, C-16), 134.6 (s, C-8), 128.5 (s, C-13), 122.9 (d, C-7), 108.7 (d, C-14), 68.0 (d, C-12), 50.4 (d, C-5)*, 50.0 (d, C-9)*, 42.2 (t, C-3), 39.1 (t, C-1), 36.7 (s, C-10), 35.5 (t, C-11), 33.1 (q, C-18), 32.9

(s, C-4), 23.9 (t, C-6), 22.7 (q, C-17), 21.8 (q, C-19), 18.7 (t, C-2), 13.6 (q, C-20), assignments marked with an asterisk may be interchanged; EIMS m/z 302 [M]⁺ (absent), 284 [M - 18]⁺ (9), 205 (28), 190 (25), 160 (78), 119 (24), 109 (61), 97 (100), 82 (47), 69 (44), 41 (52); *anal.* C 79.67%, H 9.68%, calcd for C₂₀H₃₀O₂, C 79.42%, H 10.00%.

Preparation of Mosher's Ester 17 from Alcohol 15. Alcohol **15** (8 mg, 0.025 mmol) in CH₂Cl₂ (1.0 mL) and a catalytic amount of 4-(dimethylamino)pyridine were added to a solution of (*R*)-(+)- α -methoxy- α -(trifluoromethyl)phenylacetic acid (16.0 mg, 0.07 mmol) and 1,3-dicyclohexylcarbodiimide (14.0 mg, 0.07 mmol) in CH₂Cl₂ (0.5 mL) at room temperature. The mixture was stirred for 2 h until complete consumption of alcohol. Then, the reaction mixture was filtered through a pad of Celite and evaporated under vacuum. Chromatography of the residue with hexanes-AcOEt (99:1) yielded 11 mg of pure **17** (89%): ¹H NMR (300 MHz, CDCl₃) δ 7.50–7.31 (7H, m, Ph, H-15, H-16), 6.44 (1H, m, H-14), 6.11 (1H, dd J = 11.3, 1.7 Hz, H-12), 5.39 (1H, br s, H-7), 3.44 (3H, br s, OMe), 3.19 (1H, m), 2.07 (1H, dd J = 13.9, 11.5 Hz, H_B-11), 1.70 (3H, br s, Me-17), 2.11–1.13 (11 H, m, decalin protons), 0.81 (3H, s), 0.80 (3H, s), 0.64 (3H, s).

Preparation of Mosher's Ester 18 from Alcohol 16. Alcohol **16** (13 mg, 0.043 mmol) in CH₂Cl₂ (2.0 mL) and a catalytic amount of 4-(dimethylamino)pyridine were added to a solution of (*R*)-(+)- α -methoxy- α -(trifluoromethyl)phenylacetic acid (26.0 mg, 0.13 mmol) and 1,3-dicyclohexylcarbodiimide (22.0 mg, 0.13 mmol), in CH₂Cl₂ (1 mL) at room temperature. Working as above, 21 mg (93%) of pure **18** was obtained: ¹H NMR (300 MHz, CDCl₃) δ 7.63–7.27 (7H, m, Ph, H-15, H-16), 6.28 (1H, m, H-14), 6.06 (1H, dd J = 10.0, 5.7 Hz, H-12), 5.44 (1H, m, H-7), 3.52 (3H, br s, OMe), 2.03–1.05 (12H, m), 1.77 (3H, br s, Me-17), 0.85 (3H, s), 0.82 (3H, s), 0.74 (3H, s).

General Procedure for Irradiation of 15 and 16. A solution of alcohol (100 mg) in THF (10 mL), containing ca. 1 mg of Rose Bengal, was irradiated with a 220 W tungsten lamp for 5 h. The solvent was removed under reduced pressure, and the residue was submitted to column chromatography, yielding pure compounds.

Preparation of (12*R*,15*ξ*)-12,15-dihydroxyabda-7,14-dien-16,15-olide (3) and (12*R*,16*ξ*)-12,16-dihydroxyabda-7,14-dien-15,16-olide (4). Photooxidation of **15**. Irradiation of alcohol **15** yielded, after chromatographic separation with hexanes-AcOEt (3:1), 18 mg (16%) of lactone **3** and 12 mg (11%) of lactone **4**. Compounds **3** and **4** were unstable, and correct analytical data could not be obtained.

Compound 3: ¹H NMR (300 MHz, CDCl₃) δ 7.04 (1H, m, H-14), 6.12 (1H, m, H-15), 5.42 (1H, m, H-7), 4.56 (1H, m, H-12), 1.73 (3H, s, Me-17), 0.87 (3H, s), 0.86 (3H, s), 0.73 (3H, s). Due to the instability of **3** IR and MS data could not be obtained.

Compound 4: IR (KBr) ν_{\max} 3434, 2925, 1752, 1631, 1009 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 6.24 (1H, s, H-16), 6.12 (1H, br s, H-14), 5.44 (1H, br s, H-7), 4.67 and 4.55 (1H, m, H-12), 1.73 and 1.69 (3H, br s, Me-17), 2.08–1.01 (12H, m), 0.87 (3H, s, Me-18), 0.86 (3H, s, Me-19), 0.74 and 0.73 (3H, s, Me-20); EIMS m/z 334 [M]⁺ (absent), 205 (14), 189 (10), 175 (7), 124 (36), 109 (100), 91 (22), 81 (50), 69 (26), 55 (32), 41 (27).

Preparation of (12*S*,15*ξ*)-12,15-dihydroxyabda-7,14-dien-16,15-olide (5) and (12*S*,16*ξ*)-12,16-dihydroxyabda-7,14-dien-15,16-olide (6). Photooxidation of **16**. Irradiation of alcohol **16** yielded, after chromatography with hexanes-AcOEt (3:1), 22 mg (20%) of lactone **5** and 15 mg (10%) of lactone **6**. Compounds **5** and **6** were unstable, and correct analytical data could not be obtained.

Compound 5: IR (KBr) ν_{\max} 3455, 2924, 2847, 1749, 1634, 1129 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.08 and 7.05 (1H, s, H-14), 6.15 (1H, br s, H-15), 5.45 (1H, br s, H-7), 4.55 (1H, t, J = 6.7 Hz, H-12), 1.95–1.12 (12H, m), 1.70 (3H, br s, Me-17), 0.87 (3H, s, Me-18), 0.84 (3H, s, H-19), 0.76 (3H, s, Me-20); EIMS m/z 334 [M]⁺ (absent), 316 [M - 18]⁺ (1), 298 [M - 36]⁺ (1), 205 (50), 149 (17), 124 (25), 109 (100), 81 (59), 55 (27), 41 (21).

Compound 6: IR (KBr) ν_{\max} 3435, 2920, 1754, 1631, 112 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 6.23 (1H, m, H-16), 6.07 (1H, m, H-14), 5.46 (1H, br s, H-7), 4.68 (1H, t, J = 6.2 Hz, H-12), 1.73 (3H, br s, Me-17), 2.05–1.11 (12H, m), 0.88 (3H, s, Me-18), 0.86 (3H, s, H-19), 0.79 (3H, s, Me-20); EIMS m/z 334 [M]⁺ (absent), 205 (10), 190 (80), 175 (7), 124 (40), 109 (100), 91 (18), 81 (46), 69 (20), 55 (27), 41 (21).

Preparation of (12*R*)-12-Acetoxy-15,16-epoxyabda-7,13(16),14-triene (19) from 15. Alcohol **15** (50 mg), was treated with Ac₂O-Pyr (12.0 mL, 1:2) for 24 h. Solvents were removed under reduced pressure, and the residue was filtered through a short pad of silica gel using AcOEt as solvent, yielding 55 mg (99%) of pure **19**: amorphous solid; mp 64–67 °C; [α]_D²⁵ + 15.7° (*c* 0.070, CHCl₃); IR (KBr) ν_{\max} 2923, 2847, 1725, 1361, 1240, 1020 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.40 (1H, d, J = 0.6 Hz, H-16), 7.36 (1H, t, J = 1.8 Hz, H-15), 6.38 (1H, dd, J = 1.8, 0.6 Hz, H-14), 5.93 (1H, dd, J = 10.9, 2.7 Hz, H-12), 5.41 (1H, br s, H-7), 2.06 (3H, s, OAc), 1.70 (3H, br s, Me-17), 0.88 (3H, Me-18), 0.86 (3H, s, Me-19), 0.76 (3H, s, Me-20); ¹³C NMR (50.3 MHz, CDCl₃) δ 170.7 (s, OAc), 143.2 (d, C15), 139.8 (d, C-16), 134.2 (s, C-8), 125.7 (s, C-13), 123.1 (d, C-7), 108.8 (d, C-14), 69.0 (d, C-12), 50.4 (d, C-5)*, 49.9 (d, C-9)*, 42.3 (t, C-3), 39.3 (t, C-1), 36.3 (s, C-10), 33.2 (q, C-18), 33.1 (t, C-11), 33.0 (s, C-4), 23.2 (q, C-17)*, 21.9 (q, C-19)*, 21.3 (q, OAc)*, 18.8 (t, C-2), 13.6 (q, C-20); EIMS m/z 344 [M]⁺ (absent), 284 [M - 60]⁺ (11), 190 (38), 175 (16), 160 (100), 145 (14), 119 (40), 109 (39), 97 (36), 81 (31), 43 (56); *anal.* C 76.53%, H 9.52%, calcd for C₂₂H₃₂O₃, C 76.70%, H 9.36%.

Preparation of (12*S*)-12-Acetoxy-15,16-epoxyabda-7,13(16),14-triene (20) from 16. Alcohol **16** (150 mg) was treated with Ac₂O-Pyr (18.0 mL, 1:2) for 24 h at room temperature. Solvents were removed under reduced pressure. The residue, filtered through a short pad of silica gel using AcOEt solvent, yielded 160 mg (98%) of pure **20** as a syrup: [α]_D²² -32.8° (*c* 0.090, CHCl₃); IR (film) ν_{\max} 2925, 1739, 1369, 1239, 1022 cm⁻¹; ¹H NMR (200 MHz, CDCl₃) δ 7.42 (1H, m, H-16), 7.39 (1H, t, J = 1.5 Hz, H-15), 6.41 (1H, dd, J = 1.7, 0.7 Hz, H-14), 5.87 (1H, dd, J = 8.8, 6.59 Hz, H-12), 5.42 (1H, br s, H-7), 2.02 (3H, OAc), 1.90–1.40 (12H, m), 1.80 (3H, s, Me-17), 0.84 (3H, Me-18), 0.80 (3H, s, Me-19), 0.73 (3H, s, Me-20); ¹³C NMR (50.3 MHz, CDCl₃) δ 170.3 (s, OAc), 143.2 (d, C-15), 140.9 (d, C-16), 134.3 (s, C-8), 124.5 (s, C-13), 123.0 (d, C-7), 109.2 (d, C-14), 69.9 (d, C-12), 50.2 (d, C-5), 49.9 (d, C-9), 42.1 (t, C-3), 39.2 (t, C-1), 36.8 (s, C-10), 33.0 (q, C-18), 33.0 (s, C-4), 32.3 (t, C-11), 23.8 (t, C-6), 22.7 (q, C-17), 21.8 (q, C-19), 21.3 (q, OAc), 18.7 (t, C-2), 13.6 (q, C-20); EIMS m/z 344 [M]⁺ (absent), 284 [M - 60]⁺ (11), 269 (3), 190 (26), 160 (100), 140 (25), 119 (36), 109 (48), 105 (23), 97 (50), 81 (38), 43 (74); *anal.* C 76.48%, H 9.60%, calcd for C₂₂H₃₂O₃, C 76.70%, H 9.36%.

General Procedure for Irradiation of 19 and 20. A THF solution of the corresponding acetate, containing a catalytic amount of Rose Bengal, was irradiated under direct sunlight until TLC analysis showed complete transformation of the starting material. The solvent was removed under reduced pressure, and the residue was submitted to column chromatography to yield analytically pure compounds.

Preparation of (12*R*)-12-Acetoxy-15-hydroxyabda-7,14-dien-16,15-olide (21) and (12*R*)-12-Acetoxy-16-hydroxyabda-7,14-dien-15,16-olide (22) from 19. Irradiation of **19** (75 mg) in THF (15 mL) yielded, after chromatographic separation with hexanes-AcOEt (4:1), 29 mg of the hydroxybutenolide **21** (35%) and 23 mg (33%) of its isomer **22**.

Compound 21: syrup; IR (film) ν_{\max} 3429, 2925, 2847, 1747, 1456, 1373, 1230, 1131, 1042 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 6.94 (1H, m, H-14), 6.09 (1H, m, H-15), 5.70 (1H, dd, J = 6.7, 3.8 Hz, H-12), 5.42 (1H, br s, H-7), 2.13 (3H, s, OAc), 1.75 (3H, br s, Me-17), 2.00–1.15 (12H, m), 0.87 (3H, s, Me-19), 0.72 (3H, s, Me-20); EIMS m/z 376 [M]⁺ (absent), 316 [M - 60]⁺ (23), 301 (5), 205 (9), 190 (30), 147 (28), 119 (54), 109 (100), 105 (26), 91 (28), 81 (36), 69 (27), 55 (27), 43 (52); *anal.* C 69.84%, H 8.35%, calcd for C₂₂H₃₂O₅ C 70.18%, H 8.57%.

Compound 22: syrup; IR (film) ν_{\max} 3430, 2924, 2827, 1749, 1373, 1235, 1011 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 6.20 6.03 (1H, m, H-16), 5.97 (1H, m, H-14), 5.56 (1H, br d, J = 10.6

Hz, H-12), 5.46 (1H, br s, H-7), 2.14 (3H, s, OAc), 1.71 (3H, br s, Me-17), 2.00–1.20 (12H, m), 0.88 (3H, s, Me-18), 0.86 (3H, s, Me-19), 0.75 (3H, s, H-20); EIMS m/z 376 [M]⁺ (absent), 298 [M – 60 – 18]⁺ (1), 205 (5), 190 (43), 175 (14), 124 (39), 119 (28), 109 (100), 81 (27), 69 (21), 43 (36); anal C 69.93%, H 8.38%, calcd for C₂₂H₃₂O₅ C 70.18%, H 8.57%.

Preparation of (12*S*)-12-Acetoxy-15-hydroxyabda-7,14-dien-16,15-olide (23) and (12*S*)-12-Acetoxy-16-hydroxyabda-7,14-dien-15,16-olide (24) from 20. Irradiation of 20 (120 mg), in THF (25 mL), yielded, upon chromatography with hexanes–AcOEt (4:1), 55 mg of pure hydroxybutenolide 23 (42%) and 30 mg (23%) of its isomer 24.

Compound 23: syrup; IR (film) ν_{\max} 3434, 2926, 2847, 1768, 1749, 1635, 1458, 1370, 1236, 1014 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.01, 7.00 (1H, br s, H-14), 6.12, 6.09 (1H, br s, H-15), 5.57 (1H, t, $J = 6.5$ Hz, H-12), 5.43 (1H, br s, H-7), 2.10, 2.01 (3H, s, OAc), 1.73 (3H, br s, Me-17), 2.06–0.89 (12H, m), 0.85 (3H, s, Me-18), 0.82 (3H, s, Me-19), 0.75 (3H, s, Me-20); EIMS m/z 376 [M]⁺ (absent), 316 [M – 60]⁺ (25), 301 (6), 215 (5), 190 (27), 133 (14), 119 (48), 109 (100), 105 (26), 91 (29), 81 (36), 69 (26), 55 (27), 43 (57); anal C 69.90%, H 8.46%, calcd for C₂₂H₃₂O₅ C 70.18%, H 8.57%.

Compound 24: syrup; IR (film) ν_{\max} 3430, 2926, 2862, 2847, 1747, 1638, 1458, 1367, 1232 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 6.24, 6.05 (1H, br s, H-16), 6.02 (1H, br s, H-14), 5.47 (1H, br s, H-7), 5.36 (1H, t, $J = 7.7$ Hz, H-12), 2.12, 2.09 (3H, s, OAc), 2.00–2.70 (12H, m), 1.71 (3H, s, Me-17), 0.86 (3H, s, Me-18), 0.84 (3H, s, Me-19), 0.76 (3H, s, Me-20); anal C 70.25%, H 8.50%, calcd for C₂₂H₃₂O₅ C 70.18%, H 8.57%.

Preparation of (12*R*)-12,15-Diacetoxyabda-7,14-dien-16,15-olide (25), (12*R*)-12,16-Diacetoxyabda-7,14-dien-15,16-olide (26), (12*S*)-12,15-Diacetoxyabda-7,14-dien-16,15-olide (27), and (12*S*)-12,16-Diacetoxyabda-7,14-dien-15,16-olide (28) from 21 to 24. Compounds 21–24 were treated with the mixture Ac₂O–Pyr (2 mL, 1:2) for 24 h at room temperature. Solvents were removed under reduced pressure, and the residues were purified by chromatography, using hexanes–AcOEt (9:1).

Compound 25. Following the general procedure, 24 mg (0.064 mmol) of 21 yielded 23 mg (86%) of an epimeric mixture of 25 as a syrup; $[\alpha]_D^{25} +30.9^\circ$ (c 0.330 CHCl₃); IR (film) ν_{\max} 2925, 2862, 1780, 1743, 1456, 1373, 1226, 1022, 757 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 6.89 (1H, m, H-15 minor), 6.87 (1H, t, $J = 1.6$ Hz, H-14 minor), 6.97 (2H, m, H-14 and H-15 major), 5.76 (1H, tdd, $J = 7.8, 1.6$ Hz, H-12 minor and major), 5.42 (1H, br s, H-7), 2.14 (6H, 2 × OAc major), 2.13 (3H, s, OAc, minor), 2.15 (3H, s, OAc, minor), 1.80 (3H, s, Me-17), 0.86 (6H, 2 × Me, minor), 0.85 (6H, 2 × Me, major), 0.73 (3H, Me-20, major), 0.72 (3H, Me-20 minor); EIMS m/z 418 [M⁺] (1), 358 [M⁺ – 60] (19), 298 (8), 205 (10), 190 (36), 174 (43), 119 (53), 109 (88), 105 (24), 91 (26), 81 (36), 69 (31), 43 (100); anal C 68.59%, H 7.95%, calcd for C₂₄H₃₄O₆, C 68.87%, H 8.19%.

Compound 26. Following the general procedure, 19 mg (0.051 mmol) of 22 yielded 18 mg (84%) of an epimeric mixture of 26 as a syrup; $[\alpha]_D^{24} +14.7^\circ$ (c 0.660, CHCl₃); IR (film) ν_{\max} 2962, 2920, 2847, 1800, 1747, 1653, 1455, 1373, 1261, 1226, 1026, 866 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.02 (1H, d, $J = 1.0$ Hz, H-16 minor), 6.94 (1H, d, $J = 0.9$ Hz, H-16 major), 6.12 (1H, dd, $J = 1.6, 1.1$ Hz, H-14 minor), 6.04 (1H, t, $J = 1.1$ Hz, H-14 major), 5.81 (1H, dt, $J = 11.7, 1.7$ Hz, H-12 major), 5.62 (1H, br d, $J = 13.5$ Hz, H-12 minor), 5.45 (1H, br s, H-7), 2.17 and 2.16 (3H each, s, 2 × OAc major), 2.14 and 2.10 (3H each, s, 2 × OAc minor), 1.55 (3H, s, Me-17), 2.03–1.10 (12H, m), 0.87 (3H, s, Me minor), 0.85 (3H, s, Me major), 0.85 (3H, s, Me), 0.75 (3H, s, Me minor), 0.73 (3H, s, Me major); EIMS m/z 298 [M⁺ – 60 – 60] (3), 283 (2), 255 (1), 189 (24), 175 (12), 124 (30), 119 (35), 109 (100), 81 (30), 69 (26), 55 (25), 43 (79); anal C 69.03%, H 7.88%, calcd for C₂₄H₃₄O₆, C 68.87%, H 8.19%.

Compound 27. Following the general procedure, 50 mg (0.133 mmol) of 23 yielded 51 mg (92%) of an epimeric mixture of 27 as a syrup; $[\alpha]_D^{23} -29.1^\circ$ (c 0.500, CHCl₃); IR (film) ν_{\max} 2925, 2847, 2862, 1780, 1746, 1458, 1372, 1335, 1227, 1022, 966, 797, 757 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.05 (1H, m, H-15 or H-14), 6.92 and 6.89 (1H, d, $J = 1.1$ Hz, H-14 or H-15),

5.64 (1H, t, $J = 7.1$ Hz, H-12 major), 5.62 (1H, t, $J = 7.3$ Hz, H-12 minor), 5.43 (1H, br s, H-7), 2.15, 2.14, 2.11 and 2.10 (3H each, s, OAc), 1.74 (3H, br s, Me-17), 2.03–1.13 (12H, m), 0.85 (3H, s, Me), 0.83 (3H, s, Me), 0.75 (3H, s, Me); EIMS m/z 358 [M⁺ – 60] (25), 316 (8), 298 (10), 205 (13), 190 (40), 174 (48), 119 (49), 109 (90), 91 (27), 81 (34), 69 (27), 55 (23), 43 (100); anal C 68.63%, H 8.22%, calcd for C₂₄H₃₄O₆, C 68.87%, H 8.19%.

Compound 28. Following the general procedure, 25 mg (0.06 mmol) of 24 yielded 20 mg (80%) of an epimeric mixture of 28 as a syrup; $[\alpha]_D^{24} -5.8^\circ$ (c 2.170, CHCl₃); IR (film) ν_{\max} 2962, 2920, 2850, 1797, 1744, 1455, 1372, 1260, 1024, 865, 799 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.04 (1H, m, H-16 minor), 6.97 (1H, d, $J = 1.0$ Hz, H-16 major), 6.16 and 6.07 (1H each, t, $J = 1.1$ Hz, H-14), 5.68 (1H, td, $J = 8.1, 1.20$ Hz, H-12 major), 5.46 (2H, br s, and td H-7 and H12 minor), 2.18 and 2.12 (3H each, s, 2 × OAc, major), 2.15 and 2.07 (3H, each, s, 2 × OAc minor), 2.02 (12H, m), 1.69 (3H, me-17), 0.86 (3H, s, Me), 0.84 (3H, s, Me), 0.77 and 0.75 (3H, s, Me-20); IEMS m/z 358 [M⁺ – 60] (1), 343 (1), 298 (7), 283 (5), 189 (43), 175 (17), 124 (34), 119 (34), 109 (100), 91 (17), 81 (26), 69 (19), 55 (17), 43 (53); anal C 69.12%, H 7.84%, calcd for C₂₄H₃₄O₆, C 68.87%, H 8.19%.

Acknowledgment. Financial support by the Spanish Comision Interministerial de Ciencia y Tecnologia (M.C.T., CICYT, Grant No. AGF98-0805), the Dirección General de Enseñanza Superior e Investigación Científica y Técnica (MEC-Spain), and the European Commission (M. A. Sierra, grants PB97-0323 and 2FD97-0314-CO2-02) is acknowledged. I.G. thanks the MEC (Spain) for a predoctoral fellowship. We wish to thank Prof. Mar Gómez-Gallego (UCM) for her careful editing of the manuscript and her valuable suggestions.

References and Notes

- (a) Mann, J.; Davison, R. S.; Hobbs, J. B.; Banthorpe, D. V.; Harborne, J. B. *Natural Products: Their Chemistry and Biological Significance*, Longman Group Ltd.: New York, 1994. (b) Hanson, J. R. *Nat. Prod. Rep.* **2001**, *18*, 88–94.
- Miyazawa, M.; Shimamura, H.; Nakamura, S.; Kameoka, H. *J. Agric. Food Chem.* **1995**, *43*, 3012–3015.
- Singh, M.; Pal, M.; Sharma, R. P. *Planta Med.* **1999**, *65*, 2–8.
- Zani, C. L.; Alves, T. M. A.; Queiróz, R.; Fontes, E. S.; Shin, Y. G.; Cordell, G. A. *Phytochemistry* **2000**, *53*, 877–880.
- (a) Bohlmann, F.; Zdero, C.; Gupta, R. K.; King, R. M.; Robinson, H. *Phytochemistry* **1980**, *19*, 2695–2705. (b) Bohlmann, F.; Ludwig, G.-W.; Japukov, J.; King, R. B.; Robinson, H. *Phytochemistry* **1983**, *22*, 983–986.
- (a) Rodriguez, B.; Rodriguez, B.; de la Torre, M. C.; Simmonds, M. S. J. And Blaney, W. M. *J. Nat. Prod.* **1999**, *62*, 594–600. (b) de la Torre, M. C.; Maggio, A.; Rodriguez, B. *Tetrahedron* **2000**, *53*, 8007–8017. (c) Malakov, P. Y.; Papanov, G. Y.; Rodriguez, B.; de la Torre, M. C.; Simmonds, M. S. J.; Blaney, W. M.; Boneva, I. M. *Phytochemistry* **1994**, *37*, 147–157.
- Bock, I.; Bornowski, H.; Rauff, A.; Theis, H. *Tetrahedron* **1990**, *46*, 1199–1210.
- (a) Nahm, S.; Weinreb, S. M. *Tetrahedron Lett.* **1981**, *22*, 3815–3818. (b) Shimizu, T.; Osako, K.; Nakata, T. *Tetrahedron Lett.* **1997**, *38*, 2685–2688. (c) Lucet, D.; Le Gall, T.; Moskowsky, C.; Pólux, O.; Marquet, A. *Tetrahedron Asym.* **1996**, *7*, 985–988.
- Luche, J.-L.; Gemal, A. L. *J. Am. Chem. Soc.* **1981**, *103*, 5454–5459.
- (a) Ohtani, I.; Kusumi, T.; Kasham, Y.; Kakisawa, H. *J. Am. Chem. Soc.* **1991**, *113*, 4092–4096, and references therein. (b) Kusumi, T.; Fukushima, T.; Otani, I.; Kakisawa, H. *Tetrahedron Lett.* **1991**, *32*, 2939–2942.
- Ohtani, I.; Hotta, K.; Ichikawa, Y.; Isobe, M. *Chem. Lett.* **1995**, 513–514.
- (a) Scheffold, R.; Dubs, P. *Helv. Chim. Acta* **1967**, *50*, 798–808. (b) Inoffen, H. H.; Krieser, W.; Nazir, M. *Liebigs Ann. Chem.* **1972**, *755*, 1–11. (c) Kreiser, W.; Nazir, M. *Liebigs Ann. Chem.* **1972**, *755*, 12–16. (d) Sum, F. W.; Weiler, L. *J. Org. Chem.* **1979**, *44*, 1012–1013. (e) Bourguignon, J. J.; Wermuth, C. G. *J. Org. Chem.* **1981**, *46*, 4889–4894. (f) Laugrand, S.; Guingant, A.; d'Angele, J. *J. Org. Chem.* **1987**, *52*, 4788–4790. (g) Ohta, T.; Tsuchiyama, H.; Nazoe, S. *Heterocycles* **1986**, *24*, 1137. (h) Cooper, G. K.; Dolby, L. J. *J. Org. Chem.* **1979**, *44*, 3414–3416. (i) Larcheveque, M.; Lequent, Ch.; Debal, A.; Lallemand, J. Y. *Tetrahedron Lett.* **1981**, *22*, 1595–1598.
- (a) Feringa, B. L. *Recl. Trav. Chim. Pays-Bas* **1987**, *106*, 469. (b) Wasserman, H. R.; Ives, J. L. *Tetrahedron* **1981**, *37*, 1825–1852.
- Different approaches have been developed for improving selectivity of the photooxidation reaction. See for instance: (a) Kuwajima, I.; Urabe, H. *Tetrahedron Lett.* **1981**, *22*, 5191–5194. (b) Katsumura, S.; Hori, K.; Fujimara, S.; Isoe, S. *Tetrahedron Lett.* **1985**, *26*, 4625–

4628. (c) Kernan, M. R.; Faulkner, D. J. *J. Org. Chem.* **1988**, *53*, 2772–2776. (d) Lee, G. C. M.; Syage, E. T.; Harcourt, D. A.; Holmes, J. M.; Garst, M. E. *J. Org. Chem.* **1991**, *56*, 7007–7014. (e) Hagiwara, H.; Inome, K.; Uda, K. *J. Chem. Soc., Perkin Trans. 1* **1995**, 757–764.
- (15) (a) Siddiqui, S.; Faizi, S.; Mahmood, T.; Siddiqui, B. S. *J. Chem. Soc., Perkin Trans. 1* **1986**, 1021–1025. (b) Siddiqui, B. S.; Ghiasuddin; Faiza, S.; Rasheed, M. *J. Nat. Prod.* **1999**, *62*, 1006–1009. (c) Rodriguez, B.; de la Torre, M. C.; Bruno, M.; Piozzi, F.; Vassallo, N.; Cirimina, R.; Servettaz, O. *Phytochemistry* **1997**, *45*, 383–385. (d) Siddiqui, Faizi, S.; Mahmood, T.; Siddiqui, B. S. *Tetrahedron* **1986**, *42*, 4849–4856.

NP010590U