

Two Unprecedented Dibromotyrosine-Derived Alkaloids from the Brazilian Endemic Marine Sponge *Aplysina caissara*

Beatriz M. Saeki,[†] Ana Claudia Granato,[†] Roberto G. S. Berlinck,^{*,†} Alviclér Magalhães,[‡] Alexandre B. Schefer,[‡] Antonio G. Ferreira,[‡] Ulisses S. Pinheiro,[§] and Eduardo Hajdu[§]

Instituto de Química de São Carlos, Universidade de São Paulo CP 780, CEP 13560-970, São Carlos, SP, Brazil, Departamento de Química, Universidade Federal de São Carlos CP 676, CEP 13565-905, São Carlos, SP, Brazil, and Museu Nacional, Departamento de Invertebrados, Universidade Federal do Rio de Janeiro, Quinta da Boa Vista s/n, CEP 20940-040, Rio de Janeiro, RJ, Brazil

Received November 16, 2001

Two new bromotyrosine-derived alkaloids, caissarine A (**1**) and caissarine B (**2**), along with three known biogenetically related alkaloids, aeroplysinin-1, fistularin-3, and the artifact of isolation 2-(3,5-dibromo-4-dimethoxy-1-hydroxy-2,5-cyclohexadien-1-yl)ethanamide, have been isolated from *Aplysina caissara*, an endemic species of marine sponge from the Southeastern Brazilian coast. The alkaloids have been identified by analysis of spectroscopic data. While caissarine A has a 2-hydroxyagmatine moiety in its structure, caissarin B is the first naturally occurring compound encompassing the unprecedented 1,7-diamino-3-hydroxyheptane moiety.

Sponges belonging to the order Verongida are the richest source of naturally occurring bromine-containing alkaloids, biogenetically derived from tyrosine.¹ Although the first compound of this structural class was isolated almost 40 years ago,² Verongid sponges continue to provide new bromotyrosine-derived alkaloids. Such compounds have proven to be valuable chemotaxonomic markers,^{3,4} and many of them display potent biological activities. Examples are aeroplysinin-1, a promising anticancer agent,⁵ and the bastadins from *Ianthella basta*, which display potent cytotoxic,⁶ inosine 5'-phosphate dehydrogenase inhibitory,⁷ and open Ca²⁺ channel stabilizing activities.⁸ Recent investigations on related sponges led to the isolation of completely new biologically active chemotypes, such as the antihistaminic archerine from *Aplysina archeri*,⁹ and the cytotoxic ma'edamines A and B from *Suberea* species.¹⁰ A recent study demonstrated that the Verongid sponge *Aplysina cavernicola* had a high assemblage of symbiotic bacteria.¹¹ Both *A. cavernicola* and *A. aerophoba* from the Mediterranean have associated bacteria that inhibit the growth of Gram-positive and Gram-negative bacteria, including antibiotic-resistant strains.¹²

During our ongoing search of new bioactive natural products from marine invertebrates,¹³ we have noticed that the crude methanol extract of the sponge *Aplysina caissara* (Pinheiro and Hajdu, 2001) displayed mild cytotoxic and antibacterial activities. *A. caissara* is a recently described endemic Brazilian species of marine sponge.¹⁴ A chemical investigation of the crude extract of *A. caissara* led to the isolation of three known compounds, aeroplysinin-1 (**4**),¹⁵ fistularin-3 (**5**),¹⁶ the artifact of isolation 2-(3,5-dibromo-4-ethoxy-1-hydroxy-4-methoxy-2,5-cyclohexadien-1-yl)ethanamide (**6**),¹⁷ as well as two new dibromotyrosine-derived alkaloids, caissarins A (**1**) and B (**2**), whose isolation and structure elucidation are reported herein.

The frozen sponge was extracted with methanol, and the methanol extract was concentrated to an aqueous suspen-

sion, which was partitioned with hexanes and with ethyl acetate. The ethyl acetate fraction was subjected to C₁₈ reversed-phase column chromatography and silica gel flash chromatography. The UV-absorbing fractions were further separated and purified by reversed-phase HPLC, to give, in increasing order of polarity, aeroplysinin-1, 1-acetamide-3,5-dibromo-4,4-dimethoxy-1-hydroxycyclohexa-2,5-diene, fistularin-3, caissarine B (**2**), and caissarine A (**1**).

The positive FABMS of caissarine A (**1**) displayed a molecular ion triplet at *m/z* 516, 518, and 520, indicating the presence of two bromine atoms in its structure. Analysis of the ¹H, ¹³C, gHSQC, and gHMBC NMR spectra (Table 1) indicated the presence of the 7,9-dibromo-10-hydroxy-8-methoxy-1-oxa-2-aza-spiro[4.5]deca-2,6,8-trien-3-carboxamide moiety, typically found in most of the secondary metabolites isolated from Verongida sponges. Additionally, we observed the presence of three methylene (δ_C 44.7, 25.7, and 40.2), one methine (δ_C 78.4), and a quaternary carbon (δ_C 157.5). Analysis of IR, gHSQC, ¹H–¹H gCOSY, and gHMBC indicated that the C-10 methylene (δ_C 44.7; δ_H 3.54) was attached to a carbinolic methine (C-11), in agreement with the ¹³C (78.4) and ¹H (4.45) chemical shifts. Further ¹H–¹H and long-range couplings observed between C-11 methine and the C-12 methylene (δ_C 25.7; δ_H 1.78 and 2.05), between the C-12 methylene and C-13 methylene (δ_C 40.2; δ_H 3.35), and between the C-13 methylene and the quaternary carbon (C-14) at δ 157.5 established the identity of the 2-hydroxyagmatine chain.


Considering the structure **1** proposed for caissarine A by analysis of the IR and NMR data, the *m/z* 518 peak could not be assigned to the structure established above. However, it was assigned to the deuterated form of caissarine A, in which the exchangeable hydrogens were replaced by deuterium atoms.¹⁸ The presence of deuterium atoms in **1** was consistent with the fact that the sample submitted for mass spectra was recovered from MeOH-*d*₄. In the positive FABMS we also observed a small triplet at *m/z* 511, 513, and 515, possibly indicating the presence of two deuterium atoms in the place of two exchangeable hydrogens in **1**. A high-resolution measurement on the peak at *m/z* 518 indicated the formula C₁₅H₇D₇Br₂N₅O₅ (measd 518.20911) and that on the peak at *m/z* 513 indicated the formula

* To whom correspondence should be addressed. Tel: +55-16-2739954. Fax: +55-16-2739975. E-mail: rberlinck@iqsc.sc.usp.br.

[†] Instituto de Química de São Carlos, Universidade de São Paulo.

[‡] Departamento de Química, Universidade Federal de São Carlos.

[§] Museu Nacional, Universidade Federal do Rio de Janeiro.



$C_{15}H_{12}D_2Br_2N_5O_5$ (measd 513.17885), in agreement with 1H and ^{13}C NMR data. Therefore, caissarinin A presented seven exchangeable hydrogens and seven degrees of unsaturation. Dereplication with the MarinLit database¹⁹ indicated that **1** had 16 mass units higher than purealidine L (**3**), previously isolated from the sponge *Psammaplysilla pura*.²⁰ Indeed, the 1H and ^{13}C NMR data of caissarinin A (see Table 1) are very similar to NMR data of purealidine L,²⁰ except for the 2-hydroxyagmatine chain. Several attempts to transform the guanidine group of **1** into its 2-amino-3,5-dimethylpyrimidine derivative were made, but we only obtained complex mixtures, probably due to the formation of degradation products as previously observed for aplysinamisine II (**7**).²¹ It was not possible to obtain the value of specific rotation or the circular dichroism spectrum of **1** because the sample was lost after the NMR experiments. Nevertheless, we have been able to establish the relative stereochemistry of the bicyclic moiety of caissarinin A by comparison of the 1H and ^{13}C chemical shifts of H-1, H-5, and CH₂-7 with the literature data of related compounds.²²

Caissarinin B (**2**) was also isolated as a glassy solid. The low-resolution FABMS of caissarinin B displayed a molecular ion cluster at m/z 867, 869, 871, 873, and 875. A high-resolution measurement on the peak at m/z 871 (measd 871.97930) indicated the formula $C_{27}H_{32}Br_4N_4O_9$. Considering the presence of only 17 signals in the ^{13}C NMR spectrum of **2**, we supposed that caissarinin B had two chemically equivalent bicyclic moieties. This hypothesis was further supported by the fact that the intensity of the ^{13}C signals of the bicyclic spin system was more than 3-fold the intensity of the ^{13}C signals assigned to the 1,7-diamino-3-hydroxyheptane chain.

The assignment of 1H and ^{13}C signals of the bicyclic moieties of caissarinin B was established by comparison with data reported for dihydroxyaerthionin²³ and are presented in Table 1. Although no long-range correlations were observed in the gHMBC spectrum between the hydrogens of methylenes CH₂-10 (δ 3.08 and 3.15) and CH₂-16 (δ 3.15) with their respectively attached carbonyl carbons C-9 (δ 159.2) and C-17 (δ 159.2), or between the amide hydrogens with C-10 and C-16, the presence of the 1,7-diamino-3-hydroxyheptane chain was evident in the 1H - 1H gCOSY spectrum. The resonances of the hydrogens in the 1,7-diamino-3-hydroxyheptane chain were poorly resolved, suggesting the occurrence of a conformational dynamic change. 1H - 1H correlations have been observed between CH₂-10 (δ 3.08 and 3.15), CH-11 (δ 3.55 and 3.61), and CH₂-12 (δ 1.28 and 1.40), enabling us to establish the position of the hydroxyl group. The 1H and ^{13}C chemical shifts of the methine carbinol group appeared as two signals at δ 3.55 and 3.61, certainly due to the presence of two amide rotamers in solution. Furthermore, one amide proton appeared as two triplets at δ 8.41 (major conformer) and δ 8.45 (minor conformer), and the other amide proton appeared as two triplets at δ 8.26 (major conformer) and δ 8.21 (minor conformer), providing additional evidence for the presence of two rotamers in DMSO-*d*₆. Further sequential 1H - 1H couplings observed between CH₂-12 and CH₂-13 (δ 1.60 and 1.50), between CH₂-13 and CH₂-14 (δ 3.22 and 3.28), between CH₂-14 and CH₂-15 (δ 1.65 and 1.45), and finally between CH₂-15 with CH₂-16 (δ 3.15) completed the assignments of the 1,7-diamino-3-hydroxyheptane moiety of caissarinin B. Aiming to confirm the structure of caissarinin B, a second set of 1H , ^{13}C , 1H - 1H gCOSY, gHSQC, and gHMBC NMR experiments were obtained in MeOH-*d*₄ (see Experimental Section and Supporting Information), and the results confirm the planar structure of **2**. Particularly relevant were long-range couplings observed in the gHMBC spectrum between CH₂-10 (δ 3.38 and 3.26) and C-9 (δ 163.0), as well as between CH₂-13 (δ 1.65 and 1.73) and C-11 (δ 69.5 and 71.0). The 1H spectrum of caissarinin B in MeOH-*d*₄ is even less defined than in DMSO-*d*₆, indicating that the conformational dynamic change in solution is more pronounced in MeOH-*d*₄. As in the case of caissarinin A, it was not possible to establish the absolute stereochemistry of the bicyclic moieties of caissarinin B because the sample was lost after the NMR experiments. However, its relative stereochemistry was determined as shown by comparison with literature data.²²

To solve the absolute stereochemistry of caissarinins A and B by circular dichroism analysis, we collected additional samples of *A. caissara* in April 2000. Unexpectedly, the second sponge sample was devoid of both caissarinin A and caissarinin B. A third collection of this animal is envisaged in order to investigate if occurrence of **1** and **2**

Table 1. ^1H and ^{13}C NMR Data for Caissarine A (**1**) and Caissarine B (**2**)

1			2		
position	δ $^{13}\text{C}^a$	δ ^1H (mult, J in Hz) ^a	position	δ $^{13}\text{C}^b$	δ ^1H (mult, J in Hz) ^b
CH-1	76.4	4.10 (s)	CH-1 and CH-21	74.1	3.93 (s)
C-2	115.1		C-2 and C-22	113.5	
C-3	150.2		C-3 and C-23	147.6	
C-4	123.7		C-4 and C-24	121.2	
CH-5	133.2	6.40 (s)	CH-5 and CH-25	131.7	6.57 (s)
C-6	93.4		C-6 and C-20	90.7	7.47 (dd, 8.6 and 1.5)
CH ₂ -7	41.0	3.11 (d, 18) and 3.78 (d, 18)	CH ₂ -7 and CH ₂ -19	40.0	3.12 (d, 18) and 3.62 (d, 18)
C-8	156.0		C-8 and C-18	155.0	
C-9	162.8		C-9 and C-17	159.3	
CH ₂ -10	44.7	3.54 (d, 4)	CH ₂ -10	45.7	3.08 (m) and 3.15 (m)
CH-11	78.4	4.45 (m)	CH-11	67.4 and 68.9	3.55 (m) and 3.61 (m)
CH ₂ -12	25.7	1.78 (m) and 2.05 (m)	CH ₂ -12	32.3	1.28 (m) and 1.40 (m)
CH ₂ -13	40.2	3.35 (m)	CH ₂ -13	25.5	1.50 (m) and 1.60 (m)
C-14	157.5		CH ₂ -14	36.6	3.28 (m) and 3.22 (m)
OCH ₃	61.3	3.72 (s)	CH ₂ -15	34.4	1.45 (m) and 1.65 (m)
			CH ₂ -16	39.4	3.15 (m)
			OCH ₃	60.1	3.65 (s)
			N-H		8.41 (t, 6) and 8.45 (t, 5)
			N-H		8.27 (t, 6) and 8.21 (t, 5)
			O-H		6.36 (s)

^a Taken in CD₃OD. ^b Taken in DMSO-*d*₆.

in *A. caissara* is seasonal and to determine the absolute stereochemistry of both compounds.

To the best of our knowledge, caissarine B is the first Verongid dibromotyrosine-derived alkaloid bearing a 1,7-diamino-3-hydroxyheptane chain, a diamine moiety that has no precedent among natural products. Structurally related to caissarine B are dihydroxaerotherionin from *Verongula rigida*,²³ areotherionin and homoareotherionin, which have been isolated from different species of Verongid sponges,^{24–27} and 11-hydroxaerotherionin from *Pseudocercaria durrissima*.²⁸

Experimental Section

General Experimental Procedures. IR spectra were recorded on a FT-IR Bomem MB102 infrared spectrometer. NMR spectra were run either on a Bruker AC-4.7 T spectrometer, operating at 200.1 MHz for ^1H NMR and 50.3 MHz for ^{13}C NMR spectra, or on a Bruker DRX400 9.4 T instrument, operating at 400.35 MHz for ^1H and 100.10 MHz for ^{13}C channels, respectively. All the NMR spectra were obtained at 28 °C using tetramethylsilane as internal reference. High-resolution FAB mass spectra were obtained on hybrid Kratos concept ITHQ equipment. Solvents employed for extraction and column chromatography were glass distilled prior to use. TLC analysis were performed with Aldrich precoated TLC sheets of silica gel on polyester with 254 nm fluorescent indicator eluting with two eluents: 1:1 hexanes–ethyl acetate and 9:1 CH₂Cl₂–MeOH. Plates were developed by observing at 254 nm and subsequently by spraying with phosphomolybdic acid reagent in ethanol and further heating at 120 °C.

Animal Material. Samples of *A. caissara* were collected in the São Sebastião channel, during the summer of 1999. The animals were immediately frozen. Voucher specimens are deposited in the Porifera collection of the Museu Nacional da Universidade Federal do Rio de Janeiro [MNRJ 268 (paratype), 578 (paratype), 1673 (paratype), 1675 (paratype), 1988 (holotype), 1989 (paratype)].

Extraction and Isolation. The sponge *A. caissara* (900 g, wet wt) was blended in MeOH (3 L) and filtered, and the solid residue was re-extracted with MeOH (2 L). After filtration the methanol extract was evaporated in vacuo until the alcohol was removed. The final aqueous extract (ca. 500 mL) was partitioned against hexanes, then with EtOAc (3 × 700 mL), to give 9.55 g of a brown gum of the EtOAc extract. This material was divided in portions of ca. 1 g, which were subjected to a chromatography on a C₁₈ reversed-phase Sep Pak (Waters) column, with a gradient of MeOH in H₂O. Four

fractions were obtained, with UV-absorbing compounds concentrated in fractions 2 (3.5 g) and 3 (2.9 g). Both fractions were subjected to several separations by flash chromatography (gradient of MeOH in CH₂Cl₂), yielding fractions enriched in single components. Caissarine A (24 mg, 0.0026% wet wt) was obtained as a pure compound in the last fractions of these separations. Impure compounds were purified by C₁₈ reversed-phase HPLC with a Whatman Partisil 10 ODS-3 column, using 70% MeOH for the purification of aerophysinin-1 (340 mg) and 1-acetamide-3,5-dibromo-4,4-dimethoxy-1-hydroxycyclohexa-2,5-diene (249 mg) and 65% MeOH for the purification of fistularin-3 (25 mg) and caissarine B (25 mg, 0.0026% wet wt).

Caissarine A (1): colorless, glassy solid; UV (MeOH) λ_{max} 232 (ϵ 9100), 283 (ϵ 4250); IR (neat) ν_{max} 3500–3000, 2928, 2860, 1690–1630, 1405, 1100, 605 cm⁻¹; ^1H NMR (DMSO-*d*₆, 400 MHz), see Table 1; ^{13}C NMR (DMSO-*d*₆, 400 MHz), see Table 1; FABMS (thioglycerol) m/z 518 [M + 7D – 7H]⁺ (13), 513 [M + 2D – 2H]⁺ (4), 496 (8), 478 (6), 295 (23), 279 (22), 157 (60), 71 (100); HRFABMS m/z found 518.20911 [M + 7D – 7H]⁺, calcd for C₁₅H₇D₇Br₂N₅O₅ [M + 7D – 7H]⁺ 518.20836.

Caissarine B (2): colorless, glassy solid; UV (MeOH) λ_{max} 234 (ϵ 9000), 283 (ϵ 4300); IR (neat) ν_{max} 3382, 2357, 1665, 1549, 1106, 603 cm⁻¹; ^1H NMR (DMSO-*d*₆, 400 MHz), see Table 1; ^{13}C NMR (DMSO-*d*₆, 100 MHz), see Table 1; ^1H NMR (MeOH-*d*₄, 400 MHz) δ 6.42 (2H, m, H-5 and H-25), 4.10 (2H, m, H-1 and H-21), 3.79 (1H, m, H-11) and 3.72 (1H, m, H-11), 3.77 (2H, dd, 3.6, 20.8 Hz, CH₂-7 and CH₂-19) and 3.09 (2H, dd, 2.9, 20.8 Hz, CH₂-7 and CH₂-19), 3.72 (3H, s, OCH₃), 3.40 (2H, m, CH₂-14), 3.38 (1H, m, CH₂-10) and 3.26 (1H, m, CH₂-10), 3.30 (2H, m, CH₂-16), 1.73 (1H, m, CH₂-13) and 1.65 (1H, m, CH₂-13), 1.62 (2H, m, CH₂-15), 1.55 (1H, m, CH₂-12) and 1.43 (1H, m, CH₂-12); ^{13}C NMR (MeOH-*d*₄, 100 MHz) δ 163.0 (s, C-9 and C-17), 156.2 (s, C-8 and C-18), 149.9 (s, C-3 and C-23), 135.0 (d, C-5 and C-25), 123.5 (s, C-4 and C-24), 115.0 (C-2 and C-22), 93.5 (s, C-6 and C-20), 76.0 (d, C-1 and C-21), 71.0 and 69.5 (d, C-11), 61.0 (q, OCH₃), 47.5 (t, C-10), 40.6 (t, C-16), 40.5 (t, C-7 and C-9), 37.8 (t, C-14), 35.0 (t, C-15), 32.9 (t, C-12), 27.0 (t, C-13); FABMS (thioglycerol + MeOH) m/z 875 (1), 873 (2), 871 [M]⁺ (4), 869 (2), 867 (1), 861 (1.5), 859 (3.5), 857 (5), 855 (3.5), 855 (1.5), 322 (1.5), 320 (2.5) 318 (1.5), 297 (2.5), 295 (4.5), 293 (2.5), 281 (18), 279 (37), 277 (18), 70 (100); HRFABMS m/z found 871.97930 [M]⁺, calcd for C₂₇H₃₂Br₄N₄O₉ [M]⁺ 871.97028.

Acknowledgment. The authors thank Professor Peter Northcote (Victoria University of Wellington, New Zealand) for a critical review of the manuscript, Dr. David E. Williams and Professor Raymond J. Andersen (University of British Columbia, Vancouver, Canada) for the MS analyses, and Prof.

José Carlos de Freitas and technical staff of the Centro de Biologia Marinha of the Universidade de São Paulo (CEBIMar-USP) for many facilities in sponge collection. Financial support was provided by the American Society of Pharmacognosy Foundation Research Starter Grant (1998) and a grant of the Fundação de Amparo à Pesquisa do Estado de São Paulo (96/04316-5) to R.G.S.B. A.C.G. also thanks FAPESP for a fellowship (98/11689-8).

Supporting Information Available: This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- (1) For recent reports of bromotyrosine-derived alkaloids, see: (a) Ciminiello, P.; Dell'Aversano, C.; Fattorusso, E.; Magno, S.; Pansini, M. *J. Nat. Prod.* **2000**, *63*, 263–266. (b) Ross, S. A.; Weete, J. D.; Schinazi, R. F.; Wirtz, S. S.; Tharnish, P.; Scheuer, P. J.; Hamann, M. T. *J. Nat. Prod.* **2000**, *63*, 501–503. (c) Okamoto, Y.; Ojika, M.; Sakagami, Y. *Tetrahedron Lett.* **1999**, *40*, 507–510. (d) Compagnone, R. S.; Avila, R.; Suárez, A. I.; Abrams, O. V.; Rangel, H. R.; Anvelo, F.; Piña, I. C.; Merentes, E. *J. Nat. Prod.* **1999**, *62*, 1443–1444.
- (2) (a) Sharma, G. M.; Burkholder, P. R. *J. Antibiotics* **1967**, *20*, 200–203. (b) Sharma, G. M.; Burkholder, P. R. *Tetrahedron Lett.* **1967**, 4147–4150.
- (3) Bergquist, P. R.; Wells, R. J. In *Marine Natural Products: Chemical and Biological Perspectives*; Scheuer, P. J., Ed.; Academic Press: New York, 1983; Vol. V, pp 1–50.
- (4) . Soest, R. W. M.; Braekman, J. C. *Mem. Queensland Museum* **1999**, *44*, 569–589.
- (5) . Jaspars, M. In *Advances in Drug Discovery Techniques*; Harvey, Alan L., Ed.; John Wiley & Sons: New York, 1998; pp 65–84.
- (6) Miao, S.; Andersen, R. J.; Allen, T. *J. Nat. Prod.* **1990**, *53*, 1441–1446.
- (7) Jaspars, M.; Rali, T.; Laney, M.; Schatzman, R. C.; Diaz, M. C.; Schmitz, F. J.; Pordesimo, E. O.; Crews, P. *Tetrahedron* **1994**, *50*, 7367–7374.
- (8) Chen, L.; Molinski, T. F.; Pessah, I. N. *J. Biol. Chem.* **1999**, *274*, 32603–32612.
- (9) Ciminiello, P.; Dell'Aversano, C.; Fattorusso, E.; Magno, S. *Eur. J. Org. Chem.* **2001**, 55–60.
- (10) Hirano, K.; Kubota, T.; Tsuda, M.; Watanabe, K.; Fromont, J.; Kobayashi, J. *Tetrahedron* **2000**, *56*, 8107–8110.
- (11) Friedrich, A. B.; Merkert, H.; Fendert, T.; Hacker, J.; Proksch, P.; Hentschel, U. *Mar. Biol.* **1999**, *134*, 461–470.
- (12) Hentschel, U.; Schmid, M.; Wagner, M.; Fieseler, L.; Gernert, C.; Hacker, J. *FEMS Microbiol. Ecol.* **2001**, *35*, 305–312.
- (13) Torres, Y. R.; Berlinck, R. G. S.; Magalhães, A.; Schefer, A. B.; Ferreira, A. G.; Hajdu, E.; Muricy, G. *J. Nat. Prod.* **2000**, *63*, 1098–1105, and references therein.
- (14) Pinheiro, U. S.; Hajdu, E. *Revta. Bras. Zool.* **2001**, *18*, 143–160.
- (15) Fattorusso, E.; Minale, L.; Sodano, G. *J. Chem. Soc., Chem. Commun.* **1970**, 751–753.
- (16) Gopichand, Y.; Schmitz, F. J. *Tetrahedron Lett.* **1979**, 3921–3924.
- (17) Sharma, G. M.; Vig, B.; Burkholder, P. R. *J. Org. Chem.* **1970**, *35*, 2823–2826.
- (18) We thank Professor D. John Faulkner (Scripps Institution of Oceanography, University of California at San Diego, CA) for suggesting this possibility to one of us (R.G.S.B.).
- (19) . Munro, M. H. G.; Blunt, J. W. *MarinLit—Marine Literature Database*; 1999; Updated version.
- (20) Kobayashi, J.; Honma, K.; Sasaki, T.; Tsuda, M. *Chem. Pharm. Bull.* **1995**, *43*, 403–407.
- (21) Rodríguez, A. D.; Piña, I. C. *J. Nat. Prod.* **1993**, *56*, 907–914.
- (22) Nishiyama, S.; Yamamura, S. *Bull. Chem. Soc. Jpn.* **1985**, *58*, 3453–3456.
- (23) Gunasekera, M.; Gunasekera, S. P. *J. Nat. Prod.* **1989**, *52*, 753–756.
- (24) Fattorusso, E.; Minale, L.; Sodano, G.; Moody, K.; Thomson, R. H. *Chem. Comm.* **1970**, 752–753.
- (25) Moody, K.; Thomson, R. H.; Fattorusso, E.; Minale, L.; Sodano, G. *J. Chem. Soc., Perkin Trans. 1* **1972**, 18–24.
- (26) McMillan, J. A.; Paul, I. C.; Goo, Y. M.; Rinehart, K. L.; Krueger, W. C.; Pschigoda, L. M. *Tetrahedron Lett.* **1981**, *22*, 39–42.
- (27) Venkateswarlu, Y.; Rama Rao, M.; Venkatesham, U. *J. Nat. Prod.* **1998**, *61*, 1388–1389.
- (28) Kernan, M. R.; Cambie, R. C.; Bergquist, P. R. *J. Nat. Prod.* **1990**, *53*, 615–622.

NP0105735