

Semisynthetic Analogues of the Microtubule-Stabilizing Agent Discodermolide: Preparation and Biological Activity

Sarath P. Gunasekera,* Ross E. Longley, and Richard A. Isbrucker

Division of Biomedical Marine Research, Harbor Branch Oceanographic Institution, 5600 U.S. 1 North, Fort Pierce, Florida 34946

Received July 18, 2002

A series of 12 semisynthetic discodermolide analogues, **2–13**, have been prepared using natural (+)-discodermolide (**1**) and evaluated for in vitro cytotoxicity against cultured murine P-388 leukemia and A-549 human adenocarcinoma cells. These semisynthetic analogues showed a significant variation of cytotoxicity and confirmed the importance of the C-7 through C-19 molecular fragment for potency. Specifically, these analogues suggested the importance of the C-11 and C-17 hydroxyl groups and the C-13 double bond for the potency of discodermolide. The preparation, structure elucidation, and biological activity of these new analogues are described.

Earlier, we reported the preparation, structure elucidation, biological activity, and structure–activity relationship of eight acetylated discodermolide analogues.^{1–3} Recently, our group reported the isolation, structure determination, and biological activity of five new naturally occurring discodermolide analogues from a sponge sample of *Discodermia* sp.⁴ Herein, we report the preparation, structural elucidation, biological activity, and structure–activity relationship of 12 (**2–13**) new semisynthetic analogues of natural (+)-discodermolide (**1**). Discodermolide (**1**) was first reported in 1990,⁵ and since then, studies by our group and others have indicated its immunosuppressive^{6–8} and anti-mitotic properties^{9–11} with a mechanism of action similar to that of paclitaxel.^{9,10} Discodermolide has been shown to promote the rapid polymerization of purified tubulin and to hyperstabilize the microtubule complex in cultured cells. (+)-Discodermolide inhibits the in vitro growth of several cancer cell lines, including paclitaxel-resistant ovarian and colon cancer cells.^{11–13} The Schreiber group has synthesized both antipodes of discodermolide, establishing the absolute configuration,¹⁴ and prepared a number of structural variants.¹⁵ Since then, several other groups have synthesized (+)-discodermolide,^{16,17} antipode (–)-discodermolide,^{18,19} or various fragments of discodermolide using different synthetic approaches.^{20–33} In 2001, the Paterson group³⁴ synthesized (+)-discodermolide and three epimeric discodermolides.

Results and Discussion

The starting material for this study, natural (+)-discodermolide (**1**), was isolated from the sponge *Discodermia* sp. and crystallized using the reported procedure.⁵ The preparation of various discodermolide acetates used in this study was described in our earlier publication.¹ Structurally, discodermolide possesses 13 stereogenic centers, a tetrasubstituted δ -lactone (C-1 to C-5), four secondary hydroxyl groups (C-3, C-7, C-11, and C-17), a pendant carbamate moiety (C-19), one di- (C-8, *Z*) and one trisubstituted (C-13, *Z*) double bond, and a terminal (*Z*)-diene (C-21 to C-24). Both in solid state⁵ and in solution,³⁵ discodermolide adopts a hairpin conformation where the two internal (*Z*)-olefins in the side chain act as conformational locks by minimizing (1,3) strain between their

respective substituents in concert with the avoidance of *syn*-pentane interactions. The molecular structure of discodermolide is amenable to other simple modifications; however, for this study, we focused on the four olefinic double bonds and the four hydroxy groups present in the molecule to prepare the selectively hydrogenated analogues (**2–6**), dehydroxylated analogues (**7–11**), and *seco* analogues (**12**, **13**). The ¹H NMR data of all analogues were compared to those reported for discodermolide (see Table 1) and discodermolide acetates,¹ and the chemical shifts were assigned using ¹H–¹H COSY data. DEPT, HMQC, and in some instances HMBC data were used to assign ¹³C data of all analogues and are presented in Tables 5 and 6.


HRFABMS of 8,13,21,23-octahydrodiscodermolide (**2**) supported the molecular formula C₃₃H₆₄NO₈ [(M + H)⁺, *m/z* 602.4618] as expected for the octahydro product. The ¹H NMR spectrum (see Table 1) indicated the saturation of all olefinic signals present in discodermolide. Similarly, the ¹³C spectrum (see Table 5) showed no olefinic carbons and instead indicated the presence of six additional triplets (δ 36.3, C-8; 29.0, C-9; 42.3, C-13; 23.6, C-21; 31.9, C-22 and 29.9, C-23), a doublet (δ 28.0, C-14), and a quartet (δ 14.3, C-24), which is expected for the saturation of the four double bonds in discodermolide. These data confirmed the structure of 8,13,21,23-octahydrodiscodermolide (**2**). The relative stereochemistry of the C-29 methyl group was not determined.

HRFABMS of 8,21,23-hexahydrodiscodermolide (**3**) supported the molecular formula C₃₃H₆₂NO₈ [(M + H)⁺, *m/z* 600.4478] as expected for the hexahydro product. The ¹H NMR spectrum (see Table 1) showed the presence of one olefinic doublet at δ 5.04 (*J* = 9.8 Hz). The COSY spectrum indicated the coupling of this proton to the C-12 multiplet at δ 2.54, which was in turn coupled to the C-11 hydroxymethine proton (δ 3.03, ddd, *J* = 5.2, 5.1, 5.7 Hz). The ¹³C NMR spectrum (see Table 5) showed the presence of one olefinic doublet (δ 131.5) and one olefinic singlet (δ 133.6), and these ¹H and ¹³C data established the position of the double bond in hexahydrodiscodermolide. Comparison of ¹H and ¹³C data with those of discodermolide (see Tables 1 and 5) confirmed the structure of 8,21,23-hexahydrodiscodermolide (**3**).

HRFABMS supported the molecular formula C₃₃H₆₀NO₈ [(M + H)⁺, *m/z* 598.4338] expected for tetrahydrodiscodermolide (**4**). The ¹H NMR spectrum (see Table 2) showed

* To whom correspondence should be addressed. Tel: 772-465-2400. Fax: 772-461-2221. E-mail: sgunaseker@hboi.edu.

Chart 1



the presence of two additional olefinic signals (δ 5.35, ddd, $J = 2.2, 9.1, 10.9$ Hz and δ 5.49 ddd, $J = 1.0, 10.1, 10.9$ Hz) compared to that of hexahydrodiscodermolide (**3**). The COSY spectrum revealed a coupling of the δ 5.35 olefinic signal to the H-7 hydroxymethine signal at δ 4.46 and coupling of the δ 5.49 signal to the H-10 signal at δ 2.65, which was in turn coupled to the H-11 hydroxymethine signal at δ 3.09. The ^{13}C NMR spectrum (see Table 5) showed the presence of only three olefinic doublets (δ 133.8, C-8; 133.7, C-9; 131.2, C-13) and one olefinic singlet (δ 133.8, C-14), indicating the saturation of two conjugated terminal double bonds in the discodermolide molecule. These data confirmed the structure of 21,23-tetrahydrodiscodermolide (**4**).

HRFABMS supported the molecular formula $\text{C}_{33}\text{H}_{62}\text{NO}_7$ [$(\text{M} + \text{H})^+$, m/z 584.4512] expected for the deoxy-hexahydrodiscodermolide (**5**). The ^1H NMR spectrum closely resembled that of 8,21,23-hexahydrodiscodermolide. The presence of an H-13 doublet (δ 5.03, d, $J = 9.8$ Hz) in the ^1H spectrum together with the olefinic doublet (δ 130.0, C-13) and a singlet (δ 133.2, C-14) in the ^{13}C spectrum established compound **5** to be an analogue of 8,21,23-hexahydrodiscodermolide (**3**). Comparison of the ^1H NMR spectrum of **5** with that of **3** (see Tables 1 and 2) indicated that **5** has one less hydroxymethine proton. The COSY spectrum of **5** indicated that H-5 (δ 4.30) is coupled to H-6 methylene protons (δ 1.55 and 1.65), and these were in turn coupled to another methylene group at δ 1.39 (H-7) and 1.47 (H-7). These data suggested that deoxygenation has

taken place at C-7 during the hydrogenation of discodermolide and thus confirmed the structure of **5** as 7-deoxy-hexahydrodiscodermolide.

The 7-deoxy-hexahydrodiscodermolide on acetylation furnished its triacetate **6**. HRFABMS supported the molecular formula $\text{C}_{39}\text{H}_{68}\text{NO}_{10}$ [$(\text{M} + \text{H})^+$, m/z 710.4827] expected for the triacetate (**6**). The ^1H NMR spectrum of **6** (see Table 2) when compared with that of **5** showed the presence of three acetoxy methyls (δ 2.08, 2.05, and 2.07) and downfield-shifted signals for H-3 (Δ 1.18 ppm), H-11 (Δ 1.50 ppm), and H-17 (Δ 1.51 ppm). The ^{13}C spectrum (see Table 5) when compared with that of **5** showed the presence of three additional methyl signals (δ 21.0, 20.9, 20.8) and three additional carbonyl signals (δ 171.0, 170.4, 170.3) for the presence of three acetoxy groups. These data confirmed the structure of the acetylated product of **5** as 7-deoxy-hexahydrodiscodermolide-3,11,17-triacetate (**6**).

Discodermolide-3-acetate on deacetylation under alkaline conditions gave the 3-deoxy analogue **7**. HRFABMS supported the molecular formula $\text{C}_{33}\text{H}_{54}\text{NO}_7$ [$(\text{M} + \text{H})^+$, m/z 576.3931] expected for the deoxy-discodermolide (**7**). The ^1H NMR spectrum of 3-deoxy-discodermolide-2-ene (**7**) (see Table 2) was similar to that of discodermolide (see Table 1). The ^1H NMR spectrum of **7** indicated signals for eight methyl groups as in discodermolide. However, it showed the presence of two vinylic methyl singlets compared to one vinylic methyl singlet present in discodermolide. The comparison of the ^1H NMR spectrum of **7** with that of discodermolide indicated that the downfield methyl doublet

Table 1. ¹H NMR Data of Discodermolide (**1**) and Semisynthetic Analogues **2** and **3**^a

position	1	2	3
2	2.56 (dq, 4.2, 7.2)	2.56 (dq, 4.6, 7.3)	2.57 (dd, 4.2, 7.4)
3	3.61 (ddd, 4.2, 4.2, 4.6)	3.63 (ddd, 2.3, 3.5, 4.7)	3.64 (ddd, 4.0, 4.0, 4.5)
3-OH	3.27 (4.6)	3.27 (d, 4.7)	3.31 (d, 4.5)
4	1.83 (ddq, 2.0, 4.2, 6.9)	1.87 (m)	1.87 (m)
5	4.45 (dt, 2.0, 10.1)	4.47 (dt, 2.0, 9.5)	4.48 (dt, 2.0, 10.0)
6	1.72 (m)	1.32 (m)	1.59 (m)
6	1.47 (ddd, 2.4, 10.8, 13.0)	1.61 (m)	1.65 (m)
7	4.44 (m)	3.72 (m)	3.70 (m)
7-OH	2.75 (d, 5.3)	2.77 (d, 5.2)	2.84 (d, 5.5)
8	5.38 (ddd, 2.2, 9.1, 10.9)	1.27 (m)	1.24 (m)
9	5.54 (ddd, 1, 10.1, 10.9)	1.70 (m)	1.65 (m)
		1.10 (m)	1.00 (m)
10	2.62 (m)	1.51 (m)	1.48 (m)
11	3.05 (m)	3.02 (m)	3.03 (ddd, 5.2, 5.1, 5.7)
11-OH	2.64 (d, 5.2)	2.44 (d, 5.5)	2.43 (d, 5.7)
12	2.29 (ddq, 5.2, 6.6, 10.0)	1.71 (m)	2.54 (m)
13	4.97 (d, 10.0)	1.28 (m)	5.04 (d, 9.8)
		0.97 (m)	
14		1.55 (m)	
15	1.76 (m)	1.20 (m)	2.09 (m)
15	1.63 (dd, 3.6, 12.2)	0.95 (m)	1.93 (dd, 3.6, 12.2)
16	1.76 (m)	1.80 (m)	1.93 (m)
17	3.27 (dd, 3.5, 6.2)	3.16 (ddd, 5.0, 6.1, 6.2)	3.54 (ddd, 5.2, 5.1, 5.9)
17-OH	2.59 (d, 6.5)	2.59 (d, 6.1)	2.80 (d, 5.9)
18	1.72 (m)	1.87 (m)	1.87 (m)
19	4.71 (dd, 4.1, 8.0)	4.53 (dd, 4.2, 7.3)	4.54 (dd, 4.5, 7.7)
20	3.07 (m)	1.74 (m)	1.75 (m)
21	5.42 (dd, 10.6, 10.7)	1.25 (m)	1.26 (m)
		1.09 (m)	1.01 (m)
22	6.06 (dd, 10.7, 11.0)	1.38 (m)	1.40 (m)
22		1.19 (m)	1.19 (m)
23	6.68 (ddd, 10.5, 11.0, 16.6)	1.25 (m)	1.26 (m)
		1.35 (m)	1.35 (m)
24	5.21 (d, 16.6)	0.89 (t, 6.6)	0.90 (t, 6.6)
24	5.10 (d, 10.1)		
25	1.18 (d, 7.2)	1.20 (d, 7.4)	1.21 (d, 7.4)
26	0.97 (d, 6.9)	0.97 (d, 6.6)	0.97 (d, 6.7)
27	1.00 (d, 6.9)	0.85 (d, 6.7)	0.88 (d, 6.8)
28	0.88 (d, 6.6)	0.85 (d, 6.6)	0.92 (d, 6.6)
29	1.57 (s)	0.82 (d, 6.6)	1.63 (s)
30	0.73 (d, 6.2)	0.89 (d, 6.5)	0.78 (d, 6.5)
31	0.80 (d, 6.5)	0.81 (d, 6.5)	0.90 (d, 6.6)
32	0.93 (d, 6.7)	0.78 (d, 6.7)	0.87 (d, 6.6)
NH ₂	5.05 (br s)	5.10 (br s)	5.08 (br s)

^a All spectra run at 500 MHz in CD₃CN. Chemical shifts are reported in ppm, and *J* values in Hz.

corresponding to the C-25 methyl (δ 1.18, d, *J* = 7.2 Hz, H₃-25) in discodermolide has been changed to a vinylic methyl singlet (δ 1.81, s) and has been shifted downfield (Δ 0.63 ppm) as expected. The ¹H–¹H COSY spectrum of **7** indicated allylic coupling between the vinylic methyl at 1.81 ppm and the newly formed broad olefinic singlet at 6.42 ppm. Although no coupling was seen between the olefinic singlet at 6.42 ppm and the H-4 at 2.44 ppm, the allylic nature of the H-4 established the position of the new unsaturation. The remainder of the ¹H NMR spectrum was identical to that of discodermolide. The ¹³C spectrum of **7** showed the presence of a conjugated olefinic singlet (δ 127.5) and an olefinic doublet (δ 147.0) instead of an oxygenated carbon and a methine carbon present in the ¹³C spectrum of discodermolide. These data confirmed the structure of 3-deoxy-discodermolide-2-ene (**7**).

The ¹H NMR data of the four acetylated products **8**, **9**, **10**, and **11** of 3-deoxy-discodermolide (**7**) were compared with the ¹H NMR data of discodermolide and 3-deoxy-discodermolide. Using the ¹H–¹H COSY data in combination with the HMQC data, the proton chemical shift values of these analogues were assigned (see Tables 3 and 4). The ¹H NMR spectra of these analogues indicated a significant downfield shift of the signals corresponding to the acetoxy methines (see Tables 3 and 4), and these data were used to determine the position of acetylation in each

analogue. The HRMS data of **8** and **9** are consistent with the presence of one acetoxy group in each compound and established the molecular formula C₃₅H₅₅NO₈. Comparison of the ¹H NMR spectra of **8** and **9** with that of 3-deoxy-discodermolide-2-ene (**7**) showed the presence of one acetoxy methyl in each compound and downfield shifts of H-11 (Δ 1.65 ppm) in **8** and H-17 (Δ 1.65 ppm) in **9**. The combination of these data confirmed the structures of 3-deoxy-discodermolide-2-en-11-acetate (**8**) and 3-deoxy-discodermolide-2-en-17-acetate (**9**).

HRESIMS supported the molecular formula C₃₇H₅₇NNaO₉ [(M + Na)⁺, *m/z* 682.3896] expected for the 3-deoxy-discodermolide-2-ene-diacetate (**10**). The ¹H NMR spectrum of **10** (see Table 4) when compared with that of **7** (see Table 3) showed the presence of two additional acetoxy methyls (δ 2.04 and 2.08) and a downfield shift of H-11 (Δ 1.58 ppm) and H-17 (Δ 1.64 ppm). These data are supported by the ¹³C data, which showed signals for two acetoxy groups (δ 20.9, q; 21.0, q; 171.0, s; 170.7, s). Combination of these data confirmed the structure of 3-deoxy-discodermolide-2-en-11,17-diacetate (**10**).

The molecular formula C₃₉H₅₉NO₁₀ expected for the triacetate **11** was supported by HRESIMS data [*m/z* 702.4204, for C₃₉H₆₀NO₁₀ (M + H)⁺]. The ¹H NMR spectrum of **11** (see Table 4) when compared with that of **7** (see Table 3) showed the presence of three additional acetoxy-

Table 2. ¹H NMR Data of Discodermolide Analogues **4**, **5**, and **6**^a

position	4 ^b	5 ^c	6 ^c
2	2.56 (dq, 4.2, 7.2)	2.64 (dq, 3.7, 7.4)	2.71 (dd, 3.7, 7.4)
3	3.62 (ddd, 4.2, 4.2, 4.6)	3.71 (dd, 3.7, 3.7)	4.89 (dd, 3.7, 3.7)
3-OH	3.30 (4.6)		
4	1.83 (m)	1.91 (m)	2.05 (m)
5	4.47 (dt, 2.0, 10.1)	4.30 (ddd, 2.8, 8.2, 8.2)	4.28 (ddd, 2.0, 8.5, 8.5)
6	1.72 (m)	1.55 (m)	1.53 (m)
6	1.46 (ddd, 2.4, 10.8, 13.0)	1.65 (m)	1.65 (m)
7	4.46 (m)	1.39 (m)	1.35 (m)
7		1.47 (m)	1.45 (m)
7-OH	2.80 (d, 5.3)		
8	5.35 (ddd, 2.2, 9.1, 10.9)	1.22 (m)	1.24 (m)
9	5.49 (ddd, 1.0, 10.1, 10.9)	1.70 (m)	1.65 (m)
		1.10 (m)	1.00 (m)
10	2.65 (m)	1.53 (m)	1.65 (m)
11	3.09 (m)	3.13 (dd, 4.7, 6.4)	4.63 (dd, 4.3, 7.9)
11-OH	2.64 (d, 5.2)		
12	2.38 (ddq, 5.2, 6.6, 10.0)	2.57 (ddq, 4.7, 3.4, 10.0)	2.60 (ddq, 4.7, 3.4, 10.0)
13	5.03 (d, 10.0)	5.03 (d, 9.8)	4.94 (d, 9.8)
15	1.95 (m)	1.87 (m)	1.75 (m)
15	1.75 (dd, 3.6, 12.2)	2.11 (m)	1.62 (m)
16	1.83 (m)	1.97 (m)	2.21 (m)
17	3.15 (ddd, 3.5, 6.1, 6.2)	3.29 (dd, 5.3, 5.4)	4.80 (dd, 4.2, 7.7)
17-OH	2.64 (d, 6.1)		
18	1.87 (m)	1.90 (m)	2.04 (m)
19	4.55 (dd, 4.1, 8.0)	4.58 (dd, 3.8, 7.7)	4.56 (dd, 3.8, 7.8)
20	1.74 (m)	1.72 (m)	1.75 (m)
21	1.25 (m)	1.27 (m)	1.27 (m)
	1.09 (m)	1.05 (m)	1.08 (m)
22	1.38 (dd, 10.7, 11.0)	1.38 (m)	1.40 (m)
22	1.19 (m)	1.19 (m)	1.16 (m)
23	1.25 (m)	1.25 (m)	1.26 (m)
	1.35 (m)	1.35 (m)	1.35 (m)
24	0.91 (t, 6.6)	0.87 (t, 6.6)	0.87 (t, 6.6)
25	1.19 (d, 7.4)	1.29 (d, 7.3)	1.31 (d, 7.4)
26	0.97 (d, 6.9)	1.03 (d, 6.7)	0.96 (d, 6.7)
27	0.99 (d, 6.9)	0.90 (d, 6.8)	0.81 (d, 6.8)
28	0.91 (d, 6.6)	0.95 (d, 6.8)	0.84 (d, 6.7)
29	1.61 (s)	1.65 (s)	1.63 (s)
30	0.75 (d, 6.6)	0.80 (d, 6.6)	0.72 (d, 6.5)
31	0.85 (d, 6.5)	0.94 (d, 6.7)	0.86 (d, 6.6)
32	0.84 (d, 6.7)	0.86 (d, 6.7)	0.85 (d, 6.6)
NH ₂	5.05 (br s)	4.70 (br s)	4.65 (br s)
35			2.08 (s)
37			2.05 (s)
41			2.07 (s)

^a All spectra run at 500 MHz. Chemical shifts are reported in ppm, and *J* values in Hz. ^b In CD₃CN. ^c In CDCl₃.

ethyls (δ 2.01, 2.08, 1.99) and the downfield shift of H-7 (Δ 1.20 ppm), H-11 (Δ 1.57 ppm), and H-17 (Δ 1.65 ppm) acetoxymethine protons. The ¹³C data supported the presence of three acetoxy groups in the molecule (δ 20.9, q; 21.0, q; 21.2, q; 171.0, s; 170.6, s; 169.8, s). These data confirmed the structure of 3-deoxy-discodermolide-2-en-7,11,17-triacetate (**11**).

Discodermolide (**1**) on mild oxidation with MnO₂ at room temperature gave 6,7-*seco*-discodermolide instead of the expected 7-ketodiscodermolide. HRESIMS supported the molecular formula C₂₅H₄₀NO₅ [(M + H)⁺, *m/z* 434.2911] expected for the 6,7-*seco*-discodermolide (**12**). The ¹H NMR spectrum **12** (see Table 4) was quite different from that of discodermolide (see Table 1) and showed the absence of all signals corresponding to the C-1 through C-7 fragment of the discodermolide molecule. The ¹H spectrum indicated the presence of one vinylic methyl singlet (δ 1.64, s) corresponding to the C-14 methyl group and five methyl doublets. The presence of a conjugated double bond was apparent from the downfield shift of C-8 to 5.91 ppm and C-9 to 6.64 ppm. The signals corresponding to the C-21 through C-24 diene system were present in the spectrum. The presence of the carbamate group was evident from the characteristic NH₂ signal at 4.48 ppm. The ¹³C spectrum indicated signals for 25 carbons including the α,β -unsatur-

ated δ -lactone (δ 164.0, s) and the characteristic carbamate group (δ 158.0, s). These data suggested that the C-7 allylic hydroxyl group in discodermolide had undergone a complete oxidation to an acid group and that in turn had esterified with the C-11 hydroxy group to yield the α,β -unsaturated δ -lactone. These data confirmed the structure of 6,7-*seco*-discodermolide (**12**).

The ¹H NMR spectrum of 19-*des*-aminocarbonyl-6,7-*seco*-discodermolide closely resembled that of **12** (see Table 4). The characteristic differences observed were the absence of the carbamate NH₂ functionality and the upfield shift of the C-19 oxymethine proton (Δ 1.31 ppm). HRESIMS supported the molecular formula C₂₄H₃₉O₄ [(M + H)⁺, *m/z* 391.2858] expected for the *des*-aminocarbonyl compound. The ¹³C NMR spectrum indicated the absence of the characteristic carbon signal for the carbamate group. The remaining ¹³C NMR spectral data were very similar to those of **12**. These data confirmed the structure of 19-*des*-aminocarbonyl-6,7-*seco*-discodermolide (**13**).

Biological Activity of Discodermolide (1) and Semisynthetic Analogues 2–13. Discodermolide (**1**) and its semisynthetic analogues **2–13** were tested for their in vitro cytotoxicity against cultured murine P-388 leukemia and human lung adenocarcinoma A-549 cell lines.³⁶ These compounds inhibited the in vitro proliferation of the P-388

Table 3. ¹H NMR Data of Discodermolide Analogues **7**, **8**, and **9**^a

position	7 ^b	8 ^c	9 ^c
1			
2			
3	6.42 (s)	6.33 (s)	6.32 (s)
4	2.44 (dq, 9.5, 7.3)	2.44 (dq, 9.8, 7.1)	2.43 (m)
5	4.23 (ddd, 2.5, 9.5, 9.5)	4.32 (ddd, 2.2, 9.8, 9.8)	4.32 (ddd, 2.2, 9.8, 9.8)
6	1.80, 1.60 (2H, m)	1.77, 1.70 (m)	1.80, 1.60 (m)
7	4.45 (ddd, 2, 8.1, 8.0)	4.75 (dd, 8.0, 9.0)	4.72 (ddd, 2.0, 8.0, 8.0)
8	5.38 (dd, 8.0, 10.5)	5.38 (dd, 8.0, 10.5)	5.47 (dd, 8.0, 10.5)
9	5.55 (dd, 10.5, 10.5)	5.44 (dd, 10.5, 10.5)	5.41 (dd, 10.5, 10.1)
10	2.63 (ddq, 6.1, 10.5, 6.9)	2.88 (ddq, 10.5, 6.7, 6.1)	2.73 (ddq, 10.1, 10.1, 6.6)
11	3.08 (dd, 6.1, 6.1)	4.73 (m)	3.12 (m)
12	2.27 (ddq, 6.6, 9.9, 6.1)	2.62 (ddq, 6.1, 6.7, 9.7)	2.43 (m)
13	4.97 (d, 9.7)	4.94 (d, 9.7)	5.12 (d, 9.8)
14			
15	1.75, 1.6 (m)	1.9, 1.6 (m)	1.87, 1.65 (m)
16	1.72 (m)	1.90 (m)	2.05 (m)
17	3.13 (dd, 5.1, 5.3)	3.25 (dd, 5.1, 4.9)	4.79 (dd, 5.5, 5.7)
18	1.75 (m)	1.85 (m)	2.01 (m)
19	4.72 (dd, 4.3, 7.2)	4.71 (dd, 6.0, 6.0)	4.62 (dd, 6.2, 6.1)
20	3.07 (m)	3.00 (ddq, 6.0, 6.7, 10.8)	3.13 (m)
21	5.40 (dd, 10.5, 10.9)	5.34 (dd, 10.8, 10.9)	5.33 (dd, 10.9, 10.5)
22	6.06 (dd, 10.9, 10.9)	6.01 (dd, 10.9, 11.0)	6.03 (dd, 10.9, 11.0)
23	6.66 (ddd, 10.2, 10.9, 16.3)	6.60 (ddd, 10.2, 11.0, 16.5)	6.71 (ddd, 10.2, 11.0, 16.3)
24	5.14 (d, 10.2)	5.11 (d, 10.2)	5.14 (d, 10.2)
24	5.24 (d, 16.3)	5.20 (d, 16.5)	5.21 (d, 16.3)
25	1.81 (s)	1.88 (s)	1.88 (s)
26	1.07 (d, 7.3)	1.09 (d, 7.1)	1.09 (d, 7.3)
27	1.01 (d, 7.2)	0.98 (d, 6.9)	0.98 (d, 6.6)
28	0.88 (d, 6.6)	0.87 (d, 6.7)	0.90 (d, 6.6)
29	1.60 (s)	1.61 (s)	1.62 (s)
30	0.74 (d, 6.2)	0.80 (6.0)	0.71 (d, 7.5)
31	0.75 (d, 6.8)	0.97 (d, 6.7)	0.91 (d, 6.8)
32	0.95 (d, 7.0)	0.99 (d, 6.7)	0.96 (d, 7.0)
35			2.07 (s)
37		2.02 (s)	
NH ₂	5.04 (br s)	4.53 (br s)	4.51 (br s)

^a All spectra run at 500 MHz. Chemical shifts are reported in ppm, and *J* values in Hz. ^b In CD₃CN. ^c In CDCl₃.

cell line, with IC₅₀ (nM) values of **1**, 35; **2**, >8292; **3**, 33.8; **4**, 10; **5**, 309; **6**, >7052; **7**, 33.6; **8**, 519; **9**, >8103; **10**, >7587; **11**, 7076; **12**, 3686, and **13**, >12787, and the A-549 cell line, with IC₅₀ (nM) values of **1**, 13.5; **2**, >8292; **3**, 67.7; **4**, 8.4; **5**, 377; **6**, not determined; **7**, 21.7; **8**, 3128; **9**, >8103; **10**, >7587; **11**, >7133; **12**, 6774, and **13**, >12787. The cytotoxicity data of compounds **2**–**5** indicated that double bonds play a significant role in the activity of discodermolide (**1**). The comparable cytotoxicity value of tetrahydrodiscodermolide (**4**) and discodermolide (**1**) in both cell lines suggested that the terminal conjugated diene system is not contributing significantly to the cytotoxicity of discodermolide. Further saturation of discodermolide (**1**) to hexahydrodiscodermolide (**3**) showed a 5-fold reduction in activity against the A-549 cell line but comparable activity against the P-388 cell line. These activity data indicated that the C-8 *Z*-double bond is not contributing significantly to the cytotoxicity of discodermolide (**1**). The 7-deoxy-hexahydrodiscodermolide (**5**) showed a 10-fold reduction in P-388 activity and a 5-fold reduction in A-549 activity compared to that of hexahydrodiscodermolide (**3**), suggesting the importance of the C-7 hydroxy group for cytotoxicity. A significant reduction of activity in hexahydrodiscodermolide triacetate (**6**) clearly indicated the contribution of hydroxy groups to the folding of the molecule via hydrogen bonds. A remarkable loss of activity is indicated in the fully saturated octahydrodiscodermolide (**2**). These data suggested that the C-13 *Z*-double bond, which partly controls the folding of the discodermolide molecule, is important for cytotoxicity. The series of 3-deoxy compounds **7**–**11** clearly shows the contribution of each hydroxy group toward the cytotoxicity of discodermolide (**1**). The activity

data indicated that the C-3 hydroxy group is not essential for cytotoxicity, the C-7 hydroxy group is partly contributing toward cytotoxicity, and the C-17 hydroxy group is essential for the cytotoxicity of discodermolide. The complete loss of cytotoxicity in the *seco*-compounds **12** and **13** indicated that the combination of C-11 and C-17 hydroxy groups contribute to the optimum cytotoxicity of discodermolide. These results are consistent with our earlier findings^{1,2} that acetylation at C₁₁ and more specifically at C₁₇ caused a dramatic reduction in activity and further support our earlier inference^{1,2} that the C₇ through C₁₇ moiety contributes to the overall cytotoxicity of discodermolide. The complete biological activity profile, including the effects of analogues on tubulin polymerization both within cells and using purified tubulin and G₂/M blocking activity, will be published elsewhere. The details of the P-388 and A-549 assays were described previously.^{1,2}

Experimental Section

General Experiment Procedures. UV spectra were measured with a Hitachi U-3010 spectrophotometer. IR spectra were obtained on a Midac M-1200 with Galactic GRAMS/386 software. 1D and 2D NMR spectra were measured on a Bruker AMX-500 instrument. The ¹H NMR chemical shifts were assigned using a combination of data from COSY and HMQC experiments. Similarly, ¹³C NMR chemical shifts were assigned on the basis of DEPT and HMQC experiments. The HRFABMS and HRESIMS were obtained on Finnigan MAT95Q and MAT95 Hybrid Sector mass spectrometers, respectively, at the Spectroscopic Services Group, University of Florida, Gainesville.

Preparation of 8,13,21,23-Octahydrodiscodermolide (2). Discodermolide (2.0 mg) in absolute EtOH (5.0 mL) and a

Table 4. ¹H NMR Data of Discodermolide Analogues **10**, **11**, **12**, and **13**^a

position	10	11	12	13
1				
2				
3	6.33 (s)	6.30 (s)		
4	2.46 (m)	2.42 (m)		
5	4.30 (ddd, 2.0, 9.9, 9.9)	4.04 (ddd, 2.2, 9.9, 9.9)		
6	1.77 (m)	1.90 (m)		
6	1.67 (m)	1.75 (m)		
7	4.66 (m)	5.65 (ddd, 2.0, 8.2, 9.0)		
8	5.39 (dd, 8.0, 10.5)	5.28 (dd, 9.0, 10.5)	5.92 (d, 9.3)	5.92 (d, 9.3)
9	5.49 (dd, 10.5, 10.5)	5.48 (dd, 10.5, 10.5)	6.64 (dd, 4.0, 9.3)	6.64 (dd, 4.0, 9.3)
10	2.86 (ddq, 6.1, 10.5, 6.9)	2.85 (ddq, 10.5, 6.7, 6.1)	2.54 (ddq, 4.0, 6.0, 7.4)	2.54 (4.0, 6.0, 7.4)
11	4.66 (m)	4.65 (dd, 6.1, 6.1)	3.91 (dd, 6.0, 6.0)	3.92 (dd, 6.0, 6.0)
12	2.46 (m)	2.51 (ddq, 6.1, 6.8, 10.0)	2.75 (ddq, 6.0, 6.7, 10.0)	2.78 (m)
13	4.94 (d, 10.0)	4.95 (d, 10.0)	5.18 (d, 10.0)	5.18 (d, 10.0)
14				
15	1.95, 1.65 (m)	1.8, 1.6 (m)	1.94, 1.85 (m)	1.90, 1.80 (m)
16	2.02 (m)	2.0 (m)	1.88 (m)	1.90 (m)
17	4.77 (dd, 5.1, 5.3)	4.78 (dd, 5.8, 5.7)	3.25 (dd, 5.2, 5.5)	3.42 (dd, 5.2, 5.5)
18	1.96 (m)	1.95 (m)	1.86 (m)	1.80 (m)
19	4.59 (m)	4.60 (dd, 6.0, 6.0)	4.69 (dd, 4.4, 7.7)	3.38 (dd, 4.0, 8.5)
20	3.15 (ddq, 6.0, 7.0, 10.5)	3.14 (ddq, 6.0, 6.6, 10.6)	2.98 ddq, 7.0, 7.7, 10.4)	2.74 (m)
21	5.33 (dd, 10.5, 10.9)	5.33 (dd, 10.6, 10.9)	5.33 (dd, 10.4, 10.5)	5.27 (dd, 10.4, 10.5)
22	6.04 (dd, 10.9, 11.0)	6.03 (dd, 10.9, 11.0)	6.01 (dd, 10.5, 11.0)	6.19 (dd, 10.5, 10.9)
23	6.74 (ddd, 10.2, 11.0, 16.5)	6.72 (ddd, 10.1, 11.0, 16.5)	6.60 (ddd, 10.2, 11.0, 16.3)	6.62 (ddd, 10.2, 10.9, 16.3)
24	5.16(d, 10.2)	5.15 (d, 10.1)	5.10 (d, 10.2)	5.17 (d, 10.2)
24	5.22 (d, 16.5)	5.21 (d, 16.5)	5.21 (d, 16.3)	5.21 (d, 16.3)
25	1.88 (s)	1.88 (s)		
26	1.09 (d, 7.2)	1.10 (d, 7.0)		
27	0.94 (d, 6.6)	0.96 (d, 6.6)	1.14 (d, 7.4)	1.15 (d, 7.4)
28	0.83 (d, 6.7)	0.87 (d, 6.8)	1.01 (d, 6.3)	1.02 (d, 6.3)
29	1.61 (s)	1.60 (s)	1.64 (s)	1.65 (s)
30	0.67 (d, 6.5)	0.68 (d, 6.7)	0.80 (d, 6.5)	0.82 (d, 6.5)
31	0.89 (d, 7.0)	0.90 (d, 6.7)	0.97 (d, 6.8)	0.87 (d, 6.8)
32	0.94 (d, 7.0)	0.96 (d, 6.6)	0.99 (d, 7.0)	0.96 (d, 7.0)
35	2.04 (s)	2.01 (s)		
37	2.08 (s)	2.08 (s)		
39		1.99 (s)		
NH ₂	5.04 (br s)	4.53 (br s)	4.48 (br s)	

^a All spectra run at 500 MHz in CDCl₃. Chemical shifts are reported in ppm, and *J* values in Hz.

catalytic amount of Pt(IV) oxide was allowed to shake in a hydrogen atmosphere under a balloon pressure for 12 h at room temperature. The reaction mixture was filtered, and the filtrate was evaporated under a stream of N₂ to give a colorless residue (~2 mg). The residue was subjected to HPLC on a SiO₂ gel (Lichrosorb 5μ, Phenomenex 250 × 10 mm) column using a mixture of 6% MeOH in CH₂Cl₂ to yield 8,13,21,23-octahydrodiscodermolide (**2**, 1.8 mg).

Preparation of 8,21,23-Hexahydrodiscodermolide (3), 21,23-Tetrahydrodiscodermolide (4), and 7-Deoxy-8,21,23-hexahydrodiscodermolide (5). Discodermolide (34.0 mg) in absolute EtOH (20.0 mL) and a catalytic amount of Pt(IV) oxide was maintained at 10 °C and treated with hydrogen under a balloon pressure for 0.5 h. The reaction mixture was filtered, and the filtrate was evaporated under a stream of N₂ to give a colorless residue (~34 mg). The residue was subjected to HPLC on a SiO₂ gel (Lichrosorb 5μ, 250 × 10 mm) column using a mixture of 7% MeOH in CH₂Cl₂ to yield 8,21,23-hexahydrodiscodermolide (**3**, 21.0 mg), 21,23-tetrahydrodiscodermolide (**4**, 2.0 mg), and 7-deoxy-8,21,23-hexahydrodiscodermolide (**5**, 2.3 mg).

Preparation of 7-Deoxy-8,21,23-hexahydrodiscodermolide-3,11,17-triacetate (6). 7-Deoxy-8,21,23-hexahydrodiscodermolide (1.0 mg) in pyridine (0.3 mL) and acetic anhydride (0.3 mL) was stirred overnight at 23 °C. The solvents were removed in a vacuum, and the resulting gum on purification on a Si Sep-Pak with 1% MeOH/CH₂Cl₂ gave 7-deoxy-8,21,23-hexahydrodiscodermolide-3,11,17-triacetate (**6**, 1.0 mg).

Preparation of 3-Deoxy-discodermolide-2-ene (7). Discodermolide-3-acetate¹ (2.4 mg) was treated with saturated aqueous Na₂CO₃ in ethanol (1:9, 2 mL), and the mixture was

stirred at room temperature for 24 h. The solvent was removed by distillation under reduced pressure. The residue was partitioned between EtOAc (5 mL) and H₂O (5 mL), and the EtOAc-soluble fraction on purification by HPLC (SiO₂, 5 μm, 250 × 10 mm) with 4.5% MeOH-CH₂Cl₂ gave 3-deoxy-discodermolide-2-ene (**7**) as a white solid (1.2 mg).

Preparation of 3-Deoxy-discodermolide-2-en-11-acetate (8). Discodermolide-3,7,11-triacetate¹ (3.0 mg) was treated with saturated aqueous Na₂CO₃ in ethanol (1:9, 2 mL), and the mixture was stirred at room temperature for 24 h. The solvent was removed by distillation under reduced pressure. The residue was partitioned between CH₂Cl₂ (4 mL) and H₂O (4 mL), and the CH₂Cl₂-soluble fraction on purification by HPLC (SiO₂, 5μ, 250 × 10 mm) with 2.5% MeOH-CH₂Cl₂ gave 3-deoxy-discodermolide-2-en-11-acetate (**8**) as a white solid (1.5 mg).

Preparation of 3-Deoxy-discodermolide-2-en-17-acetate (9) and 3-Deoxy-discodermolide-2-en-11,17-diacetate (10). Discodermolide-3,7,11,17-tetraacetate⁵ (**14**, 4.0 mg) was treated with saturated aqueous Na₂CO₃ in ethanol (1:9, 2 mL), and the mixture was stirred at 40 °C for 2 h. The solvent was removed by distillation under reduced pressure. The residue was partitioned between EtOAc (5 mL) and H₂O (5 mL), and the EtOAc-soluble fraction on purification by HPLC (SiO₂, 5μ, 250 × 10 mm) with 3.5% MeOH-CH₂Cl₂ gave 3-deoxy-discodermolide-2-en-17-acetate (**9**) as a white solid (1.0 mg) and 3-deoxy-discodermolide-2-en-11,17-diacetate (**10**) as a white solid (2.0 mg).

Preparation of 3-Deoxy-discodermolide-2-en-7,11,17-triacetate (11). 3-Deoxy-discodermolide-2-en-11,17-diacetate (**10**, 1.0 mg) in pyridine (0.3 mL) and acetic anhydride (0.3 mL) was stirred overnight at 23 °C. The solvents were removed

Table 5. ^{13}C NMR Data of Compounds **1–6**^a

position	1 ^b	2 ^b	3 ^b	4 ^b	5 ^c	6 ^c
1	174.7 (s)	174.8 (s)	174.9 (s)	174.6 (s)	174.2 (s)	172.9 (s)
2	44.1 (d)	44.0 (d)	44.1 (t)	44.0 (d)	43.2 (d)	40.37 (d)
3	73.2 (d)	73.2 (d)	73.2 (d)	73.1 (d)	73.4 (d)	74.9 (d)
4	36.4 (d)	37.9 (d)	38.0 (d)	37.6 (d)	37.1 (d)	35.8 (d)
5	77.6 (d)	78.3 (d)	78.3 (d)	77.6 (d)	80.2 (d)	78.7 (d)
6	42.2 (t)	42.0 (t)	41.9 (t)	42.2 (t)	33.1 (t)	33.1 (t)
7	63.5 (d)	68.1 (d)	68.1 (d)	63.6 (d)	24.9 (t)	24.4 (t)
8	133.9 (d)	36.3 (t)	37.3 (t)	133.8 (d)	30.3 (t)	30.4 (t)
9	133.8 (d)	29.0 (t)	31.8 (t)	133.7 (d)	29.2 (t)	29.2 (d)
10	36.4 (d)	36.8 (d)	36.9 (d)	36.2 (d)	35.7 (d)	34.8 (d)
11	79.8 (d)	80.1 (d)	80.7 (d)	79.6 (d)	80.8 (d)	81.5 (d)
12	37.1 (d)	32.9 (d)	36.2 (d)	36.4 (d)	35.5 (d)	34.5 (d)
13	131.2 (d)	42.3 (t)	131.5 (d)	131.2 (d)	130.0 (d)	129.1 (d)
14	133.9 (s)	28.0 (d)	133.6 (s)	133.8 (s)	133.2 (s)	133.1 (s)
15	36.3 (t)	43.6 (t)	36.9 (t)	36.9 (t)	36.5 (t)	36.4 (d)
16	34.3 (d)	32.8 (d)	33.8 (d)	34.0 (d)	33.2 (d)	31.4 (d)
17	76.0 (d)	76.3 (d)	76.9 (d)	77.0 (d)	77.2 (d)	78.3 (d)
18	38.5 (d)	35.9 (d)	35.7 (d)	35.3 (d)	34.9 (d)	32.7 (d)
19	79.3 (d)	79.4 (d)	79.6 (d)	80.0 (d)	80.2 (d)	80.8 (d)
20	34.8 (d)	36.1 (d)	36.1 (d)	36.1 (d)	35.5 (d)	33.8 (d)
21	134.3 (d)	23.6 (t)	23.6 (t)	23.6 (t)	22.9 (t)	22.9 (t)
22	130.4 (d)	31.9 (t)	130.6 (d)	31.2 (t)	31.4 (t)	31.2 (t)
23	133.3 (d)	29.9 (t)	29.9 (t)	29.9 (t)	27.1 (t)	27.1 (t)
24	118.3 (t)	14.3 (q)	13.6 (t)	14.4 (q)	14.0 (q)	14.0 (t)
25	15.8 (q)	15.7 (q)	15.7 (q)	15.7 (q)	15.8 (q)	16.6 (q)
26	13.1 (q)	12.9 (q)	13.8 (q)	13.0 (q)	12.6 (q)	12.4 (q)
27	19.7 (q)	20.7 (q)	17.6 (q)	19.4 (q)	17.1 (q)	16.9 (q)
28	17.5 (q)	16.3 (q)	16.9 (q)	17.1 (q)	16.4 (q)	16.5 (q)
29	23.3 (q)	16.6 (q)	23.6 (q)	23.4 (q)	23.4 (q)	22.9 (q)
30	15.5 (q)	14.5 (q)	14.3 (q)	14.5 (q)	13.1 (q)	12.7 (q)
31	9.2 (q)	9.8 (q)	9.7 (q)	9.3 (q)	8.7 (q)	9.5 (q)
32	18.2 (q)	13.0 (q)	16.4 (q)	16.5 (q)	15.7 (q)	15.8 (q)
33	158.4 (s)	158.4 (s)	158.4 (s)	158.5 (s)		156.9 (s)
34						171.0 (s)
35						21.0 (s)
36						170.4 (s)
37						20.9 (s)
40						170.3 (s)
41						20.8 (s)

^a All spectra run at 125.7 MHz. Chemical shifts are reported in ppm. ^b In CD_3CN . ^c In CDCl_3 .

in a vacuum, and the resulting gum on purification on a Si Sep-Pak with 1% MeOH/ CH_2Cl_2 gave 3-deoxy-discodermolid-2-en-7,11,17-triacetate (**11**, 1.0 mg) as a white solid.

Preparation of 6,7-*seco*-Discodermolide (12). Discodermolide (5.0 mg) was dissolved in EtOAc (4.0 mL) and treated with excess of MnO_2 (10 mg, Aldrich Chem. Co., 5 μm , ~85% activated). The mixture was stirred at room temperature for 48 h. The product was filtered to remove MnO_2 and evaporated to dryness using a stream of nitrogen. The residue on purification by HPLC (SiO_2 , 5 μm , 250 \times 10 mm) with 3.5% MeOH- CH_2Cl_2 gave 6,7-*seco*-discodermolide (**12**, 3.4 mg) as a white solid.

Preparation of 19-*des*-Aminocarbonyl-6,7-*seco*-discodermolide (13). 6,7-*seco*-Discodermolide (**12**, 1.0 mg) was dissolved in CH_2Cl_2 (1.0 mL) and treated with titanium isopropoxide (15 μL) at room temperature. The mixture was stirred at room temperature for two weeks. The product was partitioned between CH_2Cl_2 and H_2O . The CH_2Cl_2 -soluble layer was evaporated to dryness using a stream of nitrogen. The residue on purification by HPLC (SiO_2 , 5 μm , 250 \times 10 mm) with 1.5% MeOH- CH_2Cl_2 gave 19-*des*-aminocarbonyl-6,7-*seco*-discodermolide (**13**, 0.6 mg) as a white powder.

8,13,21,23-Octahydrodiscodermolide (2): $[\alpha]_D^{25}$ -50.0° (c 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 202 (3.25) nm; IR (neat/NaCl) ν_{max} 3365, 2967, 1720, 1602, 1389, 1237, 1028 cm^{-1} ; HRFABMS (glycerol) m/z 602.4618 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{33}\text{H}_{64}\text{NO}_8$, 602.4628).

8,21,23-Hexahydrodiscodermolide (3): $[\alpha]_D^{25}$ -29.2° (c 0.8, MeOH); UV (MeOH) λ_{max} (log ϵ) 202 (3.85) nm; IR (neat/NaCl) ν_{max} 3470, 2968, 1720, 1602, 1382, 1240 cm^{-1} ; HRFABMS (glycerol) m/z 600.4478 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{33}\text{H}_{62}\text{NO}_8$, 600.4475).

21,23-Tetrahydrodiscodermolide (4): UV (MeOH) λ_{max} (log ϵ) 202 (3.25) nm; IR (neat/NaCl) ν_{max} 3490, 2970, 1720,

Table 6. ^{13}C NMR Data of Compounds **7–12**^a

position	7 ^b	8 ^c	9 ^c	10 ^c	11 ^c	12 ^c
1	166.2 (s)	165.5 (s)	165.5 (s)	165.4 (s)	164.8 (s)	
2	127.5 (s)	127.3 (s)	127.3 (s)	127.3 (s)	127.6 (s)	
3	147.0 (d)	145.6 (d)	145.6 (d)	145.5 (d)	144.8 (d)	
4	34.8 (d)	35.0 (d)	35.0 (d)	34.5 (d)	34.1 (d)	
5	80.7 (d)	79.9 (d)	79.9 (d)	79.9 (d)	79.4 (d)	
6	41.8 (t)	40.6 (t)	40.6 (t)	40.5 (t)	38.4 (t)	
7	63.1 (d)	63.4 (d)	63.4 (d)	63.4 (d)	66.8 (d)	164.0 (s)
8	134.1 (d)	132.2 (d)	132.2 (d)	132.3 (d)	128.1 (d)	119.8 (d)
9	133.9 (d)	133.9 (d)	133.9 (t)	133.7 (d)	135.2 (d)	151.1 (d)
10	36.2 (d)	34.7 (d)	34.7 (d)	34.5 (d)	35.0 (d)	34.3 (d)
11	79.8 (d)	80.2 (d)	80.2 (d)	80.3 (d)	80.2 (d)	87.3 (d)
12	37.0 (d)	34.0 (d)	34.0 (d)	34.0 (d)	33.7 (d)	30.3 (d)
13	131.1 (d)	128.4 (d)	128.9 (d)	128.9 (d)	129.0 (d)	128.2 (d)
14	133.1 (s)	133.9 (s)	133.5 (s)	133.5 (s)	133.2 (s)	134.9 (s)
15	36.0 (t)	36.1 (t)	35.4 (t)	35.4 (t)	35.5 (t)	35.8 (t)
16	34.3 (d)	32.8 (d)	31.9 (d)	31.9 (d)	31.9 (d)	33.2 (d)
17	75.9 (d)	75.7 (d)	77.3 (d)	77.3 (d)	77.9 (d)	76.0 (d)
18	38.5 (d)	37.4 (d)	36.5 (d)	36.5 (d)	36.4 (d)	37.3 (d)
19	79.4 (d)	78.6 (d)	77.8 (d)	78.1 (d)	78.0 (d)	78.9 (d)
20	34.6 (d)	34.3 (d)	34.0 (d)	34.0 (d)	34.0 (d)	34.6 (d)
21	134.0 (d)	133.8 (d)	133.8 (d)	132.9 (d)	132.9 (d)	133.4 (d)
22	130.5 (d)	129.9 (d)	130.0 (d)	130.3 (d)	130.2 (d)	129.8 (d)
23	133.6 (d)	132.3 (d)	132.3 (d)	132.3 (d)	132.2 (d)	132.0 (d)
24	118.3 (t)	117.8 (t)	118.1 (t)	118.2 (t)	118.2 (t)	117.8 (t)
25	16.9 (q)	16.6 (q)	16.8 (q)	16.8 (q)	16.7 (q)	
26	16.7 (q)	16.3 (q)	16.6 (q)	16.6 (q)	16.3 (q)	
27	19.7 (q)	17.4 (q)	17.5 (q)	17.5 (q)	17.5 (q)	17.4 (q)
28	17.5 (q)	16.8 (q)	16.8 (q)	16.8 (q)	16.8 (q)	15.8 (q)
29	23.2 (q)	23.1 (q)	22.9 (q)	22.9 (q)	22.9 (q)	23.1 (q)
30	15.6 (q)	13.8 (q)	14.0 (q)	14.0 (q)	13.8 (q)	13.6 (q)
31	8.9 (q)	8.9 (q)	9.4 (q)	9.4 (q)	9.4 (q)	8.6 (q)
32	18.1 (q)	18.6 (q)	18.7 (q)	18.7 (q)	15.5 (q)	17.4 (q)
33	158.3 (s)	156.9 (s)	156.7 (s)	156.7 (s)	156.7 (s)	158.0 (s)
34			171.0 (s)	171.0 (s)	171.0 (s)	
35			20.9 (q)	20.9 (q)	20.9 (q)	
36		170.8 (s)		170.7 (s)	170.6 (s)	
37		21.0 (q)		21.0 (q)	21.0 (q)	
38						169.8 (s)
39						21.2 (q)

^a All spectra run at 125.7 MHz. Chemical shifts are reported in ppm. ^b In CD_3CN . ^c In CDCl_3 .

1602, 1382, 1235 cm^{-1} ; HRFABMS (glycerol) m/z 598.4338 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{33}\text{H}_{60}\text{NO}_8$, 598.4346).

7-Deoxy-8,21,23-hexahydrodiscodermolide (5): $[\alpha]_D^{25}$ -19.8° (c 0.3, MeOH); UV (MeOH) λ_{max} (log ϵ) 202 (3.25) nm; IR (neat/NaCl) ν_{max} 3480, 2970, 1720, 1602, 1372, 1240 cm^{-1} ; HRFABMS (glycerol) m/z 584.4512 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{33}\text{H}_{62}\text{NO}_7$, 584.4526).

7-Deoxy-8,21,23-hexahydroxydiscodermolide-3,11,17-triacetate (6): $[\alpha]_D^{25}$ -12.1° (c 0.01, MeOH); UV (MeOH) λ_{max} (log ϵ) 202 (3.30) nm; IR (neat/NaCl) ν_{max} 3490, 2967, 1737, 1720, 1602, 1370, 1244 cm^{-1} ; HRFABMS (glycerol) m/z 710.4827 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{35}\text{H}_{68}\text{NO}_{10}$, 710.4842).

3-Deoxy-discodermolid-2-ene (7): $[\alpha]_D^{25}$ 20.0° (c 0.04, MeOH); UV (MeOH) λ_{max} (log ϵ) 237 (3.07), 222 (3.37) nm; IR (neat/NaCl) ν_{max} 3426, 1705, 1404, 1249 cm^{-1} ; HRFABMS (glycerol) m/z 576.3931 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{33}\text{H}_{54}\text{NO}_7$, 576.3955).

3-Deoxy-discodermolid-2-en-11-acetate (8): $[\alpha]_D^{25}$ 5.1° (c 0.04, MeOH); UV (MeOH) λ_{max} (log ϵ) 237 (3.07), 220 (3.40) nm; IR (neat/NaCl) ν_{max} 3490, 2669, 1737, 1707, 1606, 1372, 1240, 1045 cm^{-1} ; HRFABMS (glycerol) m/z 618.4036 $[\text{M} + \text{H}]^+$ (calcd $\text{C}_{35}\text{H}_{56}\text{NO}_8$, 618.4005).

3-Deoxy-discodermolid-2-en-17-acetate (9): $[\alpha]_D^{25}$ 6.2° (c 0.03, MeOH); UV (MeOH) λ_{max} (log ϵ) 237 (3.07), 220 (3.40) nm; IR (neat/NaCl) ν_{max} 3480, 2970, 1736, 1707, 1605, 1374, 1243, 1045 cm^{-1} ; HRESIMS m/z 618.4037 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{35}\text{H}_{56}\text{NO}_8$, 618.4007).

3-Deoxy-discodermolid-2-en-11,17-diacetate (10): $[\alpha]_D^{25}$ 3.3° (c 0.03, MeOH); UV (MeOH) λ_{max} (log ϵ) 237 (3.07), 221 (3.41) nm; IR (neat/NaCl) ν_{max} 3490, 2973, 1736, 1707, 1374, 1243, 1045 cm^{-1} ; HRESIMS m/z 682.3896 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{37}\text{H}_{57}\text{NNaO}_9$, 682.3924).

3-Deoxy-discodermolid-2-en-7,11,17-triacetate (11): UV (MeOH) λ_{max} (log ϵ) 237 (3.07), 221 (3.41) nm; IR (neat/NaCl)

ν_{\max} 3490, 2970, 1737, 1707, 1372, 1240, 1045 cm^{-1} ; HRESIMS m/z 702.4204 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{39}\text{H}_{60}\text{NO}_{10}$, 702.4213).

6,7-*seco*-Discodermolide (12): $[\alpha]_D^{25}$ 37.5° (*c* 0.08, MeOH); UV (MeOH) λ_{\max} (log ϵ) 238 (3.94), 220 (4.29) nm; IR (neat/NaCl) ν_{\max} 3367, 2928, 1705, 1691, 1384, 1328, 1235 cm^{-1} ; HRFABMS (3-nitrobenzyl alcohol) m/z 434.2911 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{25}\text{H}_{40}\text{NO}_5$, 434.2906).

19-*des*-Aminocarbonyl-6,7-*seco*-discodermolide (13): UV (MeOH) λ_{\max} (log ϵ) 238 (3.94), 219 (4.29) nm; IR (neat/NaCl) ν_{\max} 3367, 2928, 1705, 1603, 1384, 1328, 1235 cm^{-1} ; ^{13}C NMR (CDCl_3 , 125.7 MHz) δ 164.0 (s, C-7), 119.8 (d, C-8), 151.0 (d, C-9), 34.3 (d, C-10), 87.3 (d, C-11), 30.3 (d, C-12), 128.2 (d, C-13), 134.9 (s, C-14), 35.8 (t, C-15), 32.3 (d, C-16), 76.0 (d, C-17), 37.0 (d, C-18), 78.9 (d, C-19), 34.2 (d, C-20), 133.4 (d, C-21), 129.8 (d, C-22), 132.0 (d, C-23), 117.8 (t, C-24), 17.4 (q, C-27), 15.8 (q, C-28), 23.1 (q, C-29), 13.6 (q, C-30), 8.6 (q, C-31), 17.4 (q, C-32); HRFABMS (3-nitrobenzyl alcohol) m/z 391.2858 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{24}\text{H}_{39}\text{O}_4$, 391.2840).

Acknowledgment. We thank S. A. Pomponi and J. K. Reed for the collection and identification of the sponge *Discodermia* sp. and R. K. Boeckman, Jr., University of Rochester, for helpful discussions. We also thank Patricia Linley for her assistance with the cytotoxicity assays, D. Powell of the University of Florida, Gainesville, for the HRMS measurements, and L. McNulty, Florida Atlantic University, Jupiter campus, for allowing us to use the polarimeter. This research was funded in part by a grant from the National Institutes of Health/National Cancer Institute # CA 74227. This is Harbor Branch Oceanographic Institution contribution no. 1486.

References and Notes

- Gunasekera, S. P.; Longley, R. E.; Isbrucker, R. A. *J. Nat. Prod.* **2001**, *64*, 171–174.
- Isbrucker, R. A.; Gunasekera, S. P.; Longley, R. E. *Cancer Chemother. Pharmacol.* **2001**, *48*, 29–36.
- Gunasekera, S. P.; Longley, R. E. U.S. Patent 6,127,406, 2000.
- Gunasekera, S. P.; Paul, G. P.; Longley, R. E.; Isbrucker, R. A.; Pomponi, S. A. *J. Nat. Prod.* **2002**, in press.
- Gunasekera, S. P.; Gunasekera, M.; Longley, R. E.; Schulte, G. K. *J. Org. Chem.* **1990**, *55*, 4912–4915 (correction: **1991**, *56*, 1346).
- Longley, R. E.; Gunasekera, S. P.; Faherty, D.; McLane, J.; Dumont, F. *Ann. N. Y. Acad. Sci.* **1993**, *696*, 94–107.
- Longley, R. E.; Caddigan, D.; Harmody, D.; Gunasekera, M.; Gunasekera, S. P. *Transplantation* **1991**, *52*, 650–656.
- Longley, R. E.; Caddigan, D.; Harmody, D.; Gunasekera, M.; Gunasekera, S. P. *Transplantation* **1991**, *52*, 656–661.
- ter Haar, E.; Kowalski, R. J.; Hamel, E.; Lin, C. M.; Longley, R. E.; Gunasekera, S. P.; Rosenkranz, H. S.; Day, B. W. *Biochemistry* **1996**, *35*, 243–250.
- Hung, G. T.; Chen, J.; Schreiber, S. L. *Chem. Biol.* **1996**, *3*, 287–293.
- Kowalski, R. J.; Giannakakou, P.; Gunasekera, S. P.; Longley, R. E.; Day, B. W.; Hamel, E. *Mol. Pharm.* **1997**, *52*, 613–622.
- Balachandran, R.; ter Haar, E.; Welsh, M. J.; Grant, S. G.; Day, B. W. *Anticancer Drugs* **1998**, *6*, 67–76.
- Balachandran, R.; Grant, S. G.; Welsh, M. J.; Day, B. W. *Breast J.* **1998**, *4*, 409–419.
- Nerenberg, J. B.; Hung, D. T.; Somers, P. K.; Schreiber, S. L. *J. Am. Chem. Soc.* **1993**, *115*, 12621–12622.
- Hung, D. T.; Nerenberg, J. B.; Schreiber, S. L. *J. Am. Chem. Soc.* **1996**, *118*, 11054–11080.
- Paterson, I.; Florence, G. J.; Gerlach, K.; Scott, J. P. *Angew. Chem.* **2000**, *39*, 377–380.
- Smith, A. B., III; Qui, Y.; Jones, D. R.; Kobayashi, K. *J. Am. Chem. Soc.* **1995**, *117*, 12011–12012.
- Marshall, J. A.; Johns, B. A. *J. Org. Chem.* **1998**, *63*, 7885–7892.
- Harried, S. S.; Yang, G.; Strawn, M. A.; Myles, D. C. *J. Org. Chem.* **1997**, *62*, 6098–6099.
- Clark, D. L.; Heathcock, C. H. *J. Org. Chem.* **1993**, *58*, 5878–5879.
- Paterson, I.; Wren, S. P. *J. Chem. Soc., Chem. Commun.* **1993**, 1790–1792.
- Golec, J. M. C.; Jones, S. D. *Tetrahedron Lett.* **1993**, *34*, 8159–8162.
- Evans, P. L.; Golec, J. M. C.; Gillespie, R. J. *Tetrahedron Lett.* **1993**, *34*, 8163–8166.
- Golec, J. M. C.; Gillespie, R. J. *Tetrahedron Lett.* **1993**, *34*, 8167–8168.
- Yang, G.; Myles, D. C. *Tetrahedron Lett.* **1994**, *35*, 1313–1316.
- Yang, G.; Myles, D. C. *Tetrahedron Lett.* **1994**, *35*, 2503–2504.
- Paterson, I.; Schlapbach, A. *Synlett.* **1995**, *5*, 498–500.
- Miyazawa, M.; Oonuma, K.; Maruyama, M.; Miyashita, M. *Chem Lett.* **1997**, 1191–1192.
- Marshall, J. A.; Zhi-Hui, L.; Johns, B. A. *J. Org. Chem.* **1998**, *63*, 817–823.
- Evans, D. A.; Halstead, D. P.; Allison, B. D. *Tetrahedron Lett.* **1999**, *40*, 4461–4462.
- Filla, S. A.; Song, J. J.; Chen, L.; Masamune, S. *Tetrahedron Lett.* **1999**, *40*, 5449–5453.
- Misske, A. M.; Hoffmann, H. M. R. *Tetrahedron* **1999**, *55*, 4315–4324.
- Smith, A. B., III; Kaufman, M. D.; Beachamp, T. J.; LaMarche, M. J.; Arimoyo, H. *Org. Lett.* **2000**, *2*, 1983.
- Paterson, I.; Florence, G. J.; Gerlach, K.; Scott, J. P.; Sereinig, N. *J. Am. Chem. Soc.* **2001**, *123*, 9535–9544.
- Smith, A. B., III; LaMarche, M. J.; Falcone-Hindley, M. *Org. Lett.* **2001**, *3*, 695–698.
- Gunasekera, S. P.; Longley, R. E.; Isbrucker, R. A.; Paul, G.; Pomponi, S. A.; Wright, A. E. Serial number 09/796,175, U.S. Patent filed on February 28, 2001.

NP0203234