

Marine Natural Products as Novel Antioxidant Prototypes

Satoshi Takamatsu,[†] Tyler W. Hodges,[‡] Ira Rajbhandari,[‡] William H. Gerwick,[§] Mark T. Hamann,^{†,‡} and Dale G. Nagle^{*,†,‡}

National Center for Natural Products Research, Department of Pharmacognosy, and Research Institute of Pharmaceutical Sciences, School of Pharmacy, University of Mississippi, University, Mississippi 38677-1848, and College of Pharmacy, Oregon State University, Corvallis, Oregon 97331

Received August 30, 2002

Pure natural products isolated from marine sponges, algae, and cyanobacteria were examined for antioxidant activity using a 2,2-diphenyl-1-picrylhydrazyl radical (DPPH) solution-based chemical assay and a 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA) cellular-based assay. The DCFH system detects only antioxidants that penetrate cellular membranes. Potent antioxidants were identified and the results from each system compared. The algal metabolites cymopol (**1**), avrainvilleol (**3**), and fragilamide (**4**), and the invertebrate constituent puupehenone (**5**) showed strong antioxidant activity in both systems. Several compounds were active in the DPPH assay but significantly less active in the DCFH system. The green algal metabolite 7-hydroxycymopol (**2**) was isolated from *Cymopolia barbata* and its structure determined. Compound **2** was significantly less active in the DCFH system than cymopol (**1**). The sponge metabolites (1*S*)-(+)-curcuphenol (**6**), aaptamine (**7**), isoaptamine (**8**), and curcudiol (**9**) and the cyanobacterial pigment scytonemin (**10**) showed strong antioxidant activity in the DPPH assay, but were relatively inactive in the DCFH system. Thus, cellular uptake dramatically affects the potential significance of antioxidants discovered using only the DPPH assay. The apparent "proantioxidants" hormothamnione A diacetate (**11**) and *Laurencia* monomer diacetate (**12**) require metabolic activation for antioxidant activity. Significant advantages are achieved using both a solution- and cellular-based assay to discover new antioxidants.

Reactive oxygen species (ROS) and oxidative stress play an important role in the etiology and progression of major human degenerative diseases.¹ This realization has sparked great interest in substances that act as endogenous and exogenous antioxidants. Many solution-based chemical antioxidant assay systems, such as the 2,2-diphenyl-1-picrylhydrazyl radical (DPPH[•]) assay, have been reported.² However, it is also important to evaluate the effects of antioxidants within living cells. Fluorescent technology has made it possible to evaluate antioxidants in living cells using specific probes such as 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA).³ This cell-based fluorescent method is useful to directly examine the ability of natural products to penetrate cell membranes and inhibit ROS in living human cells.

Marine organisms have proven to be rich sources of structurally novel and biologically active secondary metabolites.⁴ These natural products have served as important chemical prototypes for the discovery of new agents for use in the treatment of disease,⁵ probes in molecular pharmacology,⁶ and agrochemicals for pest management.⁷ However, relatively few attempts have been made to explore this vast resource of structurally unique chemistry for new antioxidant prototypes. Marine antioxidant research has largely focused on the antioxidant effects of crude extracts.⁸ The previously characterized marine antioxidant substances are mainly chemicals that are structurally related to plant-derived antioxidants.⁹ These marine antioxidants include pigments such as chlorophylls¹⁰ and carotenoids¹¹ and tocopherol derivatives such as vitamin E and related isoprenoids.¹² Alternatively, certain phenolic substances

produced by marine algae¹³ and UV-absorbing mycosporine-like amino acids found in marine microalgae and invertebrates have been shown to have antioxidant activity.¹⁴

Many studies of plant-derived antioxidants have examined the reduction potential or radical-scavenging effects of natural products in solution-based or TLC-based DPPH autographic chemical assays.^{2,15} Researchers have recently begun to look at the antioxidant effects of natural products in living systems.³ A cell-based method to directly examine the ability of natural products to penetrate living human cells and inhibit reactive oxygen species (ROS)-catalyzed oxidation was used to evaluate marine natural products for their ability to scavenge exogenous ROS induced by TPA (12-*O*-tetradecanoylphorbol 13-acetate) in HL-60 cells.

Results and Discussion

In the present study 130 structurally diverse pure marine natural products, isolated from the research efforts of three marine natural products research groups, were assembled and evaluated in both antioxidant evaluation systems. This panel of natural products included metabolites produced by marine Chlorophyta (green algae), Cnidaria (i.e., soft corals), Chrysophyta (golden-brown algae), Cyanobacteria (blue-green algae), Mollusca (i.e., sea hares), Phaeophyta (brown algae), Porifera (sponges), and Rhodophyta (red algae). This assemblage of secondary metabolites represented a significant level of structural-chemical diversity. Examples of the substances evaluated include the following classes of marine natural products: alkaloids (aplysinopsins, araguspongines, 1*H*-benzo[*de*]1,6-naphthyridines, bromopyrroles, bromotyrosines, brominated indole alkaloids, makaluvamines), lipopeptides (cyclic and linear, malylgamides, microcolins), furocoumarins, oxylipins (eicosanoids, prostanoids, carbocyclic and heterocyclic), peptides (simple and cyclic depsipeptides), phenolics (simple,

* To whom correspondence should be addressed. Tel: 662-915-7143. Fax: 662-915-6975. E-mail: dnagle@olemiss.edu.

[†] National Center for Natural Products Research, University of Mississippi.

[‡] Department of Pharmacognosy, University of Mississippi.

[§] College of Pharmacy, Oregon State University.


Table 1. Marine Natural Products Found to Have Antioxidant Activity

compound	sources (ref)	DPPH ^a activity	DCFH-DA ^b IC ₅₀ (μM)
cymopol (1)	<i>Cymopolia barbata</i> ¹⁶	strong	4.0
7-hydroxycymopol (2)	<i>Cymopolia barbata</i> ¹⁷	strong	>14.6
avrainvilleol (3)	<i>Avrainvillia</i> spp. ¹⁸	strong	6.1
aragilamide (4)	<i>Martensia fragilis</i> ¹⁹	moderate	11
puupehenone (5)	<i>Hyrtilis</i> spp., ²⁰ and other species	strong	27
(1 <i>S</i>)-(+)-curcuphenol (6)	<i>Didiscus</i> spp., ²² and other species	moderate	209
aaptamine (7)	<i>Aptos aptos</i> , ²³ and other species	strong	>55
isooaptamine (8)	<i>Aptos aptos</i> , ²⁴ and other species	strong	>55
(1 <i>S</i>)-curcudiol (9)	<i>Didiscus</i> spp., ²² and other species	moderate	not active
scytonemin (10)	<i>Scytonema</i> spp., ²⁵ and other species	moderate	>23
hormothamnione diacetate (11)	<i>Chrysophaeum taylori</i> ²⁶	not active	18.3
Laurencia monomer diacetate (12)	<i>Laurencia spectabilis</i> ²⁷	not active	49
hormothamnione (13)	<i>Chrysophaeum taylori</i> ²⁸	moderate	>31
carotene (α,β-mixture)		weak	>58
α-tocopherol (vitamin E)		strong	255
ascorbic acid (vitamin C)		strong	9.7

^a TLC-based 1,1-diphenyl-2-picrylhydrazyl radical scavenger antioxidant assay. ^b Cell-based 2',7'-dichlorodihydrofluorescein diacetate antioxidant bioassay.

polyphenolics, polyhalogenated, styrylchromones, quinones and hydroquinones, and terpenoids (oxygenated C₁₀–C₃₀, polyhalogenated C₁₀–C₂₀, cyclic, linear, cembranoids, steroids, and terpene quinones).

Our findings have demonstrated the potential of marine natural products to act as potent antioxidants. By using the solution-based chemical assay along with the cellular-based assay we were able to compare the results in both systems. Three interesting findings were demonstrated by this approach. Antioxidant compounds were discovered that displayed activity in both assay systems (Table 1). These include compounds such as the marine algal compounds cymopol¹⁶ (**1**), 7-hydroxycymopol¹⁷ (**2**), avrainvilleol¹⁸ (**3**), and fragilamide¹⁹ (**4**), and invertebrate metabolites such as puupehenone²⁰ (**5**). These compounds not only



act as antioxidants in solution-based antioxidant assays but can also be taken up by living cells and maintain their activity. Typical dose–response curves obtained for **1** and antioxidant standards α-tocopherol and ascorbic acid on TPA-stimulated hydrogen peroxide DCFH-DA oxidation in HL-60 cells are shown in Figure 1.

Compound **2** was isolated from a CH₂Cl₂–MeOH (1:1, v/v) extract of a Jamaican collection of *Cymopolia barbata* (L.) Lamouroux (Dasycladaceae). Compound **2** has not previously been reported to be a natural product. However, **2** has been reported as an intermediate in the synthesis of





Figure 1. Inhibition of cymopol (**1**), L-ascorbic acid, and α-tocopherol on DCFH-DA oxidation by TPA-stimulated hydrogen peroxide in HL-60 cells. The data are presented as percentages of inhibitions based upon measurement of fluorescence at 530 nm. Values are means of three independent determinations ± SD. Statistical analysis was performed using the Student's *t*-test.

7-hydroxycymopolone diacetate [4'-bromo-2',5'-diacetoxyphenyl]-*E*-(2,6-dimethyl-6-hydroxyhept-2-enyl) ketone.¹⁷ Few spectroscopic data were available to positively confirm the structure of **2** from the original synthetic work. Estrada and co-workers reported only low-field strength ¹H NMR data (60 or 90 MHz) and no ¹³C NMR spectrum from which to make an exact identification. A methoxylated analogue of **2** (3'-methoxy-7-hydroxycymopol) has recently been isolated from a Cuban collection of *C. barbata*.²¹ Since this is the first report of **2** from a natural source, the isolation, structure elucidation, and essential spectroscopic/spectrometric data are reported herein.



A detailed analysis of ¹³C NMR, ¹H, DEPT-135, and HRESIMS data of compound **2** gave the molecular formula of C₁₆H₂₃BrO₃ (molecular weight 342), with five degrees of unsaturation. The ¹³C and DEPT-135° NMR spectra indicated the presence of three methyl groups, four methylenes, three methines, and six quaternary carbons, providing a partial molecular formula of C₁₆H₂₀ and mass of 212 Da. The ¹³C and ¹H NMR spectra were quite similar to those of **1** except for the methylene carbon resonance (43.27 ppm) at position C-6, the quaternary carbon resonance (71.87 ppm) at position C-7, and two fewer sp² carbon resonances. The ¹H–¹³C HMBC spectrum showed the connectivity of the hydroxyl-bearing carbon C-7 (71.87 ppm) to the proton

resonances for the methylene at position H-6 (1.49 ppm) and the geminal methyl group resonances (1.24 ppm). The methylene carbon (43.27 ppm) at position C-6 was coupled to the methyl proton resonances (1.24 ppm) at positions H₃-8 and C-7-Me and also with the methylene proton resonance (2.07 ppm) at position H-4. The ¹H-¹H spin system (¹H-¹H COSY) established the connectivity of the methylene resonances at positions H-4, H-5, and H-6. The methylene carbon (39.97 ppm) at position C-4 was coupled to the methine proton resonance (5.29 ppm) at position H-2 and methyl protons (1.72 ppm) at position C-3-Me and also with the methylene proton (1.51 ppm) at position H-5. The quaternary carbon at C-3 (138.46 ppm) showed long-range ¹H-¹³C couplings (¹H-¹³C HMBC) with the methylene protons at positions H-1, H-4 and the protons of the methyl group attached to C-3. While **1** showed strong antioxidant activity in both assay systems, the increase in polarity associated with the 7-hydroxyl moiety significantly decreased the activity of **2** when examined using the cell-based antioxidant system.


Other marine natural products were shown to be active in the chemical assay, but had no significant activity inside cells. These include the sponge metabolites (1*S*)-(+)-curcuphenol²² (**6**), aaptamine²³ (**7**), isoaptamine²⁴ (**8**), and curcudiol²² (**9**) and the cyanobacterial UV-sunscreen pigment known as scytonemin²⁵ (**10**). This suggests that these



6

7 R = CH₃; R' = H
8 R = H; R' = CH₃

9




10

compounds do not enter the cells due to poor cellular uptake or reduced medium solubility, or perhaps lack the capacity to quench 2',7'-dichlorofluorescein (DCF) fluorescence inside the cell.

Finally, several semisynthetic marine natural product derivatives showed no activity in the solution-based assay, but showed good activity in the cellular-based assay. These include the acetylated semisynthetic styrylchromone derivative hormothamnione A diacetate²⁶ (**11**) and the acetylated red algal metabolite simply referred to as *Laurencia* monomer diacetate²⁷ (**12**). This apparent "cellular activation" is most likely due to the presence of acetylated hydroxyl groups, which upon uptake by the cell, are hydrolyzed by cellular esterases to yield free hydroxyl derivatives, which can serve as proton donors. The natural product hormothamnione A (free phenol-OH groups)²⁸ (**13**), isolated from the marine chrysophyte *Chrysosphaeum taylori*, was also evaluated in both systems. Hormothamnione A (**13**) displayed only mild antioxidant activity in the solution-based chemical assay, but did not show significant activity inside the cells. It is believed that hormothamnione A diacetate (**11**) and the *Laurencia* monomer diacetate (**12**)

act as metabolically activated antioxidants or "proantioxidants" and are more lipophilic and better absorbed into living cells.

11 R = Ac
13 R = H

12

Experimental Section

General Experimental Procedures. The UV spectrum was obtained using a Hewlett-Packard 8453 spectrophotometer. The IR spectrum was obtained using an AATI Mattson Genesis Series FTIR. NMR spectra (¹H, ¹³C, COSY, HMQC, and HMBC) of **2** were recorded in CDCl₃ on a Bruker DRX 500 spectrometer operating at 500 MHz for ¹H and 125 MHz for ¹³C, running gradients and using residual solvent peaks as internal references. Structures of other marine natural products examined were either determined as part of the original referenced research or confirmed by matching NMR spectra and MS data with that of authentic standards. The HREIMS data were acquired on a Finnigan-MAT 95 mass spectrometer (University of Minnesota Department of Chemistry, MS Service Laboratory), and the HRESIMS data were acquired on a Bruker BioAPEX 30es instrument (NCNPR, University of Mississippi). DPPH (1,1-diphenyl-2-picrylhydrazyl) was purchased from Sigma Chemical Co.

Algal Material. The marine green alga *C. barbata* (DNJ.074) was collected at Drax Cove, Jamaica, in July 1998. This sample was frozen and shipped to our (D.G.N.) laboratory storage facility (-20 °C) for storage prior to extraction. A voucher sample was placed in the UM Herbarium (Department of Biology, University of Mississippi).

Extraction and Isolation of 7-Hydroxycymopol (2) from *Cymopolia barbata*. Frozen *C. barbata* was lyophilized (286.34 g) and exhaustively extracted with 50% CH₂Cl₂ in MeOH (v/v) and dried under vacuum to yield 22.38 g of lipid extract. Analysis by 2D-TLC revealed this extract to be chemically rich with a variety of UV-absorbing secondary metabolites that produced charred products upon treatment with ethanolic H₂SO₄ (heat). A portion of crude lipid extract (5.90 g) was fractionated by Si gel vacuum-liquid chromatography with a hexanes-EtOAc-MeOH gradient. The column fraction (300 mg) that eluted with 60% EtOAc in hexanes from the Si gel vacuum-liquid chromatography was separated by Sephadex LH-20 chromatography [50% CH₂Cl₂ in MeOH (v/v)]. Fractions 15-18 were combined (110 mg), and a portion (11 mg) was chromatographed by NP-HPLC (Prodigy Si gel, 5 μm, 4.6 × 250 mm, 30% EtOAc in hexanes (v/v), 2.0 mL/min, photodiode array detection monitored at 254 nm) to yield **2** (6.6 mg, 1.1% yield).

7-Hydroxycymopol (2): oil; UV (MeOH) λ_{max} (log ε) 296 (3.50) nm; IR (film) ν_{max} 2915, 1705, 1595, 1490, 1203 cm⁻¹; ¹H NMR (CDCl₃, 500 MHz) δ 1.24 (6H, s, C-8 and C-7-Me), 1.49 (2H, m, H-6), 1.51 (2H, m, H-5), 1.72 (3H, s, C-3-Me), 2.07 (2H, m, H-4), 3.28 (2H, d, J = 7.2 Hz, H-1), 5.29 (1H, t, J = 7.2 Hz, H-2), 6.81 (1H, s, H-6'), 6.94 (1H, s, H-3'); ¹³C NMR (CDCl₃, 125 MHz) δ 16.34 (CH₃, C-3-Me), 22.61 (CH₂, C-5), 29.06 (CH₂, C-1), 29.36 (C, C-7-Me), 29.36 (C, C-8), 39.97 (CH₂, C-4), 43.27 (CH₂, C-6), 71.87 (C, C-7), 106.88 (C, C2'), 116.94 (CH, C-6'), 118.85 (CH, C-3'), 121.55 (CH, C-3'), 129.04 (C, C-5'), 138.46 (C, C-3), 146.51 (C, C-1'), 148.42 (C, C-4'); HRESIMS m/z 341.0728 [M - H]⁻ (calcd for C₁₆H₂BrO₃, 341.0753).

TLC Autographic Assay for DPPH Radical-Scavenging Effect. Si gel GF plates (10 × 20 cm; 250 μm; Uniplat) were scored to create small individual squares that could accommodate up to 96 samples of test compounds prepared

in 96-well microplates. Pure compounds (prepared in 96-well plates) were dissolved in DMSO at a concentration of 2.0 mg/mL, 4.0 μ g of each was applied in the form of a spot (4–5 mm in diameter) using a multichannel pipet, and the residual DMSO was removed under a vacuum (15–20 min). The radical-scavenging effects of the marine natural products selected were detected on the TLC plates, using a spray reagent composed of a 0.2% (w/v) solution of 1,1-diphenyl-2-picrylhydrazyl radical (DPPH) in MeOH.^{9,10} Plates were observed 30 min after spraying. Active compounds are observed as yellow spots against a purple background. Relative radical-scavenging activity was assigned as “strong” (compounds that produced an intense bright yellow zone), “medium” (compounds that produced a clear yellow spot), “weak” (compounds that produce only a weakly visible yellow spot), or “not active” (compounds that produced no sign of any yellow spot).

Microplate Assay for the Detection of Oxidative Products with DCFH-DA. This method is based on a fluorimetric assay described by Rosenkranz and co-workers.^{3b} Promyelocytic HL-60 cells (1×10^6 cells/mL, ATCC) were suspended in RPMI 1640 medium with 10% FBS and antibiotics at 37 °C in 5% CO₂–95% air. A 125 μ L aliquot of the cell suspension was added to a well of a 96-well plate. After treatment with different concentrations of the test materials for 30 min, cells were stimulated with 100 ng/mL 12-*O*-tetradecanoylphorbol 13-acetate (TPA, Sigma) for 30 min. Then 2',7'-dichlorodihydrofluorescein diacetate (5 μ g/mL) (DCFH-DA, Molecular Probes) was added to the cells, and they were incubated for another 15 min. DCFH-DA is a nonfluorescent probe that diffuses into cells. Cytoplasmic esterases hydrolyze DCFH-DA to 2',7'-dichlorodihydrofluorescein (DCFH). Reactive oxygen species (ROS) generated within HL-60 cells oxidize DCFH to the fluorescent dye 2',7'-dichlorofluorescein (DCF). The ability of the test materials to inhibit exogenous cytoplasmic ROS-catalyzed oxidation of DCFH in HL-60 cells was measured by TPA-treated control incubations with and without the test materials. Levels of DCF were measured using a CytoFluor 2350 Fluorescence Measurement System (Millipore) with excitation wavelength at 485 nm (bandwidth 20 nm) and emission at 530 nm (bandwidth 25 nm).

Acknowledgment. The Natural Resource Conservation Authority, Jamaica, and Discovery Bay Marine Laboratory are gratefully acknowledged for assistance with sample collections. This work was supported in part by the National Center for Natural Products Research (NCNPR) and Research Institute of Pharmaceutical Sciences (RIPS) [University of Mississippi], and the USDA Agricultural Research Service Specific Cooperative Agreement # 58-6408-2-0009. Efforts at OSU were supported by NIH CA52955.

References and Notes

- (a) Breimer, L. H. *Mol. Carcinog.* **1990**, *3*, 188–197. (b) Cerutti, P. A. *Science* **1985**, *227*, 375–380.
- (a) Brand-Williams, W.; Cuvelier, M. E.; Berset, C. *Food Sci. Technol.* **1995**, *28*, 25–30. (b) Cuendet, M.; Hostettmann, K.; Poterat, O.; Dyatmyko, W. *Helv. Chim. Acta* **1997**, *80*, 1144–1152. (c) Calis, I.; Kuruuzum, A.; Demirezer, O. L.; Sticher, O.; Ganci, W.; Ruedi, P. *J. Nat. Prod.* **1999**, *62*, 1101–1105.
- (a) Bass, D. A.; Parce, J. W.; Dechatelet, L. R.; Szejda, P.; Seeds, M. C.; Thomas, M. *J. Immunol.* **1983**, *130*, 1910–1917. (b) Rosenkranz, A. R.; Schmaldienst, S.; Stuhlmeier, K. M.; Chen, W.; Knapp, W.; Zlabinger, G. *J. Immunol. Methods* **1992**, *156*, 39–45. (c) Lee, S. K.; Qing, W. G.; Mar, W.; Luyengi, L.; Mehta, R. G.; Kawanishi, K.; Fong, H. H. S.; Beecher, C. W. W.; Kinghorn, A. D.; Pezzuto, J. M. *J. Biol. Chem.* **1998**, *273*, 19829–19833. (d) Chung, H. S.; Chang, L. C.; Lee, S. K.; Shamon, L. A.; van Breemen, R. B.; Mehta, R. G.; Farnsworth, N. R.; Pezzuto, J. M.; Kinghorn, A. D. *J. Agric. Food Chem.* **1999**, *47*, 36–41. (e) Rajbhandari, I.; Takamatsu, S.; Nagle, D. G. *J. Nat. Prod.* **2001**, *64*, 693–695. (f) Bedir, E.; Tatli, I. I.; Khan, R. A.; Zhao, J.; Takamatsu, S.; Walker, L. A.; Goldman, P.; Khan, I. *J. Agric. Food Chem.* **2002**, *50*, 3150–3155.
- (a) Capon, R. J. *Eur. J. Org. Chem.* **2001**, 633–645, and references therein. (b) Scheuer, P. J. *J. Chem. Educ.* **1999**, *76*, 1075–1079, and references therein. (c) Cragg, G. M.; Newman, D. J.; Snader, K. M. *J. Nat. Prod.* **1997**, *60*, 52–60.
- (a) Blunden, G. In *Pharmacognosy*, 15th ed.; Evans, W. C., Ed.; W. B. Saunders: New York, 2002; pp 115–124, and references therein. (b) Fusetani, N., Ed. *Drugs from the Sea*; S. Karger AG: New York, 2000; p 158, and references therein.
- (a) Tremblay, J.-F. *Chem. Eng. News* **2000**, Jan. 3, 14–15. (b) Coxon, A.; Bestor, T. H. *Chem. Biol.* **1995**, *2*, 119–121.
- (a) El Sayed, K. A.; Dunbar, D. C.; Perry, T. L.; Wilkins, S. P.; Hamann, M. T. *J. Agric. Food Chem.* **1997**, *45*, 2735–2739, and references therein. (b) Schrader, K. K.; Nagle, D. G.; Wedge, D. E. In *Advances in Microbial Toxin Research and its Biotechnological Exploitation*; Upadhyay, R. K., Ed.; Kluwer Academic/Plenum: New York, 2002; pp 171–195, and references therein.
- (a) Nakagawa, H.; Murata, M.; Tachibana, K.; Shiba, T. *Phycol. Res.* **1998**, *46* (Suppl.), 9–12. (b) Guzmán, S.; Gato, A.; Calleja, J. M. *Phytother. Res.* **2001**, *15*, 224–230.
- (a) Poterat, O. *Curr. Org. Chem.* **1997**, *1*, 415–440. (b) Son, B. W.; Kim, J. C.; Choi, H. D.; Kang, J. S. *Arch. Pharm. Res.* **2002**, *25*, 77–79.
- (a) Watanabe, N.; Yamamoto, K.; Ihshikawa, H.; Yagi, A.; Sakata, K.; Brinen, L. S.; Clardy, J. *J. Nat. Prod.* **1993**, *56*, 305–317. (b) Cahyana, A. H.; Shuto, Y.; Kinoshita, Y. *Biosci. Biotech. Biochem.* **1992**, *56*, 1533–1535. (c) Yamamoto, K.; Sakata, K.; Watanabe, N.; Yagi, A.; Brinen, L. S.; Clardy, J. *Tetrahedron Lett.* **1992**, *33*, 2587–2588.
- (a) Choi, J. S.; Lee, W. K.; Cho, Y. J.; Kim, D.-S.; Kim, A.; Chung, H. Y.; Jung, J. H.; Im, K. S.; Choi, W. C.; Choi, H. D.; Son, B. W. *Nat. Prod. Sci.* **2000**, *6*, 122–125. (b) Miki, W.; Otaki, N.; Yokoyama, A.; Izumida, H.; Shimidzu, N. *Experientia* **1994**, *50*, 684–686. (c) Miki, W. *Pure Appl. Chem.* **1991**, *63*, 141–146.
- (a) Kobayashi, M. *J. Chem. Res. (S)* **1994**, 494–495. (b) Yamamoto, Y.; Fujisawa, A.; Hara, A.; Dunlap, W. C. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 13144–13148.
- Fujimoto, K.; Kaneda, T. *Hydrobiologia* **1984**, *116/117*, 111–113.
- (a) Dunlap, W. C.; Shick, J. M.; Yamamoto, Y. *Redox Rep.* **1999**, *4*, 301–306. (b) Dunlap, W. C.; Yamamoto, Y. *Comp. Biochem. Physiol.* **1995**, *112B*, 105–114.
- (a) Cavin, A.; Poterat, O.; Wolfender, J.-L.; Hostettmann, K.; Dyatmyko, W. *J. Nat. Prod.* **1998**, *61*, 1497–1501. (b) Masuda, T.; Matsumura, H.; Oyama, Y.; Takeda, Y.; Jitoe, A.; Kida, A.; Hidaka, K. *J. Nat. Prod.* **1998**, *61*, 609–613. (c) Hwang, B. Y.; Kim, H. S.; Lee, J. H.; Hong, Y. S.; Ro, J. S.; Lee, K. S.; Lee, J. J. *J. Nat. Prod.* **2001**, *64*, 82–84. (d) Ohtani, I. I.; Gotoh, N.; Tanaka, J.; Higa, T.; Gyamfi, M. A.; Aniya, Y. *J. Nat. Prod.* **2000**, *63*, 676–679. (e) Westenburg, H. E.; Lee, K.-J.; Lee, S. K.; Fong, H. H. S.; van Breemen, R. B.; Pezzuto, J. M.; Kinghorn, A. D. *J. Nat. Prod.* **2000**, *63*, 1696–1698.
- Högberg, H.-E.; Thompson, R. H.; King, T. J. *J. Chem. Soc., Perkin Trans. 1* **1976**, 1697–1701.
- Estrada, D. M.; Martin, J. D.; Pérez, C. *J. Nat. Prod.* **1987**, *50*, 735–737.
- Sun, H. H.; Paul, V. J.; Fenical, W. *Phytochemistry* **1983**, *22*, 743–745.
- Kirkup, M. P.; Moore, R. E. *Tetrahedron Lett.* **1983**, *24*, 2087–2090.
- Ravi, B. N.; Perzanowski, H. P.; Ross, R. A.; Erdman, T. R.; Scheuer, P. J.; Finer, J.; Clardy, J. *Pure Appl. Chem.* **1979**, *51*, 1893–1900.
- Dorta, E.; Darias, J.; San Martín, A.; Cueto, M. *J. Nat. Prod.* **2002**, *65*, 329–333.
- (a) McEnroe, F. J.; Fenical, W. *J. Nat. Prod.* **1978**, *34*, 1661–1664. (b) Wright, A. E.; Pomponi, S. A.; McConnell, O. J.; Komoto, S.; McCarthy, P. J. *J. Nat. Prod.* **1987**, *50*, 976–978. (c) El Sayed, K. A.; Yousaf, M.; Hamann, M. T.; Avery, M. A.; Kelly, M.; Wipf, P. *J. Nat. Prod.* **2002**, *65*, 1547–1553.
- Nakamura, H.; Kobayashi, J.; Ohizumi, Y.; Hirata, Y. *Tetrahedron Lett.* **1982**, *23*, 5555–5558.
- Shen, Y. C.; Lin, T. T.; Sheu, J. H.; Duh, C. Y. *J. Nat. Prod.* **1999**, *62*, 1264–1267.
- Proteau, P.; Gerwick, W. H.; Garcia-Pichel, F.; Castenholz, R. *Experientia* **1993**, *49*, 825–829.
- (a) Gerwick, W. H.; Lopez, A.; van Duynne, G. D.; Clardy, J.; Ortiz, W.; Baez, A. *Tetrahedron Lett.* **1986**, *27*, 1979–1982. (b) Ayyangar, N. R.; Khan, R. A.; Deshpande, V. H. *Tetrahedron Lett.* **1988**, *29*, 2347–2348.
- Bernart, M. W.; Gerwick, W. H.; Corcoran, E. E.; Lee, A. Y.; Clardy, J. *Phytochemistry* **1992**, *31*, 1273–1276.
- Gerwick, W. H.; Lopez, A.; van Duynne, G. D.; Clardy, J.; Ortiz, W.; Baez, A. *Tetrahedron Lett.* **1986**, *27*, 1979–1982.