

Structure–Activity Relationships in the Fungistatic Activity against *Botrytis cinerea* of Clovanes Modified on Ring C

Athina Deligeorgopoulou,[†] Antonio J. Macías-Sánchez,[‡] Daniel J. Mobbs,[†] Peter B. Hitchcock,[†] James R. Hanson,^{*,†} and Isidro G. Collado^{*,‡}

The School of Chemistry, Physics and Environmental Science, University of Sussex, Brighton, Sussex, BN1 9QJ, United Kingdom, and Departamento de Química Orgánica, Facultad de Ciencias, Universidad de Cádiz, Apartado 40, 11510, Puerto Real, Cádiz, Spain

Received September 4, 2003

The preparation of clovanes **4**, **5**, **6**, **8**, and **9**, which bear different levels of oxidation on ring C, is described for the first time. The biotransformation of compounds **5**, **6**, and **9** by the fungus *Botrytis cinerea* is investigated, yielding compounds **10**, **11**, and **12**, which are described for the first time, together with compounds **4**–**6**, **8**, and **9**. The evaluation of the fungistatic activity against *B. cinerea* of compounds **6**, **9**, **12**, **18**, **19**, **20**, and **21** is reported. Comparison of these results with previously published data shows first that the inclusion of hydroxyl groups on ring C leads to a decrease in the biological activity and, second, that the presence of a 9 α -hydroxyl group and an alkyl chain at C-2 plays an important role in the fungistatic activity against *B. cinerea* of compounds with a clovane skeleton.


The fungus *Botrytis cinerea* is a serious plant pathogen¹ that has developed resistance to some commercial fungicides.² In addition, some fungicides, such as dicarboximide and procymidone, are persistent enough to be detected in vegetables³ and soil⁴ and in grapes, must, and wine, even several weeks after vinification.⁵ Recently, it has been shown that procymidone has effects similar to flutamide on the sexual differentiation in mice,⁶ and also it causes hepatotoxic disorders in fish.⁴ Consequently, there is interest in the development of novel rationally designed antifungal agents with activity against *B. cinerea*. We have shown⁷ that 2 β -methoxyclovan-9 α -ol (**2**) is an active fungistatic agent against *B. cinerea*. This botryane mimic is readily available by cyclization and rearrangement of caryophyllene oxide (**1**).⁸ Its detoxification by this fungus has also been studied,⁹ and this suggests that active products with the clovane skeleton should not persist for long after their application.

The development of this class of compounds as fungistatic agents against *B. cinerea* required the evaluation of a number of derivatives with clovane skeleton possessing different patterns of oxidation. In this paper we report the preparation of clovane derivatives with different levels of oxidation on ring C, which is topologically coincident with the region of botryane precursors which undergoes an oxidative cleavage to yield botrydial, dihydrobotrydial (**7**),¹⁰ and derivatives.¹¹ We also describe the metabolism of these clovanes by *B. cinerea* and the evaluation of their biological activity against *B. cinerea* in comparison with that of other clovanes.


Results and Discussion

The chemical transformations that were carried out on the clovanes are summarized in Schemes 1 and 2. 2 β -Methoxyclovan-9 α -ol (**2**), obtained from caryophyllene oxide (**1**) by cyclization and rearrangement with TCNE in methanol,⁸ was treated with Jones' reagent¹² to yield 2 β -

Scheme 1



Scheme 2




methoxyclovan-9-one (**3**) in 92% yield.⁹ This ketone was treated with sodium nitrite and boron trifluoride etherate in acetic acid to give compound **4**, in 35% yield (Scheme 1). The ¹H NMR spectrum of this product showed signals at δ_{H} 7.81 (1H, br s), 8.09 (1H, d, $J = 2.1$ Hz), and 2.1 ppm (3H, s), which were assigned to an amide proton, an olefinic proton, and the methyl group of an acetamide, respectively. The ¹³C NMR spectrum revealed a carbonyl resonance at δ_{C} 168.6 ppm and alkene resonances at δ_{C} 135.3 and 131.3 ppm. The high-resolution mass spectrum (m/z 305.1991 [M]⁺) was consistent with the formula C₁₈H₂₇NO₃. The structure for this compound was established by X-ray crystal diffraction, which showed that it was *N*-(2 β -methoxy-9-oxoclovan-10-enyl) acetamide (**4**) (Figure 1).

* To whom correspondence should be addressed. (I.G.C.) Tel: +34-956-016368. Fax: +34-956-016193. E-mail: isidro.gonzalez@uca.es. (J.R.H.) Tel: +44-01273-600755. Fax: +44-01273-677196. E-mail: j.r.hanson@sussex.ac.uk

[†] University of Sussex.

[‡] Universidad de Cádiz.

Scheme 3



Hydrolysis of the acetamide **4** with dilute hydrochloric acid in methanol gave 10-hydroxy-2 β -methoxyclovan-10-en-9-one (**5**) in 68% yield (Scheme 1). The IR spectrum of this product contained absorptions at 3429 and 1668 cm⁻¹, while the ¹H NMR spectrum contained an alkene resonance [δ_{H} 6.52 (1H, d, $J = 2.0$ Hz)] which was assigned to H-11. Reduction of compound **5** with sodium borohydride in dry methanol gave 2 β -methoxyclovan-9 β ,10 β -diol (**6**) in 91% yield (Scheme 1). The IR spectrum contained a hydroxyl absorption (3410 cm⁻¹), while the ¹H NMR spectrum lacked the alkene resonance, but contained two new methine resonances [δ_{H} 3.23 (1H, dd, $J = 6.7, 4.9$ Hz) and δ_{H} 4.17 (1H, d, $J = 4.9$ Hz)] attributed to secondary alcohols. These were assigned to H-10 α and H-9 α , respectively. The stereochemistry of the reduction was established by NOE studies. Irradiation of the signal at δ_{H} 3.23 not only enhanced the signal at δ_{H} 4.17 (4.3%) but also enhanced a signal at δ_{H} 1.14 (3.5%) (1H, d, $J = 13$ Hz), which was assigned to one of the protons at C-12. Hence H-10 was on the α face of the clovane.




Figure 1. X-ray structure of compound **4** drawn by ORTEP.

Table 1. Metabolites of Clovanes by *B. cinerea*

substrate	metabolites
5	10, 11, 7
6	10, 7
9	12, 7

The methoxyketone **3** was converted to the exocyclic epoxide **8** in 76% yield by a Corey–Chaykovsky reaction¹³ using freshly prepared trimethylsulfoxonium iodide.¹⁴ This reaction introduces a methylene group with an equatorial conformation. The product possessed an IR absorption at 916 cm⁻¹ and NMR signals [δ_{H} 2.29 and 2.70 ppm (each 1H, d, $J = 4.5$ Hz (H-17)), δ_{C} 62.92 ppm (s, C-9) and 49.45 ppm (t, C-17)], confirming the structure as 9 α ,17-epoxy-2 β -methoxyclovan (**8**).

Reduction of the epoxide **8** with lithium aluminium hydride in THF gave the C-9 tertiary alcohol **9** in 94% yield (Scheme 2). The ¹H NMR spectrum showed singlets at δ_{H} 3.28, 0.80, 0.86, 0.96, and 1.05 ppm (each 3H, s), corresponding to a methoxy group and four methyl groups. The presence of IR absorption at 3645 cm⁻¹ and a resonance at δ_{C} 74.06 ppm (s) confirmed the presence of a tertiary alcohol. Hence the reduction product had the structure 2 β -methoxy-9 β -methylclovan-9 α -ol (**9**).

The substrates 10-hydroxy-2 β -methoxyclovan-10-en-9-one (**5**), 2 β -methoxyclovan-9 β ,10 β -diol (**6**), and 2 β -methoxy-9 β -methylclovan-9 α -ol (**9**) were each incubated separately with *B. cinerea* for 7–10 days on a shaken culture. In every culture, a certain amount of the characteristic metabolite of *B. cinerea*, dihydrobotrydial (**7**), is produced. The separate feeding of compounds **5**, **6**, and **9** to liquid cultures of *B. cinerea* did not suppress the production of botryane metabolites, at the given levels. The metabolites that were isolated have been tabulated (see Table 1). Their structures were established as follows.

Compound **10** had IR absorptions at 3408 and 1708 cm⁻¹, which were consistent with the presence of an α -hydroxy ketone. The ¹H NMR spectrum showed signals at δ_{H} 3.67 (1H, d, $J = 1.1$ Hz, CHOH), 2.33 (1H, dd, $J = 14.4, 3.3$ Hz, CHHCO), and 2.44 (1H, dd, $J = 14.4, 1.5$ Hz, CHHCO), which were assigned to H-9 α , and to H-11 and H-11', respectively. An NOE effect that was observed between the signals at δ_{H} 3.67 (H-9 α) and 1.60 (1H, ddd, $J = 13.6, 1.0, 1.0$ Hz, H-12) supported the assignment of the β -stereochemistry to the hydroxyl group at C-9. The ¹³C NMR spectrum showed resonances at δ_{C} 82.3 and 212.6 ppm, consistent with the structure 9 β -hydroxy-2 β -methoxyclovan-10-one (**10**) for the metabolite.

Compound **11** possessed ¹H NMR signals at δ_{H} 3.03 and 3.10 (each 1H, d, $J = 17.4$ Hz), which were assigned to H-11 and H-11'. In the ¹³C NMR spectrum there were signals at δ_{C} 181.1 and 174.6 ppm, which were assigned to carboxyl groups at C-9 and C-10. The structure of this metabolite was established by X-ray crystallography as 2 β -methoxy-9,10-secoclovan-9,10-dioic acid (**11**) (Figure 2).

The ¹H NMR spectrum of compound **12** lacked a methoxy signal, but it contained a signal at δ_{H} 3.80 ppm (1H, dd, $J = 9.5, 5.7$ Hz), which was assigned to H-2 α . The spectrum retained four methyl group signals at δ_{H} 0.88, 0.93, 1.04, and 1.13 ppm, while the ¹³C NMR spectrum contained a tertiary alcohol signal at δ_{C} 74.28 ppm, which was assigned to C-9. The structure of this metabolite was established by X-ray crystallography as 9 β -methylclovan-2 β ,9 α -diol (**12**) (Figure 3), and it allowed, in turn, the confirmation of the stereochemistry assigned to the parent compounds **8** and **9**.

In the biotransformation of the clovanes, the methoxyl group appears to be metabolized depending upon the




Figure 2. X-ray structure of compound **11** drawn by ORTEP.




Figure 3. X-ray structure of compound **12** drawn by ORTEP.

number of hydroxyl groups present in the substrate. It did not take place when there were two hydroxyl groups or a hydroxyl and a carbonyl group in the substrate. The reduction of the carbonyl group at C-9 in the biotransformation of compound **5** to compound **10** follows the Prelog rule for the asymmetric reduction of ketones.¹⁵

The antifungal properties of compounds **6**, **9**, and **12** were determined against the growth of *B. cinerea* using the poisoned food technique.¹⁶ The commercial fungicide Euparen was used as a standard for comparison in this test. Several levels of inhibition were observed. We also report here the antifungal properties of compounds **18**, **19**,⁸ **20**,¹⁷ and **21**,⁷ which will be used in the discussion of the structure–activity relationships of clovanes versus *B. cinerea*, together with already published values for compounds **2**, **3**, **13**, **14**, **15**, **16**, and **17**.⁹

As shown in Figure 4 we observed a general decrease in the activity of compounds with the clovane skeleton when a second hydroxyl group was added to the structure even if an ether group is present at C-2. Furthermore, the closer the hydroxyl group is to position C-9 in the clovane skeleton, the lower is the activity of the compound, which decreases in the order **2** > **19** > **20** > **21** > **6**.

One of the detoxification mechanisms of 2 β -methoxyclovan-9 α -ol (**2**) is the oxidation and epimerization of the OH group at C-9.⁹ Therefore, if we can prevent this oxidation or epimerization, the activity might be prolonged. Compound **9** does not undergo biotransformation at C-9. However the biological activity of compounds **9** and **12** was less than that of the equivalent compounds (compounds **2** and **15**) which lack the methyl group at C-9 (Figure 5).

The following conclusions can be drawn from this study of structure–activity relationships. First, when compared




Figure 4. Comparison of fungal growth inhibition (*B. cinerea*) among compounds **2**, **6**, **18**, **19**, **20**, and **21** (100 ppm dose).




Figure 5. Comparison of fungal growth inhibition (*B. cinerea*) among compounds **2**, **9**, **12**, and **15** (100 ppm dose).

to 2 β -methoxyclovan-9 α -ol (**2**), the inclusion of additional hydroxyl groups, especially on ring C, leads to a decrease in biological activity. Second, the presence of a 9 α -hydroxyl group is important for biological activity. Third, the introduction of an alkyl chain at C-2 can significantly increase the potency of this class of compounds as inhibitors of the plant pathogen *B. cinerea*.¹⁸

Experimental Section

General Experimental Procedures. Melting points were measured with a Reichert-Jung Kofler block and are uncorrected. Optical rotations were determined with a Perkin-Elmer 241 polarimeter. IR spectra were recorded on a Perkin-Elmer 881 spectrophotometer. ¹H and ¹³C NMR measurements were obtained on Bruker 300 DPX and Bruker 500 AMX NMR spectrometers with SiMe₄ as internal reference. NMR assignments were made by a combination of 1D and 2D techniques and by comparison with those made for previously described compounds, where appropriate. Mass spectra were recorded on a VG 12-250 spectrometer at 70 eV. HPLC was performed with a Hitachi/Merck L-6270 apparatus equipped with a UV–vis detector (L 4250) and a differential refractometer detector (RI-71). TLC was performed on Merck Kiesegel 60 F₂₅₄, 0.2 mm thick. Silica gel (Merck) was used for column chromatography. Purification by HPLC was accomplished using a Si gel

Table 2. ^{13}C NMR Data (75.5 MHz, CDCl_3) for Compounds **4–6**, **8–11**, and **12**

carbon ^a	2	3	4	5	6	8	9	10	11	14
1	†44.1 s	†44.8 s	†41.7 s	†41.6 s	†43.0 s	†44.3 s	†44.1 s	†49.3 s	†40.7 s	†44.3 s
2	90.1 d	89.1 d	87.3 d	87.3 d	89.8 d	89.5 d	89.6 d	88.3 d	88.2 d	80.6 d
3	44.0 t	44.0 t	44.0 t	43.9 t	43.8 t	41.6 t	38.4 t	43.5 t	38.8 t	37.3 t
4	†37.0 s	†44.4 s	†38.9 s	†39.2 s	†37.4 s	†37.1 s	†37.2 s	†38.6 s	†48.9 s	†37.3 s
5	50.6 d	50.9 d	45.4 d	45.6 d	49.7 d	58.0 d	57.9 d	50.7 d	51.5 d	50.8 d
6	20.5 t	†20.5 t	†18.3 t	†18.2 t	†20.1 t	†20.4 t	†20.9 t	†18.5 t	†19.9 t	†21.1 t
7	33.1 t	†32.5 t	†33.4 t	†33.5 t	†27.1 t	†32.8 t	†33.0 t	†28.7 t	†31.3 t	†32.7 t
8	†34.7 s	†38.3 s	†48.5 s	†48.1 s	†34.6 s	†32.9 s	†37.0 s	†39.8 s	†39.8 s	†37.7 s
9	75.2 d	216.3 s	200.4 s	210.6 s	78.1 d	62.9 s	74.1 s	82.3 d	181.1 s	74.3 s
10	26.0 t	35.7 t	131.3 s	146.5 s	70.2 d	27.8 t	28.5 t	212.6 s	174.6 s	28.4 t
11	26.5 t	†34.0 t	135.3 d	122.8 d	43.1 t	†31.1 t	†32.5 t	47.4 t	43.5 t	†33.1 t
12	36.5 t	†42.7 t	33.4 t	41.7 t	38.1 t	†43.9 t	†43.8 t	41.5 t	37.8 t	†47.6 t
13	25.4 q	26.2 q	24.6 q	25.2 q	25.7 q	25.4 q	25.4 q	26.5 q	28.8 q	24.8 q
14	31.2 q	32.0 q	33.4 q	33.7 q	31.4 q	24.0 q	23.8 q	32.1 q	30.8 q	24.0 q
15	28.4 q	24.9 q	28.4 q	28.5 q	30.1 q	31.2 q	31.3 q	28.3 q	34.4 q	31.6 q
16	58.2 q	57.7 q	57.0 q	55.8 q	58.0 q	50.4 q	50.9 q	57.3 q	56.5 q	
17			168.6 s			49.4 t	24.8 q			24.8 q
18			25.6 q							

^a Notation for attached protons: s = C; d = CH; t = CH₂; q = CH₃; †, ‡ interchangeable signals.

column (Hibar 60, 7 μm , 1 cm wide, 25 cm long). X-ray diffraction data collection was carried out on a Nonius Kappa CCD area detector diffractometer. Structure was determined using the WinGX software package¹⁹ and the SHELXL-97 program.²⁰ ORTEP²¹ was used to generate Figures 1, 2, and 3.

Microorganism and Antifungal Assays. The culture of *Botrytis cinerea* employed in this work, *B. cinerea* (UCA 992), was obtained from grapes from the Domecq vineyard, Jerez de la Frontera, Cádiz, Spain. This culture of *B. cinerea* has been deposited at the Mycological Herbarium Collection (UCA), Facultad de Ciencias, Universidad de Cádiz. Bioassays were performed by measuring inhibition of radial growth on agar medium in a Petri dish. Test compounds were dissolved in EtOH to give a final compound concentration in the culture medium of 50–200 mg L⁻¹. Solutions of test compounds were added to glucose-malt-peptone-agar medium (61 g of glucose-malt-peptone-agar per L, pH 6.5–7.0). The final EtOH concentration was identical in both the control and treated cultures. The medium was poured in 6 or 9 cm diameter sterile plastic Petri dishes, and a 5 mm diameter mycelial disk of *B. cinerea* cut from an actively growing culture was placed in the center of the agar plate. Inhibition of radial growth was measured for 6 days. Every concentration was evaluated in triplicate.

General Culture Conditions. *B. cinerea* (UCA 992) was grown on shaken cultures in a 250 mL conical flask at 25 °C for 3 days on Czapek-Dox medium (70 mL per flask) comprising (per L of distilled water) glucose (40 g), yeast extract (1 g), potassium dihydrogen phosphate (5 g), sodium nitrate (2 g), magnesium sulfate (0.5 g), ferrous sulfate (10 mg), and zinc sulfate (5 mg). The substrate dissolved in EtOH was added to each flask, and the fermentation continued for a further period (see below). The mycelium was filtered and washed with brine and EtOAc. The broth was saturated with NaCl, acidified (pH 2), and extracted with EtOAc. The extracts were separated into acidic and neutral fractions with aqueous sodium hydrogen carbonate. The acid fraction was recovered in EtOAc. The extracts were dried over Na₂SO₄, the solvent was evaporated, and the residues were chromatographed on silica gel in a gradient mixture of petroleum ether–EtOAc of increasing polarity. The acidic fractions were methylated with diazomethane prior to chromatography. Chromatography of the acidic fractions yielded inseparable mixtures.

N-(2 β -Methoxy-9-oxoclov-10-enyl) Acetamide (4). Boron trifluoride complex (18 mL, 0.15 mol, 26.5 equiv, in diethyl ether) was added dropwise to a magnetically stirred solution of 2 β -methoxyclovan-9-one (**3**) (1.4 g, 5.5 mmol, 1 equiv) in acetic acid (46 mL) under nitrogen, at room temperature. NaNO₂ (7.6 g, 0.11 mol, 20 equiv) was added in portions over 3 h. After an additional 2 h, boron trifluoride (18 mL, 0.15 mol, 26.5 equiv, in diethyl ether) was added, and the solution

was stirred for 3 h and then quenched by careful addition of H₂O. The crude reaction mixture was extracted with CH₂Cl₂ and washed with H₂O, and the organic phase was dried over anhydrous Na₂SO₄. The solvent was removed in vacuo to leave a yellow oil, which was purified by column chromatography on silica gel using (3:7) EtOAc in light petroleum as eluent to give *N*-(2 β -methoxy-9-oxoclov-10-enyl) acetamide (**4**) (600 mg, 35%) as white needles (EtOAc): mp 119–121 °C; IR (Nujol) ν_{max} 3293, 1666, 1623 cm⁻¹; ¹H NMR (CDCl₃, 500 MHz) δ 8.09 (1H, d, *J* = 2.1 Hz, H-11), 7.81 (1H, br s, NH), 3.39 (1H, t, *J* = 10.0, 3.6 Hz, H-2 α), 3.35 (3H, s, OCH₃), 2.10 (3H, s, HNCOCCH₃), 1.85 (2H, s, CH₂C(CH₃)₂), 1.13 (3H, s, H-14), 1.09 (3H, s, H-15), 1.05 (3H, s, H-13); ¹³C NMR (75.5 MHz, CDCl₃) (see Table 2); HREIMS *m/z* 305.1992 [M]⁺ (calcd for C₁₈H₂₇NO₃, 305.1991).

X-ray crystal structure of *N*-(2 β -methoxy-9-oxoclov-10-enyl) acetamide (4): white crystal, obtained from EtOAc; C₁₈H₂₇NO₃; crystal size 0.4 × 0.4 × 0.4 mm. Cell parameters: *a* = 9.7008(5) Å, *b* = 12.1777(4) Å, *c* = 14.3201(8) Å, $\alpha = \beta = \gamma = 90^\circ$, *V* = 1691.7(1) Å³, *Z* = 4, *D*_{calc} = 1.20 Mg/m³, *T* = 173(2) K, $\lambda = 0.71073$ Å, space group = *P*2₁2₁2₁ (No. 19), absorption coefficient = 0.08 mm⁻¹, *F*(000) = 664, refinement with 2960 reflections (2364 > 2 σ) led to final *R* indices (all data): *R*1 = 0.0644, *wR*2 = 0.1084; and GOF = 1.017. Crystallographic data (excluding structure factors) for the structure of this compound have been deposited with the Cambridge Crystallographic Data Centre. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK [fax: +44 (0)1223 336033 or e-mail: deposit@ccdc.cam.ac.uk].

10-Hydroxy-2 β -methoxyclovan-10-en-9-one (5). HCl (32 mL, 2 N) was added dropwise to a stirred solution of acetamide **4** (600 mg, 1.96 mmol, 1 equiv) in MeOH (150 mL), and the mixture was heated to reflux for 2 h. After cooling, H₂O was added and the reaction mixture was extracted with Et₂O. The organic phase was washed with H₂O and dried over anhydrous Na₂SO₄. The solvent was evaporated under reduced pressure to give 10-hydroxy-2 β -methoxyclovan-10-en-9-one (**5**) (350 mg, 68%) as an oil: IR (Nujol) ν_{max} 3429, 1668 cm⁻¹; ¹H NMR (CDCl₃, 500 MHz) δ 6.52 (1H, d *J* = 2.0 Hz, H-11), 5.92 (1H, s, OH), 3.34 (1H, t, *J* = 9.8, 3.5 Hz, H-2 α), 3.33 (3H, s, OCH₃), 1.15 (3H, s, H-14), 1.12 (3H, s, H-15), 1.07 (3H, s, H-13); ¹³C NMR (75.5 MHz, CDCl₃) (see Table 2); HREIMS *m/z* 263.1664 [M - H]⁺ (calcd for C₁₆H₂₃O₃, 263.1647).

2 β -Methoxyclovan-9 β ,10 β -diol (6). Sodium borohydride (230 mg, 6.1 mmol, 4.5 equiv) was added to a solution of compound **5** (350 mg, 1.3 mmol, 1 equiv) in dry MeOH (25 mL) at 0 °C, and the reaction was stirred for 30 min. HOAc (2 mL) was added, and the solvent was evaporated under reduced pressure. The residue was recovered in EtOAc, and the resulting organic phase was washed with aqueous Na₂CO₃, H₂O, and brine and dried over anhydrous Na₂SO₄. The solvent

was removed in vacuo to give 2 β -methoxyclovan-9 β ,10 β -diol (**6**) (320 mg, 91%) as an oil: IR (Nujol) ν_{\max} 3410 cm⁻¹; ¹H NMR (CDCl₃, 500 MHz) δ 4.17 (1H, d, *J* 4.9 Hz, H-9 α), 3.23 (1H, dd, *J* 4.9, 6.7 Hz, H-10 α), 3.33 (3H, s, OCH₃), 3.32 (1H, t, *J* = 9.9, 5.5 Hz, H-2 α), 1.14 (1H, d, *J* = 13, H-12 α), 1.05 (3H, s, H-15), 1.03 (3H, s, H-14), 0.86 (3H, s, H-13); ¹³C NMR (75.5 MHz, CDCl₃) (see Table 2); HREIMS *m/z* 266.1871 [M - 2H]⁺ (calcd for C₁₆H₂₆O₃, 266.1881).

9 α ,17-Epoxy-2 β -methoxyclovan-9-one (8). Sodium hydride (120 mg, 60% dispersion in mineral oil) was washed three times with hexane, then dried in vacuo. Dry DMSO (40 mL) was added and the solution stirred at room temperature for 20 min under a nitrogen atmosphere. Freshly prepared trimethylsulfoxonium iodide¹⁴ (620 mg), dissolved in dry DMSO (30 mL), was added and the solution stirred at room temperature for a further 20 min. 2 β -Methoxyclovan-9-one (**3**) (350 mg), dissolved in dry DMSO (30 mL), was added, and the reaction was allowed to stir at room temperature. After 14 h, the solution was poured into H₂O (150 mL) and extracted three times with EtOAc. The combined organic layers were washed three times with brine and then dried over Na₂SO₄. The EtOAc was removed in vacuo, and the yellow oil that remained was adsorbed onto silica and purified by column chromatography over silica. Elution with (5:95) EtOAc in petroleum ether gave 9 α ,17-epoxy-2 β -methoxyclovan-9-one (**8**) (282 mg) as a colorless oil: IR (Nujol) ν_{\max} 2929, 916 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 3.26 (3H, s, OCH₃), 3.24 (1H, dd, *J* = 10.1, 5.6 Hz, H-2 α), 2.70 (1H, d, *J* = 4.5 Hz, H-17), 2.29 (1H, d, *J* = 4.5 Hz, H-17), 0.96 (3H, s, H-14), 0.80 (3H, s, H-15), 0.64 (3H, s, H-13); ¹³C NMR (75.5 MHz, CDCl₃) (see Table 2); EIMS *m/z* 264 [M]⁺ (61), 249 [M - 15]⁺ (27), 232 [M - MeOH]⁺ (100), 217 [M - 15 - MeOH]⁺ (35), 204 (23), 191 (39); HREIMS *m/z* 264.2081 (calcd for C₁₇H₂₈O₂, 264.2089).

2 β -Methoxy-9 β -methylclovan-9 α -ol (9). Compound **8** (260 mg) was taken up in dry THF (100 mL) and was added to LiAlH₄ (40 mg) under nitrogen atmosphere. The reaction was heated under reflux for 10 h and then cooled. The excess hydride was destroyed by careful addition of THF-H₂O (1:1). H₂O (100 mL) was then added, and the reaction mixture was extracted three times with EtOAc. The combined organic layers were washed three times with brine and then dried over Na₂SO₄. The ethyl acetate was removed in vacuo to afford 2 β -methoxy-9 β -methylclovan-9 α -ol (**9**) (245 mg) as a colorless oil: IR (Nujol) ν_{\max} 3456 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 3.28 (3H, s, OCH₃), 3.25 (1H, dd, *J* = 10.1, 5.6, H-2 α), 1.05 (3H, s, H-17), 0.96 (3H, s, H-14), 0.86 (3H, s, H-15), 0.80 (3H, s, H-13); ¹³C NMR (75.5 MHz, CDCl₃) (see Table 2); EIMS *m/z* 266 [M]⁺ (9), 251 [M - 15]⁺ (18), 234 [M - MeOH]⁺ (23), 216 [M - 18 - MeOH]⁺ (16), 201 [M - MeOH - 18 - 15]⁺ (15), 195 (20), 176 (57), 164 (100); HREIMS *m/z* 266.2257 (calcd for C₁₇H₃₀O₂, 266.2245).

Biotransformation of 10-Hydroxy-2 β -methoxyclovan-10-en-9-one (5) by *B. cinerea*. Compound **5** (200 mg) was distributed over 12 flasks of *B. cinerea* and the fermentation grown for a further 7 days. Chromatography of the neutral fraction with a gradient mixture of petroleum ether-EtOAc of increasing polarity gave 9 β -hydroxy-2 β -methoxyclovan-10-one (**10**) (30 mg), 2 β -methoxy-9,10-secoclovan-9,10-dioic acid (**11**) (15 mg), and dihydrobotrydial (**7**)¹⁰ (8 mg).

Biotransformation of 2 β -Methoxyclovan-9 β ,10 β -diol (6) by *B. cinerea*. Compound **6** (220 mg) was distributed over 12 flasks of *B. cinerea* and the fermentation grown for a further 7 days. Chromatography of the neutral fraction with a gradient mixture of petroleum ether-EtOAc of increasing polarity gave 2 β -methoxyclovan-9 β ,10 β -diol (**6**) (20 mg), 9 β -hydroxy-2 β -methoxyclovan-10-one (**10**) (30 mg), and dihydrobotrydial (**7**) (7 mg).

Biotransformation of 2 β -Methoxy-9 β -methylclovan-9 α -ol (9) by *B. cinerea*. Compound **9** (150 mg) was distributed over 42 flasks and the fermentation grown for a further 10 days. Chromatography of the neutral fraction with a gradient mixture of petroleum ether-EtOAc of increasing polarity gave 9 β -methylclovan-2 β ,9 α -diol (**14**) (38 mg) and dihydrobotrydial (**7**) (28 mg).

9 β -Hydroxy-2 β -methoxyclovan-10-one (10): oil; IR (Nujol) ν_{\max} 3480, 1708 cm⁻¹; ¹H NMR (CDCl₃, 500 MHz) δ 0.85 (3H, s, H-13 α), 0.94 (3H, s, H-15), 1.07 (3H, s, H-14 β), 2.33 (1H, dd, *J* = 3.3, 14.4 Hz, CHCO), 2.44 (1H, dd, *J* = 1.5, 14.4 Hz, CHCO), 3.24 (3H, s, OCH₃), 3.35 (1H, dd, *J* = 9.4, 5.4 Hz, H-2 α); ¹³C NMR (CDCl₃, 75.5 MHz) (see Table 2); HREIMS *m/z* 266.1876 [M]⁺ (calcd for C₁₆H₂₆O₃, 266.1881).

2 β -Methoxy-9,10-secoclovan-9,10-dioic acid (11): white needles (EtOAc); mp 178–180°C; IR (Nujol) ν_{\max} 1689 cm⁻¹; ¹H NMR (CDCl₃, 500 MHz) δ 1.08 (3H, s, H-13 α), 1.16 (3H, s, H-15), 1.36 (3H, s, H-14 β), 3.03 (1H, d, *J* = 17.4 Hz, CHH-COOH), 3.10 (1H, d, *J* = 17.4 Hz, CHH-COOH), 3.29 [3H, s, OCH₃]; 3.31 [1H, m, H-2 α]; ¹³C NMR (CDCl₃, 75.5 MHz) (see Table 2); HREIMS *m/z* 280.1650 [M - 18]⁺ (calcd for C₁₆H₂₄O₄ requires 280.1675).

X-ray crystal structure of 2 β -methoxy-9,10-secoclovan-9,10-dioic acid (11): white crystals, obtained from EtOAc; C₁₆H₂₆O₅; crystal size 0.4 × 0.05 × 0.05 mm. Cell parameters: *a* = 6.2588(6) Å, *b* = 13.8493(15) Å, *c* = 22.6116(18) Å, $\alpha = \beta = \gamma = 90^\circ$, *V* = 1960.0(3) Å³, *Z* = 4, *D*_{calc} = 1.30 Mg/m³, *T* = 173(2) K, $\lambda = 0.71073$ Å, space group = *P*2₁2₁ (No. 19), absorption coefficient = 0.35 mm⁻¹, *F*(000) = 816, refinement with 2614 reflections (1959 > 2 σ) led to final *R* indices (all data): *R*1 = 0.101, *wR*2 = 0.167; and GOF = 1.035. Crystallographic data (excluding structure factors) for the structure of this compound have been deposited with the Cambridge Crystallographic Data Centre. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK [fax: +44 (0)1223 336033 or e-mail: deposit@ccdc.cam.ac.uk].

9 β -Methylclovan-2 β ,9 α -diol (12): colorless needles (acetone-petroleum ether); mp 148–150°C; IR (film) ν_{\max} 3364 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 0.88 (3H, s, H-13 α), 0.93 (3H, s, H-15), 1.04 (3H, s, H-14 β), 1.04 (3H, s, H-17), 3.80 (1H, dd, *J* = 5.7, 9.5 Hz, H-2 α); ¹³C NMR (CDCl₃, 75.5 MHz) (see Table 2); HREIMS *m/z* 252.2067 (calcd for C₁₆H₂₈O₂, 252.2089).

X-ray crystal structure of 9 β -methylclovan-2 β ,9 α -diol (12): colorless crystal, obtained from acetone-petroleum ether; C₁₆H₂₈O₂; crystal size 0.3 × 0.1 × 0.1 mm. Cell parameters: *a* = 20.868(7) Å, *b* = 20.868(7) Å, *c* = 10.522(6) Å, $\alpha = \beta = \gamma = 90^\circ$, *V* = 3968(3) Å³, *Z* = 9, *D*_{calc} = 0.95 Mg/m³, *T* = 293(2) K, $\lambda = 1.5418$ Å, space group = *R*₃ (No. 146), absorption coefficient = 0.47 mm⁻¹, *F*(000) = 1260, refinement with 1836 reflections (1316 > 2 σ) led to final *R* indices (all data): *R*1 = 0.111, *wR*2 = 0.189; and GOF = 1.117. Crystallographic data (excluding structure factors) for the structure of this compound have been deposited with the Cambridge Crystallographic Data Centre. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK [fax: +44 (0)1223 336033 or e-mail: deposit@ccdc.cam.ac.uk].

Acknowledgment. This research was supported by grants from D.G.A.E.Y.C.E. (Junta de Andalucía-Spain, JACP-A22/02) and M.C.Y.T. (Spain, AGL2000-0635-C02-01). A.D. and D.M. both thank the EPSRC for a studentship.

References and Notes

- Whealer, B. E. J. *An Introduction to Plant Diseases*, John Wiley & Sons: London, 1969; pp 187–188. (b) Coley-Smith, J. R., Verhoeff, K., Jarvis, W. R., Eds. *The Biology of Botrytis*; Academic Press: London, 1980; pp 42–63.
- L. G. Copping, H. G. Hewitt *Chemistry and Mode of Action of Crop Protection Agents*, The Royal Society of Chemistry, 1998; pp 74–113. (b) Locke, T.; Hollomon, D.; Jones, D.; O'Neil, T. *Fungicide Resistance*, written on behalf of The Fungicide Resistance Action Group-UK (www.pesticides.gov.uk); August 2001. (c) Fungicide Resistance Action Committee website: <http://www.frac.info/frac.html>.
- Apladasaris, P.; Liapis, K. S.; Miliadis, G. E. *J. Agric. Food Chem.* **1994**, *42* (7), 1575–1577.
- Paris-Palacios, S.; Biagianni-Risbourg, S. Vernet, G. B. *Fr. Peche Piscic.* **1998**, (350–1), 547–562.
- Garcia Cazorla, J.; Xirau Vayreda, M. *Am. J. Enol. Viticult.* **1994**, *45*, 338–340. (b) Sala, C.; Fort, F.; Busto, O.; Zamora, F.; Arola, L.; Guasch, J. *J. Agric. Food Chem.* **1996**, *44*, 3668–3671.
- Gray, L. E.; Wolf, C.; Lambright, C.; Mann, P.; Price, M.; Cooper, R. L.; Ostby, J. *Toxicol. Ind. Health* **1999**, *15*, 94–118.
- Collado, I. G.; Hanson, J. R.; Hitchcock, P. B.; Macias-Sánchez, A. J. *J. Org. Chem.* **1997**, *62*, 1965–1969.

- (8) Collado, I. G.; Hanson, J. R.; Macías-Sánchez, A. J. *Tetrahedron* **1996**, *52*, 7961–7972.
- (9) Collado, I. G.; Hanson, J. R.; Macías-Sánchez, A. J.; Mobbs, D. J. *Nat. Prod.* **1998**, *61*, 1348–1351.
- (10) Fehlhaber, H.-W.; Geipel, R.; Mercker, H.-J.; Tschesche, R.; Welmar, K.; Schönbeck, F. *Chem. Ber.* **1974**, *107*, 1720–1730. (b) Lindner, H. J.; Gross, B. v. *Chem. Ber.* **1974**, *107*, 3332–3336.
- (11) Collado, I. G.; Hernández-Galán, R.; Prieto, V.; Hanson, J. R.; Rebordinos, L. G. *Phytochemistry* **1996**, *41*, 513–517. (b) Durán-Patrón, R.; Hernández-Galán, R.; Rebordinos, L. G.; Cantoral, J. M.; Collado, I. G. *Tetrahedron* **1999**, *55*, 2389–2400. (c) Collado, I. G.; Aleu, J.; Hernández-Galán, R.; Durán-Patrón, R. *Curr. Org. Chem.* **2000**, *4*, 1261–1286. (d) Durán-Patrón, R.; Hernández-Galán, R.; Collado, I. G. *J. Nat. Prod.* **2000**, *63*, 182–184. (e) Durán-Patrón, R.; Colmenares, A. J.; Hernández-Galán, R.; Collado, I. G. *Tetrahedron* **2001**, *57*, 1929–1933. (f) Colmenares, A. J.; Durán-Patrón, R.; Hernández-Galán, R.; Collado, I. G. *J. Nat. Prod.* **2002**, *65*, 1724–1726.
- (12) Bowden, K.; Heilbron, I. M.; Jones, E. R. H. *J. Chem. Soc.* **1946**, 39–45.
- (13) Corey, E. J.; Chaykovsky, M. *J. Am. Chem. Soc.* **1965**, *87*, 1353.
- (14) Lampman, G. M.; Koops, R. W.; Olden, C. C. *J. Chem. Educ.* **1985**, *62*, 267–268.
- (15) Prelog, V. *Pure Appl. Chem.* **1964**, *9*, 119–130.
- (16) Patil, I. S.; Kulkarni, S.; Hedge, R. K. *Pesticides* **1986**, 30–31.
- (17) Arantes, S. F.; Hanson, J. R.; Mobbs, D. J. *Phytochemistry* **1999**, *52*, 631–634.
- (18) Collado, I. G.; Hanson, J. R.; Macias-Sanchez, A. J. Spanish Patent ES 2154185, 2001; *Chem. Abstr.* **2001**, *135*, 195688.
- (19) Anon. *J. Appl. Crystallogr.* **1999**, *32*, 837–838.
- (20) Sheldrick, G. M. *SHELX-97*, Program for Crystal Structure Refinement; University of Gottingen: Germany, 1997.
- (21) Farrugia, L. J. ORTEP3 for Windows. *J. Appl. Crystallogr.* **1997**, *30*, 565.

NP030404F