

Three Cyclooctapeptides and One Glycoside from *Microtoena prainiana*

Chong-Qian Li,[†] Bo-Gang Li,[‡] Hua-Yi Qi,[‡] Qi-Lin Li,[‡] Feng-Peng Wang,^{*,†} and Guo-Lin Zhang^{*,†}

West China School of Pharmacy, Sichuan University, Chengdu 610041, People's Republic of China, and Chengdu Institute of Biology, the Chinese Academy of Sciences, Chengdu 610041, People's Republic of China

Received January 28, 2004

Three new cyclic octapeptides, microtoenins A–C (**1–3**), and a new glycoside, 3'''-*O*-methylcrenatoside (**4**), along with several known compounds, were isolated from the ethanolic extract of the stems of *Microtoena prainiana*. Their structures were determined by spectral and chemical evidence. At a concentration of 0.01 mg/mL, 3'''-*O*-methylcrenatoside (**4**), crenatoside (**5**), and isocrenatoside (**6**) inhibited angiotensin converting enzyme (ACE) activity by more than 30%.


The genus *Microtoena* comprises about 21 species, distributed from Southeast Asia to South China. *M. insuavis* (Hance) Prain is used as a folk remedy for treatment of enteritis and diarrhea. *M. prainiana* Diels (Labiatae) is a plant widely distributed in Sichuan, Yunnan, and Guizhou Provinces, China.¹ Chemical research on this genus has not been reported. In our program searching for compounds to reduce high blood pressure, it was found that the EtOH extract of *M. prainiana* exhibited ACE inhibitory activity. The EtOH extract was successively triturated with petroleum ether, CHCl₃, EtOAc, and *n*-BuOH. The EtOAc fraction was found to be the highest in activity.

Repeated chromatography of the inactive CHCl₃ fraction provided cyclic peptides, microtoenins A–C (**1–3**), and indole-3-aldehyde,² scopoletin,³ ethyl caffeate,⁴ 2-anilino-1,4-naphthoquinone,⁵ (4*E*)-2-anilino-4-(phenylimino)naphthalene-1(4*H*)-one,⁶ apigenin,⁷ luteolin,⁷ betulinic acid,⁸ pachypodol,⁹ and kumatakenin.^{10,11} Bioassay-guided isolation of the EtOAc fraction led to active compounds 3'''-*O*-methylcrenatoside (**4**), crenatoside (**5**),¹² and isocrenatoside (**6**),¹³ as well as two inactive compounds, apigenin 7-(6''-*p*-coumaroyl)glucoside¹⁴ and hesperidin.¹⁵ At a concentration of 0.01 mg/mL, compounds **4–6** inhibited ACE activity by more than 30%. Compounds **1–4** were new. The structures of all the compounds were determined on the basis of spectral and chemical methods.

Results and Discussion

Compound **1** was negative to ninhydrin but positive after it was hydrolyzed with 6 N HCl(aq). The molecular formula was assigned as C₄₇H₆₆N₈O₉ on the basis of the ion peak at *m/z* 887.5036 ([M + H]⁺) in the HRESIMS spectrum. The IR absorptions at 3435 and 1635 cm⁻¹ were consistent with amide. The ¹³C NMR spectrum showed eight amide carbonyl signals in the range δ 171.2–174.0 and eight methine signals in the range δ 48.7–62.1. The ¹H NMR spectrum showed six signals between δ 7.58 and 10.87 for amide N–H. The above evidence suggested that compound **1** was a cyclopeptide.

Standard amino acid analysis of the hydrolysate revealed the presence of 1 equiv of alanine (Ala), tyrosine (Tyr), leucine (Leu), and phenylalanine (Phe) and 2 equiv of valine (Val) and proline (Pro), which was consistent with the conclusion from ¹H–¹H COSY, TOCSY, HSQC, and HMBC experiments (Figure S1, Supporting Information).




The fragments Ala-Val¹-Pro¹-Tyr and Leu-Val²-Pro²-Phe in compound **1** were elucidated by the following HMBC correlations: 2-NH and H-2 (Ala)/C-4 (Val¹), 5-NH (Val¹)/C-9 (Pro¹), H-13 (Pro¹)/C-14 (Tyr), and 24-NH (Leu)/C-29 (Val²), 30-NH and H-30 (Val²)/C-34 (Pro²), H-38 (Pro²)/C-39 (Phe). In the ESIMS/MS experiments (Figure S2, Supporting Information), compound **1** showed fragment ions at *m/z* 218 and 472 due to the fragments of Phe-Ala and Val¹-Pro¹-Tyr-Leu, respectively. Therefore, the structure of compound **1** was determined to be cyclo-(Ala-Val¹-Pro¹-Tyr-Leu-Val²-Pro²-Phe). The proposed structure was further supported by the following NOE correlations: 15-



* Corresponding authors. (G.-L.Z.) Tel/Fax: +86-28-85225401. E-mail: zhanggl@cib.ac.cn; (F.-P.W.) Tel/Fax: +86-28-85501368. E-mail: wfp@ucms.edu.cn.

[†] Sichuan University.

[‡] Chengdu Institute of Biology.



3

4 R=CH₃ 5 R=H

6

NH (Tyr)/H-24 (Leu) and 40-NH (Phe)/H-2 (Ala) in the NOESY experiments.

The absolute configuration of the residues in **1** was characterized by acid hydrolysis and analysis of the hydrolysate by Marfey's method.¹⁶ The Marfey's derivatives of DL- and L-amino acids of all these amino acids were prepared as standards and compared with those of the hydrolysate by co-injecting on HPLC. All amino acids of compound **1** were found to have the L configuration.

Compound **2** was negative to ninhydrin but positive after it was hydrolyzed with 6 N HCl(aq). Its molecular formula was assigned as C₄₅H₅₆N₈O₈ according to the ion peak at *m/z* 837.4295 ([M + H]⁺) in the HRESIMS spectrum. IR absorptions at 3429 and 1656 cm⁻¹ suggested the presence of amide moieties. Eight signals in the range δ 169.4–174.6 for amide carbonyl groups, seven signals in the range δ 49.3–61.4 for methines, and a methylene signal at δ 44.0 were recognized in the ¹³C NMR spectrum. The ¹H NMR spectrum showed seven amide protons between δ 7.45 and 10.17. Thus, compound **2** was also a cyclopeptide.

Standard amino acid analysis of the hydrolysate prepared from **2** with 6 N HCl(aq) revealed the presence of 1 equiv of Val, Pro, and glycine (Gly), 2 equiv of Ala, and 3 equiv of Phe, which was in accordance with the results from TOCSY, HSQC, and HMBC experiments (Figure S3, Supporting Information).

The sequence of amino acids in **2** was established as cyclo-(Phe¹-Phe²-Val-Pro-Phe³-Gly-Ala¹-Ala²) by the following HMBC correlations: H-2 (Phe¹)/C-10 (Phe²), H-11 (Phe²)/C-19 (Val), H-20 (Val)/C-24 (Pro), H-28 b (Pro)/C-29 (Phe³), H-30 (Phe³)/C-38 (Gly), H-39 (Gly)/C-40 (Ala¹), 41-

NH (Ala¹)/C-43 (Ala²). In addition, the information provided by NOESY experiments (Figure S3, Supporting Information) and ESIMS/MS data (Figure S4, Supporting Information) confirmed the proposed structure. The configuration of all of the amino acids in **2** was determined to be L by analysis of its hydrolysate with Marfey's method.

Compound **3** was negative to ninhydrin but positive after it was hydrolyzed with 6 N HCl(aq). The ion peak at *m/z* 964.4916 ([M + Na]⁺) in the HRESIMS provided the molecular formula C₄₉H₆₇N₉O₁₀. IR absorptions at 3435 and 1645 cm⁻¹, nine ¹³C NMR signals in the range δ 171.5–175.1 for carbonyls, eight ¹³C NMR signals for methines in the range δ 51.0–63.1, and ¹H NMR signals for seven amide protons between δ 7.19 and 10.10 suggested that **3** was also a cyclopeptide.

Standard amino acid analysis of the hydrolysates of **3** indicated the presence of 1 equiv of Tyr, aspartic acid (Asp), Phe, Leu, and isoleucine (Ile) and 3 equiv of Pro. In the course of the analysis of ¹H–¹H COSY, TOCSY, HSQC, and HMBC data (Figure S5, Supporting Information), it became apparent that **3** incorporated an asparagine (Asn) residue instead of an Asp in view of the molecular formula and the lack of an IR band assignable to a carboxyl group.

The sequence of the amino acid residues was elucidated by the following HMBC correlations: H-2 (Pro¹)/C-6 (Tyr), H-7 (Tyr)/C-15 (Asn), H-16 (Asn)/C-19 (Phe), 20-NH (Phe)/C-28 (Pro²), H-29 (Pro²)/C-33 (Leu), H-34 (Leu)/C-39 (Pro³), H-40 (Pro³)/C-44 (Ile). Consequently, the structure of **3** was determined as cyclo-(Pro¹-Tyr-Asn-Phe-Pro²-Leu-Pro³-Ile). The ESIMS/MS data (Figure S6, Supporting Information) and NOESY experiments supported the conclusion. The amino acids in **3** were determined to have an L configuration by Marfey's method.

Compound **4** was isolated as an amorphous powder. HRESIMS gave an ion peak at *m/z* 659.1949 [M + Na]⁺, indicating a molecular formula of C₃₀H₃₆O₁₅. Alkaline treatment of **4** followed by acid hydrolysis gave glucose and rhamnose. The ¹H NMR signals at δ 7.27 (1H, d, *J* = 1.8 Hz, H-2''), 7.09 (1H, dd, *J* = 8.4, 1.8 Hz, H-6''), 6.79 (1H, d, *J* = 8.4 Hz, H-5''), and 7.56, 6.39 (each 1H, d, *J* = 15.6 Hz, H-7'', H-8'') revealed the presence of an *E*-caffeoyl (feruloyl) moiety. The protons at δ 6.73 (1H, d, *J* = 1.8 Hz, H-2), 6.69 (1H, d, *J* = 7.8 Hz, H-5), 6.61 (1H, dd, *J* = 7.8, 1.8 Hz, H-6) and 4.56 (1H, dd, *J* = 10.8, 2.4 Hz, H-7), 3.94 (1H, dd, *J* = 11.4, 2.4 Hz, H-8a), 3.50 (1H, dd, *J* = 11.4, 10.8 Hz, H-8b) were assigned to a 3,4-dihydroxyphenylethanol moiety. The above assignments were confirmed by the ¹³C NMR spectrum. The above information revealed a close structural similarity to crenatoside (**5**),² with the exception of an extra methoxy group at δ 3.80 (3H, s). This methoxy group could be located at C-3''' by a NOESY cross signal between OCH₃/H-2''' and HMBC correlation between OCH₃ (δ 3.80) and C-3''' (Figure S7, Supporting Information). Therefore, **4** incorporated a feruloyl moiety. The ¹H NMR signals at δ 4.54 (1H, d, *J* = 7.8 Hz) and 4.97 (1H, d, *J* = 1.8 Hz) were respectively assigned to the anomeric protons of β -glucopyranosyl and α -rhamnopyranosyl moieties. The linkage of the sugars and the *E*-feruloyl moiety was determined by the following HMBC correlations (Figure S7, Supporting Information): H-4' (δ 4.90) and C-9'' (δ 166.1), H-1'' (δ 4.97), and C-3' (δ 74.9). The relative configuration of C-7 was determined by the NOESY correlations (Figure S7, Supporting Information) of H-1' with H-8b and of H-7 with H-2' and H-8a. In view of the similar optical rotation of **4** and crenatoside (**5**),¹² compound **4** was thus assigned as 1,2-*O*-[2*S*-(3,4-dihydroxyphenyl)-1,2-

Table 1. ACE Inhibitory Activity of Compounds **4**, **5**, and **6** from *Microtoena prainiana*

sample	inhibition (%)		
	1.0 mg/mL	0.1 mg/mL	0.01 mg/mL
4	99.8	67.5	32.5
5	99.7	75.5	34.6
6	99.3	71.4	35.2
captopril (positive control)			97.7

ethanediy]l]-3-*O*- α -L-rhamnopyranosyl-4-*O*-feruloyl- β -D-glucopyranoside (3'''-*O*-methylcrenatoside).

At a concentration of 0.01 mg/mL, compounds **4–6** inhibited the ACE activity by 32.5%, 34.6%, and 35.2%, respectively (Table 1). Some cytotoxic and anti-HIV cyclopeptides have been reported;^{17–19} thus, cyclopeptides **1–3** were screened in vitro for cytotoxicity with human breast cancer (Bre04), human lung cancer (Lu04), and human neuroma (N04) cell lines, and they exhibited no activity (GI₅₀ > 100 μ g/mL). The antiviral test of compounds **1–6** was measured on Vero cell lines infected with Herpes simplex virus type 2 (HSV-2, strain 333); they were inactive at 250 μ g/mL.

Experimental Section

General Experimental Procedures. Melting points were determined on an XRC-1 melting point apparatus (Scientific Instruments Factory, Sichuan University). Optical rotations were measured on a Perkin-Elmer model 341 polarimeter. UV spectra and IR spectra were carried out on a Perkin-Elmer Lambda 35 UV/vis spectrometer and a Perkin-Elmer Spectrum One FT-IR spectrometer. ¹H NMR, ¹³C NMR, ¹H–¹H COSY, TOCSY, HSQC, HMBC, and NOESY spectra were performed on a Bruker Avance 600 spectrometer. Chemical shift values are in ppm (δ) with TMS as internal standard. Electrospray ionization mass spectra (ESIMS) were acquired with a Finnigan LCQ^{DECA} mass spectrometer. High-resolution electrospray ionization mass spectra (HRESIMS) were obtained on an API Q-STAR PULSAR *i* mass spectrometer. High-performance liquid chromatography (HPLC) was performed using a Perkin-Elmer series 200 pump equipped with a Perkin-Elmer series 200 UV/vis detector. Amino acid analysis was carried out on a Hitachi L-8800 amino acid analyzer. *N*₆-(2,4-Dinitro-5-fluorophenyl)-L-alaninamide (FDAA, Marfey's reagent), hippuric acid-histidine-leucine (HHL), and *O*-phthalaldehyde were purchased from Sigma-Aldrich, Inc., Germany. Silica gel (200–300 mesh) was obtained from Qingdao Ocean Chemical Company, China. MCI was purchased from Mitsubishi Company, Japan. Polyamide was acquired from Shanghai Chemical Reagent Factory, China. C-18 silica gel (RP-18 silica 60, 40–63 μ m) was obtained from Merck & Co., Inc., Germany.

Plant Material. The stems of *M. prainiana* were collected from Tianquan County, Sichuan Province, China, in October 2002 and identified by Professor Fa-Ding Fu at Chengdu Institute of Biology, the Chinese Academy of Sciences (CAS). A voucher specimen (no. TQ-6) is deposited in the Herbarium of Chengdu Institute of Biology, CAS.

Extraction and Isolation. The dried and powdered stems of *M. prainiana* (22 kg) were soaked with 95% EtOH (3 \times 160 L) at room temperature. After concentrating in vacuo, 1500 g of residue was obtained. The residue was suspended in H₂O (5 L) and then triturated successively with petroleum ether (60–90 $^{\circ}$ C) (5 \times 5 L), CHCl₃ (5 \times 3 L), EtOAc (5 \times 5 L), and n-BuOH (5 \times 5 L) to afford corresponding fractions. Each fraction was examined for ACE inhibitory activity, and the EtOAc fraction showed the strongest activity.

The EtOAc fraction (302 g) was chromatographed over a silica gel column (2500 g, CHCl₃–MeOH–H₂O, 100:1:0.02, 50:1:0.02, 20:1:0.02, 10:1:0.02, 5:1:0.02, 0:1:0.02, each 5000 mL, $\emptyset \times l = 12 \times 55$ cm) to afford four subfractions. 3'''-*O*-Methylcrenatoside (**4**) (304 mg, 0.0014% dry weight) and

compound **6** (30 mg, 0.00014%) were isolated from fraction 1 (30 g) using a silica gel column (350 g, CHCl₃–MeOH, 10:1, 3000 mL, 7.5 \times 14 cm) and were each purified by recrystallization from acetone. Fraction 2 (30 g) was chromatographed over silica gel (200 g, EtOAc–MeOH, 20:1, 2000 mL, 6.5 \times 15 cm) to afford **5** (800 mg, 0.0036%). Apigenin 7-(6''-*p*-coumaroyl)glucoside (5 mg, 0.00002%) was obtained from fraction 3 (300 mg) using a silica gel column (15 g, CHCl₃–MeOH–H₂O, 15:1:0.02, 350 mL, 2.5 \times 20 cm). Fraction 4 (3 g) was chromatographed over a polyamide column (70 g, acetone–H₂O, 1:5, 1000 mL, 5 \times 10 cm) to afford hesperidin (3 mg, 0.000014%).

The CHCl₃ fraction (100 g) was chromatographed over MCI gel (500 g) using MeOH–H₂O (7:3, 8:2, 9:1) to afford fraction 5. Fraction 5 (38 g) was chromatographed over silica gel, resulting in nine subfractions (fractions 6–14). Indole-3-aldehyde (20 mg, 0.00009%) and scopoletin (15 mg, 0.00007%) were obtained by recrystallization of fraction 6 (100 mg) and fraction 7 (80 mg) from acetone, respectively. Ethyl caffeate (30 mg, 0.00014%) was isolated from fraction 8 (400 mg). Fraction 9 (240 mg) was separated over a silica gel column, yielding 2-anilino-1,4-naphthoquinone (3 mg, 0.00001%) and (4*E*)-2-anilino-4-(phenylimino)naphthalene-1(4*H*)-one (2 mg, 0.000009%). The separation of fraction 10 (150 mg) using a C₁₈ silica gel column (30 g, MeOH–H₂O, 1:1) afforded apigenin (38 mg, 0.00017%) and luteolin (15 mg, 0.00007%). Fraction 11 (2 g) was chromatographed over silica gel to give betulinic acid (27 mg, 0.00012%). Microtoenin A (**1**) (58 mg, 0.00026%) and microtoenin B (**2**) (30 mg, 0.00014%) were isolated from fraction 12 (2.4 g) using a silica gel column [120 g, petroleum ether (60–90 $^{\circ}$ C)–EtOAc–MeOH, 5:1:0.2, 2000 mL, 5.6 \times 10 cm] and were further purified by recrystallization from acetone. Fraction 13 (35 mg) was chromatographed over silica gel to afford pachypodol (8 mg, 0.00004%) and kumatakenin (3 mg, 0.00001%). Microtoenin C (**3**) (58 mg, 0.00026%) was isolated from fraction 14 (2 g) over a silica gel column [60 g, petroleum ether (60–90 $^{\circ}$ C)–acetone, 1:1, 2000 mL, 5 \times 6.5 cm] and was further purified by recrystallization from acetone.

Microtoenin A (1): white amorphous powder; mp 280–282 $^{\circ}$ C; $[\alpha]_D^{20} -104.8^{\circ}$ (*c* 0.23, MeOH); UV (MeOH) λ_{max} (log ϵ) 206 (4.31) nm; IR (KBr) ν_{max} 3435, 2927, 1635, 1517, 1454 cm⁻¹; ¹H NMR (C₅D₅N, 600 MHz) δ 5.34 (1 H, m, H-2), 1.49 (3H, d, *J* = 6.0 Hz, H-3), 8.94 (1H, d, *J* = 8.4 Hz, 2-NH), 4.69 (1H, dd, *J* = 8.4, 6.0 Hz, H-5), 2.51 (1H, m, H-6), 1.01 (9H, d, *J* = 6.6 Hz, H-7, H-8, H-33), 7.58 (1H, d, *J* = 8.4 Hz, 5-NH), 4.30 (1H, d, *J* = 7.8 Hz, H-10), 2.26 (1H, m, H-11a), 1.04 (1H, overlapped, H-11b), 1.50 (4H, m, H-12, H-37), 3.72 (1H, m, H-13a), 3.55 (2H, m, H-13b, H-38b), 4.95 (1H, overlapped, H-15), 3.20 (2H, m, H-16a, H-41a), 2.81 (1H, m, H-16b), 7.00 (2H, d, *J* = 8.4 Hz, H-18, H-22), 7.09 (2H, d, *J* = 8.4 Hz, H-19, H-21), 11.55 (1H, brs, 20-OH), 10.87 (1H, brs, 15-NH), 5.38 (1H, m, H-24), 2.00 (1H, m, H-25a), 1.79 (2H, m, H-25b, H-26), 0.85 (3H, d, *J* = 6.0 Hz, H-27), 0.91 (3H, d, *J* = 6.0 Hz, H-28), 8.62 (1H, d, *J* = 8.4 Hz, 24-NH), 4.74 (1H, t, *J* = 8.4 Hz, H-30), 2.45 (1H, m, H-31), 1.04 (3H, d, *J* = 6.6 Hz, H-32), 8.11 (1H, d, *J* = 9.0 Hz, 30-NH), 4.00 (1H, d, *J* = 7.8 Hz, H-35), 2.21 (1H, m, H-36a), 0.73 (1H, m, H-36b), 3.65 (1H, m, H-38a), 4.86 (1H, m, H-40), 2.95 (1H, m, H-41b), 7.04 (2H, d, *J* = 7.2 Hz, H-43, H-47), 7.25 (2H, dd, *J* = 7.3, 7.2 Hz, H-44, H-46), 7.20 (1H, overlapped, H-45), 10.83 (1H, brs, 40-NH); ¹³C NMR (C₅D₅N, 150 MHz) δ 174.0 (C, C-1), 48.7 (CH, C-2), 17.8 (CH₃, C-3), 171.2 (C, C-4), 61.3 (CH, C-5), 31.1 (CH, C-6), 19.8 (CH₃, C-7), 17.7 (CH₃, C-8), 171.7 (C, C-9), 62.1 (CH, C-10), 30.5 (CH₂, C-11), 21.9 (CH₂, C-12), 46.4 (CH₂, C-13), 172.2 (C, C-14), 53.8 (CH, C-15), 37.6 (CH₂, C-16), 126.2 (C, C-17), 130.9 (CH, C-18, C-22), 116.6 (CH, C-19, C-21), 158.2 (C, C-20), 173.5 (C, C-23), 51.5 (CH, C-24), 42.6 (CH₂, C-25), 25.2 (CH, C-26), 22.3 (CH₃, C-27), 23.0 (CH₃, C-28), 171.5 (C, C-29), 62.0 (CH, C-30), 31.3 (CH, C-31), 19.9 (CH₃, C-32), 18.5 (CH₃, C-33), 171.3 (C, C-34), 61.9 (CH, C-35), 30.5 (CH₂, C-36), 21.8 (CH₂, C-37), 46.3 (CH₂, C-38), 171.6 (C, C-39), 54.1 (CH, C-40), 38.3 (CH₂, C-41), 136.1 (C, C-42), 129.7 (CH, C-43, C-47), 129.1 (CH, C-44, C-46), 127.6 (CH, C-45); ESIMS *m/z* 909 ([M + Na]⁺, 100), 887 ([M + H]⁺, 15), 885 ([M – H][–], 100); ESIMS/MS on [M + H]⁺ *m/z* 887 (8), 869 (10), 859 (45), 740 (15), 691 (100), 669 (14), 620 (18), 578 (13), 528

(11), 415 (11), 310 (3); ESIMS/MS/MS on $[M + H - \text{Phe}]^+ m/z$ 740 (20), 712 (100), 472 (10), 359 (11), 267 (20), 218 (17), 196 (90), 147 (18); HRESIMS m/z 887.5036 ($[M + H]^+$, $\text{C}_{47}\text{H}_{67}\text{N}_8\text{O}_9$ requires 887.5031).

Microtoenin B (2): white amorphous powder; mp 288–290 °C; $[\alpha]_D^{20} -68.3^\circ$ (c 0.12, MeOH); UV (MeOH) λ_{max} (log ϵ) 211 (4.61) nm; IR (KBr) ν_{max} 3429, 2927, 1656, 1534, 1453 cm^{-1} ; ^1H NMR ($\text{C}_5\text{D}_5\text{N}$, 600 MHz) δ 5.43 (1H, m, H-2), 3.23 (1H, m, H-3a), 3.38 (1H, m, H-3b), 7.45 (3H, d, $J = 7.2$ Hz, H-5, H-9, 20-NH), 7.28 (4H, m, H-6, H-8, H-14, H-18), 7.15 (3H, m, H-7, H-15, H-17), 7.64 (1H, d, $J = 7.2$ Hz, 2-NH), 4.64 (1H, m, H-11), 3.20 (1H, m, H-12a), 3.32 (1H, m, H-12b), 7.37 (3H, m, H-16, H-34, H-36), 7.93 (1H, brs, 11-NH), 4.08 (1H, t, $J = 9.6$ Hz, H-20), 1.96 (1H, m, H-21), 0.88 (3H, d, $J = 6.6$ Hz, H-22), 0.78 (3H, d, $J = 6.6$ Hz, H-23), 3.78 (1H, d, $J = 7.8$ Hz, H-25), 2.04 (1H, q, $J = 6.0$ Hz, H-26a), 0.98 (1H, m, H-26b), 1.52 (1H, m, H-27a), 1.43 (1H, m, H-27b), 3.64 (1H, m, H-28a), 3.53 (1H, t, $J = 10.2$ Hz, H-28b), 4.68 (1H, m, H-30), 3.27 (1H, t, $J = 12.6$ Hz, H-31a), 3.16 (1H, m, H-31b), 7.30 (2H, d, $J = 8.4$ Hz, H-33, H-37), 7.25 (1H, m, H-35), 10.17 (1H, brs, 30-NH), 4.61 (1H, m, H-39a), 3.85 (1H, dd, $J = 16.8, 5.4$ Hz, H-39b), 8.57 (1H, brs, 39-NH), 5.02 (1H, m, H-41), 1.72 (3H, d, $J = 7.2$ Hz, H-42), 8.54 (1H, d, $J = 8.4$ Hz, 41-NH), 4.60 (1H, m, H-44), 1.49 (3H, d, $J = 7.2$ Hz, H-45), 9.03 (1H, d, $J = 3.6$ Hz, 44-NH); ^{13}C NMR ($\text{C}_5\text{D}_5\text{N}$, 150 MHz) δ 171.7 (C, C-1), 53.6 (CH, C-2), 39.2 (CH₂, C-3), 137.2 (C, C-4), 130.7 (CH, C-5, C-9), 128.3 (CH, C-6, C-8), 126.7 (CH, C-7), 171.3 (C, C-10), 58.4 (CH, C-11), 37.4 (CH₂, C-12), 136.7 (C, C-13), 129.7 (CH, C-14, C-18), 129.03 (CH, C-15, C-17), 127.3 (CH, C-16), 173.6 (C, C-19), 58.6 (CH, C-20), 27.0 (CH, C-21), 20.6 (CH₃, C-22), 18.7 (CH₃, C-23), 172.0 (C, C-24), 61.4 (CH, C-25), 30.9 (CH₂, C-26), 21.9 (CH₂, C-27), 46.5 (CH₂, C-28), 174.6 (C, C-29), 53.7 (CH, C-30), 38.2 (CH₂, C-31), 136.5 (C, C-32), 129.8 (CH, C-33, C-37), 129.09 (CH, C-34, C-36), 127.6 (CH, C-35), 169.4 (C, C-38), 44.0 (CH₂, C-39), 173.7 (C, C-40), 49.3 (CH, C-41), 18.0 (CH₃, C-42), 173.1 (C, C-43), 52.1 (CH, C-44), 17.0 (CH₃, C-45); ESIMS m/z 859 ($[M + \text{Na}]^+$, 100), 835 ($[M - \text{H}]^-$, 100); ESIMS/MS on $[M - \text{H}]^- m/z$ 835(3), 817 (20), 791 (7), 710 (5), 693 (15), 660 (3), 634 (17), 606 (11), 558 (100), 540 (2), 492 (6), 345 (5), 288 (2); ESIMS/MS/MS on $[M - 3\text{H} - \text{Phe} - \text{Gly} - \text{Ala}]^- m/z$ 558 (2), 540 (60), 513 (10), 470 (10), 430 (62), 362 (52), 345 (100), 327 (15), 302 (5), 274 (7), 215 (11), 200 (11), 195 (3); HRESIMS m/z 837.4295 ($[M + H]^+$, $\text{C}_{45}\text{H}_{57}\text{N}_8\text{O}_8$ requires 837.4299).

Microtoenin C (3): white amorphous powder; mp 256–258 °C; $[\alpha]_D^{20} -93.8^\circ$ (c 0.13, MeOH); UV (MeOH) λ_{max} (log ϵ) 207 (4.36) nm; IR (KBr) ν_{max} 3435, 2961, 1645, 1517, 1452 cm^{-1} ; ^1H NMR ($\text{C}_5\text{D}_5\text{N}$, 600 MHz) δ 3.87 (1H, m, H-2), 1.74 (2H, m, H-3a, H-42a), 2.33 (2H, m, H-3b, H-36), 1.91 (2H, m, H-4a, H-48b), 1.84 (1H, m, H-4b), 3.41–3.52 (7H, m, H-5a, H-8b, H-17b, H-21a, H-32a, H-43), 3.79 (1H, t, $J = 8.4$ Hz, H-5b), 4.63 (1H, m, H-7), 4.05 (1H, t, $J = 6.6$ Hz, H-8a), 7.28 (2H, d, $J = 7.8$ Hz, H-10, H-14), 7.11–7.19 (5H, m, H-11, H-13, H-24, H-25, H-26), 11.43 (1H, s, 12-OH), 7.19 (1H, overlapped, 7-NH), 5.15 (1H, m, H-16), 3.90 (1H, m, H-17a), 8.57, 7.90 (each 1H, s, 18-NH₂), 10.10 (1H, s, 16-NH), 5.12 (1H, m, H-20), 3.59 (1H, d, $J = 12.6$ Hz, H-21b), 7.34 (2H, d, $J = 7.2$ Hz, H-23, H-27), 8.63 (1H, d, $J = 8.4$ Hz, 20-NH), 4.32 (1H, d, $J = 7.8$ Hz, H-29), 2.41 (1H, q, $J = 6.0$ Hz, H-30a), 1.64 (2H, m, H-30b, H-42b), 1.35 (1H, m, H-31a), 0.76 (1H, m, H-31b), 3.34 (1H, t, $J = 10.2$ Hz, H-32b), 5.10 (1H, m, H-34), 2.04 (1H, m, H-35a), 1.79 (1H, m, H-35b), 0.79 (3H, d, $J = 6.0$ Hz, H-37), 1.11 (3H, d, $J = 6.0$ Hz, H-38), 8.04 (1H, brs, 34-NH), 4.77 (1H, t, $J = 7.8$ Hz, H-40), 2.28 (1H, m, H-41a), 2.13 (1H, m, H-41b), 4.48 (1H, brs, H-45), 2.60 (1H, brs, H-46), 1.07 (3H, d, $J = 6.0$ Hz, H-47), 1.52 (1H, m, H-48a), 0.84 (3H, t, $J = 7.2$ Hz, H-49), 8.91 (1H, brs, 45-NH); ^{13}C NMR ($\text{C}_5\text{D}_5\text{N}$, 150 MHz) δ 174.3 (C, C-1), 61.0 (CH, C-2), 29.5 (CH₂, C-3), 27.2 (CH₂, C-4), 49.1 (CH₂, C-5), 175.1 (C, C-6), 63.1 (CH, C-7), 36.1 (CH₂, C-8), 139.6 (C, C-9), 132.5 (CH, C-10, C-14), 117.3 (CH, C-11, C-13), 158.3 (C, C-12), 171.8 (C, C-15), 54.6 (CH, C-16), 38.1 (CH₂, C-17), 171.51 (C, C-18), 171.58 (C, C-19), 59.7 (CH, C-20), 39.9 (CH₂, C-21), 130.8 (C, C-22), 130.6 (CH, C-23, C-27), 129.8 (CH, C-24, C-26), 128.0 (CH, C-25), 171.54 (C, C-28), 62.4 (CH, C-29), 32.0 (CH₂, C-30), 22.8 (CH₂, C-31), 48.0 (CH₂, C-32), 171.65 (C, C-33), 51.0 (CH,

C-34), 46.8 (CH₂, C-35), 26.3 (CH, C-36), 25.2 (CH₃, C-37), 22.5 (CH₃, C-38), 172.79 (C, C-39), 60.1 (CH, C-40), 28.5 (CH₂, C-41), 26.2 (CH₂, C-42), 48.6 (CH₂, C-43), 172.71 (C, C-44), 61.3 (CH, C-45), 35.3 (CH, C-46), 16.9 (CH₃, C-47), 26.2 (CH₂, C-48), 11.2 (CH₃, C-49); ESIMS m/z 964 ($[M + \text{Na}]^+$, 100), 940 ($[M - \text{H}]^-$, 100); ESIMS/MS on $[M - \text{H}]^- m/z$ 940 (2), 922 (32), 843 (20), 730 (7), 664 (100), 646 (40), 619 (5), 517 (25), 422 (32); ESIMS/MS/MS on $[M - \text{Tyr} - \text{Asn}]^- m/z$ 664 (2), 646 (100), 619 (15), 517 (20), 454 (5), 420 (3), 357 (3), 260 (3), 209 (5); HRESIMS m/z 964.4916 ($[M + \text{Na}]^+$, $\text{C}_{49}\text{H}_{67}\text{N}_9\text{O}_{10}\text{Na}$ requires 964.4908).

3''-O-Methylcrenatoside (4): white amorphous powder; mp 224–226 °C; $[\alpha]_D^{20} -53.2^\circ$ (c 0.37, MeOH); UV (MeOH) λ_{max} (log ϵ) 328 (4.18), 290 (3.96), 231 (4.03), 219 (4.07), 205 (4.31) nm; IR (KBr) ν_{max} 3403, 2957, 1719, 1638, 1603, 1517, 1448, 1302, 1274, 1243, 1174, 1122, 1064, 1042, 1018 cm^{-1} ; ^1H NMR (DMSO-*d*₆, 600 MHz) δ 6.73 (1H, d, $J = 1.8$ Hz, H-2), 8.90 (1H, s, 3-OH), 8.89 (1H, s, 4-OH), 6.69 (1H, d, $J = 7.8$ Hz, H-5), 6.61 (1H, dd, $J = 7.8, 1.8$ Hz, H-6), 4.56 (1H, dd, $J = 10.8, 2.4$ Hz, H-7), 3.94 (1H, dd, $J = 11.4, 2.4$ Hz, H-8a), 3.50 (1H, dd, $J = 11.4, 10.8$ Hz, H-8b), 4.54 (1H, d, $J = 7.8$ Hz, H-1'), 3.38 (1H, m, H-2'), 4.03 (1H, t, $J = 9.6$ Hz, H-3'), 4.90 (1H, t, $J = 9.6$ Hz, H-4'), 3.70 (1H, t, $J = 7.8$ Hz, H-5'), 3.43 (2H, m, H-6'a, H-5''), 3.40 (1H, m, H-6'b), 4.86 (1H, t, $J = 6.0$ Hz, 6'-OH), 4.97 (1H, d, $J = 1.8$ Hz, H-1''), 3.53 (1H, brs, H-2''), 3.24 (1H, m, H-3''), 3.09 (1H, t, $J = 9.6$ Hz, H-4''), 1.02 (3H, d, $J = 6.0$ Hz, H-6''), 4.53 (2H, overlapped, 2''-OH, 4''-OH), 4.45 (1H, d, $J = 5.4$ Hz, 3''-OH), 7.27 (1H, d, $J = 1.8$ Hz, H-2'''), 9.61 (1H, s, 4''-OH), 6.79 (1H, d, $J = 8.4$ Hz, H-5'''), 7.09 (1H, dd, $J = 8.4, 1.8$ Hz, H-6'''), 7.56 (1H, d, $J = 15.6$ Hz, H-7'''), 6.39 (1H, d, $J = 15.6$ Hz, H-8'''), 3.80 (3H, s, OCH₃); ^{13}C NMR (DMSO-*d*₆, 150 MHz) δ 128.6 (C, C-1), 114.2 (CH, C-2), 145.8 (C, C-3), 145.7 (C, C-4), 115.9 (CH, C-5), 117.8 (CH, C-6), 76.7 (CH, C-7), 71.6 (CH₂, C-8), 97.5 (CH, C-1'), 81.0 (CH, C-2'), 74.9 (CH, C-3'), 69.3 (CH, C-4'), 76.8 (CH, C-5'), 61.0 (CH₂, C-6'), 100.9 (CH, C-1''), 71.0 (CH, C-2''), 70.9 (C-3''), 72.0 (CH, C-4''), 69.4 (CH, C-5''), 18.5 (CH₃, C-6''), 126.2 (C, C-1'''), 111.8 (CH, C-2'''), 148.5 (C, C-3'''), 150.1 (C, C-4'''), 116.1 (CH, C-5'''), 123.8 (CH, C-6'''), 146.5 (CH, C-7'''), 114.5 (CH, C-8'''), 166.1 (C, C-9''), 56.3 (CH₃, OCH₃); ESIMS m/z 659 ($[M + \text{Na}]^+$, 85), 675 ($[M + \text{K}]^+$, 10), 1295 ($[2M + \text{Na}]^+$, 100), 635 ($[M - \text{H}]^-$, 100), 671 ($[M + \text{Cl}]^-$, 20), 1271 ($[2M - \text{H}]^-$, 25); HRESIMS m/z 659.1949 ($[M + \text{Na}]^+$, $\text{C}_{30}\text{H}_{36}\text{O}_{15}\text{Na}$ requires 659.1951).

Amino Acid Analysis of 1–3. Microtoenins A (1) (5.4 mg), B (2) (5.7 mg), and C (3) (5.4 mg) were treated separately with 6 N HCl (10.8, 11.4, and 10.8 mL, respectively) at 110 °C for 30 h in a sealed tube, after which the reaction mixtures were concentrated in vacuo to dryness and redissolved in 0.02 N HCl (2.7, 2.9, and 2.7 mL, respectively) and then subjected to amino acid analysis.

Acid Hydrolysis of Cyclopeptides. 1–3 (each 0.5 mg) were separately treated with 6 N HCl (each 1 mL) at 110 °C for 30 h. The reaction mixture was concentrated in vacuo to dryness. These acid hydrolysates were then subjected to Marfey's derivatization.¹⁶

Marfey's Derivatization and Analysis. To each acid hydrolysate was added 50 μL of 1 M sodium bicarbonate and 1 mL of 0.1% FDAA in acetone. The solution was stirred at 40 °C for 60 min. Then the mixture was neutralized with 25 μL of 2 N HCl, and the derivatized sample was concentrated in vacuo to dryness and redissolved in H₂O (each 1 mL). As reference DL- and L-amino acids were derivatized in the same manner. HPLC analysis involved elution of 20 μL of the derivatized solution through a C₁₈ column (Lichrospher 100, RP-18e, 5 μm , 4 \times 250 mm, Merck KGaA, Germany) with a flow rate of 1 mL/min in 60 min linear gradient of CH₃CN in H₂O (0%–60%) containing 0.05% TFA and detection at 340 nm.

Retention times (t_R , min) of Marfey's derivatives of authentic amino acids: 32.31 (L-Asp), 32.96 (D-Asp), 34.34 (Gly), 36.14 (L-Ala), 37.69 (D-Ala), 36.78 (L-Pro), 37.80 (D-Pro), 39.33 (L-Tyr), 40.35 (D-Tyr), 41.84 (L-Val), 45.23 (D-Val), 45.13 (L-Ile), 48.68 (D-Ile), 46.09 (L-Leu), 49.71 (D-Leu), 46.32 (L-Phe), 48.59 (D-Phe).

Retention times (t_R , min) of the Marfey's derivatives of the acid hydrolysate of microtoenin A (1): 36.50 (L-Ala), 37.08 (L-Pro), 39.77 (L-Tyr), 42.23 (L-Val), 46.48 (L-Leu), 46.73 (L-Phe).

Retention times (t_R , min) of the Marfey's derivatives of the acid hydrolysate of microtoenin B (2): 35.08 (Gly), 36.71 (L-Ala), 37.20 (L-Pro), 42.19 (L-Val), 46.52 (L-Phe).

Retention times (t_R , min) of the Marfey's derivatives of the acid hydrolysate of microtoenin C (3): 32.03 (L-Asp), 36.88 (L-Pro), 39.33 (L-Tyr), 45.30 (L-Ile), 46.03 (L-Leu), 46.36 (L-Phe).

Alkaline Treatment of Compound 4 followed by Acid Hydrolysis. Compound 4 (10 mg) in 2% NaOH solution was kept overnight under N_2 at room temperature. The reaction mixture was acidified with diluted HCl(aq) and extracted with Et_2O . The aqueous layer was extracted with *n*-BuOH. The *n*-BuOH layer was evaporated to dryness. The residue was dissolved in 1% H_2SO_4 solution and heated on a H_2O bath (95 °C) for 1 h and then cooled. The mixture was extracted with Et_2O . The aqueous layer was neutralized with $Ba(OH)_2$ (aq), and the precipitate was filtered off. The filtrate was evaporated to dryness. Glucose and rhamnose were detected in the residue by TLC comparison with authentic samples.

Angiotensin I Converting Enzyme Assay. ACE activity was determined by the method of Carmel.²⁰ Briefly, 10 μ L of ACE extract (containing 10 μ g of total protein) was incubated with 10 μ L of samples of various concentration for 30 min at 37 °C. Then 120 μ L of HHL was added and incubated for 15 min at 37 °C. Afterward, 40 μ L of 1 M NaOH and 10 μ L of 20 g/L *O*-phthaldialdehyde were added, and the reaction was terminated after 10 min by the addition of 20 μ L of 3 M HCl(aq). The fluorescence intensity of the product His-Leu was measured at 405 nm (excitation) and 535 nm (emission) with a fluorescence spectrophotometer (Wallac Victor2). For each assay, the blank control, negative control, and positive control were prepared. If the letters *N*, *S*, and *B* respectively represent the measured fluorescence intensity values of the reacting systems of negative control, detected sample, and blank control, then the ACE inhibitory rate (%) = $100\% \times (N - S) / (N - B)$. At least three separate determinations were conducted for each sample. In this assay protocol, the positive control (captopril) completely inhibited ACE activity at the concentration of 40 nM with an IC_{50} of 20.3 nM. These data are in accordance with those reported,²¹ indicating the reliability of this assay protocol.

In Vitro Cytotoxicity Assay. Cytotoxic assays were performed by the sulforhodamine B (SRB) method.²² The cell lines used were human breast cancer (Bre04), human lung cancer (Lu04), and human neuroma (N04).

Antiherpetic Activity Assay. Inhibition of Herpes simplex virus type 2 (HSV-2) replication was evaluated in the Vero cell line using assays of cytopathic effect (CPE).²³ Acyclovir (ACV) was used as positive drug control.

Supporting Information Available: Figures S1–S7, selected 2D NMR, and mass fragmentation data for compounds 1–4. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- (1) Delectis Florae Reipublicae Popularis Sinicae Agendae Academiae Sinicae Edita. *Flora Reipublicae Popularis Sinicae*, Science Press: Beijing, 1977; Vol. 66, pp 46–64.
- (2) Mei, S. X.; Pan, X. H.; Hou, Z. F.; Li, Y. *Chin. Pharm. J.* **1999**, *34*, 366–367.
- (3) Aбышев, A. Z.; Zmeikov, V. P. *Chem. Nat. Compd.* **1982**, *18*, 270–276.
- (4) Borges, M. F. M.; Pinto, M. M. M. *J. Liq. Chromatogr.* **1994**, *17*, 1125–1139.
- (5) Sadtler Research Laboratories, Inc. *Nuclear Magnetic Resonance Spectra*; Sadtler Research Laboratories, Inc. Publishers: Philadelphia, 1978; Vol. 43–46, 27798M.
- (6) Boulos, L. S.; Arsanious, M. H. N. *Synth. Commun.* **2002**, *32*, 2779–2790.
- (7) Berghofer, B.; Holzl, J. *Planta Med.* **1987**, *53*, 216–217.
- (8) Siddiqui, S.; Hafeez, F.; Begum, S.; Siddiqui, B. S. *J. Nat. Prod.* **1988**, *51*, 229–233.
- (9) Itokawa, H.; Suto, K.; Takeya, K. *Chem. Pharm. Bull.* **1981**, *29*, 254–256.
- (10) Valesi, A. G.; Rodriguez, E.; Velde, G. V.; Mabry, T. J. *Phytochemistry* **1972**, *11*, 2821–2826.
- (11) Wang, Y.; Hamburger, M.; Gueho, J.; Hostettmann, K. *Phytochemistry* **1989**, *28*, 2323–2327.
- (12) Afifi, M. S.; Lahloub, M. F.; El-khayaat, S. A.; Anklin, C. G.; Ruegger, H.; Sticher, O. *Planta Med.* **1993**, *59*, 359–362.
- (13) Murayama, T.; Yanagisawa, Y.; Kasahara, A.; Dnoder, K. I.; Kurimoto, M.; Ikeda, M. *Nat. Med.* **1998**, *52*, 455–458.
- (14) Nawwar, M. A. M.; El-mousallamy, A. M. D.; Barakat, H. B.; Buddrus, J.; Linscheid, M. *Phytochemistry* **1989**, *28*, 3201–3206.
- (15) Markham, K. R.; Ternai, B. *Tetrahedron* **1976**, *32*, 2607–2612.
- (16) Marfey, P. *Carlsberg Res. Commun.* **1984**, *49*, 591–596.
- (17) Li, W. L.; Yi, Y. H.; Wu, H. M.; Xu, Q. Z.; Tang, H. F.; Zhou, D. Z.; Lin, H. W.; Wang, Z. H. *J. Nat. Prod.* **2003**, *66*, 146–148.
- (18) Hallock, Y. F.; Sowder, R. C.; Pannell, L. K.; Hughes, C. B.; Johnson, D. G.; Gulakowski, R.; Cardellina, J. H.; Boyd, M. R. *J. Org. Chem.* **2000**, *65*, 124–128.
- (19) Zampella, A.; D'Auria, M. V.; Paloma, L. G.; Casapulla, A.; Minale, L.; Debitus, C.; Henin, Y. *J. Am. Chem. Soc.* **1996**, *118*, 6202–6209.
- (20) Carmel, A.; Yaron, A. *Eur. J. Biochem.* **1978**, *87*, 265–273.
- (21) Duncan, A. C.; Jager, A. K.; Staden, J. *J. Ethnopharmacol.* **1999**, *68*, 63–70.
- (22) Skehan, P.; Storeng, R.; Scudiero, D.; Monks, A.; McMahon, J.; Vistica, D.; Warren, J. T.; Bockesch, H.; Kenny, S.; Boyd, M. R. *J. Natl. Cancer Inst.* **1990**, *82*, 1107–1112.
- (23) Mahmoud, H.; Vladimir, I.; Jacov, T.; Shoshana, M. A. *J. Biochem. Biophys. Methods* **2002**, *50*, 189–200.

NP040021X